



## References

- Abbate, A, (2003):** Increased myocardial apoptosis in patients with unfavorable left ventricular remodeling and early symptomatic postinfarction heart failure. *J. Am. Coll. Cardiol.* 41, 753–760.
- Acehan, D, Jiang, X, Morgan, DG, Heuser, JE, Wang, X and Akey, CW (2002):** "Three-dimensional structure of the apoptosome: implications for assembly, procaspase-9 binding, and activation." *Mol Cell* 9(2): 423-432.
- Adams RH, Porras A, Alonso G, Jones M, Vintersten K, Panelli S, Valladares A, Perez L, Klein R, Nebreda AR (2000):** Essential role of pSSalpha MAP kinase in placental but not embryonic cardiovascular development. *Mol Cell*, 6:109-116.
- Ahmad, M. and Shi, Y. (2000):** TRAIL-induced apoptosis of thyroid cancer cells: potential for therapeutic intervention. *Oncogens* 19, 3363-3371.
- Ahn EY, Pan G, Vickers SM, McDonald JM (2002):** IFN-gamma upregulates apoptosis-related molecules and enhances Fas-mediated apoptosis in human cholangio-carcinoma. *Int J Cancer*;100(4):445-451.
- Alarcon RM, Rupnow BA, Graeber TG, Knox SJ. Giaccia AJ (1996):** Modulation of c-myc. activity and apoptosis in vivo. *Cancer Res* 56: 4315-4319.
- Alles A.J. , and sulik, K. K. (1990):** Retinoic acid-induced spina bifida , evidence for apathological mechanism. *Development (Cambridge, UK)* 108,73-81.
- Allison. J., Thomas. H. E., Catterall, T., (2005):** Transgenic expression of dominant-negative Fas-associated death domain protein in beta cells protects against Fas ligand-induced apoptosis and reduces spontaneous diabetes in nonobese diabetic mice. *Journal of Immunology*, 175, 293-301.
- Alnemri ES, Livigston DJ, Nicholson DW (1996):** Human ICE/CED-3 protease nomenclature. *Cell*; 187:171.
- Ameisen, JC (2002):** "On the origin, evolution, and nature of programmed cell



- death: a timeline of four billion years." *Cell Death Differ* 9(4): 367-393.
- Anderson, A.J., Pike, C.J., and Cotman, C.W. (1995):** Differential induction of IEGs in response to Ab peptides in cultured neurons: association between c-Jun immunoreactivity and Ab-mediated apoptosis. *J. Neurochem.* 65:1487–1498.
- Anderson, A.J., Su, J.H., and Cotman, C.W. (1996):** DNA damage and apoptosis in Alzheimer's disease: colocalization with c-jun immunoreactivity, relationship to brain area, and effect of postmortem delay. *J. Neurosci.* 16:1710-1719.
- Appel, S.H. (1981):** A unifying hypothesis for the cause of amyotrophic lateral sclerosis, parkinsonism, and Alzheimer disease. *Ann. Neurol.* 10:499-505.
- Arscott, P.L. et al (1999):** Fas (CD95) expression is up-regulated on papillary thyroid carcinoma. *J. Clin.Endocrinol.Metab.* 84,4246-4252.
- Arscott, PL. et al. (1997):** Fas (APO-1, CD95) mediated apoptosis in thyroid cells is regulated by a labile protein inhibitor. *Endocrinology* 138, 5019-5027.
- Ashkenazi, A. and Dixit, V M. (1999):** Apoptosis control by death and decoy receptors. *Curr. Opin. Cell Biol.* 11,255-260.
- Ashwell JD, Lu FW, Vacchio MS (2000):** Glucocorticoids in T cell development and function. *Annu Rev Immunol*, 18:309-345.
- Askew DS. Ashmun RA, Simmons BC. Cleveland JL (1991):** Constitutive c-myc expression in an IL-3-dependent myeloid cell line suppresses cell cycle arrest and accelerates apoptosis. *Oncogene* 6: 1915 1922.
- Attila AL Moss SF, Walters JRF., (1995):** Increased small bowel epithelial apoptosis reflects celiac sprue activity. *Gastroenterology*;108:A271.
- Babic, I., Cherry, E., & Fujita, D. (2006):** SUMO modification of Sam68 enhances its ability to repress cyclin D1 expression and inhibits its ability to induce apoptosis. *Oncogene*, 25, 4955-4964.
- Babu, S. R., Bao, F., Roberts, C., (2003):** Caspase 7 is a positional candidate gene for IDDM 17 in a Bedouin Arab family. *Annals of the New York Academy of Sciences*, 1005, 340–343.



- Baker SJ, Preisinger AC, Jessup JM, (1990):** p53 gene mutations occur in combination with 17p allelic deletions as late events in colorectal tumorigenesis. *Cancer Res*;50:7717-7722.
- Baker, J.R., Jr (1995) :** The immune response to papillary thyroid cancer. *J.Clin. Endocrinol Metab.* 80,3419-3420.
- Bantel H, Lugerling A, Poremba C., (2001):** Caspase activation correlates with the degree of inflammatory liver injury in chronic hepatitis C virus infection. *Hepatology*;34(4 Part 1):758-767.
- Barazzone C, Horowitz S, Donati YR., (1998):** Oxygen toxicity in mouse lung: Pathways to cell death. *Am J Respir Cell Mol Biol* 1998; 19:573-581.
- Barbas-Filho JV, Ferreira MA, Sesso A. (2001):** Evidence of type II pneumocyte apoptosis in the pathogenesis of idiopathic pulmonary fibrosis (IPF)/usual interstitial pneumonia (UIP). *J Clin Pathol*;54:132-138.
- Barnes PJ, Shapiro SD, Pauwels RA (2003):** Chronic obstructive pulmonary disease: Molecular and cellular mechanisms. *Eur Respir J*;22:672-688.
- Batteux, F. et al (2000):** Transgenic expression of Fas ligand on thyroid follicular cells prevents autoimmune thyroiditis. *J. Immunol* 164, 1681-1688.
- Bauer, MK, Schubert, A, Rocks, O and Grimm, S (1999):** "Adenine nucleotide translocase-1, a component of the permeability transition pore, can dominantly induce apoptosis." *J Cell Biol* 147(7): 1493-1502.
- Bäumler CB, Böhler T, Hen- I, Benner A, Krammer PH, Debatin K-M (1996):** Activation of the CD95 (APO-1/Fas) system in T cells from human immunodeficiency virus type-1-infected children. *Blood* ;88(5):1741-1746.
- Baylin S, Bestor TH (2002):** Altered methylation patterns in cancer cell genomes: cause or consequence? *Cancer Cell*;1(4):299-305.
- Behringer R.R., Finegold M.J. and cate R.L. (1994):** Mullerian – inhibiting substance function during mammalian sexual development . cell (Cambridge , mass) 79,415 – 425 .
- Behrmann I, Walczak H, Krammer PH (1994):** Structure of the human APO-1 gene. *J Immunol*;24:3057-3062.



- Benliar. M & Stamler. .1. (2005):** S A central role for S-nitrosylation in apoptosis  
Nat Cell Biol 7. 645-646.
- Berg LJ, Kang J (2001):** Molecular determinants of TCP expression and  
selection. Curr Opin Immunol, 13:232-241.
- Berke G 1995:** The CTLs kiss of death. Cell;81:9-12.
- Beutner, G, Ruck, A, Riede, B and Brdiczka, D (1998):** "Complexes between  
porin, hexokinase, mitochondrial creatine kinase and adenylate  
translocator display properties of the permeability transition pore.  
Implication for regulation of permeability transition by the kinases."  
Biochim Biophys Acta 1368(1): 7-18.
- Bissonnette RP, Echeverri F, Mahboubi A, Green DR (1992):** Apoptotic cell  
death induced by c-myc is inhibited by bcl-2. Nature 359: 552-554.
- Blandino G, Scardigli R, Rizzo MG, Crescenzi M, Soddu S, Sacchi A (1995):**  
Wild-type p53 modulates apoptosis of normal, IL-3 deprived,  
hematopoietic cells. Oncogene;10(4):731-737.
- Blum, D., Wu, Y., Nissou, M.F., Arnaud, S., Benabid, A.L., and Verna, J.M.  
(1997):** p53 and Bax activation in 6-hydroxydopamine-induced apoptosis  
in PC12 cells. Brain Res. 751:139-142.
- Borgerson, K.L. et al (1999):** The role of Fas-mediated apoptosis in thyroid  
autoimmune disease. Autoimmunty 30, 251-264.
- Borner, C (2003):** "The Bcl-2 protein family: sensors and checkpoints for life-or-  
death decisions." Mol Immunol 39(11): 615-647.
- Bortezomib, Nat. Rev., Drug Discov. (2004) :** S8– S9.
- Bosca, L., & Hortelano, S. (1999):** Mechanisms of nitric oxide-dependent  
apoptosis: Involvement of mitochondria! mediators. Cell Signal, 17,239-  
244.
- Botla R, Spivey JR, Aguilar H, Bronk SF, Gores GJ (1995):** Ursodeoxycholate  
(UDCA) inhibits the mitochondrial membrane permeability transition  
induced by glycochenodeoxycholate: a mechanism of UDCA cytoprotec-  
tion. J Pharmacol Exp Ther 1995;272(2):930-938.



- Bouillet P, Metcalf D, Huang DC, Tarlinton DM, Kay TW, Kontgen F, Adams JM, Strasser A (1999):** Proapoptotic Bcl-2 relative Bim required for certain apoptotic responses, leukocyte homeostasis, and to preclude autoimmunity. *Science*, 286:1735-1738.
- Bouillet P, Purton JF, Godfrey DL, Zhang LC, Coultas L, Puthalakath H, Pellegrini M, Cory S, Adams JM, Strasser A (2002):** BH3-only Bcl-2 family member Bim is required for apoptosis of autoreactive thymocytes. *Nature*, 415:922-926.
- Bouillet, P and Strasser, A (2002):** "BH3-only proteins - evolutionarily conserved proapoptotic Bcl-2 family members essential for initiating programmed cell death." *J Cell Sci* 115(Pt 8): 1567-1574.
- Brctz, J.D, et al (1999):** TRAIL dcatii pathway expression and induction in thyroid follicular cells. *J. Biol. Chem.* 274, 23627-23632.
- Bretz, J.D. et al (1999):** Inflammatory cytokine regulation of Fas-me dialed apoptosis in thyroid follicular cells. *J. Biol Chem.* 274, 25433-25438.
- Brewer JA, Kanagawa O, Sleckman BP, Muglia LJ (2002):** Thymocyte apoptosis induced by T cell activation is mediated by glucocorticoids in vivo. *J Immunol*, 169:1837-1843.
- Broussard-Diehl C, Bauer SR, Scheuermann RH (1996):** A role forc-myc in the regulation of thymocyte differentiation and possibly positive selection. *J Immunol* 156: 3141-3150.
- Budinger GR, Tso M, McClintock DS, (2002):** Hyperoxia-induced apoptosis does not require mitochondrial reactive oxygen species and is regulated by Bcl-2 proteins. *J Biol Chem* ;277:15654-15660.
- Budinger GR, Tso M, McClintock DS., (2002):** Hyperoxia-induced apoptosis does not require mitochondrial reactive oxygen species and is regulated by Bcl-2 proteins. *J Biol Chem*;277:15654-15660.
- Burris H.A, (2006):** 3rd Dual kinase inhibition in the treatment of breast cancer : initial experience with the EGFR/ErbB-2 inhibitor lapatinib, *Oncologist* 9 (Suppl. 3) , 10– 15.



- Butler, A., Jang, J., Gurlo, T., (2004):** Diabetes due to a progressive defect in beta-cell mass in rats transgenic for human islet amyloid polypeptide (HIP Rat): A new model for Type 2 diabetes. *Diabetes*, 53, 1509-1516.
- Bykov, V.J., Selivanova, G., Wiman, K.G., (2003):** Small molecules that reactivate mutant p53. *Eur. J. Cancer* 39, 1828–1834.
- Cameron J.S., (1988):** the long term outcome of the glomerular disease . In *disease of the kidney* edited by R. Schrier, Gottschalk, Goston, Little Browne P. 2127.
- Carini R, De Cesaris MG, Splendore R, et al (2001):** Signal pathway involved in the development of hypoxic preconditioning in rat hepatocytes. *Hepatology*;33:131-9.
- Carnevali S, Petruzzelli S, Longoni B, et al (2003):** Cigarette smoke extract induces oxidative stress and apoptosis in human lung fibroblasts. *Am J Physiol Lung Cell Mol Physiol*;284:L955-63.
- Celli BR, MacNee W, (2004):** ATS/ERS Task Force. Standards for the diagnosis and treatment of patients with COPD: A summary of the ATS/ERS position paper. *Eur Respir J*;23:932-46.
- Chandler D., Naggar A.K., Brisbay S., Redline R.W., Redline, Mc Donnell T.J., (1994):** Apoptosis and expression of the bcl-2 proto-oncogene in the fetal and adult human kidney : Evidence for the contribution of bcl-2 expression to renal carcinogenesis . *Hum pathol* 25:789-796.
- Chang, I., Cho, N., Kiin, S., et al. (2004):** Role of calcium in pancreatic islet cell death by IFN-gamma/TNF-alpha. *Journal of Immunology*, 172, 7008-7014.
- Chen Q, Galleano M, Cederbaum AI (1997):** Cytotoxicity and apoptosis produced by arachidonic acid in Hep G2 cells overexpressing human cytochrome P4502E1. *J Biol Chem*;272:14532-14541.
- Chen, Y.N. et al. (2002) :** Selective killing of transformed cells by cyclin/cyclin-dependent kinase 2 antagonists. *Proc. Natl. Acad. Sci. U. S. A.* 96, 4325-4329.



- Chen. W., Salojin, K., Mi, Q., et al. (2004):** Insulin-like growth factor (IGF)-I/IGF-binding protein-3 complex: Therapeutic efficacy and mechanism of protection against Type 1 diabetes. *Endocrinology*, 145, 627-638.
- Choi DW. (1992):** Excitotoxic cell death . Department of Neurology , washington University School of medicine , ST. Louis, Missouri 63110. *J Neurobiol*, Nov, 23:9,1261-76.
- Christ, B., Huang, R., Scaal, M., (2004):** Formation and differentiation of the avian sclerotome. *Anat. Embryol. (Berl)* 208, 333–350.
- Chung H.M; Neff A. W. and Malacinski G. M, (1989):** Autonomous cell death of amphibian (*Xenopus Laevis*) Cranial Myotomes . *J. Exp. Zool* . 251,290-299.
- Claessens YE, Bouscary D, Dupont JM, Picard F, Melle J, Gis-selbrecht S, et al (2002):** In vitro proliferation and differentiation of ery-throid progenitors from patients with myelodysplastic syndromes: evidence for Fas-dependent apoptosis. *Blood*;99(5):1594-601.
- Clark JM, Lambertson CJ (1971) :** Pulmonary oxygen toxicity: A review. *Pharmacol Rev*;23:37-133.
- Clayton E, Bardi G, Bell SE, Chantry D, Downes CP, Gray A, Humphries LA, Rawlings D, Reynolds H, Vigorito E et al (2002):** A crucial role for the p110delta subunit of phosphatidylinositol 3-kinase in B cell development and activation. *J Exp Med*, 196:753-763.
- Clayton LK, Ghendler Y, Mizoguchi E, Patch RJ, Ocain TD, Orth K, Bhan AK, Dixit VM, Reinherz EL (1997):** T-cell receptor ligation by peptide/MHC induces activation of a caspase in immature thymocytes: the molecular basis of negative selection. *EMBO J*, 16:2282-2293.
- Clermont, O., Burlet, P., Burglen, L., Lefebvre, S., Pascal, F., McPherson, J. et al. (1994):** Use of genetic and physical mapping to locate the spinal muscular atrophy locus between two new highly polymorphic DNA markers. *Am. J. Hum. Genet.* 54:687-694.



- CM. Matter, C.E. Chadjichristos, P. Meier, T. von Lukow-icz, C. Lohmann, P.K. Schuler, D. Zhang. B. Odermatt, E. Hofmann, T. Brunner, B.R. Kwak, T.F. Luscher et al., (2006):** Role of endogenous Fas (CD95/Apo-1) ligand in balloon-induced apoptosis, inflammation, and neointima formation, *Circulation* 113 , 1879-1887.
- Cnop, M., Welsh, N., Jonas, J., et al. (2005):** Mechanisms of pancreatic  $\beta$ -cell death in Type 1 and Type 2 diabetes: Many differences, few similarities. *Diabetes*, 54, S97-S107.
- Cohen E.E., F. Rosen, W.M. Stadler, W. Recant, K. Stenson, D. Huo, E.E. Vokes, (2005):** Phase II trial of ZD1839 in recurrent or metastatic squamous cell carcinoma of the head and neck, *J. Clin. Oncol.* 21, 1980–1987.
- Cohen PL, Caricchio R, Abraham V, et al (2002):** Delayed apoptotic cell clearance and lupus-like autoimmunity in mice lacking the c-met membrane tyrosine kinase. *J Exp Med* 196(1): 135-40.
- Cohen PL, Eisenberg RA. Lpr and gld (1991):** single gene models of systemic autoimmunity and lymphoproliferative disease. *Annu Rev Immunol*;9:243-269.
- Contreras, J., Smyth, C., Bilbao, G, (2002):** Simvastatin induces activation of the serine-threonine protein kinase AKT and increases survival of isolated human pancreatic islets. *Transplantation*, 74, 1063-1069.
- Copp A. J. (1978):** Interaction between inner cell mass and trophoblast of the mouse blastocyst. A study of cellular proliferation. *J. Embryol. Exp. Morphol.* 48, 109-125.
- Cory, S and Adams, JM (2002):** "The Bcl2 family: regulators of the cellular life-or-death switch." *Nat Rev Cancer* 2(9): 647-656.
- Counism M.F.; chaudin E; Courtois Y. and allinquant B. (1991):** Dnase activities in embryonic chicken lens: In epithelial cells or in differentiating fibers where chromatin is progressively cleaved. *Biol. cell* 72, 231-238.



- Coutinho, P., Parsons, M.J., Thomas, K.A., Hirst, E.M., Saude, L., Campos, I., Williams, P.H., Stemple, D.L., (2004):** Differential requirements for COPI transport during vertebrate early development. *Dev. Cell* 7, 547–558.
- Coyle, J.T., and Puttfarcken, P. (1993):** Oxidative stress, glutamate, and neurodegenerative disorders. *Science* 262:689-695.
- Crapo JD, Barry BE, Foscue HA, (1980):** Structural and biochemical changes in rat lungs occurring during exposure to lethal and adaptive doses of oxygen. *Am Rev Respir Dis*:122:123-143.
- Creagh, EM and Martin, SJ (2001):** "Caspases: cellular demolition experts." *Biochem Soc Trans* 29(Pt 6): 696-702.
- Cullen, S.P.,and Martin, S.J., (2008):** Mechanisms of granule-dependent killing. *Cell Death Differ.* 15, 251–262.
- Cummins J, Ward T.H., E.C. LaCasse, M. Dawson, O. Denny, D.C. McManus, J.P. Durkin, M. Ranson, C. Dive (2004):** Validation of pharmacodynamic assays to evaluate the clinical efficacy of an antisense targeted to the XIAP inhibitor of apoptosis, *Br. J. Cancer* 91,s75.
- Cuppige F., Huseman R. and Chapman A, (1980):** ultrastructure and function of cysts from human adult polycystic kidneys. *kidney Int*; 17:372-381 .
- D. Verzola, M.T. Gandolfo, F. Ferrario, M.P. Rastaldi, B. Villaggio, F. Gianiorio, M. Giannoni, L. Rimoldi, F. Lauria, M. Miji, G. Deferrari, G. Garibotto, et al., (2007):** Apoptosis in the kidneys of patients with type II diabetic nephropathy, *Kidney Int.* 72 , 1262–1272.
- D'Agostini F, Balansky RM, Izzotti A, (2001):** Modulation of apoptosis by cigarette smoke and cancer chemopreventive agents in the respiratory tract of rats. *Carcinogenesis* 22:375-380.
- D'amato, R.J., Lipman, Z.P., and Snyder, S.H. (1986):** Selectivity of the parkinsonian neurotoxin MPTP: toxic metabolite MPP<sup>+</sup> binds to neuromelanin. *Science* 231:987-989.



- Damon AJ, Nicholson DW, Bleakley RC (1995) :** Activation of the apoptotic protease CPP32 by cytotoxic T-cell-derived granzyme B. *Nature*;377:446-448.
- Datto MB, Li Y, Panus JF. Howe DJ, Xiong Y, Wang XF (1995):** Transforming growth factorp induces the cyclin-dependent kinase inhibitor p21 through ap53-independent mechanism. *Proc Natl Acad Sci USA* 92: 5545-5549.
- De Maria R. Boirivant M, Cifone MG, (1997):** Functional expression of Fas and Fas ligand on human gut lamina propria T lymphocytes. *J Clin Invest*;97:316-322.
- De Maria R. Fais S, Silvestri L, (1993):** Continuous in vivo activation and transient hyporesponsiveness to TCR/CD3 triggering of human gut lamina propria T lymphocytes. *Eur J Immunol*;23:3104-3108.
- DeFranco AL (1997):** The complexity of signaling pathways activated by the BCR. *Curr Opin Immunol*, 9:296-308.
- Demedts IK, Brusselle GG, Bracke KR, (2005):** Matrix metalloproteinases in asthma and COPD. *Curr Opin Pharmacol*;5:257-263.
- Demierre M.F., Merlino G, (2004):** Chemoprevention of melanoma, *Curr. Oncol. Rep.* 6 , 406-413.
- Denault, JB and Salvesen, GS (2002):** "Caspases: keys in the ignition of cell death." *Chem Rev* 102(12): 4489-4500.
- Deng, G., Pike, C.J., and Cotman, C.W. (1996):** Alzheimer-associated presenilin-2 confers increased sensitivity to apoptosis in PC 12 cells. *FEBS Lett.*397:50-54.
- Deng, H.-X., Hentati, A., Tainer, J.A., Iqbal, Z., Cayabyab, A., Hung ,W.-Y, (1993):** Amyotrophic lateral sclerosis and structural defects in Cu, Zn superoxide dismutase. *Science* 261:1047-1051.
- Dewson, G., Kratina, T., Sim, H.W., Puthalakath, H., Adams, J.M., Colman, P.M., Kluck, R.M., (2009):** To trigger apoptosis, Bak exposes its BH3 domain and homodimerizes via BH3: groove interactions. *Mol. Cell* 30, 369–380.



- Dias N., Stein C.A., (2006):** Antisense ligonucleotides: basic concepts and mechanisms, *Mol. Cancer Ther.* 1 , 347– 355.
- Dini L, Lentini A, Diez GD, (1995):** Phagocytosis of apoptotic bodies by liver endothelial cells. *J Cell Sci*:108:967-973.
- Do RK, Hatada E, Lee H, Tourigny MR, Hilbert D, Chen-Kiang S (2000):** Attenuation of apoptosis underlies B lymphocyte stimulator enhancement of humoral immune response. *J Exp Med*, 192:953-964.
- Doctor R., Bacallao R., and mandel L (1994):** Method for recovering ATP content and mitochondrial function after chemical anoxia in renal cell cultures *A. M. J. physiol*, 266:1803-1811.
- Doerfler P, Forbush KA, Perlmutter RM (2000):** Caspase enzyme activity is not essential for apoptosis during thymocyte development. *J Immunol*, 164:4071-4079.
- Dolinay T, Wu W, Kaminski N, (2008):** Mitogen-activated protein kinases regulate susceptibility to ventilator-induced lung injury. *PLoS ONE*;3:e1601.
- Dong C, Davis RJ, Flavell RA (2002):** MAP kinases in the immune response. *Annu Rev Immunol*, 20:55-72.
- Dror Y, Freedman MH (2001):** Shwachman-Diamond syndrome marrow cells show abnormally increased apoptosis mediated through the Fas pathway. *Blood*:97(10):3010-3016.
- Du, C, Fang, M, Li, Y, Li, L and Wang, X (2000):** "Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition." *Cell* 102(1): 33-42.
- Du, Y., Bales, K.R., Dodel, R.C., Hamilton-Byrd, E., Horn, H.W., Czilli, D.L, (1997):** Activation of a caspase 3-related cysteine protease is required for glutamate-mediated apoptosis of cultured cerebellar granule neurons. *Proc. Natl. Acad. Sci. USA* 94:11,657-662.
- Dumez H., Debiec-Rychter M., I. Judson, B. Wasag, J. Verweij, M. Brown, S. Dimitrijevic, R. Sciot, M. Stul, H. Vranck, M. Scurr, A. Hagemeijer,**



- M. van Glabbeke, A.T. van Oosterom, (2004):** Use of c-KIT/ PDGFRA mutational analysis to predict the clinical response to imatinib in patients with advanced gastrointestinal stromal tumours entered on phase I and II studies of the EORTC Soft Tissue and Bone Sarcoma Group, Eur. J. Cancer 40 , 689–695.
- Durand E, A. Scoazec, A. Lafont, J. Boddaert, A. Al Hajzen, F. Addad, M. Mirshahi, M. Desnos, A. Tedgui, Z Mallat (2004):** In vivo induction of endothelial apoptosis leads to vessel thrombosis and endothelial denudation: a clue to the understanding of the mechanisms of thrombotic plaque erosion. Circulation 109 2503-2506.
- Editorial, (2002) :** Scrip's Review of 2001. Scrip Magazine 109, 73.
- Eguchi Y; Ewert D.L. and Tsujimoto Y. (1992):** Isolation and characterization of chicken bcl-2 gene , expression in a variety of tissues including lymphoid and neuronal organs in adult and embryo. Nucleic Acids Res . 20, 4187-4192.
- Eisenman RN, Cooper JA (1995):** Beating apathy to myc. Nature 378: 438-439.
- Elghazi, L., Balcazar, N., & Bernal-Mizrachi, E. (2006):** Emerging role of protein kinase B/Akt signaling in pancreatic p-cell mass and function. International Journal Biochemistry and Cell Biology, 38, 689-695.
- Enari M., Hug H., Nagata S., (1995):** Involvement of an ICE- Like protease in fas mediated apoptosis . Nature 375:78-81.
- Eramo. A. et al (2004):** CD95 death-inducing signaling complex formation and internalization occur in lipid rafts of type I and type II cells Eur . J Immunol 34. 1930 - 1940.
- Estevez, A.G., Spear, N., Manuel, S.M., Radi, R., Henderson, C.E., Barbeito, L., and Beckman, J.S. (1998):** Nitric oxide and superoxide contribute to motor neuron apoptosis induced by trophic factor deprivation. J. Neurosci. 18:923-931.
- Evan GI, Wylhe AH, Glibert CS, Littlewood TD, Land H, Brooks M, Waters CM, Penn LZ, Hancock DC (1992):** Induction of apoptosis in fibroblasts by c-myc protein. Cell 69: 119-128.



- Fadeel, B., Orrenius, S., (2005):** Apoptosis: a basic biological phenomenon with wideranging implications in human disease. *J. Intern. Med.* 258, 479–517.
- Faitova J, Krekac D, R. Hrstka, B. Vojtesek, (2006):** Endoplasmic reticulum stress and apoptosis, *Cell. Mol. Biol. Lett.* 11 , 488–505.
- Federici, M., Hribal, M., Perego, L, (2001):** High glucose causes apoptosis in cultured human pancreatic islets of Langerhans: A potential role for regulation of specific Bel family genes toward an apoptotic cell death program. *Diabetes*, 50, 1290-1301.
- Ferrari R, Agnoletti L, Comini L, Gaia G, Bachetti T, Cargnoni A, Ceconi C, Curello S, Visioli O (1998):** Oxidative stress during myocardial ischemia and heart failure. *Eur Heart J*;19:B2–B11.
- Fiucci G, Ruberti G (1994):** Detection of polymorphisms within the FAS cDNA gene sequence by GC-clamp denaturing gradient gel electrophoresis. *Inimmunogenetics*;39:437-439.
- Formigli L, Papucci L, Tani A, (2000):** Aponecrosis: Morphological and biochemical exploration of a syncretic process of cell death sharing apoptosis and necrosis. *J Cell Physiol*; 182:41-49.
- Fox CK, Furtwaengler A, Nepomuceno RR, Martinez OM, Krams SM (2001):** Apoptotic pathways in primary biliary cirrhosis and autoimmune hepatitis. *Liver*;21(4):272-279.
- Freeman B, Crapo JD (1981):** Hyperoxia increases oxygen radical production in rat lungs and lung mitochondria. *J Biol Chem*:256:10986-10992.
- Fukuda K, Kojiro M, Chiu JF (1993):** Demonstration of extensive chromatin cleavage in transplanted Morris Hepatoma 7777 tissue: Apoptosis or necrosis. *Am J Pathol*; 142:935–946.
- Gajate C, Alonso MT, Schimmang T, Mollinedo F (1996):** *c-fos* is not essential for apoptosis. *Biochem Biophys Res Commun* 218: 267-272.
- Gallucci BB; Sale GE, McDonald GB, (1982):** The fine structure of human rectal epithelium in acute graft-versus-host disease. *Am J Surg Pathol*:6:293-305.



- Garcia-Martinez J.M., Pe´ rez-Navarro E., Xifro´ , J.M. Canals, M. Diaz-Herna´ ndez, Y. Trioulier, E. Brouillet, J.J. Lucas, J. Alberch, (2007):** BH-3 only proteins Bid and BimEL are differentially involved in neuronal dysfunction in mouse models of Huntington's disease, *J. Neurosci. Res.* 85 2756–2769.
- George, S., Rochford, J., Wolfrum, C., (2004):** A family with severe insulin resistance and diabetes due to a mutation in AKT2. *Science*, 304, 1325–1328.
- Ghadge, G.D., Lee, J.P., Bindokas, V.P., Jordan, J., Ma, L., Miller, R.J., and Roos, R.P. (1997):** Mutant superoxide dismutase-1-linked familial amyotrophic lateral sclerosis-molecular mechanisms of neuronal death and protection. *J. Neurosci.* 17:8756-8766.
- Gil, J.M., Rego, A.C., (2008):** Mechanisms of neurodegeneration in Huntington's disease. *Eur. J. Neurosci.* 27, 2803–2820.
- Goldberg, Y.R., Nicholson, D.W., Rasper, D.M., Kalchman, M.A., Koide, H.B., Graham, R.K, (1996):** Cleavage of huntingtin by apopain, a proapoptotic cysteine protease, is modulated by the polyglutamine tract. *Nature Genet.* 13:442-449.
- Gong Q, Cheng AM, Akk AM, Alberola-Ila J, Gong G, Pawson T, Chan AC (2001):** Disruption of T cell signaling networks and development by Grb2 haploid insufficiency. *Nat Immun*, 2:29-36.
- Goodrich DW, Lee WH (1992):** Abrogation by c-myc of G1 phase arrest induced by RB protein but not by p53. *Nature* 360: 177-179.
- Graeber TG, Osmanian C, Jacks T, (1996):** Hypoxia-mediated selection of cells with diminished apoptotic potential in solid tumors. *Nature*; 379:88-91.
- Gramzinski R.A; Parchment R.E. and Pierce G.B. (1990):** Evidence linking programmed cell death in the blastocyst to polyamine oxidation . *Differentiation (Berlin)* 43,59,65.



- Grassme. H., Cremesti, A., Kolesnick. R. & Gulbms, E (2005):** Ceramide-mediated clustering is required for CD95-DISC formation. *Oncogene* 22. 5457-5470.
- Greenlund LJS, Deckwerth TL, Johnson EM Jr (1995):** Superoxide dismutase delays neuronal apoptosis: A role for reactive oxygen species in programmed neuronal death. *Neuron* ;14:303–315.
- Griffin, W.S.T., Stanley, L.C., Ling, C., White, L., MacLeod, V., Perrot, L.J, (1989):** Brain interleukin 1 and S-100 immunoreactivity are elevated in Down syndrome and Alzheimer disease. *Proc. Natl. Acad. Sci. USA* 86:7611 -7615.
- Griffith C.M. and Hay E.D (1992):** Epithelial – mesenchymal transformation during platal fusion , carboxyfluorescein traces cells at light and electron microscopic levels . *Development (Cambridge, UK)* 116,1087-1099.
- Grimm, S, Bauer, MK, Baeuerle, PA and Schulze-Osthoff, K (1996):** "Bcl-2 down-regulates the activity of transcription factor NF-kappaB induced upon apoptosis." *J Cell Biol* 134(1): 13-23.
- Guo Z, Clydesdale G, Cheng J, Kim K, Gan L, McConkey DJ, Ullrich SE, Zhuang Y, Su B (2002):** Disruption of Mekk2 in mice reveals an unexpected role for MEKK2 in modulating T-cell receptor signal transduction. *Mol Cell Biol*, 22:5761-5768.
- Guo, Q., Sopher, B.L., Pham, D.G., Furukawa, K., Robinson, N., Martin, G.M., and Mattson, M.P. (1997):** Alzheimer's presenilin mutation sensitizes neural cells to apoptosis induced by trophic factor withdrawal and amyloid b-peptide: involvement of calcium and oxyradicals. *J. Neurosci.* 17:4212-4222.
- Habit FM, Springal DR, Davies GJ, Oakley CM, Yacub MN, Polak JM (1996):** Tumor necrosis factor and inducible nitric oxide synthase in dilated cardiomyopathy. *Lancet*;347:1151–1154.
- Haddad JJ, Land SC (2000):** The differential expression of apoptosis factors in the alveolar epithelium is redox sensitive and requires NF-kappaB (RelA)-selective targeting. *Bio-chem Biophys Res Commun*:271:257-67.



- Haldar S, Negrini M, Monne M, (1994):** Down-regulation of bcl-2 by p53 in breast cancer cells. *Cancer Res*;54:2095-2097.
- Hamaguchi, T., Matsuoka, Y., Bechberger, J., Ohnishi, T., Fujita, K., Naus, C.C., Kusunoki, M., Tsubura, A., Tsuda, H., (2006):** Establishment of an apoptosis-sensitive rat mammary carcinoma cell line with a mutation in the DNA-binding region of p53. *Cancer Lett.* 232, 279-288.
- Hara H, Takeda A, Takeuchi M, Wakeham AC, Itie A, Sasaki M, Mak TW, Yoshimura A, Nomoto K, Yoshida H (2002):** The apoptotic protease-activating factor 1-mediated pathway of apoptosis is dispensable for negative selection of thymocytes. *J Immunol*, 168:2288-2295.
- Harade K, Ozaki S, Gershwin ME, Nakanuma Y (1997):** Enhanced apoptosis relates to bile duct loss in primary biliary cirrhosis. *Hepatology*;26(6):1399-1405.
- Harrington EA, Bennett MR, Fanidi A, Evan GI (1994):** C-myc-induced apoptosis in fibroblasts is inhibited by specific cytokines. *EMBO J* 13:3286-3295.
- Harrison D.J. , (1988):** Cell death in the diseased glomerulus . *Histopathology* 12:679-683.
- Hawkins, CJ, Yoo, SJ, Peterson, EP, Wang, SL, Vernooy, SY and Hay, BA (2000):** "The Drosophila caspase DRONC cleaves following glutamate or aspartate and is regulated by DIAP1, HID, and GRIM." *J Biol Chem* 275(35): 27084-27093.
- Hegewisch S, Weh HJ, Hossfeld DK (1990):** TNF-induced cardiomyopathy. *Lancet* 2:294–295.
- Hengartner, MO (1999):** "Programmed cell death in the nematode *C. elegans*." *Recent Prog Horm Res* 54: 213-22; discussion 222-224.
- Henson PM, Vandivier RW, Douglas IS (2006):** Cell death, remodeling, and repair in chronic obstructive pulmonary disease? *Proc Am Thorac Soc*;3:713-717.



- Herbst R.S.,and Frankel S.R, (2004):** Oblimersen sodium (Genasense bcl-2 antisense oligonucleotide): a rational therapeutic to enhance apoptosis in therapy of lung cancer, Clin. Cancer Res. 10 , 4245s-4248s.
- Hermeking H, Eick D (1994):** Mediation of c-myc-induced apoptosis by p53. Science 265: 2091-2093.
- Hermiston ML, Gordon JI (1995):** Inflammatory bowel disease and adenomas in mice expressing a dominant negative N-cadherin. Science;270:1203-1207.
- Hess P, Pihan G, Sawyers CL, Flavell RA, Davis RJ (2002):** Survival signaling mediated by c-Jun NH2-terminal kinase in transformed B lymphoblasts. Nat Genet, 32:201-205.
- Hickey, M.A. and Chesselet, M.F. (2003):** Apoptosis in Huntington's disease. Prog. Neuropsychopharmacol. Biol. Psychiatry 27, 255–265.
- Higuchi H, Bronk SF, Takikawa Y, (2001):** The bile acid glycochenodeoxycholate induces trail-receptor 2/DR5 expression and apoptosis. J Biol Chem;276(42):38610-38618.
- Hockenbery D.M. , Zutter W., Hickey W., Korsmeyer S.J. , Nahm M., (1991):** bcl-2 Protein is topographically restricted in tissues characterized by apoptotic cell death. Proc Natl Acad Sci, USA 88:6861-6965.
- Hodge S, Hodge G, Holmes M, (2005):** Increased airway epithelial and T-cell apoptosis in COPD remains despite smoking cessation. Eur Respir J;25:447-454.
- Hodge S, Hodge G, Scicchitano R, (2003):** Alveolar macrophages from subjects with chronic obstructive pulmonary disease are deficient in their ability to phagocytose apoptotic airway epithelial cells. Immunol Cell Biol;81:289-296.
- Hodge SJ, Hodge GL, Reynolds PN, (2003):** Increased production of TGF-beta and apoptosis of T lymphocytes isolated from peripheral blood in COPD. Am J Physiol Lung Cell Mol Physiol;285:L492-499.



- Hoglen, N.C. et al. (2004):** Characterization of the caspase inhibitor IDN-1965 in a model of apoptosis-associated liver injury. *J. Pharmacol. Exp. Ther.* 297, 811–818.
- Hoglund, P., Mintern, J., Waltzinger, C, (1999):** Initiation of autoimmune diabetes by developmentally regulated presentation of islet cell antigens in the pancreatic lymph nodes. *Journal of Experimental Medicine*, 189, 331-339.
- Hokanson, J., Kinney, G., Cheng, S., (2006):** Susceptibility to type 1 diabetes is associated with ApoCIII gene haplotypes. *Diabetes*, 55, 834–838.
- Hong, J.R., Lin, G.H., Lin, C.J., Wang, W.P., Lee, C.C., Lin, T.L., Wu, J.L., (2004):** Phosphatidylserine receptor is required for the engulfment of dead apoptotic cells and for normal embryonic development in zebrafish. *Development* 131, 5417–5427.
- Hoozemans JJ, van Haastert ES, Eikelenboom P, de Vos RA, Rozemuller JM, Scheper W, (2007):** *Biochem Biophys Res Commun* 2007;354(3):707-711.
- Horikoshi, M., Hara, K., Ohashi, J., (2006):** A polymorphism in the AMPKalpha2 subunit gene is associated with insulin resistance and type 2 diabetes in the Japanese population. *Diabetes*, 55, 919–923.
- Hotchkiss RS, Chang KC, Grayson MH, (2003):** Adoptive transfer of apoptotic splenocytes worsens survival, whereas adoptive transfer of necrotic splenocytes improves survival in sepsis. *Proc Natl Acad Sci USA*;100:6724–6729.
- Hu, J., Ferreira, A., and Van Eldik, L., (1997):** S100B induces neuronal cell death through nitric oxide release from astrocytes. *J. Neurochem.* 69:2294-2301.
- Hu, Y. et al. (2003) :** Antisense oligonucleotides targeting XIAP induces apoptosis and enhance chemotherapeutic activity against human lung cancer cell in vitro and in vivo. *Clin. Cancer Res.* 9, 2826-2836.



- Huang, Y, Park, YC, Rich, RL, Segal, D, Myszka, DG and Wu, H (2001):** "Structural basis of caspase inhibition by XIAP: differential roles of the linker versus the BIR domain." *Cell* 104(5): 781-790.
- Ikeda, H., Yamaguchi, M., Sugai, S., Aze, Y., Narumiya, S., and Kakizuka, A. (1996):** Expanded polyglutamine in the Machado-Joseph disease protein induces cell death in vitro and in vivo. *Nature Genet.*13:196-202.
- Ikeguchi M, Ueda T, Sakatani T, Hirooka Y, Kaibara N, (2002):** Expression of survivin messenger RNA correlates with poor prognosis in patients with hepatocellular carcinoma. *Diagn Mol Pathol*;11(1):33-40.
- Imai K, Mercer BA, Schulman LL, (2005):** Correlation of lung surface area to apoptosis and proliferation in human emphysema. *Eur Respir J*;25:250-258.
- Imai Y, Parodo J, Kajikawa O, (2003):** Injurious mechanical ventilation and end-organ epithelial cell apoptosis and organ dysfunction in an experimental model of acute respiratory distress syndrome. *JAMA*;289:2104-2112.
- Inaba T, Inukai T, Yoshihara T, Seyschab RA, Canman CE, Laken SJ, Kastan MB, Look AT (1996):** Reversal of apoptosis by the leukaemia-associated E2A-HLF chimaeric transcription factor. *Nature* 382: 541-544.
- Ishihara, H., Sasaoka, T., Kagawa, S., (2003):** Association of the polymorphisms in the 5'-untranslated region of PTEN gene with type 2 diabetes in a Japanese population. *FEBS Letters*, 554, 450–454.
- Ishii T, Matsuse T, Igarashi H, I (2001):** Tobacco smoke reduces viability in human lung fibroblasts: Protective effect of glutathione 5-transferase PL *Am J Physiol Lung Cell Mol Physiol*;280:L1189-1195.
- Ishizaki Y; Voyuvodic J.T; Burne J.F. and Raff M.C. (1993):** Control of lens epithelial cell survival . *J. cell Biol.* 121, 899-908.
- Ismail M, Gibson FM, Gordon-Smith EC, Rutherford TR (2001):** Bcl-2 and Bcl-x expression in the CD34<sup>+</sup> cells of aplastic anemia patients relationship with increased apoptosis and upregulation of Fas antigen. *Br J Haematol*;113(3):706-712.



- Isner JM, Kearney M, Bortman S, Passeri J (1995):** Apoptosis in human atherosclerosis and restenosis. *Circulation*;91:2703–2711.
- Iwahashi, H., Eguchi, Y., Yasuhara, N., Hanafusa, T., Matsuzawa, Y., and Tsujimoto, Y. (1997):** Synergistic anti-apoptotic activity between Bcl-2 and SMN implicated in spinal muscular atrophy. *Nature* 390:413-417.
- Jacks T, Weinberg RA (1996):** Cell cycle control and its watchman. *Nature*;381:643-644.
- Jascur, T., Brickner, H., Salles-Passador, I., Barbier, V., El Khissiin, A., Smith, B., Fotedar, R., Fotedar, A., (2005):** Regulation of p21(WAF1/CIP1) stability by WISp39, a Hsp90 binding TPR protein. *Mol. Cell* 17, 237-249.
- Javitch, J.A., D'amato, R.J., Strittmatter, S.M., and Snyder, S.H. (1985):** Parkinsonism-inducing neurotoxin, Nmethyl-4-phenyl-1,2,3,6-tetrahydropyridine: uptake of the metabolite N-methyl-4-phenylpyridine by dopamine neurons explains selective toxicity. *Proc. Natl. Acad. Sci. USA* 82:2173-2177.
- Jen J, Kim H, Piantadosi S, (1994):** Allelic loss of chromosome 18q and prognosis in colorectal cancer. *N Engl J Med*;331:213- 221.
- Jeremias I, Kupatt C, Martin-Villalba A, (2000):** Involvement of CD95/Apo1/Fas in cell death after myocardial ischemia. *Circulation*; 102:915-920.
- Jin Y, Kim HP, Chi M, (2008):** Deletion of caveolin-1 protects against oxidative lung injury via up-regulation of heme oxygenase-1. *Am J Respir Cell Mol Biol*;39:171-179.
- Jirsova Z . and vernerova Z. (1993):** Involution of the wolffian duct in the rat. *Funct . Dev. Morphol .* 3, 205-506.
- Joe M.K., Lee H.J., Suh Y.H., Han K.L., Lim J.H., J. Song, J.K. Seong, M.H. Jung, (2008):** Crucial roles of neuronatin in insulin secretion and high glucose-induced apoptosis in pancreatic beta-cells, *Cell. Signal.* 20 , 907–915.



- Johannesen, J., et al. (2001):** Linkage of the human inducible nitric oxide synthase gene to type 1 diabetes. *Journal of Clinical Endocrinology and Metabolism*, 86, 2792–2796.
- Johnson, J., Han, Z., Otani, K., (2004):** RyR2 and calpain-10 delineate a novel apoptosis pathway in pancreatic islets. *Journal of Biological Chemistry*, 279, 24794-24802.
- Johnstone RW, Ruefli AA, Lowe SW (2002):** Apoptosis: a link between cancer genetics and chemotherapy. *Cell*;108(2):153-164.
- Jones NL, Yeger H, Cutz E, (1996):** Helicobacter pylori induces apoptosis of gastric antral epithelial cells in vivo. *Gastroenterology*;110:A933.
- Joyce, D.E. et al, (2001) :** Gene expression profiling of antithrombotic protein C defines new mechanisms modulating inflammation and apoptosis. *J. Biol. Chem.* 276, 11199–11203
- Juntti-Berggren, L., Refai, E., Appelskog, I., (2004):** Apolipoprotein CIII promotes Ca<sup>2+</sup>-dependent beta cell death in type 1 diabetes. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 10090-10094.
- Kajstura J, Cheng W, Reiss K, Clark WA, Sonnenblick EH, Krajewski S, Reed JC, Olivetti G, Anversa P (1996):** Apoptotic and necrotic myocyte cell deaths are independent contributing variables of infarct size in rats. *Lab Invest*;74:86–107.
- Kakudo, Y., Shibata, H., Otsuka, K., Kato, S., Ishioka, C., (2005):** Lack of correlation between p53-dependent transcriptional activity and the ability to induce apoptosis among 179 mutant p53s. *Cancer Res.* 65, 2108–2114.
- Kang, E., Kim, H., Nam, M., (2006):** A novel 111/121 diplotype in the Calpain-10 gene is associated with type 2 diabetes. *Journal of Human Genetics* (Epub ahead of print).
- Karin, M and Lin, A (2002):** "NF-kappaB at the crossroads 'of life and death.'" *Nat Immunol* 3(3): 221 -227.



- Kasahara Y, Tudor RM, Cool CD, (2001):** Endothelial cell death and decreased expression of vascular endothelial growth factor and vascular endothelial growth factor receptor 2 in emphysema. *Am J Respir Crit Care Med*;163:737-744.
- Kasler HG, Victoria J, Duramad O, Winoto A (2000):** ERK5 is a novel type of mitogen-activated protein kinase containing a transcriptional activation domain. *Mol Cell Biol*, 20:8382-8389.
- Kawaguchi, Y., Okamoto, T., Taniwaki, M., Aizawa, M., Inoue, M., Katayama, S, (1994):** CAG expansions in a novel gene for Machado-Joseph disease at chromosome 14q32.1. *Nature Genet.* 8:221-227.
- Keane, M.M. et al, (1999) :** Chemotherapy augments TRAIL-induced apoptosis in breast cell lines. *Cancer Res.* 59, 734-741.
- Keller, R., Davidson, L.A., Shook, D.R., (2003):** How we are shaped: the biomechanics of gastrulation. *Differentiation* 71, 171–205.
- Kennedy W.A. , Stenberg G. and Lackgre T (1994):** Renal tubular apoptosis after partial uretral obstruction . *J urol*, 152:568-664.
- Kerr JFR, Wyllie AH, Currie AR (1972) :** Apoptosis: A basic biological phenomenon with wide ranging implications in tissue kinetics. *Br J Cancer* 26:239-257.
- Kim H, Liu X, Kobayashi T, (2004):** Reversible cigarette smoke extract induced DNA damage in human lung fibroblasts. *Am J Respir Cell Mol Biol*;31:483-490.
- Kim, K.W., Kim, Y.S., Ha, K.Y., Woo, Y.K., Park, J.B., Park, W.S., An, H.S., (2005):** An autocrine or paracrine Fas-mediated counterattack: a potential mechanism for apoptosis of notochordal cells in intact rat nucleus pulposus. *Spine* 30, 1247–1251.
- Komarov, P.G., Komarova, E.A., Kondratov, R.V., Christov-Tselkov, K., Coon, J.S., Chernov, M.V., Gudkov, A.V., (2003):** A chemical inhibitor of p53 that protects mice from the side effects of cancer therapy. *Science* 285, 1733–1737.



- Kopp C.W, and R. de Martin (2004):** Gene therapy approaches for the prevention of restenosis, *Curr. Vase. Pharmacol.* 2, 183-189.
- Korsmeyer S.J., (1992):** bcl-2 initiates anew category of oncogenes regulators of cell death . *blood* 8:879-886.
- Koseki C; Her Zlinger D. and Al-Awqati Q. (1992):** Apoptosis in metanephric development . *J. cell Biol.* 119,1327-1333.
- Kotani, X et al , (1995):** Apoptosis in thyroid tissue from patients with Hashimoto's tliyroiditis. *Auto immunity* 20, 231-236.
- Kothakota S, Asuma T, Reinhard C (1997):** Caspase-3-generated fragment of gelsolin: effector of morphological change in apoptosis. *Science*;278:294-298.
- Kovalchuk LV, Pavliuk AS, Kasparov AA, Shiglenko NA, Kasparova EA, Sinuhin VN, Chirun NV (2000):** Analysis of pharmacological preparations on the model of apoptosis of human lymphocytes in vitro in norm and immunopathology. *Allergology Immunol*;1(1):24-30.
- Krajweski M, Moss SF, Krajewski S, (1996):** Elevated expression of Bcl-X and reduced Bak in primary coloreetal adenocarcinomas. *Cancer Res*;56:2422-2427.
- Kroemer, G and Reed, JC (2000):** "Mitochondrial control of cell death." *Nat Med* 6(5): 513-9. Kroemer, G, Zamzami, N and Susin, SA (1997). "Mitochondrial control of apoptosis." *Imnuinol Today* 18(1): 44-51.
- Kuang AA, Cado D, Winoto A (1999):** Nur77 transcription activity correlates with its apoptotic function in vivo. *Eur J Immunol*, 29:3722-3728.
- Kuida, K, Zheng, TS, Na, S, Kuan, C, Yang, D, Karasuyama, H, Rakic, P and Flavell, RA (1998):** "Decreased apoptosis in the brain and premature lethality in CPP32-deficient mice." *Nature* 384(6607): 368-372.
- Kuo WH. Chen JH, Lin HH, (2005):** Induction of apoptosis in the lung tissue from rats exposed to cigarette smoke involves p38/JNK MAPK pathway. *Chem Biol Interact*; 155:31-42.



- Kuwano K. (2008):** Involvement of epithelial cell apoptosis in interstitial lung diseases. *Intern Med*;47:345-353.
- Labi, V., Erlacher, M., Kiessling, S., Villunger, A., (2006):** BH3-only proteins in cell death initiation, malignant disease and anticancer therapy. *Cell Death Differ.* 13, 1325–1338.
- LaFerla, F.M., Tinkle, B.T., Bieberich, C.J., Haudenschield, C.C., and Jay, G. (1995):** The Alzheimer's AB peptide induces neurodegeneration and apoptotic cell death in transgenic mice. *Nature Genet.* 9:21-30.
- Lane DP, Midgley CA, Hupp TR, (1995):** On the regulation of the p53 tumour suppressor, and its role in the cellular response to DNA damage. *Philos Trans R Soc Lond Biol*;347:83- 87.
- Larrick J.W and wright S.C (1990):** Cytotoxic mechanisms of tumor necrosis factor- $\alpha$  *FASEB J.*4, 3215-3223.
- Lau HT, Yyu M, Fontana A, Stoeckert Jr CJ (1996):** Prevention of islet allograft rejection with engineered myoblasts expressing FasL in mice. *Science*;273:109-112.
- Lawson K.A; Meneses J.J and pedersen R.A. (1986):** Cell fate and cell lineage in the endoderm of the presomite mouse embryo, studied with an intracellular tracer . *Dev. Biol.* 115, 325-339.
- Le, W.D., Colom, L.V., Wie, W.J., Smith, R.G., Alexianu, M., and Appel, S.H. (1995):** Cell death induced by Bamyloid 1-40 in MES 23.5 hybrid clone: the role of nitric oxide and NMDA-gated channel activation leading to apoptosis. *Brain Res.* 686:49-60.
- Lee FD (1994):** Drug-related pathological lesions of the intestinal tract. *Histopathology*;25:303-308.
- Lee SH, Shin MS, Lee HS, (2001):** Expression of Fas and Fas-related molecules in human hepatocellular carcinoma. *Hum Pathol*;32(3):250-256.
- Lee, S., Ko, G., Li, J., (2001):** Factors predicting the age when Type 2 diabetes is diagnosed in Hong Kong Chinese subjects. *Diabetes Care*, 24, 646-649.



- Lefebvre, S., Burglen, L., Reboullet, S., Clermont, O., Burlet, P., Viollet, L, (1995):** Identification and characterization of a spinal muscular atrophy-determining gene. *Cell* 80:155-165.
- Leist, M and Jaattela, M (2001):** "Four deaths and a funeral: from caspases to alternative mechanisms." *Nat. Rev. Mol. Cell Biol.* 2(8): 589-598.
- Leist, M., and Nicotera, P. (1998):** Apoptosis, excitotoxicity, and neuropathology. *Exp. Cell Res.* 239:183-201.
- Leitges M, Schmedt C, Guinamard R, Davoust J, Schaal S, Stabel S, Tarakhovsky A (1996):** Immunodeficiency in protein kinase cbeta-deficient mice. *Science*, 273:788-791.
- Li H, Kolluri SK, Gu J, Dawson MI, Cao X, Hobbs PD, Lin B, Chen G, Lu J, Lin F et al (2000):** Cytochrome c release and apoptosis induced by mitochondrial targeting of nuclear orphan receptor TR3. *Science*, 289:1159-1164.
- Li LF, Liao SK, Ko YS, (2007):** Hyperoxia increases ventilator-induced lung injury via mitogen-activated protein kinases: A prospective, controlled animal experiment. *Crit Care*; 11R25.
- Li LF, Liao SK, Lee CH, (2005):** Ventilation-induced neutrophil infiltration and apoptosis depend on apoptosis signal-regulated kinase 1 pathway. *Crit Care Med*;33:2118-2119.
- Li LY, Luo X, Wang X (2001):** Endonuclease G is an apoptotic DNase when released from mitochondria. *Nature*, 412:95-99.
- Li Y, Arita Y, Koo HC, Davis JM, (2003):** Inhibition of c-Jun N-terminal kinase pathway improves cell viability in response to oxidant injury. *Am J Respir Cell Mol Biol* ; 29:779-783.
- Li YQ, Fan CY, O'Connor PJ, (1992):** Target cells for the cytotoxic effects of carcinogens in the murine small bowel. *Carcinogenesis*;13:361-368.
- Li, J., Miller, E., Ninomiya-Tsuji, J., (2005):** AMP-activated protein kinase activates p38 mitogen-activated protein kinase by increasing recruitment of p38 MAPK to TAB1 in the ischemic heart. *Circulation Research*, 97, 872-879.



- Lieberthal W. Traica V. , and Levine J. (1996):** Mechanisms of death induced by cisplatin in proximal tubular epithelial cells : Apoptosis VS necrosis . A m .J. physiol. 270 (Renal fluid Electrolyte physiol) ; 39 : F 700- F708.
- Lin, C., Gurlo, T, Haataja, L., (2005):** Activation of peroxisome proliferator-activated receptor-gamma by rosiglitazone protects human islet cells against human islet amyloid polypeptide toxicity by a phosphatidylinositol 3'-kinase-dependent pathway. Journal of Clinical Endocrinology and Metabolism, 90, 6678-6686.
- Liston, P., Roy, N., Tamai, K., Lefebvre, C., Baird, S., Cherton-Horvat, G, (1996):** Suppression of apoptosis in mammalian cells by NAIP and a related family of IAP genes. Nature 379:349-353.
- Liston, P., Young, S.S., Mackenzie, A.E., and Korneluk, R.G. (1997):** Life and death decisions: the role of the IAPS in modulating programmed cell death. Apoptosis 2:423-441.
- Liu X, Conner H, Kobayashi T, Kim H, Wen F, Abe S, (2005):** Cigarette smoke extract induces DNA damage but not apoptosis in human bronchial epithelial cells. Am J Rcspir Cell Mol Biol;33:121-129.
- Liu Y, Cigola E, ChengW, Kajstura J, Olivetti G, Hintze TH, Anversa P (1995):** Myocyte nuclear mitotic division and programmedmyocyte nuclearmitotic division and programmed myocyte cell death characterize the cardiac myopathy induced by rapid ventricular pacing in dogs. Lab Invest;73:771–787.
- Liu, Z.M., Chen, G.G., Ng, E.K., Leung, W.K., Sung, J.J., Chung, S.C., (2004):** Upregulation of heme oxygenase-1 and p21 confers resistance to apoptosis in human gastric cancer cells. Oncogene 23, 503–513.
- Loeffler, M and Kroemer, G, (2000):** "The mitochondrion in cell death control: certainties and incognita." Exp Cell Res 256(1): 19-26.
- Loo, D.T., Copani, A., Pike, C.J., Whittemore, E.R., Walencewicz, J.A., and Cotman, C.W. (1993):** Apoptosis is induced by B-amyloid in cultured central nervous system neurons. Proc. Natl. Acad. Sci. USA 90:7951-7955.



- Lotem J, Sachs L (1995):** A mutant p53 antagonizes the deregulated c-myc-mediated enhancement of apoptosis and decrease in leukemogenicity. *Proc Natl Acad Sci USA* 92: 9672-9676.
- Love, S. (2003):** Apoptosis and brain ischaemia. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 27, 267-282
- Lowin B., Hahne M., Mattmann C., Ischop J., (1994):** Cytolytic Tcell Cytotoxicity is mediated through perforin and fas lytic pathways. *Nature* 370:650-652.
- Luo C, P. Laaja (2004):** Inhibitors of JAKs/STATs and the kinases: a possible new cluster of drugs, *Drug Discov. Today* 9,268–275.
- Luo, X, Budihardjo, I, Zou, H, Slaughter, C and Wang, X, (1998):** "Bid, a Bcl2 interacting protein, mediates cytochrome c release from mitochondria in response to activation of cell surface death receptors." *Cell* 94(4): 481-490.
- Lupi, R., Dotta, F, Marselli, L., (2002):** Prolonged exposure to free fatty acids has cytostatic and pro-apoptotic effects on human pancreatic islets: Evidence that beta-cell death is caspase mediated, partially dependent on ceramide pathway, and Bcl-2 regulated. *Diabetes*, 51, 1437-1443.
- MacNee W, Rahman I (1999):** Oxidants and antioxidants as therapeutic targets in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*; 160:858-865.
- Maedler, K., Spinas, G., Lehmann, R., (2001):** Glucose induces beta-cell apoptosis via upregulation of the Fas(CD95) receptor in human islets. *Diabetes*, 50, 1683-1690.
- Mahida YR, Makh S. Hyde S, (1995) :** Human intestinal epithelial cells produce interleukin-8 (IL-8) and undergo apoptosis following cell detachment induced by *Clostridium difficile* toxin A. *Gastroenterology* ;108:A868.
- Majumder M, Ghosh AK, Steele R, (2002):** Hepatitis C virus NS5A protein impairs TNF-mediated hepatic apoptosis, but not by an anti-FAS antibody, in transgenic mice. *Virology*;294:94-105.



- Mantell LL, Lee PJ (2000):** Signal transduction pathways in hyperoxia-induced lung cell death. *Mol Genet Metab*;71:359-370.
- Mantell LL, Shaffer TH, Horowitz S, (2002):** Distinct patterns of apoptosis in the lung during liquid ventilation compared with gas ventilation. *Am J Physiol* 2002;283: LSI-41.
- Marchetti, P, Decaudin, D, Macho, A, Zamzami, N, Hirsch, T, Susin, SA and Kroemer, G (1997):** "Rcdox regulation of apoptosis: impact of thiol oxidation status on mitochondrial function." *Eur J Immunol* 27(1): 289-296.
- Marchetti, P., Del Guerra, S., Marselli, L, (2004):** Pancreatic islets from type 2 diabetic patients have functional defects and increased apoptosis that are ameliorated by metformin. *Journal of Clinical Endocrinology and Metabolism*, 89, 5535-5541.
- Marsden VS, O'Connor L, O'Reilly LA, Silke J, Metcalf D, Ekert PG, Huang DC, Cecconi F, Kuida K, Tomaselli KJ, (2002):** Apoptosis initiated by Bcl-2-regulated caspase activation independently of the cytochrome c/Apaf-1/caspase-9 apoptosome. *Nature*, 419:634-637.
- Marsden, VS, O'Connor, L, O'Reilly, LA, Silke, J, Metcalf, D, Ekert, PG, Huang, DC, Cecconi, F, Kuida, K, Tomaselli, KJ, Roy, S, Nicholson, DW, Vaux, DL, Bouillet, P, Adams, JM and Strasser, A (2002):** "Apoptosis initiated by Bcl-2-regulated caspase activation independently of the cytochrome c/Apaf-1/caspase-9 apoptosome." *Nature* 419(6907): 634-637.
- Martin P, Duran A, Minguet S, Gaspar ML, Diaz-Meco MT, Rennert P, Leitges M, Moscat J (2002):** Role of zeta PKC in B-cell signaling and function. *EMBO J*, 21:4049-4057.
- Masiello, P. (2006):** Animal models of Type 2 diabetes with reduced pancreatic (i-cell mass. *International Journal of Biochemistry and Cell Biology*, 38, 873-893.
- Matsumura S, Van De Water J, Kita H, (2002):** Contribution to antimito-chondrial antibody production: cleavage of pyruvate dehydrogenase complex-E2 by apoptosis- related proteases. *Hepatology*;35(1):14 22.



- Mattson, M.P. (1998):** Modification of ion homeostasis by lipid peroxidation: role in neuronal degeneration and adaptive plasticity. *Trends in Neurol. Sci.* 21:53-57.
- Mattson, M.P., Cheng, B., Davis, D., Bryant, K., Lieberburg, I., and Rydel, R.E, (1992):** B-amyloid peptides destabilize calcium homeostasis and render human cortical neurons vulnerable to excitotoxicity. *J. Neurosci.* 12:376-389.
- McDonnell TJ, Deane N, Platt FM, Nunez G, Jaeger U, McKearn JP, (1989):** Bcl-2-immunoglobulin transgenic mice demonstrate extended B cell survival and follicular lymphoproliferation. *Cell*;57(1):79-88.
- McGrath-Morrow SA, Stahl J (2001) :** Growth arrest in A549 cells during hyperoxic stress is associated with decreased cyclin B1 and increased p21(Waf1/Cip1/Sd1) levels. *Biochim Biophys Acta*;1538:90-97.
- McGregor, A.M. et al, (1980):** Carbimazole and the autoimmune response in Graves' disease. *New Engl .J. Med.* 303,302-307
- McMurtry MS, Archer SL, Altieri DC, (2005):** Gene therapy targeting survivin selectively induces pulmonary vascular apoptosis and reverses pulmonary arterial hypertension. *J Clin Invest ;* 115:1479-1491.
- McVicker BL, Casey CA (1999):** Effects of ethanol on receptor-mediated endocytosis in the liver. *Alcohol*;19:255-260.
- Mehal WZ, Juedes AE, Crispe IN (1999):** Selective retention of activated CD8+ T cells by the normal liver. *J Immunol*;163(6):3202-3210.
- Merino, D., Giam, M., Hughes, P.D., Siggs, O.M., Heger, K., O'Reilly, L.A., Adams, J.M., Strasser, A., Lee, E.F., Fairlie, W.D., Bouillet, P., (2009):** The role of BH3-only protein Bim extends beyond inhibiting Bcl-2-like prosurvival proteins. *J. Cell Biol.* 186, 355–362.
- Merritt A J, Potten CS, Kemp CJ, (1994):** The role of p53 in spontaneous and radiation-induced apoptosis in the gastrointestinal tract of normal and p53-deficient mice. *Cancer Res*;54:614-617.



- Metzstein MM, Hengartner MO, Tsung N, Ellis RE, Horvitz HR (1996):** Transcriptional regulator of programmed cell death encoded by *Caenorhabditis elegans* gene *ces-2*. *Nature* 382: 545-547.
- Milligan C.E; Oppenheim Row. And Schwart Z. L.M. (1994):** Motoneurons deprived of trophic support in vitro require of trophic support in vitro require new gene expression to undergo programmed . cell death. *J. Neurobiol* 25, 1005 -1016.
- Missero C, Calautti E, Eckner R, Chin J, Tsai LII, Livingston DM, Dotto GP (1995):** Involvement of the cell-cycle inhibitor Cipl/WAFI and the E1A-associated p300 protein in terminal differentiation. *Proc Natl Acad Sci USA* 92: 5451-5455.
- Mitsiades, N. et al (2001):** Ewing's sarcoma family tumors are sensitive to tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) and express DR4 and DR5 receptors. *Cancer Res.* 61, 2704-2712.
- Mitsiades, N. et al, (1999):** Fas ligand expression in thyroid carcinomas: a potential mechanism of immune evasion.,/. *Clin. Endocrinol. Metab.* 84, 2924-2932.
- Mitsiades, N. et al, (1998):** Fas/Fas ligand up-regulation and Bcl-2 down-regulation may be significant in the pathogenesis of Hashimoto's thyroiditis. \*/. *Clin. Endocrinol. Metab.* 83, 2199-2203.
- Mitsiades, N. et al, (2000):** Fas ligand expression in thyroid follicular cells from patients with thionamide-treated Graves' disease. *Thyroid* 10, 527-532.
- Mitsiades, N. et al, (2000):** Thyroid carcinoma cells are resistant to FAS-mediated apoptosis but sensitive to tumor necrosis factor-related apoptosis-inducing ligand. *Cancer Res.* 60, 4122-4129.
- Miyamoto A, Nakayama K, Imaki H, Hirose S, Jiang Y, Abe M, Tsukiyama T, Nagahama H, Ohno S, Hatakeyama S, (2002):** Increased proliferation of B cells and auto-immunity in mice lacking protein kinase C $\delta$ . *Nature*, 416:865-869.



- Miyashita T, Harigal M, Hanada M, (1994):** Identification of a p53- dependent negative response element in the bcl-2 gene. *Cancer Res*;54(12):3131-3135.
- Mody N, Leitch J, Armstrong C, Dixon J, Cohen P (2001):** Effects of MAP kinase cascade inhibitors on the MKK5/ERK5 pathway. *FEBS Lett*, 502:21-24.
- Morgan, R., Nalliah, A., Morsi El-Kadi, A.S., (2004):** FLASH, a component of the FAS–CASPASE8 apoptotic pathway, is directly regulated by Hoxb4 in the notochord. *Dev. Biol.* 265, 105–112.
- Morse D, Otterbein LE, Watkins S, (2003):** Deficiency in the c-Jun NH2-terminal kinase signaling pathway confers susceptibility to hyperoxic lung injury in mice. *Am J Physiol Lung Cell Mol Physiol*;285:L250-257.
- Mu, X., He, J., Anderson, D.W. Trojanowski, J.Q., and Springer, J.E. (1996):** Altered expression of bcl-2 and bax mRNA in amyotrophic lateral sclerosis spinal cord motor neurons. *Ann. Neurol.* 40:379-386.
- Mund, T, Gewies, A, Sctioenfeld, N, Bauer, MK and Grimm, S (2003):** "Spike, a novel BII3-only protein, regulates apoptosis at the endoplasmic reticulum." *Faseb J* 17(6): 696-8.
- Murphy AN, Fiskum G, Beal MF (1999):** Mitochondria in neurodegeneration: Bioenergetic function in cell life and death. *J Cereb Blood Flow Metab*; 19:231-245.
- Muschen M, Re D, Braunmger A, Wolf J, Hansmann ML, Diehl V, (2000):** Somatic mutations of the CD95 gene in Hodgkin and Reed-Sternberg cells. *Cancer Res*;60(20):5640- 5643.
- Muschen, M. et al, (2000):** Defining CD95 as a tumor suppressor gene. *J. Mot. Med.* 78, 312-325.
- Muzio M. Arul M, Chinnaiyan FC, (1996):** FLICE, A Novel FADD homologous ICE/CED-3-like protease, is recruited to the CD95 (Fas/Apo-1) death-inducing signalling complex. *Cell* :85:817-827.
- Myc, A. et al, (1999) :** Characterization of FAP-1 expression and function in thyroid follicular cells. *Endocrinology* 140, 5431-5434.



- Nagata S, Goldstein P. *The Fas death factor* (1995):** Science;267:1449–1456.
- Naismith, JH and Sprang, SR, (1998):** "Modularity in the TNF-receptor family." Trends Biochem Sci 23(2): 74-79.
- Nakayama K., Wegishi I., Kuida . K., Sawa H., Loh D.Y., (1994):** Targeted disruption of bcl-2 in mice ; occurrence of grey hair, PKD and lymphocytopenia. Proc Natl Acad sci USA 91:3700-3704.
- Natori, S. et al. (2003):** The caspase inhibitor IDN-6556 prevents caspase activation and apoptosis in sinusoidal endothelial cells during liver preservation injury. Liver Transpl. 9, 278–284.
- Newton K, Harris AW, Bath ML, Smith KGC, Strasser A, (1998):** Adominant interfering mutant of FADD/MORT1 enhance deletion of autoreactive thymocytes and inhibits proliferation of mature T lymphocytes. EMBO J, 17:706-718.
- Nishina H, Radvanyi L, Raju K, Sasaki T, Kozieradzki I, Penninger JM (1998):** Impaired TCR-mediated apoptosis and Bel-XL expression in T cells lacking the stress kinase activator SEK1/MKK4 J Immunol, 161:3416-3420.
- Nolsoe, R., Hamid, Y., Pociot, F., (2006):** Association of a microsatellite in FASL to type II diabetes and of the FAS-670G>A genotype to insulin resistance. Genes and immunity (Epub ahead of print).
- Nzeako UC, Guicciardi ME, Yoon JH, Bronk SF, Gores GJ (2002):** COX-2 inhibits Fas-mediated apoptosis in cholan-giocarcinoma cells. Hepatology;35(3):552-559.
- O'Connell J, O'Sullivan G, Collins JK, (1996):** The fas counterattack- another mode of lymphoepithelial communication. Gastroenterology ;110:A981.
- Odin JA, Huebert RC. Casciola-Rosen L, LaRusso NF, Rosen A (2001):** Bcl-2-dependent oxidation of pyruvate dehy-drogenase-E2, a primary biliary cirrhosis autoantigen, during apoptosis. J Clin Invest;108(2):223-232.
- Offen, D., Ziv, I., Panel, H., Wasserman, L., Stein, R., Melamed, E., and Barzilai, A, (1997):** Dopamineinduced apoptosis is inhibited in PC12 cells



- expressing Bcl-2. *Cell. Mol Neurobiol.* 17:289-304.
- Ofner D, Riehemann K, Maier H, (1995) :** Immunohistochemically detectable bcl-2 expression in colorectal carcinoma: Correlation with tumour stage and patient survival. *Br J Cancer*;72:981- 985.
- Ogasawara, J. et al, (1993) :** Lethal effect of the anti-Fas antibody in mice. *Nature* 364, 806-809.
- Ohnishi T, Wang X, Ohnishi K, Matsumoto II, Takahashi A (1996):** p53-dependent induction of WAF 1 by heat treatment in human glioblastoma cells. *J Biol Chem* 271: 14510-14513.
- Okaro AC, Fennell DA, Corbo M, Davidson BR (2002):** Cotter FE. Pk11195, a mitochondrial benzodiazepine receptor antagonist, reduces apoptosis threshold in Bcl-X(L) and Mcl-1 expressing human cholangiocarcinoma cells. *Gut*;51(4):556-561.
- Olivetti G, Abbi R, Quaini F, Kajstura J, ChengW, Nitahara JA, Quaini E, Loreto CD, Beltrami CA, Krajewski S, Reed JC, Anversa P (1997):** Apoptosis in the failing human heart. *N Engl J Med*;336:1131–1141.
- O'Reilly MA, Staversky RJ, Watkins RH, (2001):** The cyclin-dependent kinase inhibitor p21 protects the lung from oxidative stress. *Am J Respir Cell Mol Biol* 2001;24:703-710.
- Osborn B (1996):** Apoptosis and the maintenance of homeostasis in the immune system. *Curr Opin Immunol*;8:245-254.
- Otsuka M, Kato N, Taniguchi H, (2002):** Hepatitis C virus core protein inhibits apoptosis via enhanced Bcl-xL expression. *Virology*;296:84-93.
- Ow, Y.P., Green, D.R., Hao, Z., Mak, T.W., (2008):** Cytochrome c: functions beyond respiration. *Nat. Rev. Mol. Cell Biol.* 9, 532–542.
- Owen-Schaub, L.B. et al, (1998):** Fas and Fas ligand interactions suppress melanoma lung metastasis. *J. Exp. Med.* 188, 1717-1723.
- Packer M, (1995):** Is tumor necrosis factor an important neurohormonal mechanism in chronic heart failure? *Circulation*;95:1379–1382.



- Pages G, Guerin S, Grail D, Bonino F, Smith A, Anjuere F, Auberger P, Pouyssegur J (1999):** Defective thymocyte maturation in p44 MAP kinase (Erk 1) knockout mice. *Science*, 286:1374-1377.
- Pan J, Duan LX, Sun BS, Feitelson MA (2001):** Hepatitis B virus X protein protects against anti-Fas-mediated apoptosis in human liver cells by inducing NF-kappa B. *J Gen Virol* ;82:171-182.
- Paradis, E., Douillard, H., Koutroumanis, M., Goodyer, C., and LeBlanc, A. (1996):** Amyloid B peptide of Alzheimer's disease downregulates Bcl-2 and upregulates Bax expression in human neurons. *J. Neurosci.* 16:7533-7539.
- Pardridge, W.M, (2002):** Neurotrophins, neuroprotection and the blood–brain barrier. *Curr. Opin. Investig. Drugs* 3, 1753–1757.
- Park JW, Kim HP, Lee SJ, (2008):** Protein kinase C {alpha} and {zeta} differentially regulate death-inducing signaling complex formation in cigarette smoke extract-induced apoptosis. *J Immunol*; 180:4668-4678.
- Pastorino JG, Hoek JB, (2000):** Ethanol potentiates tumor necrosis factor-alpha cytotoxicity in hepatoma cells and primary rat hepatocytes by promoting induction of the mitochondrial permeability transition. *Hepatology*,31:1141-1152.
- Paulson, H.L., Perez, M.K., Trotter, Y., Trojanowski, J.Q., Subramony, S.H., Das, S.S, (1997b):** Intranuclear inclusions of expanded polyglutamine protein in spinocerebellar ataxia type 3. *Neuron* 19:333-344.
- Pekarsky Y, Hallas C, Palamarchuk A, Koval A, Bullrich F, Hirata Y, Bichi R, Letofsky J, Croce CM (2001):** Akt phosphorylates and regulates the orphan nuclear receptor Nur77. *Proc Natl Acad Sci USA*, 98:3690-3694.
- Pesce M; Farrace M.G; Piecentini M; Dolci S, and De Felici M, (1993):** Stem cell factor and leukemia inhibitory factor promote primordial germ cell survival by suppressing programmed cell death (apoptosis). *Development (Cambridge , UK)* 118, 1089-1094.



- Petrache 1, Natarajan, V, Zhen L, (2005):** Ceramide upregulation causes pulmonary cell apoptosis and emphysema-like disease in mice. *Nat Med* 11:491-498.
- Petrache L Choi ME, Otterbein LE, (1999):** Mitogen-activated protein kinase pathway mediates hyperoxia-induced apoptosis in cultured macrophage cells. *Am J Physiol*;277: L589-595.
- Petrovsky, N., Silva, D., Socba, L., (2002):** The role of Fas(CD95) ligand in beta cell destruction in autoimmune diabetes of NOD mice. *Annals of the New York Academy of Sciences*, 958, 204-208.
- Pexieder T, (1975):** Cell death in the morphogenesis and teratogenesis of the heart. *Adv. Anat. Embryol. Cell Biol* 51, 1-100.
- Peyrou M., Hanna P.E., Cribb A.E., Cisplatin, (2007):** Gentamicin, and p-aminophenol induce markers of endoplasmic reticulum stress in the rat kidneys, *Toxicol. Sci.* 99 , 346–353.
- Piazza F, Gurrieri C, Pandolfi PP. The theory of APL (2001):** *Oncogene*;20(49): 7216-7222.
- Pinkovski M, Hobman M, Heibin J (1998):** Entry and trafficking of granzyme B during apoptosis. *Blood*;92:1044-1054.
- Poelmann R.F and Vermeij-Keers C, (1976):** Cell degeneration in the mouse embryo: A prerequisite for normal development. In "Progress in Differentiation Research" (N. Müller-Bérat, ed.), North-Holland Pub., Amsterdam pp 93-102.
- Ponnio T, Burton Q, Pereira FA, Wu DK, Conneely OM (2002):** The nuclear receptor Nor-1 is essential for proliferation of the semicircular canals of the mouse inner ear. *Mol Cell Biol*, 22:935-945.
- Potten CS, Li YQ, O'Connor PJ, (1992):** A possible explanation for the differential cancer incidence in the intestine, based on distribution of the cytotoxic effects of carcinogens in the murine large bowel. *Carcinogenesis*;13:2305-2312.
- Pryhuber GS, O'Brien DP, Baggs R, (2000):** Ablation of tumor necrosis factor receptor type 1 (p55) alters oxygen-induced lung injury. *Am J Physiol*;278:L 1082-1090.



- Puthalakath H, Huang DC, O'Reilly LA, King SM, Strasser A, (1999):** The proapoptotic activity of the Bcl-2 family member Bim is regulated by interaction with the dynein motor complex. *Mol Cell*, 3:287-296.
- Puthalakath, H and Strasser, A, (2002):** "Keeping killers on a tight leash: transcriptional and post-translational control of the pro-apoptotic activity of BH3-only proteins." *Cell Death Differ* 9(5): 505-512.
- Qu, H., Bharaj, B., Liu, X., (2005):** Assessing the validity of the association between the SUMO4 M55V variant and risk of type 1 diabetes. *Nature Genetics*, 37, 112.
- Rabizadeh, S., Bitler, C.M., Butcher, L.L., and Bredesen, D.E. (1994):** Expression of the low-affinity nerve growth factor receptor enhances B-amyloid peptide toxicity. *Proc. Natl. Acad. Sci. USA* 91:10703-10706.
- Rabizadeh, S., Gralla, E.B., Borchelt, D.R., Gwinn, R., Valentine, J.S., Sisodia, S, (1995):** Mutations associated with amyotrophic lateral sclerosis convert superoxide dismutase from an antiapoptotic gene to a proapoptotic gene: studies in yeast and neural cells. *Proc. Natl. Acad. Sci. USA* 92:3024-3028.
- Racusen L., Fivush B. and statniky, (1991):** Dissociation of tubular cell detachment and tubular cell death in clinical and experimental acute tubular necrosis *Lab. Invest*; 69:456-555.
- Ramenghi U, Bonisconi S, Migliaretti G, DeFranco S, Bottarel F, Gambaruto C, (2000):** Deficiency of the Fas apoptosis pathway without Fas gene mutations is a familial trait predisposing to development of autoimmune diseases and cancer. *Blood*;95(10): 3176-3182.
- Rathmell JC, Lindsten T, Zong WX, Cinalli RM, Thompson CB (2002):** Deficiency in Bak and Bax perturbs thymic selection and lymphoid homeostasis. *Nat Immun*, 3:932-939.
- Raveendran M, Wang J, Senthil D, (2005):** Endogenous nitric oxide activation protects against cigarette smoking induced apoptosis in endothelial cells. *FEES Lett*; 579:733-740.



- Reich 'K, Lingnau F, Williams RM, (1996):** Corticosteroids downregulate bcl-2 and induce apoptosis in CD4+ LPL in Crohn's disease. *Gastroenterology*, 110:A999.
- Riachy, R., Vandewalle, B., Moerman, E., (2006):** 1,25 Dihydro-vitamin D3 protects human pancreatic islets against cytokine-induced apoptosis via down-regulation of the Fas receptor. *Apoptosis*, 11, 151-159.
- Richardson, H and Kumar, S (2002):** "Death to flies: Drosophila as a model system to study programmed cell death." *J Immunol Methods* 265(1-2): 21-38.
- Ritzel, R., & Butler, P. (2003):** Replication increases beta-cell vulnerability to human islet amyloid polypeptide-induced apoptosis. *Diabetes*, 52,1701-1708.
- Rivero M, Crespo J, Fabrega E, (2002):** Apoptosis mediated by the Fas system in the fulminant hepatitis by hepatitis B virus. *J Viral Hepatitis*;9:107-113.
- Robles MS, Leonardo E, Criado LM, Izquierdo M, Martinez AC, (2002):** Inhibitor of apoptosis protein from *Ornyia pseudotsugata* nuclear polyhedrosis virus provides a costimulatory signal required for optimal proliferation of developing thymocytes. *J Immunol*, 168:1770-1779.
- Romashko J, Horowitz S, Franek WR, (2003):** MAPK pathways mediate hyperoxia-induced oncotic cell death in lung epithelial cells. *Free Radic Biol Med* ;35: 978-993.
- Rothstein, J.F., Van Kammen, M., Levey, A.I., Martin, L.J., and Kuncel, R.W. (1995):** Selective loss of glial glutamate transporter GLT-1 in amyotrophic lateral sclerosis. *Ann. Neurol.* 38:73-84.
- Rouvier E, Luciani M-F, Golstein P, (1993):** Fas involvement in  $Ca^{2+}$ -independent T cell-mediated cytotoxicity. *J Exp Med*; 177:195-200.
- Roy, N., Mahadevan, M.S., McLean, M., Shutler, G., Yaraghi, Z., Farahani, R, (1995):** The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy. *Cell* 80:167-178.
- Rudin C.M, M. Kozloff, P.C. Hoffman, M.J. Edelman, R. Karnauskas, R. Tomek, L. Szeto, E.E. Vokes, (2004):** Phase I study of G3139, a bcl-2



- antisense oligonucleotide, combined with carboplatin and etoposide in patients with small-cell lung cancer, *J. Clin. Oncol.* 22 (2004) 1110–1117.
- Rumora, L., Hadzija, M., Barisic, K., (2002):** Amylin-induced cytotoxicity is associated with activation of caspase-3 and MAP kinases. *Biological Chemistry*, 383, 1751-1758.
- Ryan KM, Birnie GD (1996):** Myc oncogenes: the enigmatic family. *Biochem J* 314: 713-721.
- S. Pryshchep, K. Sato, J.J. Goronzy, C.M. Weyand , (2006):** Tcell recognition and killing of vascular smooth muscle cells in acute coronary syndrome, *Circ. Res.* 98 , 1168-1 176.
- Sabapathy K, Kallunki T, David JP, Graef I, Karin M, Wagner EF (2001):** c-Jun NH2-terminal kinase (JNK)1 and JNK2 have similar and stage-dependent roles in regulating T cell apoptosis and proliferation. *J Exp Med*, 193:317-328.
- Saikumar P, Dong Z, Weinberg JM, (1998):** Mechanisms of cell death in hypoxia/reox-ygenation injury. *Oncogene*; 17:3341-3349.
- Sakao S, Taraseviciene-Stewart L, Lee JD, (2005):** Initial apoptosis is followed by increased proliferation of apoptosis-resistant endothelial cells. *FASEB J*;19:1178-1180.
- Saldeen, J. (2000):** Cytokines induce both necrosis and apoptosis via a common Bcl-2-inhibitable pathway in rat insulin-producing cells. *Endocrinology*, 141, 2003-2010.
- Salvesen OS, Duckett CS (2002):** IAP proteins: blocking the road to death's door. *Nat Rev Mol Cell Biol*;3(6):401-410.
- Salvesen, GS and Duckett, CS (2002a):** "IAP proteins: blocking the road to death's door." *Nat Rev Mol Cell Biol* 3(6): 401-410.
- Salvesen, GS and Renatus, M (2002b):** "Apoptosome: the seven-spoked death machine." *Dev Cell* 2(3): 256-257.



- Sanders E.J. and wride MA, (1995):** Programmed cell death in development, International review of cytology, 163, 105:173.
- Sanders E.J; Khare M.K; Oai V.C, and Bellairs R. (1986):** An experimental and morphological analysis of the tail bud mesenchym of the chick embryo. Anat, embryol, 174, 179-185.
- Saraste, A and Fulkki, K (2000):** "Morphologic and biochemical hallmarks of apoptosis." Cardiovasc Res 45(3): 528-537.
- Savill J (1995):** Apoptosis and renal injury Curr, Opin. Nephrol, Hypertns.;4 (3); 263-269.
- Savill J. Mooney A, and Hughes J, (1996):** what role does apoptosis play in progression of renal disease. Curr. Opin, NephroL Hypertens; 5(4): 364-369.
- Savitz, S.I., and Rosenbaum, D.M, (1998):** Apoptosis in neurological diseases. Neurosurgery 42:555-574.
- Sawyers C.L., E.A. Sausville, Overview (2002):** the art of cancer drug screening: molecular target versus milieu-based screens, Curr. Opin. Investig. Drugs 3 (2002) 478– 481.
- Scaffidi, C, Fulda, S, Srinivasan, A, Friesen, C, Li, F, Tomaselli, KJ, Dcbatin, KM, Krammcr. PH and Peter, ME (1998):** "Two CD95 (APO-1/Fas) signaling pathways." Embo J 17(6): 1675-1687.
- Schimmer AD, Hedley DW, Pemi LZ, Minden MD (2001):** Receptor and mitochondrial-mediated apoptosis in acute leukemia: a translational view. Blood;98(13):3541-3553.
- Schirmer M, Vallejo AN, Weyand CM, Goronzy JJ, (1998):** Resistance to apoptosis and elevated expression of Bcl-2 in clonally expanded CD4<sup>+</sup>CD28<sup>-</sup> T cells from rheumatoid arthritis patients. J Immunol: 161:1018-1025.
- Schrank, B., Gotz, R., Gunnersen, J.M., Ure, J.M., Toyka, K.V., Smith, A.G., and Sendtner M. (1997):** Inactivation of the survival motor neuron gene, a candidate gene for human spinal muscular atrophy, leads to massive cell



- death in early mouse embryos. *Proc. Natl. Acad. Sci. USA* 94:9920-9925.
- Schumer M, Colombel MC, Sawczuk IS, Gobe G, Connor J, O'Toole KM, Wise GJ, Buttyan R (1992):** Morphologic, biochemical & molecular evidence of apoptosis during the reperfusion phase after brief periods of renal ischemia. *Am J Pathol*;140:831–838.
- Schwartz LM, Smith SW, Jones MEE, (1993):** Do all programmed cell deaths occur via apoptosis? *PNAS* 90:980-984.
- Segura-Valdez L, Pardo A, Gaxiola M, (2000):** Upregulation of gelatinases A and B, collagenases 1 and 2, and increased parenchymal cell death in COPD. *Chest*; 117: 684-694.
- Selman M, King TE, Pardo A, (2001):** American Thoracic Society, European Respiratory Society, American College of Chest Physicians. Idiopathic pulmonary fibrosis: Prevailing and evolving hypotheses about its pathogenesis and implications for therapy. *Ann Intern Med*; 134:136-151.
- Server A.C. and Mobley W.C (1991):** Neuronal cell death and the role of apoptosis In "apoptosis: the molecular basis of cell death" (L.D, Tomei and F.O. Cope, eds.) Cold spring harbor lab. Press, Plainview, NY, pp. 263-278.
- Shah N.P., Tran C., Lee F.Y., Chen P., Norris D., Sawyers C .L, (2004):** Overriding imatinib resistance with a novel ABL kinase inhibitor, *Science* 305 , 399- 401.
- Shapiro SD (2004):** Smoke gets in your cells. *Am J Respir Cell Mol Biol*;31:4812.
- Shaw P, Bovey R, Tardy S, (1992):** Induction of apoptosis by wildtype p53 in a human colon tumour-derived cell line. *Proc Natl Acad Sci USA*;89:4495-4499.
- Shetty V, Hussaini S, Broady-Robinson L, Allampallam K, Mundle S, Borok R, (2000):** Intramedullary apoptosis of hematopoietic cells in myelodysplastic syndrome patients can be massive: apoptotic cells recovered from high-density fraction of bone marrow aspirates. *Blood*;96(4): 1388-1392.



- Shi, Y. (2002) :** Mechanisms of caspase activation and inhibition during apoptosis. *Mol. Cell* 9, 459-470.
- Shibue, T., Takeda, K., Oda, E., Tanaka, H., Murasawa, H., Takaoka, A., Morishita, Y., Akira, S., Taniguchi, T., Tanaka, N., (2003):** Integral role of Noxa in p53-mediated apoptotic response. *Genes Dev.* 17, 2233–2238.
- Shimizu A. and yamanaka N (1993):** Apoptosis and cell desquamation in repair process of ischemic tubular necrosis. *Curr. Opin. Nephrol. Hypertens*;8(2): 171-180.
- Shimizu A., Kitamura H., Masudav., Ishizaki M., Sugisaki Y., Yamanaka N., (1995):** Apoptosis in repair process of experimental proliferative glomerulonephritis . *Kidney Int* 47:114-121.
- Siegel RM, Chan FK, Chun HJ, Lenardo MJ (2000):** The multifaceted role of Fas signaling in immune cell homeostasis and autoimmunity. *Nat Immunol* ;1(6):469-474.
- Simunovic F, Yi M, Wang Y, Macey L, Brown LT, Krichevsky AM, Andersen SL, Stephens RM, Benes FM, Sonntag Kc, (2009):** *Brain*;132(Pt 7):1795-1809.
- Slebos DJ, Ryter SW. van der Toorn M, (2006):** Mitochondria! localization and function of heme oxygenase-1 in cigarette smoke-induced cell death. *Am J Respir Cell Mol Biol* ;36:409-417.
- SmeyneRJ, Vendrell M, Hayward M. Baker SJ, Miao GG, Schilling K, Robertson LM.CurranT, Morgan JI (1993):** Continuous *c-fos* expression precedes programmed cell death in vivo. *Nature* 363: 166-169.
- Sohn SJ, Sarvis BK, Cado D, Winoto A (2002):** ERK5 MAP kinase regulates embryonic angiogenesis and acts as a hypoxia-sensitive repressor of VEGF expression. *J Biol Chem*, 277:43344-43351.
- Solary E, Droin N, Bettaieb A, Corcos L, Dimanche-Boitrel MT, Gamdo C (2000):** Positive and negative regulation of apoptotic pathways by cytotoxic agents in hematological malignancies. *Leukemia*:14:1833-1849.



- Stahnke K, Hecker S, Kohne E, Debatin K-M (1998):** CD95 (APO-1/FAS) mediated apoptosis in cytokine-activated hematopoietic cells. *Exp Hematol*;26:844-850.
- Stassi, G. D., De Maria, R., Trucco, G., (1997):** Nitric oxide primes pancreatic beta cells for Fas(CD95)-mediated destruction in insulin-dependent diabetes mellitus. *Journal of Experimental Medicine*, 186, 1193-1200.
- Stassi, G. et al, (2000) :** Control of target cell survival in thyroid autoimmunity by T helper cytokines via regulation of apoptotic proteins. *Nat. Immunol.* 1. 483-488.
- Stegh AH, Barnhart BC, Volkland J, (2002):** Inactivation of caspase-8 on mitochondria of Bcl-xL-expressing MCF7-Fas cells: Role for the bifunctional apoptosis regulator protein. *JBiol Chem*;277:435I- 4360.
- Steinberg KP, Hudson LD (2000) :** Acute lung injury and acute respiratory distress syndrome: The clinical syndrome. *Clin Chest Med*;21:402-417.
- Stemple, D.L., (2005):** Structure and function of the notochord: an essential organ for chordate development. *Development* 132, 2503–2512.
- Sterling, J., Binzer, J., Andersson, A., (2005):** Nitric oxide contributes to cytokine-induced apoptosis in pancreatic beta cells via potentiation of INK activity and inhibition of Akt. *Diabetologia*, 48, 2039-2050.
- Strand, S. et al, (1996):** Lymphocyte apoptosis induced by CD95 (APO-1/Fas)ligand-expressing tumor cells-a mechanism of immune evasion? *Nat.Med.* 2, 1361-1366.
- Strasser A, Harris AW, Bath ML, Cory S (1990):** Novel primitive lymphoid tumours induced in transgenic mice by cooperation between myc and bcl-2. *Nature*;348(6299):331-333.
- Strasser, A, O'Connor, L and Dixit, VM (2000):** "Apoptosis signaling." *Annu Rev Biochem* 69: 217-245.
- Straus SE, Jaffe ES, Puck JM, Dale JK, Elkon KB, Rosen-Wolff A, (2001):** The development of lymphomas in families with autoimmune lymphoproliferative syndrome with germline Fas mutations and defective lymphocyte apoptosis. *Blood*;98(1):194-200.



- Strittmatter, W.J., Saunders, A.M., Schmechel, D., Pericak-Vance, M., Enghild, J., Salvesen, G.S., and Roses A.D. (1994):** Apolipoprotein E: high-avidity binding to beta-amyloid and increased frequency of type 4 allele in late-onset familial Alzheimer disease. *Proc. Natl. Acad. Sci. USA* 90:1977-1981.
- Su, J.H., Deng, G., and Cotman, C.W. (1997):** Bax protein expression is increased in Alzheimer's brain: correlations with DNA damage, Bcl-2 expression and brain pathology. *J. Neuropathol. Exp. Neurol.* 56:86-93.
- Suda T., and Nagata S., (1994):** Purification and characterization of the fas ligand that induces apoptosis. *J Exp Med* 179: 873-879.
- Sun E, Gao Y, Chen J, (2004):** Allograft tolerance induced by donor apoptotic lymphocytes requires phagocytosis in the recipient. *Cell Death Differ*;11:1258–1264.
- Sun W, Kesavan K, Schaefer BC, Garrington TP, Ware M, Johnson NL, Gelfand EW, Johnson GL, (2001):** MEKK2 associates with the adapter protein Lad/RIBP and regulates the MEK5-BMK1/ ERK5 pathway. *J Biol Chem*, 276:5093-5100.
- Suzuki A, Yamaguchi MT, Ohteki T, Sasaki T, Kaisho T, Kimura Y, Yoshida R, Wakeham A, Higuchi T, Fukumoto M, (2001):** T cell-specific loss of Pten leads to defects in central and peripheral tolerance. *Immunity*, 14:523-534.
- Suzuki, Y, Nakabayashi, Y, Nakata, K, Reed, JC and Takahashi, R (2001):** "X-linked inhibitor of apoptosis protein (XIAP) inhibits caspase-3 and -7 in distinct modes." *J Biol Chem* 276(29): 27058-27063.
- Swat W, Fujikawa K, Ganiatsas S, Yang D, Xavier RJ, Harris NL, Davidson L, Ferrini R, Davis RJ, Labow MA (1998):** SEK1/MKK4 is required for maintenance of a normal peripheral lymphoid compartment but not for lymphocyte development. *Immunity*, 8:625-634.
- Takahashi T, Tanaka M, Brannan CI (1994):** Generalized lymphoproliferative disease in mice, caused by a point mutation in the Fas ligand. *Cell*;76:969–976.



- Takahashi, R, Deveraux, Q, Tamm, I, Welsh, K, Assa-Munt, N. Salvesen, GS and Rood, JC (1998):** "A single SIR domain of XIAP sufficient for inhibiting caspases." *J Biol Chem* 273(14): 7787-7790.
- Takayama S, Sato T, Krajewski S, (1995):** Cloning and functional analysis of BAG-1: A novel Bcl-2 binding protein with anti-cell death activity. *Cell*;80:279-284.
- Takayama S., Sato T., Krajewski S., Kochel K., Irie S., Millan J.A., (1995):** Cloning and functional analysis of BAG -1 : A novel bcl-2 binding protein with anti-cell death activity . *cell* 80:279-284.
- Takemura T., Katsumi M., Hirofumi M., (1995):** Expression of fas antigen and bcl-2 in human glomerulonephritis . *Kidney Int* 48:1886-1892.
- Tan, J., Zhuang, L., Leong, H.S., Iyer, N.G., Liu, E.T., Yu, Q., (2005):** Pharmacologic modulation of glycogen synthase kinase-3 $\beta$  promotes p53-dependent apoptosis through a direct Bax-mediated mitochondrial pathway in colorectal cancer cells. *Cancer Res.* 65, 9012–9020.
- Tanaka M, Ito H, Adachi S, Akimoto H, Nishikawa T, Kasajima T, Marumo F, Hiroe M, (1994):** Hypoxia induces apoptosis with enhanced expression of Fas antigen messenger RNA in cultured neonatal rat cardiomyocytes. *Circ Res*;75: 426–433.
- Tanaka N, Kamanaka M, Enslen H, Dong C, Wysk M, Davis RJ, Flavell RA (2002):** Differential involvement of p38 mitogen-activated protein kinase kinases MKK3 and MKK6 in T-cell apoptosis. *EMBO Rep*, 3:785-791.
- Taraseviciene-Stewart L, Scerbavicius R, Choe KH, (2006):** Simvastatin causes endothelial cell apoptosis and attenuates severe pulmonary hypertension. *Am J Physiol Lung Cell Mol Physiol*;291 :L668-676.
- Tatton, N.A., and Kish, S.J. (1997):** In situ detection of apoptotic nuclei in the substantia nigra compacta of MPTP-treated mice using TdT labeling and acridine orange staining. *Neurosci.* 77:1037-1048.
- Taylor, R.C., Cullen, S.P., Martin, S.J., (2008):** Apoptosis: controlled demolition at the cellular level. *Nat. Rev. Mol. Cell Biol.* 9, 231–241.



- Thallinger C, M.F. Wolschek, H. Maierhofer, H. Skvara, H. Pehamberger, B.P. Monia, B. Jansen, V. Wacheck, E. Selzer , (2004):** Mcl-1 is a novel therapeutic target for human sarcoma: synergistic inhibition of human sarcoma xenotransplants by a combination of Mcl-1 antisense oligonucleotides with low-dose cyclophosphamide, Clin. Cancer Res. 10 , 4185– 4191.
- Thiagarajan RR, Winn RK, Harlan JM (1997):** The role of leukocyte and endothelial adhesion molecules in ischemia-reperfusion injury. Thromb Haemost;78:310 4.
- Thompson CB (1996):** A fate worse than death. Nature 382: 492-493.
- Thompson CB, 1995:** Apoptosis in the pathogenesis and treatment of disease. Science;267:1456-1462.
- Tominaga T, Kure S, Narisawa K, Yoshimoto T (1993):** Endonuclease activation following focal ischemic injury in the rat brain. Brain Res;608:21–26.
- Tompkins, MM., Basgall, E.J., Zamrini, E., and Hill, W.D, (1997):** Apoptotic-like changes in Lewy-body-associated disorders and normal aging in substantia nigral neurons. Am. J. Pathol. 150:119-131.
- Tovar, C., Rosinski, J., Filipovic, Z., Higgins, B., Kolinsky, K., Hilton, H., Zhao, X., Vu, B.T., Qing, W., Packman, K., Myklebost, O., Heimbrook, D.C., Vassilev, L.T., (2006):** From the cover: smallmolecule MDM2 antagonists reveal aberrant p53 signaling in cancer: implications for therapy. Proc. Natl. Acad. Sci. (USA) 103, 1888–1893.
- Troncoso, J.C., Sukhov, R.R., Kawas, C.H., and Koliatsos, V.E, (1996):** In situ labeling of dying cortical neurons in normal aging and in Alzheimer's disease: correlations with senile plaques and disease progression. J. Neuropathol. Exp. Med. 55:1134-1142.
- Trudeau, .1., Dutz, J., Arany, E., (2000):** Neonatal beta-cell apoptosis: A trigger for autoimmune diabetes? Diabetes, 49, 1-7.



- Tsujiim and Dubois R. (1995):** Alterations in cellular adhesion and apoptosis in epithelial cells over expressing prostaglandin endoperoxide synthase 2. *cell*; 83:493-501.
- Tsujimoto Y, Finger LR, Yunis J, Nowell PC, Croce CM (1984):** Cloning of the chromosome breakpoint of neoplastic B cells with the t (14;18) chromosome translocation. *Science*;226(4678):1097-1099.
- Ueki, K., Okada, T, Hu, J., (2006):** Total insulin and 1GF-I resistance in pancreatic beta cells causes overt diabetes. *Nature Genetics*, 38, 583-588.
- Van Cruchten, S and Van Den Broeck, W, (2002):** "Morphological and biochemical aspects of apoptosis, oncosis and necrosis." *Anat Histol Embryol* 31 (4): 214-223.
- Van der Hoven F; Schimmang T; volkmann A; mattei M.G; Kyewski B. and Rüther U. (1994):** Programmed cell death is affected in the novel mouse mutant fused toes (Ft). *Development (Cambridge , uk)* 120,2601-2607.
- van Loo, G, Saelens, X, van Gurp, M, MaeFarlane, M, Martin, SJ and Vandenabeele. P, (2002):** "The role of mitochondria! factors in apoptosis: a Russian roulette with more than one bullet." *Cell Death Differ* 9(10): 1031-1042.
- Varfolomeev, EE, Sehuehmann, M, Luria, V, Chiannikulchai, N, Bcckmann, JS, Mett, IL, Rebrikov, D, Brodianski, VM, Kemper, OC, Kollet, O, Lapidot, T, Softer, D, Sobe, T, Avraham, KB, Goncharov, T, Holtmann, H, Lonai, P and Wallach, D (1998):** "Targeted disruption of the mouse Caspase 8 gene ablates cell death induction by the TNF receptors, Fas/Apol, and DR3 and is lethal prcnatally." *Immunity* 9(2): 267-276.
- Vaschetto R, Kuiper JW, Chiang SR, (2008):** Inhibition of poly(adenosine diphosphate-ribose) polymerase attenuates ventilator-induced lung injury. *Anesthesiology* : 108:261-268.
- Verhagen, AM, Silke, J, Ekert, PG, Pakusch, M, Kaufmann, H, Connolly, LM, Day, CL, Tikoo, A, Burke, R, Wrobel, C, Moritz, RL, Simpson, RJ and Vaux, DL (2002):** "HtrA2 promotes cell death through its serine protease



- activity and its ability to antagonize inhibitor of apoptosis proteins." *J Biol Chem* 277(1): 445-454.
- Verheul H.M, Pinedo H.M. (2007):** Vascular endothelial growth factor and its inhibitors, *Drugs Today (Barc)* 39 (Suppl. C) , 81–93.
- Viitala K, Makkonen K, Israel Y, (2000):** Autoimmune responses against oxidant stress and acetaldehyde-derived epitopes in human alcohol consumers. *Alcohol Clin Exp Res*;24(7):1103-1109.
- Vis J.C., Schipper R.T., Van B, Huizen, M.M. Verbeek, R.M.W. De Waal, P. Wesseling, H.J. Ten Donkelaar, B. Kremer, (2005):** Expression pattern of apoptosis-related markers in Huntington's disease, *Acta Neuropathol.* 109 ,321–328.
- Vogelstein B, Lane D, Levins AJ (2000):** Surfing the p53 network. *Nature*:408:307-10.
- Volpe, R. (1994):** Evidence that the immunosuppressive effects of anti thyroid drugs are mediated through actions on the thyroid cell, modulating thyrocyte-immunocyte signaling: a review. *Thyroid*4, 217-223.
- von der Thusen J.H., van Vlijmen B.J., Hoebe R.C, MM. Kockx, L.M. Havekes, T.J. van Berkel, E.A. Biessen, (2002):** Induction of atherosclerotic plaque rupture in apolipoprotein E-mice after adenovirus-mediated transfer of p53. *Circulation* 105 , 2064-2070.
- W.L. Chan, N. Pejnovic, H. Hamilton, T.V. Liew, D. Popadic, A. Poggi, S.M. Khan, (2005):** Atherosclerotic abdominal aortic aneurysm and the interaction between autologous human plaque-derived vascular smooth muscle cells, type 1 NKT, and helper T cells, *Circ. Res.* 96 , 675–683.
- Walczak, H. et al, (1999):** Tumoricidal activity of tumor necrosis factor-related apoptosis-inducing ligand in vivo. *Nat. Med.* 5,157-163.
- Waltzer W.C., Bach Varoff R.J. and Rasivach A.D, (1984):** Immunological maintenance in patients with end stage renal disease. *J. clin . Immunol*;5:369-368.



- Wang J, Walsh K, (1996):** Resistance to apoptosis conferred by cdk inhibitor during myocyte differentiation. *Science* 273: 359-361.
- Wang J, Zheng L, Lobilo A, Chan FK, Dale J, Sneller M, (1999):** Inherited human caspase 10 mutations underlie defective lymphocyte and dendritic cell apoptosis in autoimmune lymphoproliferative syndrome type II. *Cell*;98(1):47-58.
- Wang J. Wilcken DE, Wang XL. (2001):** Cigarette smoke activates caspase-3 to induce apoptosis of human umbilical venous endothelial cells. *Mol Genet Metab*;72:82-88.
- Wang JY (2000):** Regulation of cell death by the Abl tyrosine kinase. *Oncogene*;19(49):5643-5650.
- Wang R, Ibarra-Sunga O, Verlinski L, (2000):** Abrogation of bleomycin-induced epithelial apoptosis and lung fibrosis by captopril or by a caspase inhibitor. *Am J Physiol Lung Cell Mol Physiol*;279:L143-151.
- Wang X, Ryter S W, Dai C, (2003) :** Necrotic cell death in response to oxidant stress involves the activation of the apoptogenic caspase-8/bid pathway. *J Biol Chem* 2003;278: 29184-191.
- Wang X, Wang Y, Kim HP, (2007):** Carbon monoxide protects against hyperoxia-induced endothelial cell apoptosis by inhibiting reactive oxygen species formation. *J Biol Chem*;282:1718-1726.
- Wang X, Wang Y, Kim HP, (2007):** FLIP inhibits endothelial cell apoptosis during hyperoxia by suppressing Bax. *Free Radic Biol Med*;42:1599-609.
- Wang X, Wang Y, Zhang J, (2005):** FLIP protects against hypoxia/reoxygenation-induced endothelial cell apoptosis by inhibiting Bax activation. *Mol Cell Biol*;25:4742-4751.
- Wang X, Zhang J, Kim HP, (2004):** Bel-XL disrupts death-inducing signal complex formation in plasma membrane induced by hypoxia/reoxygenation, *FASEB J*; 18: 1826-1833.
- Wang X, Zhou Y, Kim HP, (2004):** Hepatocyte growth factor protects against hypoxia/ reoxygenation-induced apoptosis in endothelial cells. *J Biol Chem*;279:5237-5243.



- Wang Y, Prives C (1995):** Increased and altered DNA binding of human p53 by S and G2/M but not G1 cyclin-dependent kinases. *Nature* 376: 88-91.
- Wang, S, Miura, M, Jung, YK, Zhu, H, Li, E and Yuan, J (1998):** "Munne easpase-11, an ICK-mteracting protease, is essential for the activation of ICE." *Cell* 92(4): 501-509.
- Wang, X (2001):** "The expanding role of mitochondria in apoptosis." *Genes Dev* 15(22): 2922-33. Wang, XW (1999). "Role of p53 and apoptosis in carcinogenesis." *Anticanoer Res* 19(6A): 4759-4771.
- Watson AIM, Askew JN, Benson RSP (1995):** Poly(Adenosine diphosphate ribose) polymerase inhibition prevents necrosis induced by H<sub>2</sub>O<sub>2</sub> but not apoptosis. *Gastroenterology*;109: 472-482.
- Watson AJM, Merritt A J, Jones LS, (1996):** Evidence for reciprocity of bcl-2 and p53 expression in human colorectal adenomas and carcinomas. *Br J Cancer*;76:889-994.
- Weetman, A.P. and McGregor, A.M, (1994) :** Autoimmune thyroid disease: further developments in our understanding. *Endocr. Rev.* 15,788-830.
- Wei, C.L., Wu, Q., Vega, V.B., Chiu, K.P., Ng, P., Zhang, T., Shahab, A., Yong, H.C., Fu, Y., Weng, Z., Liu, J., Zhao, X.D., Chew, J.L., Lee, Y.L., Kuznetsov, V.A., Sung, W.K., Miller, L.D., Lim, B., Liu, E.T., Yu, Q., Ng, H.H., Ruan, Y., (2006):** A global map of p53 transcription-factor binding sites in the human genome. *Cell* 124, 207-219.
- Wei, MC, Zong, WX, Cheng, EH, Lindsten, T, Panoutsakopoulou, V, Ross, AJ, Roth, KA, MacGregor, GR, Thompson, CB and Korsmeyer, SJ (2001):** "Proapoptotic BAX and BAK: a requisite gateway to mitochondria! dysfunction and death." *Science* 292(5517): 727-730.
- Weil, M, Jacobson, MD, Coles, US, Davies, TJ, Gardner, RL, Raff, KD and Raff, MC (1996):** "Constitutive expression of the machinery for programmed cell death." *J Cell Biol* 133(5): 1053-1059.
- Welsh A.O. and Enders A.C. (1987):** Trophoblast . decidua cell interactions and the establishment of maternal blood circulation in parietal yolk sac



- placenta of the rat. *A nat . Rec .* 217,203-219.
- Werlen G, Hausmann B, Palmer E, (2000):** A motif in the alphabets T-cell receptor controls positive selection by modulating ERK activity. *Nature*, 406:422-426.
- Wickenden JA, Clarke MC, Rossi AG, (2003):** Cigarette smoke prevents apoptosis through inhibition of caspase activation and induces necrosis. *Am J Respir Cell Mol Biol*;29:562-570.
- Wickremasinghe RG, Hoffbrand AV (1999):** Biochemical and genetic control of apoptosis: relevance to normal hematopoiesis and hematological malignancies. *Blood*,93(11):3587-3600.
- Williams GT, Smith CA, Spooner E, Dexter TM, Taylor DR (1990):** Haematopoietic colony stimulating factors promote cell survival by suppressing apoptosis. *Nature*;343:376.
- Winoto A, Littman DR (2002):** Nuclear hormone receptors in T lymphocytes. *Cell*, 109(Suppl):S57-S66.
- Wisniewski, T., Ghiso, J., and Frangione, B, (1994):** Alzheimer's disease and soluble A beta. *Neurobiol. Aging* 15:143-152.
- Wolozin, B., Iwasaki, K., Vito, P, Ganjei, J.K., Lacan, E., Sunderland, T, (1996):** Participation of presenilin 2 in apoptosis: enhanced basal activity conferred by an Alzheimer mutation. *Science* 274:1710-1713.
- Wong, P.C., Pardo, C.A., Borchelt, D.R., Lee, M.K., Copeland, N.G., Jenkins N.A, (1995):** An adverse property of a familial ALS-linked SOD1 mutation causes motor neuron disease characterized by vacuolar degeneration of mitochondria *Neuron* 14:1105-1116.
- Wong SC, Fukuchi M, Melnyk P, Rodger I, Giaid A, (1998):** Induction of cyclooxygenase-2 and activation of nuclear factor-kappa B in myocardium of patients with congestive heart failure. *Circulation*;98:100–103.
- Woo D, (1995):** Apoptosis and loss of renal tissue in polycystic kidney disease . *N. Eng. J. med*; 333:18-25.



- Wood KJ, Sakaguchi S, (2003):** Regulatory T cells in transplantation tolerance. *Nat Rev Immunol*;3:199–210.
- Workman P, (2004):** Inhibiting the phosphoinositide 3-kinase pathway for cancer treatment, *Biochem. Soc. Trans.* 32 (2004) 393-396.
- Wu CH, Lin HH, Yan FP, (2005):** Immunohistochemical detection of apoptotic proteins, p53/Bax and JNK/FasL cascade, in the lung of rats exposed to cigarette smoke. *Arch Toxicol* ;9:1-9.
- Wu M, Arsura M, Bellas RE, Fitzgerald MJ, Lee H, Schauer SL, Sherr DH, Sonenshein GE, (1996):** Inhibition of c-myc expression induces apoptosis of WEHI murine B cells. *Mol Cell Biol* 16: 5015-5025.
- Y.H. Huang, M. Lu, H. Wu (2004):** Antagonizing XIAP-mediated caspase-3 inhibition: Achilles' heel of cancers? *Cancer Cell* 5 , 1-2.
- Yang, Y, Fang, S, Jensen, JP, Weissman, AM and Ashwell, JD, (2000):** "Ubiquitin protein ligase activity of IAPs and their degradation in proteasomes in response to apoptotic stimuli." *Science* 288(5467): 874-877.
- Yao M, Keogh A, Spratt P, dos Remedios CG, Kiebling PC. Elevated Dnase I levels in human idiopathic dilated cardiomyopathy (1996):** An indicator of apoptosis? *J Mol Cell Cardiol*;28:95–101.
- Yokohori N, Aoshiba K, Nagai A, (2004):** Increased levels of cell death and proliferation in alveolar wall cells in patients with pulmonary emphysema. *Chest* ;125:626-632.
- Yoshino T, Kishi H, Nagata T, Tsukada K, Saito S, Muraguchi A (2001):** Differential involvement of p38 MAP kinase pathway and Bax translocation in the mitochondria-mediated cell death in TCR- and dexamethasone-stimulated thymocytes. *Eur J Immunol*, 31:2702-2708.
- Youle, R.J.,and Strasser, A, (2008):** The BCL-2 protein family: opposing activities that mediate cell death. *Nat. Rev. Mol. Cell Biol.* 9, 47–59.
- Zhang X, Shan P, Otterbein LE, (2003):** Carbon monoxide inhibition of apoptosis during ischemia-reperfusion lung injury is dependent on the p38 mitogen-activated protein kinase pathway and involves caspase 3. *J Biol Chem*;278:1248-1258.



- Zhao L., Ackerman S.L., (2006):** Endoplasmic reticulum stress in health and disease, *Curr. Opin. Cell Biol.* 18 444–452.
- Zhao, B., Chrest, F.J., Horton, Jr. W.E., Sisodia, S.S., and Kusiak, J.W. (1997)** : Expression of mutant amyloid precursor proteins induces apoptosis in PC12 cells. *J. Neurosci. Res.* 47:253-263.
- Zhihe Li, Bing OHL, Long X, Robinson KG, Lakatta EG (1997):** Increased cardiomyocyte apoptosis during the transition heart failure in the spontaneously hypertensive rat. *Am J Physiol*;272:H2313–H2319.
- Ziv, 1., Offen, D., Barzilai, A., Haviv, R., Stein, R., Zilkha, R., (1997):** Modulation of control mechanisms of dopamine-induced apoptosis-a future approach to the treatment of Parkinson's disease. *J. Neural Transm.* 849:195-202.
- Zuzarte-Luis, V., Hurle, J.M., (2005):** Programmed cell death in the embryonic vertebrate limb. *Semin. Cell Dev. Biol.* 16, 261–269.
- Zychlinsky A, Preuost MC, Sansonetti PJ 1992 :** *Shigella flexneri* induces apoptosis in infected macrophages. *Nature*;358:167- 169.