

References

A

Abdel Aziz MT, Zayed M, Al-Asmar MF et al . (1998)

Detection of HPV infection in Egyptian women by PCR in situ hybridization.

Proceedings of the first Pan African Conference on Biochemistry and Molecular Biology. Nairobi, Kenya 163–170 .

Abdel Aziz MT , Abdel Aziz MZ , Atta HM et al. (2006)

Screening for human papillomavirus (HPV) in Egyptian women by the second-generation hybrid capture (HC II) test.

Med Sci Monit, 12(7): 43-49.

Abdel-Hady ES, Martin-Hirsch P, Duggan-Keen M et al. (2001)

Immunological and viral factors associated with the response of vulval intraepithelial neoplasia to photodynamic therapy.

Cancer Res 61:192–196.

Adam MB, Auerbach JD, Buehler JW et al. (2001)

Workshop summary: scientific evidence on condom effectiveness for sexually transmitted diseases prevention. Herndon, VA: National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services, 1-12 .

Adams F (1886)

The genuine works of Hippocrates: Translated from the Greek with a preliminary discourse and annotations, vol 2. William Wood, New York
Quoted from : Rosenblatt A & De Campos Guidi HG (2009) Human Papillomavirus History and Epidemiology , in: Rosenblatt A & De Campos Guidi HG (eds.) Human Papillomavirus , A Practical Guide for Urologists , Springer-Verlag Berlin Heidelberg , ch: 1 ,pp: 3-15 .

Ahmed AM, Madkan V & Tyring SK (2006)

Human Papillomaviruses and Genital Disease .

Dermatol Clin 24 : 157 – 165 .

Ahn WS, Huh SW, Bae SM et al., (2003)

A major constituent of green tea, EGCG, inhibits the growth of a human cervical cancer cell line, CaSki cells, through apoptosis, G(1) arrest, and regulation of gene expression.

DNA Cell Biol 22: 217–224 .

Alam M & Stiller M (2001)

Direct medical costs for surgical and medical treatment of condylomata acuminata.

Arch Dermatol 137: 337–341 .

Alberico S, Pinzano R, Comar M et al., (1996)

Maternal-fetal transmission of human papillomavirus.

Minerva Ginecol 48: 199–204 .

Allen AL & Siegfried EC (2000)

What's new in human papillomavirus infection.

Curr Opin Pediatr 12:365–369 .

American Academy of Pediatrics (1999)

Task Force on Circumcision. Circumcision policy statement.

Pediatrics 103:686–693.

Arany I, Evans T & Tyring SK (1998)

Tissue specific HPV expression and downregulation of local immune responses in condylomas from HIV seropositive individuals.

Sex Transm Infect 74:349–353.

Agrawal SK, Bhattacharya SN & Singh N (2004)

Pearly penile papules: a review.

Int J Dermatol 43:199-201.

Armstrong LR, Preston EJ, Reichert M et al. (2000)

Incidence and prevalence of recurrent respiratory papillomatosis among children in Atlanta and Seattle.

Clin Infect Dis 1:107–109.

Ault KA (2007)

Effect of prophylactic human papillomavirus L1 virus-like-particle vaccine on risk of cervical intraepithelial neoplasia grade 2, grade 3 , and

adenocarcinoma in situ: A combined analysis of four randomised clinical trials.

Lancet 369: 1861–1868 .

Aynaud O, Buffet M, Roman P et al. (2008)

Study of persistence and recurrence rates in 106 patients with condyloma and intraepithelial neoplasia after CO2 laser treatment.

Eur J Dermatol 18: 153–158 .

B

Bafverstedt B (1967)

Condylomata acuminata-past and present.

Acta Derm Venereol 47: 376–381 .

Baken LA, Koutsky LA, Kuypers J et al. (1995)

Genital human papillomavirus infection among male and female sex partners: prevalence and typespecific concordance.

J Infect Dis 171:429–432.

Baldwin SB, Wallace DR, Papenfuss MR et al. (2003)

Human papillomavirus infection in men attending a sexually transmitted disease clinic.

J Infect Dis 187: 1064–1070 .

Bandieramonte G, Colecchia M, Mariani L et al. (2008)

Peniscopically controlled CO2 laser excision for conservative treatment of in situ and T1 penile carcinoma: report on 224 patients.

Eur Urol 54: 875–882 .

Bar-Am A & Niv J (2007)

The role of HPV DNA in the evaluation and follow-up of asymptomatic male sexual partners of females with CIN3.

Eur J Gynaecol Oncol 28: 207–210 .

Barr E & Tamms G (2007)

Quadrivalent Human Papillomavirus Vaccine

Clinical Infectious Diseases 45:609–617 .

Barnabas RV, Laukkanen P, Koskela P et al. (2006)

Epidemiology of HPV 16 and cervical cancer in Finland and the potential impact of vaccination: Mathematical modelling analyses.
PLoS Med 3: 138 -143 .

Baseman JG & Koutsky LA (2005)

The epidemiology of human papillomavirus infections.
Journal of Clinical Virol . 32: 16-24.

Bechara FG, Huesmann M, Stucker M, et al. (2002)

An exceptional localization of angiokeratoma of Fordyce on the glans penis.
Dermatology 205:187-188.

Bell B (1793)

Treatise on gonorrhoea virulenta and lues venerea, vol 1. Watson Mudie, Edinburg . Quoted from : Rosenblatt A & De Campos Guidi HG (2009) Human Papillomavirus History and Epidemiology , in: Rosenblatt A & De Campos Guidi HG (eds.) Human Papillomavirus , A Practical Guide for Urologists , Springer-Verlag Berlin Heidelberg , ch: 1 ,pp: 3-15 .

Bennett SR, Carbone FR, Karamalis F et al. (1997)

Induction of a CD8+ cytotoxic T lymphocyte response by cross-priming requires cognate CD4+ T cell help.
J Exp Med 186:65-70.

Beutner KR, Ferenczy A. (1997)

Therapeutic approaches to genital warts.
Am J Med 102:28–37.

Beutner KR, Richwald GA, Wiley DJ et al.(1998)

External genital warts: report of the American Medical Association Consensus Conference. AMA Expert Panel on External Genital Warts.
Clin Infect Dis 27:796–806 .

Beutner KR, Wiley DJ, Douglas JM, et al. (1999)

Genital warts and their treatment.
Clin Infect Dis 28(Suppl 1): 37–56.

Bhattacharjee P & Brodell RT (2003)

Podophyllin in Brodell RT & Johnson SM (ed): Warts , Martin Dunitz, a member of the Taylor & Francis Group ch :7 , pp: 59-69 .

Bleeker MC, Hogewoning CJ, Voorhorst FJ et al. (2005)

HPV-associated flat penile lesions in men of a non-STD hospital population: less frequent and smaller in size than in male sexual partners of women with CIN.

Int J Cancer 113: 36–41 .

Bleeker MC, Snijders PF, Voorhorst FJ et al. (2006)

Flat penile lesions: the infectious “invisible” link in the transmission of human papillomavirus.

Int J Cancer 119: 2505–2512 .

Bohle A, Doehn C, Kausch I et al. (1998)

Treatment of recurrent penile condylomata acuminata with external application and intraurethral instillation of bacillus calmetteguerin.

J Urol 160: 394–396 .

Böhle A, Büttner H, and Jocham D (2001)

Primary treatment of condylomata acuminata with viable bacillus Calmette- Guérin.

J Urol 165: 834–836.

Bonnez W (2009)

Introduction in: Bonnez W (ed .) Guide to Genital HPV Diseases and Prevention , Informa Healthcare USA, Inc. p: xi – xii .

Bonnez W & Toy EP (2009)

External genital warts in Guide to Genital HPV Diseases and Prevention ed: Bonnez W Informa Healthcare USA, Inc. ch : 3 , p:29-34 .

Bonnez W , Brown DR & Rand CM (2009)

Prevention, in Bonnez W (ed): Guide to Genital HPV Diseases and Prevention, Informa Healthcare USA, Inc. ch: 7 , pp:87-127 .

Bosch FX (2009)

Broad-Spectrum Human Papillomavirus Vaccines: New Horizons but One Step at a Time .

JNCI Vol. 101, Issue 11 : pp:771-773 .

Bosch FX, Castellsague X, Munoz N, et al. (1996)

Male sexual behavior and human papillomavirus DNA: key risk factors for cervical cancer in Spain.

J Natl Cancer Inst 88:1060–1067.

Bosch FX, Lorincz A, Munoz N et al. (2002)

The causal relation between human papillomavirus and cervical cancer.
J Clin Pathol 2002; 55:244–265 .

Bosch FX & De Sanjosé SS (2003)

Human papillomavirus and cervical cancer-burden and assessment of causality.
J Nat Cancer Inst Monogr 31:3–13.

Brodell LA, Mercurio MG & Brodell RT (2007)

The diagnosis and treatment of HPV-mediated genital lesions.
Cutis, 79(4 Suppl), 5-10 .

Brotherton JM, Gold MS, Kemp AS et al. (2008)

Anaphylaxis following quadrivalent human papillomavirus vaccination.
CMAJ 179: 525–533 .

Brown DR, Schroeder JM, Bryan JT et al. (1999)

Detection of multiple human papillomavirus types in Condylomata acuminata lesions from otherwise healthy and immunosuppressed patients.
J Clin Microbiol 37: 3316–3322 .

Brown D (2007)

HPV type 6/11/16/18 vaccine: First analysis of cross-protection against persistent infection, cervical intraepithelial neoplasia (CIN), and adenocarcinoma in situ (AIS) caused by oncogenic HPV types in addition to 16/18 In: 47th Interscience Conference on Antimicrobial Agents and Chemotherapy, Chicago

Brown RE, Breugelmans JG, Theodoratou D et al. (2006)

Costs of detection and treatment of cervical cancer, cervical dysplasia and genital warts in the UK.
Curr Med Res Opin 22:663-670.

Buechner SA (2002)

Common skin disorders of the penis.
BJU Int 90: 498–506 .

Bunker CB & Neill SM (2004)

The genital, perianal and umbilical regions. In: Burns DA, Breathnach SM, Cox NH, Griffiths CEM (eds .): Rook's textbook of dermatology, 7th ed, vol. 4. Oxford: Blackwell Publishing. pp. 53-68.

Burchell AN, Richardson H, Mahmud SM et al. (2006)

Modeling the sexual transmissibility of human papillomavirus infection using stochastic computer simulation and empirical data from a cohort study of young women in Montreal, Canada.
Am J Epidemiol 163: 534–543 .

Burns JE & Maitland NJ (2005)

Human papilloma virus and cancer.
Microbiology today aug.2005 117-120 .

C

Carr J & Gyorfı T (2000)

Human papillomavirus. Epidemiology, transmission and pathogenesis.
Clin Lab Med 20: 235-255 .

Carrasco D, Van Der Straten M & Tying SK (2002)

Treatment of anogenital warts with imiquimod 5% cream followed by surgical excision of residual lesions.
J Am Acad Dermatol 47: S212–S216 .

Carter JJ & Galloway DA (1997)

Humoral immune response to human papillomavirus infection.
Clin Dermatol 15:249-259.

Carter JJ, Koutsky LA, Hughes JP et al. (2000)

Comparison of human papillomavirus types 16, 18, and 6 capsid antibody responses following incident infection.
J Infect Dis 181: 1911–1919 .

Carter JJ, Madeleine MM, Shera K, et al. (2001)

Human papillomavirus 16 and 18 L1 serology compared across anogenital cancer sites.
Cancer Res 61(5): 1934–1940 .

Calista D (2000)

Topical cidofovir for severe cutaneous human papillomavirus and molluscum contagiosum infections in patients with HIV/ AIDS. A pilot study.

J Eur Acad Dermatol Venereol 14:484–488.

Callen JP, Greer DE, Hood AF, et al. (1993)

Color Atlas of Dermatology. Philadelphia, WB Saunders, pp 327,328.quoted from: Walsh PC, (ed.) Campbell's Urology, Philadelphia: WB Saunders; 2002:731-742.

Casas A & Batlle A (2002)

Rational design of 5-aminolevulinic acid derivatives aimed at improving photodynamic therapy.

Curr Med Chem Anticancer Agents 2: 465–475 .

Cason J, Kaye JN, Jewers RJ et al. (1995)

Perinatal infection and persistence of human papillomavirus types 16 and 18 in infants.

J Med Virol 47: 209–218 .

Castellsague X, Bosch FX, Munoz N, et al. (2002)

Male circumcision, penile human papillomavirus infection, and cervical cancer in female partners.

N Engl J Med 346:1105–1112.

CDR (2006)

Trends in anogenital warts and anogenital herpes simplex virus infection in the United Kingdom: 1996 to 2005.

In: Communicable Disease Report Weekly (CDR Weekly), vol 16. Health Protection Agency .

CDC (2006)

Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines 2006.

MMWR , vol .55:62–67.

CDC (2007)

Centers for Disease Control and Prevention , Quadrivalent Human Papillomavirus Vaccine , Recommendations of the Advisory Committee on Immunization Practices (ACIP) .

MMWR vol. 57:1-32 .

CDC (2010)

Centers for Disease Control and Prevention. Sexually transmitted diseases treatment guidelines 2010.

MMWR (Mortality and Morbidity Weekly Report) vol.59 :69-73 .

Chen K, Chang BZ, Ju M et al. (2007)

Comparative study of photodynamic therapy vs. CO2 laser vaporization in treatment of condylomata acuminata: a randomized clinical trial.

Br J Dermatol 156: 516–520 .

Chesson HW, Blandford JM, Gift TL et al. (2004)

The estimated direct medical cost of sexually transmitted diseases among American youth, 2000.

Perspect Sex Reprod Health 36: 11–19 .

Chiao EY, Giordano TP, Palefsky JM et al. (2006)

Screening HIV-infected individuals for anal cancer precursor lesions: a systematic review.

Clin Infect Dis 43:223–233.

Cogliano V, Baan R, Straif K et al. (2005)

Carcinogenicity of human papillomaviruses.

Lancet Oncol 6: 204 -209 .

Coleman N, Birley HD, Renton AM et al. (1994)

Immunological events in regressing genital warts.

Am J Clin Pathol 102:768-773.

Conley LJ, Ellerbrock TV, Bush TJ et al. (2002)

HIV-1 infection and risk of vulvovaginal and perianal condylomata acuminata and intraepithelial neoplasia: a prospective cohort study.

Lancet 359:108–113.

Cook L, Koutsky L & Holmes K (1994)

Circumcision and sexually transmitted diseases.

Am J Public Health 84:197–201.

Cox T, Buck HW, Kinney W et al. (2001)

HPV types: natural history and epidemiology. Human papillomavirus (HPV) and cervical cancer . Clinical proceedings . Washington , DC:

Association of Reproductive Health Professionals; 2001. Available from: <http://www.arhp.org/healthcareprovider/onlinepublications/clinicalproceedings/cphpv/types.cfm?ID=149>. Accessed on :2009-09-12 .

Crawford LV & Crawford EM (1963)

A comparative study of polyoma and papilloma viruses.
Virology 21: 258–263 .

Crowe MA (2010)

Molluscum contagiosum from: www.emedicine.com . Topic no.910570 ,
last update : Apr. 2010. accessed: dec , 2010 .

D

Daling JR, Madeleine MM, Johnson LG, et al. (2005)

Penile cancer: importance of circumcision, human papillomavirus and smoking in in situ and invasive disease.
Int J Cancer 116:606–616 .

Day PM, Roden RB, Lowy DR et al. (1998)

The papillomavirus minor capsid protein, L2, induces localization of the major capsid protein, L1, and the viral transcription/replication protein, E2, to PML oncogenic domains.
J Virol 72: 142–150 .

De Carvalho JJ, Syrjanen KJ, Jacobino M et al. (2006)

Prevalence of genital human papillomavirus infections established using different diagnostic techniques among males attending a urological clinic.
Scand J Urol Nephrol 40: 138–143 .

De Villiers EM, Fauquet C, Broker TR et al. (2004)

Classification of papillomaviruses.
Virology 324: 17–27 .

Dias D, Van Doren J, Schlottmann S, et al. (2005)

Optimization and validation of a multiplexed luminex assay to quantify antibodies to neutralizing epitopes on human papillomaviruses 6, 11, 16, and 18.
Clin Diagn Lab Immunol 12(8): 959–969 .

Dillner J , Andersson-Ellström A , Hagmar B et al. (1999)

High risk genital papillomavirus infections are not spread vertically. Rev. med. Virol., 9, 23–29 .

Dinh TH, Sternberg M, Dunne EF et al. (2008)

Genital warts among 18- to 59-year-olds in the United States, National Health and Nutrition Examination Survey, 1999–2004. Sex Transm Dis 35: 357–360 .

Doorbar J (2005)

The papillomavirus life cycle.
J Clin Virol 32 (suppl 1): S7–15.

Doorbar J & Sterling JC (2001)

The biology of human papillomaviruses. In: Sterling JC, Tyring SK (eds) Human papillomaviruses: Clinical and scientific advances. Arnold, London, ch: 1 , pp 10–23 .

Douglas JM Jr , Eron LJ, Judson FN et al. (1990)

A randomized trial of combination therapy with intralesional interferon alpha 2b and podophyllin versus podophyllin alone for the therapy of anogenital warts.
J Infect Dis 162: 52–59 .

Dunne EF & Lauri E (2006)

Genital Human Papillomavirus Infection
Clinical Infectious Diseases 43:624–629 .

Dunne EF, Unger ER, Sternberg M et al. (2007)

Prevalence of HPV infection among females in the United States.
JAMA 297: 813–819 .

Dupin N (2004)

Genital warts .
Clinical Dermatology . 22:81-86.

E

Edwards L, Ferenczy A, Eron L et al. (1998)

Selfadministered topical 5% imiquimod cream for external anogenital warts.

Arch Dermatol 134:25–30.

Einstein MH , Schiller JT , Viscidi RP et al. (2009)

Clinician's guide to human papillomavirus immunology: knowns and unknowns .

Lancet Infect Dis 9: 347–356 .

El Mejjad A (2003)

Le condylome acuminé géant-tumeur de Buschke Loewenstein (à propos de 3 cas).

Prog Urol 13(3): 513–517.

Evans DM, Hudson EA, Brown CL (1986)

Terminology in gynaecological cytopathology: report of the working party of the British Society for Clinical Cytology.

J Clin Pathol 39:933-942.

F

FDA (2010 a)

Food and Drug Administration. Product approval information—licensing action [package insert]. Gardasil (quadrivalent human papillomavirus types 6,11,16,18). Merck & Co., Whitehouse Station, NJ.available at: www.fda.gov/cber/products/gardasil.htm , accessed in May 29, 2010 ,last updated april 2010

FDA (2010 b)

Food and Drug Administration. Product approval-prescribing information [package insert]. Cervarix [human papillomavirus bivalent (types 16 and 18) vaccine, recombinant], GlaxoSmithKline Biologicals: Food and Drug Administration Available at :

<http://www.fda.gov/biologicsbloodvaccines/vaccines/approvedproducts/ucm186957.htm>. Accessed in : May 25, 2010.

Fehrman, F & Laimins, LA (2003)

Human papillomaviruses: Targeting differentiating epithelial cells for malignant transformation.

Oncogene, 22, 5201–5207 .

Ferenczy A, Bergeron C & Richart RM (1990)

Carbon dioxide laser energy disperses human papillomavirus deoxyribonucleic acid onto treatment fields.

Am J Obstet Gynecol 163: 1271–1274 .

Field S & Buckley C (2009)

Perianal Verrucous Epidermal Nevus Mimics a Cutaneous Sign of Child Abuse.

J Pediatr 2009;154:306 .

Fleischer AB Jr, Parrish CA, Glenn R et al. (2001)

Condylomata acuminata (genital warts): patient demographics and treating physicians.

Sex Transm Dis 28:643–647.

Forbes CD & Jackson WF (2003)

Color Atlas and Text of Clinical Medicine, 3rd ed. London, Mosby, p:337

Forhan SE, Gottlieb SL, Sternberg MR et al. (2008)

Prevalence of sexually transmitted infections and bacterial vaginosis among Female adolescents in the United States: Data from the National Health and Nutritional Examination Survey (NHANES) 2003–2004. In: National STD Prevention Conference, Chicago, IL

Franceschi S, Rajkumar R, Snijders, et al. (2005)

Papillomavirus infection in rural women in southern India.

Br. J. Cancer, 92, 601–607

Frattoni MG, Lim HB & Laimins LA (1996)

In vitro synthesis of oncogenic human papillomaviruses requires episomal genomes for differentiation-dependent late expression.

Proc Natl Acad Sci USA 93:3062–3067.

Frazer IH, Cox JT and Mayeaux EJ (2006)

Advances in prevention of cervical cancer and other HPV-related diseases.

Pediatric Infectious Diseases Journal 25 :65-81.

Frega A, Cenci M, Stentella P et al. (2003)

Human papillomavirus in virgins and behaviour at risk.

Cancer Lett 194: 21–24 .

Frimberger D, Schneede P, Hungerhuber E et al. (2006)

Autofluorescence and 5-aminolevulinic acid induced fluorescence diagnosis of penile carcinoma — new techniques to monitor Nd:YAG laser therapy.

Urol Res 30:295-300 .

G

Gall SA (2001)

Female genital warts: global trends and treatments.

Infect Dis Obstet Gynecol, 9, 149- 54.

Garcia-Pineros AJ, Hildesheim A, Herrero R, et al. (2006)

Persistent human papillomavirus infection is associated with a generalized decrease in immune responsiveness in older women.

Cancer Res 66: 11070–11076 .

Gearhart PA (2009)

Human papilloma virus from: www.emedicine.com . Topic no. 1037 , last update : Jan 2009. accessed in: May , 15,2009

Gerberding JL (2004)

Report to Congress: prevention of genital human papillomavirus infection. Centers for Disease Control and Prevention, Department of Health and Human Services .

Gilbert DM & Cohen SN (1987)

Bovine papilloma virus plasmids replicate randomly in mouse fibroblasts throughout S phase of the cell cycle.

Cell 50:59–68 .

Gillard P, Vanhooteghem O, Richert B et al. (2005)

Tumor de Buschke-Loewenstein.

Ann Dermatol Venereol 132: 98–99.

Gissmann L & Zur Hausen H (1980)

Partial characterization of viral DNA from human genital warts (Condylomata acuminata).

Int J Cancer 25: 605–609 .

Giuliano AR, Harris R, Sedjo RL et al. (2002)

Incidence, prevalence, and clearance of type-specific human papillomavirus infections: the Young Women's Health study.
J Infect Dis 2002; 186:462–469.

Giuliano AR, Lu B, Nielson CM et al. (2008)

Age-specific prevalence, incidence, and duration of human papillomavirus infections in a cohort of 290 US men.
J Infect Dis 198: 827–835 .

Goh BT (2005)

Syphilis in adults.
Sex Transm Inf 81:448-452 .

Goldman L (2007)

Human Papillomavirus and Genital Warts : From Nelson AL and Woodward JA (eds) Current Clinical Practice: Sexually Transmitted Diseases: A Practical Guide for Primary Care .Humana Press Inc. p:21-46

Gross G & Barrasso R (1997)

Human papillomavirus infection. A Clinical Atlas. Ullstein- Mosby, Berlin, Wiesbaden, 415-432.

Gross G, Meyer KG, Pres H et al. (2007)

A randomized, double-blind, four-arm parallelgroup, placebo-controlled Phase II/III study to investigate the clinical efficacy of two galenic formulations of Polyphenon E in the treatment of external genital warts.
J Eur Acad Dermatol Venereol 21: 1404–1412 .

Guidelines for the Management of Genital Warts in Australia and New Zealand , (2002)

Published by: Professional Advisory Board (PAB) of the Australia and New Zealand HPV Project.3rd Edition , p: 6-11 .

Gunter J (2003)

Genital and perianal warts: new treatment opportunities for human papillomavirus infection.
Am J Obstet Gynecol 189:S3–S11.

H

Habif TP (2004)

Genital warts. In: Habif TP(ed) Clinical Dermatology: a Color Guide to Diagnosis And Therapy 4th ed. St. Louis, Mo: Mosby .210:224.

Hagensee ME , Yaegashi, N & Galloway DA (1993)

Self-assembly of human papillomavirus type 1 capsids by expression of the L1 protein alone or by coexpression of the L1 and L2 capsid proteins. J. Virol., 67, 315–322 .

Hajak ZF (1956)

Contribution to the etiology of laryngeal papilloma in children. J. Laryngol. Otol., 70, 166–168 .

Hallez S, Simon P, Maudoux F et al. (2004)

Phase I/II trial of immunogenicity of a human papillomavirus (HPV) type 16 E7 protein-based vaccine in women with oncogenic HPV-positive cervical intraepithelial neoplasia. Cancer Immunol Immunother 53: 642–650 .

Hallmo P & Naess O (1991)

Laryngeal papillomatosis with human papillomavirus DNA contracted by a laser surgeon. Eur Arch Otorhinolaryngol 248: 425–427 .

Hand E (1949)

Circumcision and venereal disease. Arch Dermatol Syphilol 60:341–346.

Handley J, Hanks E, Armstrong K et al. (1997)

Common association of HPV 2 with anogenital warts in prepubertal children. Pediatr Dermatol 14: 339–343 .

Hariri S, Dunne EF, Sternberg M et al. (2008)

Seroepidemiology of human papillomavirus type 11 in the United States: Results from the third National Health and Nutrition Examination Survey, 1991–1994. Sex Transm Dis 35: 298–303 .

Harper D, Gall S, Naud P et al. (2008)

Sustained immunogenicity and high efficacy against HPV-16/18 related cervical neoplasia: long-term follow up through 4 years in women vaccinated with CervarixTM (GSK's HPV 16/18 AS04 candidate vaccine). 36th annual meeting of the Society of Gynecologic Oncologists; Tampa, FL, USA; March 9–12, 2008 .

Harro CD, Pang YY, Roden RB et al. (2001)

Safety and immunogenicity trial in adult volunteers of a human papillomavirus 16 L1 virus-like particle vaccine.
J Natl Cancer Inst 93: 284–292 .

Health Protection Agency (2007)

Trends in genital warts and genital herpes diagnoses in the United Kingdom. In: Health Protection Weekly Report, vol 1. Health Protection Agency, UK .

Hebner CM, Wilson R, Rader J et al. (2006)

Human papillomaviruses target the double-stranded RNA protein kinase pathway.
J Gen Virol 87: 3183–3193.

Hengge UR & Tietze G (2000)

Successful treatment of recalcitrant condyloma with topical cidofovir.
Sex Transm Infect 76: 143-150 .

Hernandez BY, Wilkens LR, Zhu X et al. (2008)

Circumcision and human papillomavirus infection in men: a site-specific comparison.
J Infect Dis 197:787–794 .

Herrero R , Castle PE , Schiffman M et al. (2005)

Epidemiologic profile of type-specific human papillomavirus infection and cervical neoplasia in Guanacaste, Costa Rica.
J. infect. Dis., 191, 1796–1807 .

Hicheri J , Jaber K, M. R. Dhaoui MR et al.(2006)

Giant condyloma
Acta Dermatoven APA Vol 15, 2006, No 4 ,pp:181-183 .

Hildesheim A, Herrero R, Wacholder S et al. (2007)

Effect of human papillomavirus 16/18 L1 viruslike particle vaccine among young women with preexisting infection: A randomized trial. JAMA 298: 743–753 .

Hillemanns P, Breugelmans JG, Giesecking F et al. (2008)

Estimation of the incidence of genital warts and the cost of illness in Germany: A cross-sectional study. BMC Infect Dis 8: 76-80 .

Ho GY, Bierman R, Beardsley L et al. (1998)

Natural history of cervicovaginal papillomavirus infection in young women. N Engl J Med 338:423–428 .

Hoekx L & Wyndaele JJ(1998)

Angiokeratoma: A cause of scrotal bleeding. Acta Urol Belg 66:27-28.

Hollier LM & Workowski K (2008)

Treatment of Sexually Transmitted Infections in Women , Infect Dis Clin N Am 22 : 665–691 .

Hoyme UB, Hagedorn M, Schindler AE et al. (2002)

Effect of adjuvant imiquimod 5% cream on sustained clearance of anogenital warts following laser treatment. Infect Dis Obstet Gynecol 10: 79–88 .

Hubert WG (2003)

Papillomaviruses: biology and analysis in: Brodell RT & Johnson SM (ed):Warts Diagnosis and Management: An Evidence-based Approach . Martin Dunitz, a member of the Taylor & Francis Group .ch :1, pp: 6-22

Huh WK & Roden RB (2008)

The future of vaccines for cervical cancer. Gynecol Oncol 109: S48–S56 .

Hutchinson J (1855)

On the influence of circumcision in preventing syphilis. Med Times Gaz 2:542–3. Quoted from : Rosenblatt A & De Campos Guidi HG (2009) Human Papillomavirus History and Epidemiology , in: Rosenblatt A & De Campos Guidi HG (eds.) Human Papillomavirus , A

Practical Guide for Urologists , Springer-Verlag Berlin Heidelberg , ch: 1 ,pp: 3-15 .

I

IARC (2007)

International Agency for Research on Cancer IARC Monographs on the Evaluation of Carcinogenic Risks to Humans ,Human papillomaviruses. 90: 61-79 .

Insinga RP, Dasbach EJ & Myers ER (2003)

The health and economic burden of genital warts in a set of private health plans in the United States.
Clinical Infectious Diseases. 36(11):1397–1403 .

J

Jones V, Smith SJ & Omar HA (2007)

Nonsexual transmission of anogenital warts in children: A retrospective analysis.
Scientific WorldJournal 7: 1896–1899 .

Joura EA, Leodolter S, Hernandez-Avila M et al. (2007)

Efficacy of a quadrivalent prophylactic human papillomavirus (types 6, 11, 16, and 18) L1 viruslike- particle vaccine against high-grade vulval and vaginal lesions: A combined analysis of three randomised clinical trials.
Lancet 369: 1693–1702 .

K

Khawaja HT

Podophyllin versus scissor excision in the treatment of perianal condylomata acuminata: a prospective study.
Br J Surg 1989; 76:1067–1068.

Knowles G, O'Neil BW and Campo MS(1996)

Phenotypical characterization of lymphocytes infiltrating regressing papillomas.
J Virol 70:8451-8458.

Kodner CM & Nasraty S (2004)

Management of genital warts.
American Family Physician 70:2335–2342.

Konya J & Dillner J (2001)

Immunity to oncogenic human papillomaviruses.
Adv Cancer Res 82: 205–238

Körber A & Dissemond J (2009)

Pearly penile papules.
CMAJ , 108: 379.

Koshiol JE, Laurent SA & Pimenta JM (2004)

Rate and predictors of new genital warts claims and genital warts-related healthcare utilization among privately insured patients in the United States.
Sex Transm Dis 31: 748–752 .

Koutsky L (1997)

Epidemiology of genital human papillomavirus infection.
Am J Med 102: 3–8.

Krantz I & Ahlberg B. (1995)

Circumcision and HIV.
Lancet 345:730-734 .

Kumar B & Gupta S (2001)

The acetowhite test in genital human papillomavirus infection in men: what does it add?
J Eur Acad Dermatol Venereol 15: 27–29 .

Kurman RJ & Solomon D (1994)

The Bethesda System for reporting cervical/vaginal cytologic diagnoses. Definitions, criteria, and explanatory notes for terminology and specimen adequacy. Springer-Verlag, New York, 1994: 1-267.

L

Lacey CJ, Goodall RL, Tennvall GR et al. (2003)

Randomised controlled trial and economic evaluation of podophyllotoxin solution, podophyllotoxin cream, and podophyllin in the treatment of genital warts.

Sex Transm Infect 79: 270–275 .

Laumann E, Masi C, Zuckerman M (1997)

Circumcision in the United States.

JAMA 277:1052–1057.

Leggatt GR & Frazer IH (1999)

Imunobiology of HPV Infection in: Tindle RW (ed) Vaccines for Human Papillomavirus Infection and Anogenital Disease R.G. Landes Company Austin, Texas, U.S.A. pp: 13-32 .

Leman JA & Benton EC (2000)

Verrucas. Guidelines for management.

Am J Clin Dermatol 1:143-149.

Loh CS, MacRobert AJ, Bedwell J et al. (1993)

Oral versus intravenous administration of 5-aminolaevulinic acid for photodynamic therapy.

Br J Cancer 68: 41–51 .

Lorincz A & Anthony J (2001)

Advances in HPV detection by hybrid capture.

Papillomavirus Rep 145: 154-157

Lowry DR & Androphy EJ (2003)

Warts . In : Freedberg IM, Eisen AZ, Klaus W et al.(ed.): Fitzpatrick's dermatology in general medicine , vol s. 2. 6th ed New York: McGraw-Hill Inc . 2119 – 2131

Luna J & Saah A (2007)

Safety, efficacy, and immunogenicity of quadrivalent HPV vaccine (Gardasil) in women aged 24–45.In: 24th International Papillomavirus Congress, Beijing .

M

Madani TA (2006)

Sexually transmitted infections in Saudi Arabia.
BMC Infect Dis, 10, 3-10 .

Majewski S, Marczak M, Mlynarczyk B et al. (2005)

Imiquimod is a strong inhibitor of tumor cell-induced angiogenesis.
Int J Dermatol 44: 14–19 .

Mansur CP (2002)

Human papillomaviruses. In: Tyring SK, Yen-Moore A (eds)
Mucocutaneous manifestations of viral diseases. Marcel Dekker, New
York, ch:12, pp 247–294 .

Mao C, Hughes JP, Kiviat N et al. (2003)

Clinical findings among young women with genital human
papillomavirus infection.
Am J Obstet Gynecol 188: 677–684 .

Margolis DJ & Wein AJ (2002)

Color atlas of genital dermatology. In: Walsh PC, (ed.) Campbell's
Urology, Philadelphia: WB Saunders; 2002:731-742.

Matteelli A, Beltrame A, Graifemberghi S et al. (2001)

Efficacy and tolerability of topical 1% cidofovir cream for the treatment
of external anogenital warts in HIV-infected persons.
Sex Transm Dis 28: 343–346 .

McCance D (2004)

Papillomaviruses in: Zuckerman AJ, Banatvala JE , Pattison JR et al.
(eds) Principles and Practice of Clinical Virology, Fifth Edition. 2004
John Wiley & Sons Ltd, pp: 261-673 .

McClean H & Shann S (2005)

A cross-sectional survey of treatment choices for anogenital warts.
Int J STD AIDS 16: 212–216 .

McMillan A & Ogilvie MM (2004)

Human papilloma virus infection. In McMillan A & Scott GR (ed.): Clinical practice in sexually transmissible infections; 1st. ed. Elsevier. Ch. 5 :71-73 .

McMillan NA, Payne E, Frazer IH et al. (1999)

Expression of the alpha6 integrin confers papillomavirus binding upon receptornegative B-cells.
Virology 261:271–279.

Metaweia B, El-Nashar AR, Kamel I et al. (2005)

Application of viable bacille calmette-guerin topically as a potential therapeutic modality in condylomata acuminata: A placebo-controlled study.
Urology 65: 247–250 .

Miller RD (2000)

Local anesthetics. In: Miller RD, ed. Anesthesia. 5th ed. Philadelphia: Churchill-Livingstone, Harcourt Brace & Company, p:2877.

Miller & James (2002)

Angiokeratoma of Fordyce as a cause of red scrotum.
Cutis 69:50-51.

Moberg M, Gustavsson I & Gyllensten U (2004)

Type-specific associations of human papillomavirus load with of developing cervical carcinoma in situ.
Intl J Cancer 112(5): 854–859 .

Monsonogo J, Breugelmans JG, Bouee S et al.(2007)

Anogenital warts incidence, medical management and costs in women consulting gynaecologists in France.
Gynecol Obstet Fertil 2007, 35:107-113.

Moore RA, Edwards JE, Hopwood J et al. (2001)

Imiquimod for the treatment of genital warts: a quantitative systematic review.
BMC Infect Dis 1: 3-11 .

Moscicki AB (2005)

Impact of HPV infection in adolescent populations.
J Adolesc Health 37: S3–S9 .

Mroueh AM, Seoud MA, Kaspar HG et al. (2002)

Prevalence of genital human papillomavirus among Lebanese women.
Eur J Gynaecol Oncol 2002, 23:429-432.

Munoz N, Bosch FX, de Sanjose S et al. (2003)

Epidemiologic classification of human papillomavirus types associated with cervical cancer.
N Engl J Med 348:518–527.

N

Narang T & Kanwar A (2006)

Verrucous epidermal naevus on penis.
Indian J Dermatol vol 51, pp 222–223 .

Newman L, Datta D, Ahrens K et al. (2008)

Monitoring genital wart disease in STD clinics through the STD Surveillance Network (SSuN). In: National STD Prevention Conference, Illinois, Chicago .

Nickoloff BJ & Turka LA (1993)

Keratinocytes: Immunocytes of the integument.
Am J Pathol 143:325-331.

Nielsen A, Kjaer SK, Munk C et al. (2008)

Type-specific HPV infection and multiple HPV types: prevalence and risk factor profile in nearly 12,000 younger and older Danish women.
Sex Transm Dis 35: 276–282 .

Nielson CM, Flores R, Harris RB et al. (2007)

Human papillomavirus prevalence and type distribution in male anogenital sites and semen.
Cancer Epidemiol Biomarkers Prev 16: 1107–1114 .

O

Oriel JD (1971)

Natural history of genital warts .
Br J Vener Dis 47:1-13 .

P

Paavonen J, Jenkins D, Bosch FX et al. (2007)

Efficacy of a prophylactic adjuvanted bivalent L1 virus-like-particle vaccine against infection with human papillomavirus types 16 and 18 in young women: An analysis of a phase III double-blind, randomised controlled trial.

Lancet 369: 2161–2170 .

Padilla-Ailhaud A (2006)

Carbon dioxide laser vaporization of condyloma acuminata.

J Low Genit Tract Dis 10: 238–241 .

Palefsky JM, Holly EA, Hogeboom CJ et al. (1998)

Virologic, immunologic, and clinical parameters in the incidence and progression of anal squamous intraepithelial lesions in HIV-positive and HIV-negative homosexual men.

J Acquir Immune Defic Syndr Hum Retrovirol 17: 314–319 .

Parker S, Stewart A, Wren M et al. (1983)

Circumcision and sexually transmitted diseases.

Med J Aust 2:288–290.

Parkin DM (2006)

The global health burden of infection-associated cancers in the year 2002.

Int J Cancer 118:3030–3044.

Peate I (2006)

Nursing care and treatment of the patient with HPV.

Br J Nurs, 15, 1063-1069 .

Perez G, Lazcano-Ponce E, Hernandez-Avila M et al. (2008)

Safety, immunogenicity, and efficacy of quadrivalent human papillomavirus (types 6, 11, 16, 18) L1 virus-like-particle vaccine in Latin American women.

Int J Cancer 122: 1311–1318 .

Petersen CS, Bjerring P, Larsen J et al. (1991)

Systemic interferon alpha-2b increases the cure rate in laser treated patients with multiple persistent genital warts: a placebo-controlled study.

Genitourin Med 67: 99–102 .

Petersen CS & Weismann K (1995)

Quercetin and kaempferol: an argument against the use of podophyllin?
Genitourin Med 71(2):92 –93.

Petry KU, Scheffel D, Bode U et al. (1994)

Cellular immunodeficiency enhances the progression of human papillomavirus-associated cervical lesions.
Int J Cancer 57:836-840.

Pierson D, Bandel C, Ehrig T et al. (2003)

Benign epithelial tumors and proliferations. In: Bologna JL, Jorizzo JL & Rapini RP (eds) Dermatology. Mosby, Edinburgh, pp 1697–1720.

R

Renzi A , Giordano P, et al. (2006)

Buschke-Löwenstein tumor successful treatment by surgical excision alone: A case report.
Surg Inn 13(1):69-72.

Ricord P (1838)

Traité pratique des maladies vénériennes. J. Rouvier and E. le Bouvier, Paris . Quoted from : Rosenblatt A & De Campos Guidi HG (2009) Human Papillomavirus History and Epidemiology , in: Rosenblatt A & De Campos Guidi HG (eds.) Human Papillomavirus , A Practical Guide for Urologists , Springer-Verlag Berlin Heidelberg , ch: 1 ,pp: 3-15 .

Rogozinski TT & Janniger CK (1988)

Bowenoid papulosis.
Am Fam Physician 38:161– 164.

Rombaldi RL, Serafini EP, Mandelli J et al. (2008)

Transplacental transmission of human papillomavirus.
Virol J 5: 106-108 .

Rose RC & Stoler MH (2009)

Biology of HPV in: Bonnez W (ed) Guide to Genital HPV Diseases and Prevention, Informa Healthcare USA, Inc 2009 ch :1 , p: 1- 14.

Rosenblatt A & De Campos Guidi HG (2009)

Human Papillomavirus History and Epidemiology ,Human Papillomavirus and External Genital Lesions in: Rosenblatt A & De Campos Guidi HG (eds):Human Papillomavirus A Practical Guide for Urologists, Springer-Verlag Berlin Heidelberg , pp: 3-15 , 33-73 .

Routh HB, Bhowmik KR & Parish LC (1997)

Myths, fables and even truths about warts and human papillomavirus.
Clin Dermatol 15: 305–307 .

Rupp R, Stanberry LR & Rosenthal SL (2004)

New biomedical approaches for sexually transmitted infection prevention: vaccines and microbicides.
Adolesc Med Clin 15:393–407.

S

Sánchez-Alvarado JP, Franco-Morales A, Rosas-Nava JE et al. (2008)

Giant condyloma acuminatum (Buschke-Löwenstein tumor): a case report
Mex Urol ,68(6):348-350 .

Sarkola M, Rintala M, Grenman S et al. (2008)

Human papillomavirus DNA detected in breast milk.
Pediatr Infect Dis J 27: 557–558 .

Satterwhite CL (2008)

Up, up, and away? Trends in chlamydial infections in the U.S. In: National STD Prevention Conference, Chicago, IL .

Scheinfeld N & Lehman DS (2006)

An evidence-based review of medical and surgical treatments of genital warts.
Dermatol Online J 12: 5 – 9 .

Schneede P, Schlenker B and Hungerhuber E (2006).

Condylomata acuminata: an indication for prophylactic HPV vaccination?
Urologe A, 45, 1514-1520.

Schoen E (1997)

Benefits of newborn circumcision: is Europe ignoring medical evidence?
Arch Dis Child 77:258–260.

Schofer H, Van Ophoven A, Henke U et al. (2006)

Randomized, comparative trial on the sustained efficacy of topical imiquimod 5% cream versus conventional ablative methods in external anogenital warts.
Eur J Dermatol 16: 642–648 .

Schroder JM (1995)

Cytokine networks in the skin.
J Invest Dermatol 105:20S-24S.

Schwartz DB, Greenberg MD, Daoud Y et al. (1988)

Genital condylomas in pregnancy: use of trichloroacetic acid and laser therapy.
AmJ Obstet Gynecol 158:1407–1416.

Schwartz RA & Janniger CK (1991)

Bowenoid papulosis.
J Am Acad Dermatol 24:261-264.

Schwarz TF & Dubin G (2007)

Human papillomavirus (HPV) 16/18 L1 AS04 virus-like particle (VLP) cervical cancer vaccine is immunogenic and well-tolerated 18 months after vaccination in women up to age 55 years.
J Clin Oncol 25: 3007-3008 .

Sellers JW, Karwalajtys TL, Kaczorowski J, et al. (2003)

Incidence, clearance and predictors of human papillomavirus infection in women.
CMAJ 168:421–425 .

Sellers JW & Sankaranarayanan R (2003)

Colposcopy and Treatment of Cervical Intraepithelial Neoplasia: pub.:International Agency for Research on Cancer pp:72-88 .

Sherrard J & Riddell L (2007)

Comparison of the effectiveness of commonly used clinic-based treatments for external genital warts.
Int J STD AIDS 18: 365–366 .

Silverberg E, Boring C, and Squire T(1990)

Cancer statistics.

CA Cancer J Clin 40: 9–26.

Simms I & Fairley CK (1997)

Epidemiology of genital warts in England and Wales: 1971 to 1994.
Genitourin Med 73(5):365 –367.

Singer A & Monaghan J (1999)

Lower genital tract precancer In: Singer A, Monaghan J (eds.) ∴ Colpos Pathol Treat. 2nd ed . Blackwell Science , pp.339-344.

Silverberg MJ, Ahdieh L, Munoz A et al.(2002)

The impact of HIV infection and immunodeficiency on human papillomavirus type 6 or 11 infection and on genital warts.
Sex Transm Dis 29:427–435 .

Silverberg MJ, Thorsen P, Lindeberg H et al. (2003)

Condyloma in pregnancy is strongly predictive of juvenile-onset recurrent respiratory papillomatosis.
Obstet Gynecol 101:645–652.

Skinner SR, Garland SM, Stanley MA et al. (2008)

Human papillomavirus vaccination for the prevention of cervical neoplasia: Is it appropriate to vaccinate women older than 26?
Med J Aust 188: 238–242 .

Smetana Z, Malik Z, Orenstein A et al. (1997)

Treatment of viral infections with 5-aminolevulinic acid and light.
Lasers Surg Med 21: 351–358 .

Smith JS, Moses S, Hudgens MG et al. (2007)

Human papillomavirus detection by penile site in young men from Kenya.
Sex Transm Dis 34: 928–934 .

Sonnex C, Strauss S & Gray JJ (1999)

Detection of human papillomavirus DNA on the fingers of patients with genital warts.
Sex Transm Infect 75: 317–319

Sonnex C & Vrotsou K (2007)

Treatment of ano-genital warts: the effect of an educational event on practitioner choice.
Int J STD AIDS 18: 531–537 .

St Louis ME , Icenogle JP Manzila T et al. (1993)

Genital types of papillomavirus in children of women with HIV-1 infection in Kinshasa, Zaire.
Int. J. Cancer, 54, 181–184.

Stanley M (2003)

Genital Human Papillomavirus Infections-Current and Prospective Therapies, Journal of the National Cancer Institute Monographs No. 31, PP:117-225 .

Stanley M (2006)

Immune responses to human papillomavirus.
Vaccine 24 (suppl 1): S16–22.

Stanley M, Lowy DR & Frazer I (2006)

Prophylactic vaccines: underlying mechanisms.
Vaccine 24:S106–S113.

Steele JC, Stankovic T & Gallimore PH (1993)

Production and characterization of human proliferative T-cell clones specific for human papillomavirus type 1 E4 protein.
J Virol 67:2799-2806.

Steele JC , Mann CH , Rookes, S et al. (2005)

T-cell responses to human papillomavirus type 16 among women with different grades of cervical neoplasia.
Br. J. Cancer, 93:248–259 .

Sterling JC, Handfield-Jones S & Hudson PM (2001)

British Association of Dermatologists. Guidelines for the management of warts.
Br J Dermatol 144:4-11.

Stockfleth E, Beti H, Orasan R et al. (2008)

Topical Polyphenon E in the treatment of external genital and perianal warts: a randomized controlled trial.
Br J Dermatol 158: 1329–1338 .

Stone KM, Becker TM, Hadgu A et al. (1990)

Treatment of external genital warts: a randomised clinical trial comparing podophyllin, cryotherapy, and electrodesiccation.
Genitourin Med 66: 16–19 .

Stone KM, Karem KL, Sternberg MR et al. (2002)

Seroprevalence of human papillomavirus type 16 infection in the United States.
J Infect Dis 186: 1396–1402 .

Strauss MJ, Shaw EW, et al. (1949)

Crystalline virus-like particles from skin papillomas characterized by intranuclear inclusion bodies.
Proc Soc Exp Biol Med 72: 46–50 .

Syed TA, Lundin S, Cheema KM et al. (1994)

Human leukocyte interferonalpha in cream, for the treatment of genital warts in Asian women: a placebo-controlled, double-blind study.
Clin Investig 72: 870–873 .

Syrjanen S & Puranen M (2000)

Human papillomavirus infections in children: The potential role of maternal transmission.
Crit Rev Oral Biol Med 11: 259–274 .

Syrjoen K & Syrjoen S (2000)

Papillomavirus infections in human pathology. J Wiley & Sons, New York, 1-615.

Syrjoen KJ (1997)

Condyloma acuminatum and other HPV-related squamous cell tumours of the genito-anal area. In: Gross G, von Krogh G, eds. Human papillomavirus infections in dermatovenereology. CRC Press, Boca Raton, FL, 151-180.

Syrjon KJ (2000)

Terminology used in cytopathology.
CME J Gynecol Oncol 5:12-17.

T

Tabrizi SN & Garland SM (1996)

Is cryotherapy treating or infecting?
Med J Aust 164:263-266 .

Tang, CK , Shermeta DW & Wood C (1978)

Congenital condylomata acuminata.
Am. J. Obstet. Gynecol., 131, 912–913 .

Teichman JM , Thompson IM & Elston DM(2010)

Noninfectious Penile Lesions
Am Fam Physician. 81(2):167- 174.

Thomas JO , Herrero R , Omigbodun AA et al. (2004)

Prevalence of papillomavirus infection in women in Ibadan, Nigeria:
Apopulation-based study.
Br. J. Cancer, 90, 638–645 .

Tjalma WA, Arbyn M, Paavonen J et al. (2004)

Prophylactic human papillomavirus vaccines: the beginning of the end of
cervical cancer.
Int J Gynecol Cancer 14:751–761.

Tobian AA, Serwadda D, Quinn TC et al. (2009)

Male circumcision for the prevention of HSV-2 and HPV infections and
syphilis.
N Engl J Med 360: 1298-1309 .

Trottier H & Franco EL (2006)

The epidemiology of genital human papillomavirus infection.
Vaccine 24(Suppl 1): S1–S15 .

Tsambaos D (1997)

Treatment of condylomata acuminata with oral isotretinoin.
J Urol 158(5):1810-1013 .

V

Vaccarella S, Lazcano-Ponce E, Castro-Garduno JA et al. (2006)

Prevalence and determinants of human papillomavirus infection in men attending vasectomy clinics in Mexico.

Int J Cancer 119:1934–1949.

Vambutas A, DeVoti J, Pinn W et al. (2001)

Interaction of human papillomavirus type 11 E7 protein with TAP-1 results in the reduction of ATP-dependent peptide transport.

Clin Immunol 101: 94–99.

Villa LL, Costa RL, Petta CA, et al. (2006)

High sustained efficacy of a prophylactic quadrivalent human papillomavirus types 6/11/16/18 L1 virus-like particle vaccine through 5 years of follow-up.

Br J Cancer 95:1459–1466.

Viscidi RP, Ahdieh-Grant L, Clayman B et al. (2003)

Serum immunoglobulin G response to human papillomavirus type 16 virus-like particles in human immunodeficiency virus (HIV)-positive and risk-matched HIV-negative women.

J Infect Dis 187: 194–205 .

Von Krogh G (2001)

Management of anogenital warts (condylomata acuminata).

Eur J Dermatol 11: 598–603 .

Von Krogh G, Lacey CJ, Gross G et al. (2000)

European course on HPV associated pathology: guidelines for primary care physicians for the diagnosis and management of anogenital warts.

Sex Transm Infect 76:162–168.

W

Wang XL, Wang HW, Wang HS et al. (2004)

Topical 5-aminolaevulinic acid-photodynamic therapy for the treatment of urethral condylomata acuminata.

Br J Dermatol 151:880-885.

Waszak S (1978)

The historic significance of circumcision.
Obstet Gynecol 51:499–501.

Watts DH , Koutsky LA , Holmes KK et al. (1998)

Low risk of perinatal transmission of human papillomavirus: Results from a prospective cohort study.
Am. J. Obstet. Gynecol., 178, 365–373 .

Weaver BA, Feng Q, Holmes KK, et al.(2004)

Evaluation of genital sites and sampling techniques for detection of human papillomavirus DNA in men.
J Infect Dis 189(4):677– 685.

Wen LM, Estcourt CS, Simpson JM et al. (2007)

Risk Factors for HPV in Men .
JID 196:200 -207 .

Wentzensen N, Vinokurova S & Von Knebel DM (2004)

Systematic review of genomic integration sites of human papillomavirus genomes in epithelial dysplasia and invasive cancer of the female lower genital tract.
Cancer Res 64: 3878–3884 .

Wikstrom A (1995)

Clinical and serological manifestations of genital human papillomavirus infection.
Acta Derm Venereol Suppl (Stockh) 193: 81–85 .

Wiley DJ & Beutner KR (2000)

Genital warts.
Clin Evidence 3:764–774.

Wiley DJ, Douglas J & Beutner K (2002)

External genital warts: diagnosis, treatment, and prevention.
Clin Infect Dis 35: S210–S224 .

Wilson R (1947)

Circumcision and venereal disease. Can Med Assoc J 1947;56: 54–6.
quoted from: Lavreys L , Rakwar JB , Thompson ML et al.,
Circumcision and venereal disease .
The Journal of Infectious Diseases 1999;180:330–336 .

Winer RL, Feng Q, Hughes JP et al . (2008)

Risk of female human papillomavirus acquisition associated with first male sex partner.

J Infect Dis 197: 279–282 .

Winer RL, Hughes JP, Feng Q, et al(2006)

Condom use and the risk of genital human papillomavirus infection in young women.

N Engl J Med 354:2645–2654.

Winer RL, Lee SK, Hughes JP et al. (2003)

Genital human papillomavirus infection: Incidence and risk factors in a cohort of female university students.

Am J Epidemiol 157: 218–226 .

Winer, RL & Koutsky, LA (2004)

Delivering reassurance to parents: Perinatal human papillomavirus transmission is rare.

Sex Transm. Dis., 31, 63–64 .

Workowski KA & Berman SM (2007)

Centers for Disease Control and Prevention sexually transmitted diseases treatment guidelines.

Clin Infect Dis 44 Suppl 3: S73–S76 .

www. cervical-vaccine.com (2010)

http://cervical-vaccine.com/trust_gardasil.html , accessed in may 2010 .

www.gsk.fr (2010)

<http://gsk.fr/gsk/medicament/photo/cervarix.jpg> accsed in may 2010 .

Y

Yan J, Chen SL, Wang HN et al. (2006)

Meta-analysis of 5% imiquimod and 0.5% podophyllotoxin in the treatment of condylomata acuminata.

Dermatology 213: 218–223 .

Z

Zabawski EJ & Cockerell CJ (1998)

Topical and intralesional cidofovir: A review of pharmacology and therapeutic effects.

J Am Acad Dermatol 39:741–745.

Zur Hausen H (1976)

Condylomata acuminata and human genital cancer.

Cancer Res 36: 794 -801 .

Zur Hausen H (1991)

Human papillomaviruses in the pathogenesis of anogenital cancer.

Virology 184: 9–13 .

Zur Hausen H (2000)

Papillomaviruses causing cancer: Evasion from host-cell control in early events in carcinogenesis.

J Natl Cancer Inst 92: 690–698 .