
References

- Aaltonen R, Heikkinen T, Hakala K, Laine K, Alanen A (2005):** Transfer of proinflammatory cytokines across term placenta. *Obstet Gynecol.*; 106(4):802-807.
- Abbas, A. K., Lichtman, A. H., and Pober, J. S.,(1994):** Cellular and Molecular Immunology, 2nd ed., W. B. Saunders, Philadelphia, , 239-260.
- Adam Bitner, Jarosław Kalinka (2010).** IL- β ;, IL-6 promoter, TNF- α ; promoter and IL-1RA gene polymorphisms and the risk of preterm delivery due to preterm premature rupture of membranes in a population of Polish women *Arch Med Sci* 2010; 6, 4: 552-557
- Akira, S., Isshiki, H., Sugita, T., et al. (1990b).** A nuclear factor for IL-6 expression (NF-IL6) is a member of a C/EBP family *EMBO J.* 9, 1897-2005
- Allaire AD, Ballenger KA, Wells SR, McMahon MJ, Lessey BA(2000):** Placental apoptosis in preeclampsia. *Obstet Gynecol* 2000, 96:271-276.
- Al-Othman S, Omu AE, Diejomaoh FM, Al-Yatama M, Al-Qattan F(2001):** Differential levels of interleukin 6 in maternal and cord sera and placenta in women with pre- eclampsia. *Gynecol Obstet Invest*, 52:60-65
- Andrews WW, Hauth JC, Cliver SP, Conners M, GoldenbergRL(2000).** Endometrial microbial colonization and plasma cell endometritis are common three months postpartum (abstract). *Am J Obstet Gynecol.*;182:S53.
- Andrews AL, Nasir T, Bucchieri F, Holloway JW, Holgate ST, Davies DE(2006).**IL-13 receptor alpha 2: a regulator of IL-13 and IL-4 signal transduction in primary human fibroblasts. *J Allergy Clin Immunol*; 118:858–865.
-

-
- Arck PC. Stress and pregnancy loss(2000):** role of immune mediators, hormones and neurotransmitters. *Am J Reprod Immunol*; 46:117–23.
- Ashkar AA, Di Santo JP, Croy BA.** Interferon gamma contributes to initiation of uterine vascular modification, decidual integrity, and uterine natural killer cell maturation during normal murine pregnancy. *J Exp Med*; 92:259–70
- Ashkar S, Weber GF, Panoutsakopoulou V, Sanchirico ME, Jansson M, Zawaideh S, Rittling SR, Denhardt DT, Glimcher MJ, Cantor H. (2000).** Eta-1 (osteopontin): an early component of type-1 (cell-mediated) immunity. *Science* 287: 860-864 .
- Ashkenazi A. (2002)** Targeting death and decoy receptors of the tumournecrosis factor superfamily. *Nat Rev Cancer*;2:420 – 30.
- Atreya, R., Mudter, J., Finotto, S., Mullberg, J., et al ., (2000)** Blockade of interleukin-6 trans signaling suppresses T-cell resistance against apoptosis in chronic intestinal inflammation: evidence in Crohn’s disease and experimental colitis in vivo. *Nature Med* 6,583-588
- Austgulen, R., Lien, E., Liabakk, N.B., Jacobsen, G., Arntzen, K.J., 1994.** Increased levels of cytokines and cytokine activity modifiers in normal pregnancy. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 57, 149–155.
- Azizieh F, Raghupathy R, Makhseed M(2005).** Maternal cytokine production patterns in women with pre-eclampsia. *Am J Reprod Immunol.*; 54: 30-7.
- Bahar AM, Ghalib HW, Moosa RA, Zaki ZM, Thomas C, Nabri OA(2003).** Maternal serum interleukin-6, interleukin-
-

- 8, tumor necrosis factor-alpha and interferon-gamma in preterm labor. *Acta Obstet Gynecol Scand.*;82:/543_9.
- Barakoyi A, Polgar B, Szekeres-Bartho J(1999).** The role of Tcell receptor-positive cells in pregnancy :partII. *AmJReprod Immuol*;42:83-7
- Barata H, Thompson M, Zielinska W, Han YS, Mantilla CB, Prakash YS, Feitoza S, Sieck G, Chini EN(2004).** The role of cyclic-ADP-ribose-signaling pathway in oxytocin-induced Ca²⁺ transients in human myometrium cells. *Endocrinology*; 145: 881–889.
- Baud V, Karin M(2001).** Signal transduction by tumor necrosis factor and its relatives. *Trends Cell Biol*; 11: 372–377.
- Bauer S, Pollheimer J, Hartmann J, Husslein P, Aplin JD, Knofler M(2004):** Tumor necrosis factor-alpha inhibits trophoblast migration through elevation of plasminogen activator inhibitor-1in first-trimester villous explant cultures. *J Clin Endocrinol Metab*, 89:812-822
- Baxter, G. T., Kuo, R. C., Jupp, O. J., Vandenabeele, P. and MacEwan, D. J. (1999)** Tumor necrosis factor-alpha mediates both apoptotic cell death and cell proliferation in a human hematopoietic cell line dependent on mitotic activity and receptor subtype expression. *J. Biol. Chem.* 274, 9539-9547.
- Ben-Hur H, Gurevich P, Elhayany A, Avinoach I, Schneider DF, Zusman I(2005).** Transport of maternal immunoglobulins through the human placental barrier in normal pregnancy and during inflammation. *Int J Mol Med.*; 16(3):401-407.
- Benyo DF, Smarason A, Redman CW, Sims C, Conrad KP(2001):** Expression of inflammatory cytokines in placentas from women with preeclampsia. *J Clin Endocrinol Metab*, 86:2505-2512

- Beutler, B(1992).** Tumour Necrosis Factor: the Molecules and their Emerging Role in Medicine (Raven, New York,).
- Bodnar LM, Ness RB, Harger GF, Roberts JM(2005).** Inflammation and triglycerides partially mediate the effect of prepregnancy body mass index on the risk of preeclampsia. *Am J Epidemiol.*; 162(12)1198–1206
- Bose P, Black S, Kadyrov M, Weissborn U, Neulen J, Regan L, Huppertz B(2005):** Heparin and aspirin attenuate placental apoptosis in vitro: implications for early pregnancy failure. *Am J Obstet Gynecol*, 192:23-30.
- Bouwmeester T, Bauch A, Ruffner H, Angrand PO, Bergamini G, Croughton K, Cruciat C, Eberhard D, Gagneur J, Ghidelli S, Hopf C, Huhse B, Mangano R, Michon AM, Schirle M, Schlegl J, Schwab M, Stein MA, Bauer A, Casari G, Drewes G, Gavin AC, Jackson DB, Joberty G, Neubauer G, Rick J, Kuster B, Superti-Furga G(2004).** A physical and functional map of the human TNF-alpha/NF-kappa B signal transduction pathway. *Nat Cell Biol.* Feb;6(2):97-105.
- Bowen, J.M., Chamley, L., Keelan, J.A., Mitchell, M.D., (2002)** Cytokines of the placenta and extra-placental membranes: roles and regulation during human pregnancy and parturition. *Placenta* 23, 257–273.
- Breard, J., Costa, O., and Kordon, C.,(1995)** Organization and functional relevance of neuroimmune networks, in *Stress, the Immune System and Psychiatry*, Leonard, B. E. and Miller, K., Eds., John Wiley & Sons, Chichester, U.K., 1995, 17-45.
- Brewster JA, Orsi NM, Gopichandran N, Ekbote UV, Walker JJ(2007).** Gestational effects on host inflammatory response in normal and preeclamptic pregnancies. *Eur J Obstet Gynecol Reprod Biol*; in press.

- Burton GJ, Jauniaux E. (2004): Placental oxidative stress**From miscarriage to preeclampsia. *J Soc Gynecol Investig.* 11(6):342-352.
- Caniggia I, Winter J, Lye SJ, Post M(2000):** Oxygen and placental development during the first trimester: implications for the pathophysiology of pre-eclampsia.*Placenta*,21(Suppl A):25-30.
- Carp H, Torchinsky A, Fein A,(2001).** Hormones, cytokines and fetal anomalies in habitual abortion. *Gynecol Endocrinol*; 15:472–83.
- Casart CY, Tarrazzi K, Camejo MI(2007).** Serum levels of interleukin-6, interleukin-1b and human chorionic gonadotropin in pre-eclamptic and normal pregnancy. *Gynecol Endocrinol*;23:300–303.
- Castro-Rendo'n WA, Castro-Alvarez JF, Guzman-Martinez C, Bueno-Sanchez JC(2006).** Blastocyst-endometrium interaction: intertwining a cytokine network. *Braz J Med Biol Res*; 39: 1373–1385.
- Catalano RD, Yanaihara A, Evans AL, Rocha D, Prentice A, Saidi S, Print CG, Charnock-Jones DS, Sharkey AM & Smith SK 2003** The effect of RU486 on the gene expression profile in an endometrial explant model. *Molecular Human Reproduction* 9 465–473.
- Catherine L. Haggerty, Robert E. Ferrell, Carl A. Hubel., (2005):** Association between allelic variants in cytokine genes and preeclampsia. *A JObstet Gynecol.* ;193:209-15.
- Cedergren MI(2004).** Maternal morbid obesity and the risk of adverse pregnancy outcome. *Obstet Gynecol*;103:219-24.
- Chaouat G, Cayol V, Mairovitz V, Dubanchet S(1999).** Localization of the Th2 cytokines IL-3, IL-4, IL-10 at the fetomaternal interface during human and murine pregnancy and lack of requirement for Fas / Fas ligand interaction for a

- successful allogeneic pregnancy. *Am J Reprod Immunol*; 42: 1–13.
- Chaouat G, Ledee-Bataille N, Dubanchet S(2004),.** TH1/TH2 paradigm in pregnancy: paradigm lost? Cytokines in pregnancy/early abortion: reexamining the TH1/TH2 paradigm. *Int Arch Allergy Immunol*; 134:93–119.
- Chavarria ME, Lara-Gonzalez L, Gonzalez-Gleason A, Sojo I, and Reyes A (2002):** Maternal plasma cellular fibronectin concentrations in normal and preeclamptic pregnancies: a longitudinal study for early prediction of preeclampsia, *Am J Obstet Gynecol* 187:595-601.
- Chegini N, Ma C, Roberts M, Williams RS & Ripps BA (2002)** Differential expression of interleukins (IL) IL-13 and IL-15 throughout the menstrual cycle in endometrium of normal fertile women and women with recurrent spontaneous abortion. *Journal of Reproductive Immunology* 56 93–110.
- Chen, G., Goeddel, D.V. 2002.** TNF-R1 signaling: a beautiful pathway. *Science*. 296:1634-1635.
- Chiaramonte MG, Mentink-Kane M, Jacobson BA, Cheever AW, Whitters MJ, Goad ME, Wong A, Collins M, Donaldson DD, Grusby MJ. (2003).** Regulation and function of the interleukin 13 receptor alpha 2 during a T helper cell type 2-dominant immune response. *Journal of Experimental Medicine* 197 687-701
- Chrousos, G. P(1995),** The hypothalamic-pituitary-adrenal axis and the immune/inflammatory reaction, *N. Engl. J. Med.*, 332, 1351, 1995.
- Chung CP, Oeser A, Solus JF, Gebretsadik T, Shintani A, Avalos I, Sokka T, Raggi P, Pincus T, Stein CM(2008).** Inflammation-associated insulin resistance: differential effects in rheumatoid arthritis and systemic lupus erythematosus define potential mechanisms *Arthritis Rheum*. Jul;58(7):2105-12.

- Clark DA, Arck PC, Chaouat G(1999).** Why did your mother reject you? Immunogenetic determinants of the response to environmental selective pressure expressed at the uterine level. *Am J Reprod Immunol*; 41:5–22
- Conde-Agudelo A, Villar J, and Lindheimer M (2004):** World Health Organization systematic review of screening tests for preeclampsia. *Obstet Gynecol Dec*; 104(6):1367-1391.
- Cooper MA, Fehniger TA and Caligiuri MA (2001).** "The biology of human natural killer cell subsets." *Trends Immunol* 22(11): 633-40.
- Cope, A., Ettinger, R., McDevitt, H., 1997.** The role of TNF alpha and related cytokines in the development and function of the autoreactive T-cell repertoire. *Res. Immunol.* 148 (5), 307–312
- Culig Z, Bartsch G, Hobisch A (2003).** "Interleukin-6 regulates androgen receptor activity and prostate cancer cell growth". *Mol. Cell. Endocrinol.* 197 (1-2): 231–8.
- Cunningham FG, Leveno KJ, Hauth JC, Hauth LC, Gilstrap III LC, and Wenstrom KD (2005):** Hypertension disorders in pregnancy, *Williams Obstetrics 22nd edition.* Chapter 34:761 - 808.
- Curry AE, Vogel I, Skogstrand K, Drews C, Schendel DE, Flanders WD, Hougaard DM, Thorsen P(2008).** Maternal plasma cytokines in early- and mid-gestation of normal human pregnancy and their association with maternal factors. *J Reprod Immunol*; 77: 152–160
- Curry AE, Thorsen P, Drews C, Schendel D, Skogstrand K, Flanders WD, Hougaard D, Olsen J, Vogel I.(2009)** First-trimester maternal plasma cytokine levels, pre-pregnancy body mass index, and spontaneous preterm delivery. *Acta Obstet Gynecol Scand.* 2009;88(3):332-42.

- Dan-Sohkawa, M., M. Morimoto, and H. Kaneko. 1995.** In vitro, reactions of coelomocytes against sheep red blood cells in the solitary ascidian *Halocynthia roretzi*. *Zool. Sci.*12:411–417.
- Davis JR, Giardina JB, Green GM, Alexander BT, Granger JP, Khalil RA(2002).** Reduced endothelial NO-cGMP vascular relaxation pathway during TNF- α -induced hypertension in pregnant rats. *Am J Physiol Regul Integr Comp Physiol.*; 282: R390–R399.
- De Cesaris P, Starace D, Riccioli A, Padula F, Filippini A, Ziparo E(1998).** Tumor necrosis factor-alpha induces interleukin-6 production and integrin ligand expression by distinct transduction pathways. *J Biol Chem.*; 273: 7566–7571.
- Del Prete G, Maggi E, Romagnani S(1994).** Human Th1 and Th2 cells: Functional properties, mechanisms of regulation, and role in disease. *Lab Invest.*; 70(3):299-306.
- Demerath EW, Reed D, Rogers N, Sun SS, Lee M, Choh AC, Couch W, Czerwinski SA, Chumlea WC, Siervogel RM et al. (2008)**Visceral adiposity and its anatomical distribution as predictors of the metabolic syndrome and cardiometabolic risk factor levels. *American Journal of Clinical Nutrition* **88** 1263–1271.
- Denney JM, Nelson EL, Wadhwa PD, Waters TP, Mathew L, Chung EK, Goldenberg RL, Culhane JF.(2010)** Longitudinal modulation of immune system cytokine profile during pregnancy.Cytokine. 2010 Nov 29.
- Denison, F.C., Riley, S.C., Wathen, N.C. et al. (1998)** Differential concentrations of monocyte chemotactic protein-1 and interleukin-8 within the fluid compartments present during the first trimester of pregnancy. *Hum. Reprod.*, 13, 2292–2295.

- Desai TR, Leeper NJ, Hynes KL, Gewertz BL(2002):** Interleukin-6 causes endothelial barrier dysfunction via the protein kinase C pathway. *J Surg Res*, 104:118-23
- Dinarello CA(2007).** Historical insights into cytokines. *Eur J Immunol* ;37 (Suppl. 1): S34–S45
- Ding ZQ, Rowe J, Ng B, Sinosich MJ, Gallery ED(2002):** Modulation of prostacyclin and thromboxane secretion by cytotrophoblasts from normal and pre-eclamptic human pregnancies. *Placenta*, 23:594-599.
- Djurovic S, Clausen T, Wergeland R, Brosstad F, Berg K, Henriksen T(2002).** Absence of enhanced systemic inflammatory response at 18 weeks of gestation in women with subsequent pre-eclampsia. *BJOG*. Jul;109(7):759-64.
- Dosiou C, Giudice LC(2005):** Natural killer cells in pregnancy and recurrent pregnancy loss: endocrine and immunologic perspectives. *Endocrine Rev*, 26:44-62.
- Dubiński A, Zdrojewicz Z (2007).** "[The role of interleukin-6 in development and progression of atherosclerosis]". *Pol. Merkur. Lekarski* 22 (130): 291–4
- Dunn CL, Kelly RW, Critchley HO(2003):** Decidualization of the human endometrial stromal cell: an enigmatic transformation. *Reprod Biomed* , 7:151-161
- Durum SK and Oppenheim JJ (1993).** Proinflammatory cytokines and immunity. In: Paul WE, editor. *Fundamental immunology*, 3rd ed. New York:Raven Press, Ltd, 801–835.
- Durum, S. K. and Muegge, K(1995).** Cytokines linking the immune and inflammatory systems: IL1, TNF, IL6, IFN $\alpha\beta$ and TGF β . In Rich, R. (Ed.): Clinical Immunology. St. Louis, Mosby, pp. 350-362
- Ellis J, Wennerholm UB, Bengtsson A, Lilja H, Pettersson A, Sultan B, Wennergren M, Hagberg H(2001):** Levels of dimethylarginines and cytokines in mild and severe preeclampsia. *Acta Obstet Gynecol Scand*, 80:602-608

- Engert S, Rieger L, Kapp M, Becker JC, Dietl J, Kammerer U(2007).** Profiling chemokines, cytokines and growth factors in human early pregnancy decidua by protein array. *Am J Reprod Immunol*; 58: 129–137
- Fagiolo, U., Cossarizza, A., Scala, E., Fanales-Belasio, E., Ortolani, C., Cozzi, E., Monti, D., Franceschi, C., Paganelli, R., (1993).** Increased cytokine production in mononuclear cells of healthy elderly people. *Eur. J. Immunol.* 23, 2375–2378.
- Fain JN, Madan AK, Hiler ML, Cheema P & Bahouth SW 2004** Comparison of the release of adipokines by adipose tissue, adipose tissue matrix, and adipocytes from visceral and subcutaneous abdominal adipose tissues of obese humans. *Endocrinology* **145** 2273–2282.
- Fichorova RN, Anderson DJ(1999).** Differential expression of immunobiological mediators by immortalized human cervical and vaginal epithelial cells. *Biol Reprod*; 60: 508–514.
- Florio P, Reis FM, Pezzani I, Luisi S, Severi FM, and Petraglia F (2003):** The addition of activin A and inhibin A measurement to uterine artery Doppler velocimetry to improve the early prediction of pre-eclampsia, *Ultrasound Obstet Gynecol.* Feb;21(2):165-169.
- Flynn, R.M. and Palladino, M.A. (1992).** TNF and TGF-J3: the opposite sides of the avenue? In "Tumor Necrosis Factors. The Molecules and their Emerging Role in Medicine" (ed. B. Beutler), pp. 131-144. Raven Press, New York.
- Founds SA, Powers RW, Patrick TE, Ren D, Harger GF, Markovic N,(2008).** A comparison of circulating TNF-alpha in obese and lean women with and without preeclampsia. *Hypertens Pregnancy*;27(1):39–48.

- Frances M. Stewart, Dilys J. Freeman, Jane E. Ramsay, Ian A. Greer, Muriel Caslake and William R(2007).** Ferrell Longitudinal Assessment of Maternal Endothelial Function and Markers of Inflammation and Placental Function throughout Pregnancy in Lean and Obese Mothers The Journal of Clinical Endocrinology & Metabolism Vol. 92, No. 3 969-975
- Freeman DJ, McManus F, Brown(2004).** Short- and long-term changes in plasma inflammatory markers associated with preeclampsia. Hypertension; 43: 708-714
- Friebe-Hoffmann U, Chiao JP, Rauk PN(2001).** Effect of IL-1beta and IL-6 on oxytocin secretion in human uterine smooth muscle cells. Am J Reprod Immunol.; 46(3):226-231.
- Gammill HS, Roberts JM(2007):** Emerging concepts in preeclampsia investigation. Front Biosci, 12:2403-2411
- Gdrard, C., Bruyns, C., Marchant, A., et al. (1993).** Interleukin 10 reduces the release of tumor necrosis factor and prevents lethality in experimental endotoxemia. J. Exp. Med. 177, 547-550.
- Goepfert AR, Goldenberg RL, Andrews WW, Hauth JC, Mercer B, Iams J,(2001).** The Preterm Prediction Study: association between cervical interleukin 6 concentration and spontaneous preterm birth. National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network. Am J Obstet Gynecol.;184:/483_8.
- Goldenberg RL, Iams JD, Mercer BM, Meis PJ, Moawad A, Das A, et al. (2001)**The Preterm Prediction Study: toward a multiple-marker test for spontaneous preterm birth. Am J Obstet Gynecol.;185:/643_51
- Gonzalez F, Minium J, Rote NS, Kirwan JP(2006).** Altered tumor necrosis factor alpha release from mononuclear cells

- of obese reproductive-age women during hyperglycemia. *Metabolism*; 55(2)271–276
- Green NS, Damus K, Simpson JL, Iams J, Reece EA, Hobel CJ, et al. (2005)** Research agenda for preterm birth: recommendations from the March of Dimes. *Am J Obstet Gynecol*.
- Guarini S., Tagliavini S., Bazzani C., Bertolini A., Ferrari W. (1991)** Nicotine reverses hemorrhagic shock in rats. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 343:427–430
- Gurevich¹ Pavel, Asher Elhayany² Andrey P. Milovanov³ Reuvit Halperin⁴ Ella Kaganovsky⁵ Itzhak Zusman⁶ Herzel Ben-Hur · (2007)** The Placental Barrier in Allogenic Immune Conflict in Spontaneous Early Abortions: Immunohistochemical and Morphological Study *American Journal of Reproductive Immunology* Volume 58, Issue 5, pages 460–467, November .
- Guven MA et al.(2009)** Association of Maternal Serum CRP, IL-6, TNF- α , Homocysteine, Folic Acid and Vitamin B12 Levels with the Severity of Preeclampsia and Fetal Birth Weight. *Hypertension in Pregnancy*; 28:190–200.
- Guzeloglu-Kayisli O, Kayisli UA & Taylor HS 2009** The role of growth factors and cytokines during implantation: endocrine and paracrine interactions. *Seminars in Reproductive Medicine* 27 62–79.
- Gwozdziewiczová, S., R. Lichnovská, J. Hřebíček and F. Cesk,(2004).** Tumor Necrosis Factor alpha (TNF alpha) and insulin resistance. *Cesk Fysiol.*, 53: 167-175. PMID: 15704741
- Haddad R, Tromp G, Kuivaniemi H, Chaiworapongsa T, Kim YM, Mazor M, Romero R(2006).** Human spontaneous labor without histologic chorioamnionitis is characterized by an acute inflammation gene expression signature. *Am J Obstet Gynecol*; 195: 394 e1–394 e24

- Hadden, D.R. and C. McLaughlin,(2009).** Normal and abnormal maternal metabolism during pregnancy. *Semin Fetal Neonatal Med.*, 14: 66-71. PMID: 18986856
- Hager, K., Machein, U., Krieger, S., Platt, D., Seefried, G., Bauer, J. (1994).** Interleukin-6 and selected plasma proteins in healthy persons of different ages. *Neurobiol. Aging* 15, 771–772.
- Hamai Y, Fujii T, Yamashita T, Miki A, Hyodo H, Kozuma S, Geraghty DE, Taketani Y(1999):** The expression of human leukocyte antigen-G on trophoblasts abolishes the growth-suppressing effect of interleukin-2 towards them. *Am J Reprod Immunol*, 41:153-158.
- Hanna J, Goldman-Wohl D, Hamani Y, Avraham I, Greenfield C, Natanson-Yaron S, Prus D, Cohen-Daniel L, Arnon TI, Manaster I, Gazit R, Yutkin V, Benharroch D, Porgador A, Keshet E, Yagel S, Mandelboim O(2006).** Decidual NK cells regulate key developmental processes at the human fetal-maternal interface. *Nat Med*; 12:1065–1074.
- Hannan NJ, Salamonsen LA(2007).** Role of chemokines in the endometrium and in embryo implantation. *Curr Opin Obstet Gynecol*; 19: 266–272
- Hardardóttir I, Grünfeld C, Feingold KR(1994):** Effects of endotoxin and cytokines on lipid metabolism. *Curr Opin Lipidol* , 5:207-215.
- Hauguel-de Mouzon S & Guerre-Millo M (2006)** The placenta cytokine network and inflammatory signals. *Placenta* 27 794–798.
- Hayashi M, Ueda Y, Ohkura T, Inaba N(2005):** Interleukin-6 concentrations in the placenta and blood in normal pregnancies and preeclampsia. *Horm Metab Res*, 37:419-424
- Heikkinen J, Mottonen M, Komi J, Alanen A and Lassila O (2003).** "Phenotypic characterization of human decidual macrophages." *Clin Exp Immunol* 131(3): 498-505

- Heinrich PC, Behrmann I, Haan, S, Hermanns, HM, Müller-Newen G, Schaper, F (2003).** "Principles of interleukin-6-type cytokine signalling and its regulation". *Biochem. J.* 374: 1–20. doi:10.1042/BJ20030407.
- Heiskanen J, Romppanen EL, Hiltunen M, Iivonen S, Mannermaa A, Punnonen K, Heinonen S(2002).** Polymorphism in the tumor necrosis factor-alpha gene in women with preeclampsia. *J Assist Reprod Genet.*; 19: 220–223.
- Herrera JA, Chaudhuri G, Lopez-Jaramillo P(2001).** Is infection a major risk factor for preeclampsia? *Med Hypotheses*; 57: 393-397.
- Hiby SE, Walker JJ, O'Shaughnessy KM, Redman CWG, Carrington M, Trowsdale J, Moffett A(2004):** Combinations of maternal KIR and fetal HLA-C genes influence the risk of preeclampsia and reproductive success. *J Exp Med*, 200:957-965.
- Hirohata, S., Tanimoto, K. and Ito, K. (1993).** Elevation of cerebrospinal fluid interleukin-6 activity in patients with vasculitides and central nervous system involvement. *Clin.Immunol. Immunopathol.* 66, 225-229.
- Homburger, F. (1945).** A plasma fibrinogen-increasing factor obtained from sterile abscesses in dogs. *J. Clin. Invest.* 24,43-45.
- Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM(1995):** Increased adipose tissue expression of tumor necrosis factor- α in human obesity and insulin resistance. *J Clin Invest*, 95:2409-2415.
- Hotamisligil GS 2006** Inflammation and metabolic disorders. *Nature* 444 860–867.
- Hu W, Wang H, Wang Z, Huang H, Dong M(2007).** Elevated serum levels of interleukin-15 and interleukin-16 in preeclampsia. *J Reprod. Immunol.*; 73:166-71.

- Huang SJ, Schatz F, Masch R, Rahman M, Buchwalder L, Niven- Fairchild T, Tang C, Abrahams WM, Krikun G, Lockwood CJ. : (2006) : Regulation of chemokine production in response to pro-inflammatory cytokines in first trimester decidual cells. J Reprod Immunol ; 72: 60-73.**
- Huber J, Kiefer FW, Zeyda M, Ludvik B, Silberhumer GR, Prager G, Zlabinger GJ & Stulnig TM 2008** CC chemokine and CC chemokine receptor profiles in visceral and subcutaneous adipose tissue are altered in human obesity. *Journal of Clinical Endocrinology and Metabolism* **93** 3215–3221.
- Hunt JS (2006).** "Stranger in a strange land." Immunol Rev 213(36-47
- Ishihara K, Hirano T (2003).** "IL-6 in autoimmune disease and chronic inflammatory proliferative disease". Cytokine Growth Factor Rev. 13 (4-5): 357–68.
- Jacobs-Helber, S. M., K. H. Roh, D. Bailey, E. N. Dessypris, J. J. Ryan, J. Chen, A. Wickrema, D. L. Barber, P. Dent, S. T. Sawyer.(2003).** Tumor necrosis factor- α expressed constitutively in erythroid cells or induced by erythropoietin has negative and stimulatory roles in normal erythropoiesis and erythroleukemia. Blood 101: 524-531.
- James, J. L., Whitley, G. S. and Cartwright, J. E. (2010),** Pre-eclampsia: fitting together the placental, immune and cardiovascular pieces. The Journal of Pathology, 221: 363–378.
- Jilka et al., (1992).** Increased osteoclast development afterestrogen loss: mediation by interleukin-6. Science. 257, 88-91.
- Johnson, C.A., 2008.** Glucose homeostasis during canine pregnancy: Insulin resistance, ketosis and hypoglycemia. Theriogenology, 70: 1418-1423.

- Jones SA(2005):** Directing transition from innate to acquired immunity: defining a role for IL-6. *J Immunol*, 175:3463-3468
- Jonsson Y, Matthiesen L, Berg G, Ernerudh J, Nieminen K, Ekerfelt C(2005).** Indications of an altered immune balance in preeclampsia: a decrease in in vitro secretion of IL-5 and IL-10 from blood mononuclear cells and in blood basophil counts compared with normal pregnancy. *J Reprod Immunol.*; 66: 69-84.
- Jonsson Y, Rubèr M, Matthiesen L, Berg G, Nieminen K, Sharma S, Ernerudh J, Ekerfelt C(2006).** Cytokine mapping of sera from women with preeclampsia and normal pregnancies. *J Reprod Immunol.* Jun;70(1-2):83-91. Epub Jan 18.
- Kaaja RJ, Poyhonen-Alho MK(2006).** Insulin resistance and sympathetic overactivity in women. *J Hypertens*; 24(1)131–141
- Kac G, Dos Santos Vaz J, Schlüssel MM, Moura AS.(2010)** C-reactive protein and hormones but not IL-6 are associated to body mass index in first trimester of pregnancy.*Arch Gynecol Obstet.* Oct 26.
- Karumanchi SA, Bdolah Y(2004).** Hypoxia and sFlt-1 in preeclampsia: The "chicken-and-egg" question. *Endocrinology.*; 145(11):4835-4837.
- Kaufmann P, Black S, and Huppertz B (2003):** Endovascular trophoblast invasion: implication for the pathogenesis of intrauterine growth retardation and preeclampsia. *Biol Reprod* Mar;69:1-7.
- Kawamura K, Kawamura N, Kumagai J, Fukuda J, Tanaka T(2007).** Tumor necrosis factor regulation of apoptosis in mouse preimplantation embryos and its antagonism by

- transforming growth factor alpha / phosphatidylinositol 3-kinase signaling system. *Biol Reprod*; 76: 611–618
- Keelan JA, Mitchell MD(2007).** Placental cytokines and preeclampsia. *Front Biosci.*; 12: 2706-27.
- Kelly RW, Carr GG, Critchley HO (1997).** A cytokine switch induced by human seminal plasma: an immune modulation with implications for sexually transmitted disease. *Hum Reprod*; 12: 677–681.
- Keogh RJ, Harris LK, Freeman A, Baker PN, Aplin JD, Whitley GS, Cartwright JE(2007).** Fetal-derived trophoblast use the apoptotic cytokine tumor necrosis factor-alpha-related apoptosis-inducing ligand to induce smooth muscle cell death. *Circ Res*; 100: 834–841
- Kern PA, Saghizadeh M, Ong JM, Bosch RJ, Deem R, Simsolo RB(1995):** The expression of tumor necrosis factor in human adipose tissue; Regulation by obesity, weight loss, and relationship to lipoprotein lipase. *J Clin Invest*, 95:2111-2119.
- Kharfi A, Giguère Y, Sapin V, Massé J, Dastugue B, Forest JC(2003):** Trophoblastic remodeling in normal and preeclamptic pregnancies: implication of cytokines. *Clin Biochem*, 36:323-331.
- Kheshtchin N, Gharagozloo M, Andalib A, Ghahiri A, Maracy MR, Rezaei A. (2010)** The expression of Th1- and Th2-related chemokine receptors in women with recurrent miscarriage: the impact of lymphocyte immunotherapy. *Am J Reprod Immunol*. Aug 1;64(2):104-12.
- Kim JW, Back SH, Kim V, Ryu I, and Jang SK (2005):** Sequestration of TRAF2 into stress granules interrupts tumor necrosis factor signaling under conditions. *Mol Cell Biol Mar*; 25(6): 2450-2462.

- King AE, Kelly RW, Sallenave JM, Bocking AD, Challis JR(2007).** Innate immune defences in the human uterus during pregnancy. *Placenta*; 28: 1099–1106.
- Kishimoto, T(2005).** Interleukin-6: From basic science to medicine, 40 years in immunology. *Annu. Rev. Immunol.* 23:1-21.
- Kocyigity Y, Atamer Y, Atamer A, Tuzcu A, Akkus Z(2004).** Changes in serum levels of leptin, cytokines and lipoprotein in preeclamptic and normotensive pregnant women. *Gynecol Endocrinol*; 19: 267-273
- Koga K, Osuga Y, Yoshino O, Hirota Y, Ruimeng X, Hirata T, Takeda S, Yano T, Tsutsumi O, Taketani Y(2003):** Elevated serum soluble vascular endothelial growth factor receptor 1 (sVEGFR-1) levels in women with preeclampsia. *J Clin Endocrinol Metab*, 88:2348-2351.
- Kozma N, Halasz M, Palkovics T, Szekeres-Bartho J(2006):** The progesterone-induced blocking factor modulates the balance of PKC and intracellular Ca. *Am J Reprod Immunol*, 55:122-129.
- Krahl'ckova' M, Sima P, Rokyta Z(2005).** Role of the leukemia-inhibitory factor gene mutations in infertile women: the embryo-endometrial cytokine cross talk during implantation – a delicate homeostatic equilibrium. *Folia Microbiol*; 50: 179–186.
- KRIKOVSKY D, V&S&RHELYI B, T&TH-HEYN P, K&RNER A, TULASSAY T, MAD&CSY L (2002)** Association between G-308A polymorphism of the tumor necrosis factor-alpha gene and 24-hour ambulatory blood pressure val&es in type 1 diabetic adolescents. *Clin Genet* 62: 474-477.
- Kristiansen OP, Mandrup-Poulsen T (2006).** "Interleukin-6 and diabetes: the good, the bad, or the indifferent?". *Diabetes* 54 Suppl 2: S114–24.

- Kruessel JS, Biefield P, Polan ML, Simo'n C(2003).** Regulation of embryonic implantation. *Eur J Obstet Gynecol Reprod Biol*; 110 (Suppl. 1): S2–S9 .
- Kurpisz M and Fernandez N (1995).** "Immunology of human reproduction." BIOS, Oxford, UK.
- Lago F, Dieguez C, Gomez-Reino J & Gualillo O 2007** The emerging role of adipokines as mediators of inflammation and immune responses. *Cytokine and Growth Factor Reviews* **18** 313–325.
- Laird SM, Tuckerman EM, Li TC(2006).** Cytokine expression in the endometrium of women with implantation failure and recurrent miscarriage. *Reprod Biomed Online*; 13: 13–23.
- Lala PK, Chakraborty C(2003):** Factors regulating trophoblast migration and invasiveness: possible derangements contributing to pre-eclampsia and fetal injury. *Placenta*, 24:575-587.
- Lam C, Lim K-H, and Karumanchi SA (2005):** Circulating angiogenic factors in the pathogenesis and prediction of preeclampsia, *Hypertension* 46(5):1077-1085.
- Langhoff-Roos J, Kesmodel U, Jacobsson B, Rasmussen S, Vogel I(2006).** Spontaneous preterm delivery in primiparous women at low risk in Denmark: population based study. *BMJ*./ 332:/937_9.
- Lash GE, Otun HA, Innes BA, Kirkley M, De Oliveira L, Searle RF, Robson SC, Bulmer JN(2006).** Interferon-gamma inhibits extravillous trophoblast cell invasion by a mechanism that involves both changes in apoptosis and protease levels. *FASEB J*; 20: 2512–2518.
- Laskin CA, Bombardier C, Hannah ME, Mandel FP, Ritchie JW, Farewell V, Farine D, Spitzer K, Fielding L, Soloninka CA, Yeung M(1997):** Prednisone and aspirin in women with autoantibodies and unexplained recurrent fetal loss. *N Engl J Med*, 337:148-153.

- Lea RG(2001)** Proliferation, differentiation and apoptosis in pregnancy and cancer. In *Cancer and Pregnancy*, pp 216–228. Eds ER Barnea, E Jauniaux & PE Schwartz. London: Springer Verlag.
- Leal AM, Poon LC, Frisova V, Veduta A, Nicolaides KH (2009)**First-trimester maternal serum tumor necrosis factor receptor-1 and pre-eclampsia. *Ultrasound Obstet Gynecol.* Feb;33(2):135-41.
- Ledingham MA, Thomson AJ, Jordan F, Young A, Crawford M, Norman JE(2001).** Cell adhesion molecule expression in the cervix and myometrium during pregnancy and parturition. *Obstet Gynecol*; 97: 235–242.
- Lee HY, Sherwood OD(2005).** The effects of blocking the actions of estrogen and progesterone on the rates of proliferation and apoptosis of cervical epithelial and stromal cells during the second half of pregnancy in rats. *Biol Reprod*; 73: 790–797.
- Leung N, Zhang J, Lau TK, Chan LY, and Lo YM (2001):** Increased maternal plasma fetal DNA concentrations in women who eventually develop preeclampsia. *Clin Chem* 47:137-139.
- Levine RJ, Qian C, Leshane ES, Yu KF, England LJ, Schisterman EF, Wataganara T, Romero R, Bianchi DW(2004):** Two-stage elevation of cell-free fetal DNA in maternal sera before onset of preeclampsia. *Am J Obstet Gynecol*, 190:707-713 .
- Lewis SH, Perrin E (eds) (1999) .** *Pathology of the Placenta*. 2nd Edition
- Lin H, Mosmann TR, Guilbert L, Tuntipopipat S & Wegmann TG (1993)**Synthesis of T helper 2-type cytokines at the maternal-fetal interface. *Journal of Immunology* 151 4562–4573.

- Locksley RM, Killeen N, Lenardo MJ(2001).** The TNF and TNF receptor superfamilies: integrating mammalian biology. *Cell*;104:487–501.
- Loudon JA, Sooranna SR, Bennett PR, Johnson MR(2004).** Mechanical stretch of human uterine smooth muscle cells increases IL-8 mRNA expression and peptide synthesis. *Mol Hum Reprod*; 10: 895–899.
- Lu, H.L., H.W. Wang, Y. Wen, M.X. Zhang and H.H. Lin,(2006).** Roles of adipocyte derived hormone adiponectin and resistin in insulin resistance of type 2 diabetes. *World J. Gastroenterol.*, 12: 1747-1751. PMID: 16586545 .
- Ludmir J, Sehdev HM(2000).** Anatomy and physiology of the uterine cervix. *Clin Obstet Gynecol*; 43: 433–439.
- Luppi P, Haluszczak C, Betters D, Richard CA, Trucco M and DeLoia JA (2002a).** "Monocytes are progressively activated in the circulation of pregnant women." *J Leukoc Biol* 72(5): 874-84.
- Lyall F(2005):** Priming and remodelling of human placental bed spiral arteries during pregnancy – a review. *Placenta*, (Suppl A);31-36.
- Madazli R, Aydin S, Uludag S, Vildan O, Tolun N(2003):** Maternal plasma levels of cytokines in normal and preeclamptic pregnancies and their relationship with diastolic blood pressure and fibronectin levels. *Acta Obstet Gynecol Scand*, 82:797-802 .
- Madazli R, Kuseyrioglu B, Uzun H, Uludag S, and Ocak V (2005):** Prediction of preeclampsia with maternal mid-trimester placental growth factor, activin A, fibronectin and uterine artery Doppler velocimetry. *Int J Gynaecol Obstet Jun*; 89(3):251-257.
- Majumdar A, Cruz D, Asamoah N, et al (2007):** Activation of microglia acidifies lysosomes and leads to degradation of

- alzheimer amyloid fibrils. *Molecular Biology of the Cell* 18, 1490–1496 .
- Makhseed, M., Raghupathy, R., El-Shazly, S., Azizieh, F., Al-Harmi, J.A., Al-Azemi, M.M., (2003).** Pro-inflammatory maternal cytokine profile in preterm delivery. *Am. J. Reprod. Immunol.* 49, 308–318.
- Makrigiannakis A, Minas V(2007).** Mechanisms of implantation. *Reprod Biomed Online*; 14: 102–109.
- Makris A, Xu B, Yu B, Thornton C, Hennessy A(2006).** Placental deficiency of interleukin-10 (IL-10) in preeclampsia and its relationship to an IL10 promoter polymorphism. *Placenta*.; 27: 445-51.
- Malek A(2003).** Ex vivo human placenta models: Transport of immunoglobulin G and its subclasses. *Vaccine*.; 21(24):3362-3364.
- Malmstrom E, Sennstrom M, Holmberg A, Frielingsdorf H, Eklund E, Malmstrom L, Tufvesson E, Gomez MF, Westergren-Thorsson G, Ekman- Ordeberg G, Malmstrom A(2007).** The importance of fibroblasts in remodeling of the human uterine cervix during pregnancy and parturition. *Mol Hum Reprod*; 13: 333–341.
- Maria-Jesus Dominguez-Santallaa,Lourdes Loidib, Celsa Quinteirob, Luis F. Perezc(2004)** Relation of tumor necrosis factor (TNF) gene polymorphisms with serum concentrations and in vitro production of TNF-alpha and interleukin-8 in heavy drinkers: *Alcohol*, Volume 34, Issue 2, Pages 273-277 (October).
- Marvin KW, Keelan JA, Eykholt RL, Sato TA and Mitchell MD (2002).** "Use of cDNA arrays to generate differential expression profiles for inflammatory genes in human gestational membranes delivered at term and preterm." *Mol Hum Reprod* 8(4): 399-408.

- Mastorakos G, Ilias I (2007).** "Interleukin-6: a cytokine and/or a major modulator of the response to somatic stress". *Ann. N. Y. Acad. Sci.* 1088: 373–81.
- Matthiesen L, Berg G, Ernerudh J, Ekerfelt C, Jonsson Y, Sharma S.(2005)** Immunology of preeclampsia. *Chem Immunol Allergy.*; 89:49-61.
- Maury E & Brichard SM (2010)**Adipokine dysregulation, adipose tissue inflammation and metabolic syndrome. *Molecular and Cellular Endocrinology* **314** 1–16.
- McLachlan, K.A., D. O'Neal, A. Jenkins and F.P. Alford, (2006).** Do adiponectin, TNF- α , leptin and CRP relate to insulin resistance in pregnancy? Studies in women with and without gestational diabetes, during and after pregnancy. *Diabetes Metab. Res. Rev.*, 22: 131-138. PMID: 16170833.
- Milne F, Redman C, Walker J, et al(2005).** The pre-eclampsia community guideline (PRECOG): How to screen for and detect onset of pre-eclampsia in the community. *BMJ.*; 330(7491):576-580.
- Miyaura H, Iwata M(2002):** Direct and indirect inhibition of Th1 development by progesterone and glucocorticoids. *J Immunol*, 168:1087-1094.
- Montagnana M, Lippi G, Albiero A, Salvagno GL, Franchi M, Guidi GC(2008).** Serum pro-inflammatory cytokines in physiological and pre-eclamptic pregnancies *Gynecol Endocrinol.* Mar;24(3):113-6.
- Moore KL and Persaud TVN (2003).** "The developing human. 7th ed." Saunders, Philadelphia, USA .
- Morrison JJ,RennieJM.(1997)** scientific and ethical aspects of fetal and neonatal care at extremely preterm periods of gestation .*Br J Obstet Gynaecol.*104;1341-1350
- Munno, L. M. Chiechi, G. Lacedra, C. Berardesca, C. Patimo, L. Marcuccio, P. Nardelli and P. Loizzi (1999)** Evaluation of Nonspecific Immunity and Plasma Levels of Interferon- γ ,

- Interleukin-6 and Tumor Necrosis Factor- α in Preeclampsia
Immunopharmacology and Immunotoxicology, Vol. 21, No. 3, Pages 551-564
- Muttukrishna S, Hyett J, Paine M, Moodley J, Groome N, and Rodeck C (2006):** Uterine vein and maternal urinary levels of activin A and inhibin A in pre-eclampsia patients. Clin Endocrinol (Oxf). Apr;64(4):469-473.
- Nadar SK, Karalis I, Al Yemeni E, Blann AD, Lip GY(2005):** Plasma markers of angiogenesis in pregnancy induced hypertension. Thromb Haemost, 94:1071-1076 .
- National Heart, Lung, and Blood Institute(1998).** Clinical guidelines of the identification, evaluation, and treatment of overweight and obesity in adults. NIH Publication No. 98-4083. Bethesda, MDUSA: Department of Health and Human Services, National Institutes of Health; .
- Nauts HC, Fowler GA, Bogatko FH(1953).** A review of the influence of bacterial infection and of bacterial products (Coley's toxins) on malignant tumors in man; a critical analysis of 30 inoperable cases treated by Coley's mixed toxins, in which diagnosis was confirmed by microscopic examination selected for special study. Acta Med Scand Suppl 1953;276:1–103.
- Newman KC, Riley EM(2007).** Whatever turns you on: Accessory-cell-dependent activation of NK cells by pathogens. Nat Rev Immunol.; 7(4):279-291.
- Norman JE, Bollapragada S, Yuan M, Nelson SM(2007).** Inflammatory pathways in the mechanism of parturition. BMC Pregnancy Childbirth; 7 (Suppl. 1): S7.
- Ohsugil Y (2007):** Recent advances in immunopathophysiology of interleukin-6: An innovative therapeutic drug, tocilizumab (recombinant humanized anti-human interleukin-6 receptor antibody), Unveils the mysterious etiology of immune-

- mediated inflammatory diseases *biol pharm bull* 30, 11: 2001—2006.
- Okada H, Nakajima T, Yoshimura T, Yasuda K & Kanzaki H (2003)** Microarray analysis of genes controlled by progesterone in human endometrial stromal cells in vitro. *Gynecological Endocrinology* 17 271–280.
- Oppenheim, J. J. and F. W. Ruscetti. (1997).** Cytokines. In: Stites, D. P., Terr, A. I. and Parslow, T. G. (Eds.), *Medical Immunology*. Appleton & Lange, Stamford, CT., pp: 146-168.
- Osman I, Young A, Ledingham MA, Thomson AJ, Jordan F, Greer IA and Norman JE (2003).** "Leukocyte density and pro-inflammatory cytokine expression in human fetal membranes, decidua, cervix and myometrium before and during labour at term." *Mol Hum Reprod* 9(1): 41-5.
- Panagiotakos DB, Pitsavos C, Yannakoulia M, Chrysoshoou C, Stefanadis C.(2005)** The implication of obesity and central fat on markers of chronic inflammation: The ATTICA study. *Atherosclerosis*; 183(2)308–315.
- Papanicolaou, M.D.(1998)** Interleukin-6: The Endocrine Cytokine *The Journal of Clinical Endocrinology & Metabolism* Vol. 85, No. 3 1331-1333.
- Parr, E.L., Chen, H.L., Parr, M.B., Hunt, J.S.,(1995).** Synthesis and granular localization of tumor necrosis factor-alpha in activated NK cells in the pregnant mouse uterus. *J. Reprod. Immunol.* 28, 31–40.
- Paruk F, Moodley J(2000).** Maternal and neonatal outcome in early- and late-onset pre-eclampsia. *Semin Neonatol.*; 5(3):197-207.
- Patni S, Flynn P, Wynen LP, et al(2007).** An introduction to toll-like receptors and their possible role in the initiation of labour. *BJOG.*; 114(11):1326-1334.

- Peristats (2007) an interactive perinatal data resource [database on the Internet]. March of Dimes Birth Defects Foundation Perinatal Data Center. 2007.**
- Peter M.E mmerPM,Nelen WLOM,Steegers EAP,Hendriks JM,Veerhook M,Joosten I(2000).**Peripheral Natural Killer cytotoxicity and CD(+)CD16(−)cells increase during early pregnancy in women with a history of recurrent spontaneous abortion .Human Repro.15:1163-1169;
- Petersen and Bente Klarlund Pedersen(2005)** Centre of Inflammation and Metabolism at The Copenhagen Muscle Research Centre and The Department of Infectious Diseases, Rigshospitalet, University of Copenhagen, Copenhagen, DenmarkJ Appl Physiol 98: 1154-1162,
- Piccinni MP (2006)** T cells in normal pregnancy and recurrent pregnancy loss. Reproductive Biomedicine Online 13 840–844.
- Piguet PF, Vesin C.(2002)** Modulation of platelet caspases and life-span by anti-platelet antibodies in mice May;68(5):253-61eur J haematol.
- Pujol-Borrell,T,Toodd,I,Doshi, , M., Bottazzo, G.F., Sutton, , R., Gray, , D., Adolf, G.R. & Feldmann , M. (1987)** HLA class II induction in human islet cells by interferon-γ plus tumor necrosis factor or lymphotoxin. Nature, 326, 304.
- Quenby S, Mountfield S, Cartwright JE, Whitley GS, Vince G(2004):** Effects of low-molecular-weight and unfractionated heparin on trophoblast function. Obstet Gynecol, 104:354-361
- Quenby S, Kalumbi C, Bates M, Farquharson R, Vince G(2005):** Prednisolone reduces preconceptual endometrial natural killer cells in women with recurrent miscarriage. Fertil Steril, 84:980-984
- Raghupathy R, Makhseed M, Azizieh F, Omu A, Gupta M, Farhat R(2000).** Cytokine production by maternal

- lymphocytes during normal human pregnancy and in unexplained recurrent spontaneous abortion. *Hum Reprod*; 15(3):713-718.
- Ramos-Casals M, Brito-Zeron P, Munoz S, Soto MJ(2008).** BIOGEAS STUDY Group. A systematic review of the off-label use of biological therapies in systemic autoimmune diseases. *Medicine*;87:345–64.
- Ramsey JE, Clark P, Walker ID, Sattar N, Greer IA(2004).** Short and long term changes in plasma inflammatory markers associated with preeclampsia. *Hypertension*;43:708–714.
- Ray, A., Sassone-Corsi, P. and Sehgal, P.B. (1989).** A multiple cytokine- and second messenger-responsive element in the enhancer of the human interleukin-6 gene: similarities with cfos gene regulation. *Mol. Cell. Biol.* 9, 5537-5547.
- Rebel, Sheila Hartnett, Geoffrey R. Hill, Suzan B. Lazo-Kallanian,§ James L.M. Ferrara, and Colin A. Sieff (1999)** Essential Role for the p55 Tumor Necrosis Factor Receptor in Regulating Hematopoiesis at a Stem Cell Level *J. Exp. Med.* The Rockefeller University Press • Volume 190,□□ Number 10, November 15, 1493–1503
- Redman CW, Sargent IL(2005).** Latest advances in understanding preeclampsia. *Science*; 308(5728):1592-1594.
- Reis FM, D'Antona D, and Petraglia F (2002):** Prediction value of hormone measurement in maternal and fetal complication of pregnancy. *Endocrine Reviews* 23(2): 230-257.
- Robertson SA, Mayrhofer G, Seamark RF(1996).** Ovarian steroid hormones regulate granulocyte-macrophage colony-stimulating factor synthesis by uterine epithelial cells in the mouse. *Biol Reprod*; 54: 183 196.
- Robertson SA, Bromfield JJ, Tremellen KP(2003).** Seminal ‘priming’ for protection from pre-eclampsia-a unifying hypothesis. *J Reprod Immunol*; 59: 253–265.

- Robertson SA, O’Leary S, Armstrong DT(2006).** Influence of semen on inflammatory modulators of embryo implantation. *Soc Reprod Fertil Suppl*; 62: 231–245.
- Roederer, M., Staal, F.J.T., Raju, P.A., Herzenberg, L.A. and Herzenberg, L.A. (1992).** The interrelationship of tumor necrosis factor, glutathione, and AIDS. In "Tumor Necrosis Factor: Structure-function Relationship and Clinical Application" (ed T. Osawa and B. Bonavida), pp. 215-229. Karger, Basel.
- Rogers MS, Wang CC, Tam WH, Li CY, Chu KO, and Chu CY (2006):** Oxidative stress in midpregnancy as a predictor of gestational hypertension and pre-eclampsia. *BJOG*. Sep; 113(9):1053-1059.
- Rohra DK and Qidwai N (2005):** Biochemical screening for the prediction of preeclampsia. *JPMMA Feb*; 59(2):79-82.
- Rusten, L.S., F.W. Jacobsen, W. Lesslauer, H. Loetscher, E.B. Smeland, and S.E. Jacobsen. (1994).** Bifunctional effects of tumor necrosis factor alpha (TNF alpha) on the growth of mature and primitive human hematopoietic progenitor cells: involvement of p55 and p75 TNF receptors. *Blood*. 83:3152–3159.
- Saito S, Sakai M(2003):** Th1/Th2 balance in preeclampsia. *J Reprod Immunol*, 59:161-173.
- Saji F, Samejima Y, Kamiura S, Koyama M(1999).** Dynamics of immunoglobulins at the feto-maternal interface. *Rev Reprod.*; 4(2):81-89.
- Salem M, Delwel R, Touw I, Mahmoud LA, Elbasousy EM, Löwenberg B(1990).** Modulation of colony stimulating factor-(CSF) dependent growth of acute myeloid leukemia by tumor necrosis factor. *Leukemia*. Jan;4(1):37–43.
- Salomon LJ, Benattar C, Audibert F, Fernandez H, Duyme M, Taieb J, and Frydman R (2003):** Severe preeclampsia is

associated with high inhibin A levels and normal leptin levels at 7 to 13 weeks into pregnancy. *Am J Obstet Gynecol* 189:1517-1522.

Samuel O. Olusi, MBBS, FRCPath, PhD; Michael Diejomaoh, MBBS, FRCOG; Alexander Omu, MBBS, FRCOG; Amal Abdulaziz, BM, ChB, CABOG; Kinatamitath Prabha, MBBS, MD; Sunila George, MSc(2000) INTERLEUKINS IN PREECLAMPSIA *Annals of Saudi Medicine*, Vol 20, No 1,7

Savino, R., Lahm, A., Giorgio, M., Cabibbo, A., Tramontano, A and Ciliberto, G. (1993). Saturation mutagenesis of the human interleukin 6 receptor-binding site: implications for its three-dimensional structure. *Proc. Natl Acad. Sci. USA* 90,4067-4071

Schwantner, Andreas; Dingley Andrew J, Ozbek Suat, Rose-John Stefan, Grötzinger Joachim (Jan. 2004). "Direct determination of the interleukin-6 binding epitope of the interleukin-6 receptor by NMR spectroscopy". *J. Biol. Chem. (United States)* 279 (1): 571-6.

Sehgal, P.B., Zilberstein, A., Ruggieri, R.M. et al. (1986) Human chromosome 7 carries the beta-2 interferon gene. *Proc. Natl Acad. Sci.*, 83, 5219-5222.

Seki H, Matuoka K, Inooku H, Takeda S(2007). TNF-alpha from monocyte of patients with pre-eclampsia-induced apoptosis in human trophoblast cell line. *J Obstet Gynaecol Res*; 33: 408- 416.

Sennström MB, Ekman G, Westergren-Thorsson G, Malmström A, Byström B, Endresen U, Mlambo N, Norman M, Stabi B and Brauner A (2000). "Human cervical ripening, an inflammatory process mediated by cytokines." *Mol Hum Reprod* 6(4): 375-81

Serin Ibrahim Serdar, Bulent Ozcelik, Mustafa Basbug, Huseyin Kiliç, Derya Okur, Rusen Erez(2002) Predictive

- value of tumor necrosis factor alpha (TNF- α) in preeclampsia *European Journal of Obstetrics & Gynecology and Reproductive Biology* 5 August (Vol. 104, Issue 1, Page 86)
- Sharkey DJ, Macpherson AM, Tremellen KP & Robertson SA (2007)** Seminal plasma differentially regulates inflammatory cytokine gene expression in human cervical and vaginal epithelial cells. *Molecular Human Reproduction* 13 491–501.
- Sharma A, Satyam A, Sharma JB(2007).** Leptin, IL-10 and inflammatory markers (TNF-alpha, IL-6 and IL-8) in pre-eclamptic, normotensive pregnant and healthy non-pregnant women. *Am J Reprod Immunol*; 58: 21-30.
- Sharon Phelan(2010)**Obesity in the American population: calories, cost, and culture *American Journal of Obstetrics and Gynecology*, Volume 203, Issue 6, December, Pages 522-524
- Shirai, T., Yamaguchi, H., Ito, H., Todd, C.W. and Wallace R.B. (1985).** Cloning and expression in *Escherichia coli* of the gene for human tumour necrosis factor. *Nature* 313 803-806
- Sibai BM, Dekker G, Kupferminic M(2005):** Pre-eclampsia. *Lancet*, 365:785-799
- Sibai B et al(2009).** Maternal plasma concentrations of the soluble tumour necrosis factor receptor 2 are increased prior to the diagnosis of preeclampsia. *Am J Obstet Gynecol*;200:630.e1–630.e8.
- Smith, E. M.(1992),** Hormonal activities of cytokines, in *euroimmunoendocrinology*, 2nd ed., Blalock, J. E., Ed., Karger, Basel, Switzerland, , 154-169.
- Soleymanlou N, Jurisica I, Nevo O, Ietta F, Zhang X, Zamudio S, Post M, Caniggia I(2005):** Molecular evidence of

- placental hypoxia in preeclampsia. *J Clin Endocrinol Metab*, 90:4299-4308
- Statistics Canada(2005)**. Adult obesity in Canada: measured height and weight. *adults-adultes/8060-eng.htm*. Accessed January 6, 2010.
- Sullivan K.E,^{1*} A. B. M. Reddy,¹ K. Dietzmann,¹ A. R. Suriano,¹ V. P. Kocieda,¹ M. Stewart,² and M. Bhatia;(2007)** Epigenetic Regulation of Tumor Necrosis Factor Alpha Molecular and Cellular Biology, July, p. 5147-5160, Vol. 27, No. 14
- Svensson L, Arvola M, Sallstrom MA, Holmdahl R, Mattsson R(2001)**. The Th2 cytokines IL-4 and IL- 10 are not crucial for the completion of allogeneic pregnancy in mice. *J Reprod Immunol.*; 51(1):3-7.
- Sverremark Ekstrom E, Nilsson C, Holmlund U, et al(2002)**. IgE is expressed on, but not produced by, fetal cells in the human placenta irrespective of maternal atopy. *Clin Exp Immunol.*; 127(2):274-282.
- Szarka A, Rigo JJ, Lazar L, Beko G, Molvarec A.(2010)**Circulating cytokines, chemokines and adhesion molecules in normal pregnancy and preeclampsia determined by multiplex suspension array.*BMC Immunol*. Dec 2;11(1):59.
- TacchinAi-Cottier F, Vesin C, Redard M, Buurman W, and Piguet PF (1998)**. Role of TNF receptors I and II in TNF-induced platelets consumption in mice. *J Immunol* **160**: 6182–6186.
- Tavakol Afshari J, Ghomian N, Shameli A, Shakeri MT, Fahmidehkar MA, Mahajer E, et a(2005)l**. Determination of interlukin-6 and tumor necrosis factor-alpha concentrations in Iranian-Khorasanian patients with preeclampsia. *BMC Pregnancy and Childbirth*; 5: 14.

- Teppa RJ, Ness RB, Crombleholme WR, et al. (2000):** Free leptin is increased in normal pregnancy and further increased in preeclampsia. *Metabolism* 49:1043-1048.
- Teran E, Escudero C, Moya W, Flores M, Vallance P, Lopez-Jaramillo P(2001)** Elevated C-reactive protein and pro-inflammatory cytokines in Andean women with preeclampsia. *IntJGynecol Obstet*, 75:243- 9
- Thadhani R, Mutter WP, Wolf M, Levine RJ, Taylor RN, Sukhatme VP, Ecker J, Karumanchi SA(2004):** First trimester placental growth factor and soluble fms-like tyrosine kinase 1 and risk for preeclampsia. *J Clin Endocrinol Metab*, 89:770-775
- Thellin O, Heinen E(2003).** Pregnancy and the immune system: between tolerance and rejection. *Toxicology*; 185:179–84.
- Tordjman J, Poitou C, Hugol D, Bouillot JL, Basdevant A, Bedossa P, Guerre-Millo M & Clement K 2009** Association between omental adipose tissue macrophages and liver histopathology in morbid obesity: influence of glycemic status. *Journal of Hepatology* 51 354–362.
- Tornblom Sa, Klimaviciute A, Bystrom B et al(2005).** Non-infected preterm parturition is related to increased concentrations of IL- 6, IL-8 and MCP-1 in human cervix. *Reprod Biol Endocrinol.*;25;(3):39
- Tosun M, Celik H, Avci B, Yavuz E, Alper T, Malatyalioglu E(2010)** Maternal and umbilical serum levels of interleukin-6, interleukin-8, and tumor necrosis factor-alpha in normal pregnancies and in pregnancies complicated by preeclampsia *J Matern Fetal Neonatal Med.* Aug;23(8):880-6
- Trinchieri, G.(1994),** Interleukin-12: a cytokine produced by antigen-presenting cells with immunoregulatory functions in the generation of T-helper cells type 1 and cytotoxic lymphocytes, *Blood*, 84, 4008,.

- Vassiliadis, S., Ranella, A., Papadimitriou, L., Makrygiannakis, A., Athanassakis, I., (1998).** Serum levels of pro- and anti-inflammatory cytokines in non-pregnant women, during pregnancy, labour and abortion. *Mediators Inflamm.* 7, 69–72.
- Vesce F, Buzzi M, Ferretti ME, Pavan B, Bianciotto A, Jorizzo G, Biondi C(1998)** Inhibition of amniotic prostaglandin E release by ampicillin. *Am J Obstet Gynecol*, 178:759-764.
- Vesce F, Pavan B, Buzzi M, Pareschi MC, Bianciotto A, Jorizzo G, Biondi C(1999):** Effect of different classes of antibiotics on amniotic prostaglandin E release. *Prostaglandins Other Lipid Mediat*, 57:207-218.
- Vesce, F., Scapoli, C., Giovannini, G. et al. (2001)** Plasminogen activator system in serum and amniotic fluid of euploid and aneuploid pregnancies. *Obstet. Gynecol.*, 97, 404– 408.
- Vesce F, Scapoli C, Giovannini G, Tralli L, Gotti G, Valerio A, Piffanelli A(2002):** Cytokine imbalance in pregnancies with fetal chromosomal abnormalities. *Hum Reprod*, 17:803-808.
- Vesce F, Pavan B, Lunghi L, Giovannini G, Scapoli C, Piffanelli A, Biondi C(2004):** Inhibition of amniotic interleukin-6 and prostaglandin E2 release by ampicillin. *Obstet Gynecol*, 103:108-113.
- Vgontzas AN, Bixler EO, Lin HM, et al. (2005).** "IL-6 and its circadian secretion in humans". *Neuroimmunomodulation* 12 (3): 131–40.
- Von Dadelszen P, Magee LA, Kraiden M, et al(2003).** Levels of antibodies against cytomegalovirus and chlamydothila pneumoniae are increased in early onset pre-eclampsia. *BJOG.*; 110(8):725- 730.
- Vural P(2002).** Nitric oxide/endothelin-1 in preeclampsia. *Clinica Chimica Acta* 2002; 317: 65-70.
- Vural, P., Degirmencioglu, S., Saral, N. Y., Demirkan, A., Akgul, C., Yildirim, G., Issever, H. and Eroglu, H.**

- (2010), Tumor necrosis factor α , interleukin-6 and interleukin-10 polymorphisms in preeclampsia. *Journal of Obstetrics and Gynaecology Research*, 36: 64–71.
- Walker JJ.,(2000)**. Pre-eclampsia. *Lancet.*; 356(9237):1260-1265
- Walther T, Wessel N, Malberg H, Voss A, Stepan H, and Faber R (2006)**: A combined technique for predicting pre-eclampsia: concurrent measurement of uterine perfusion and analysis of heart rate and blood pressure variability. *J Hypertens*. Apr; 24(4):747-750.
- Wataganara T and Bianchi DW (2004)**: Fetal cell-free nucleic acids in the maternal circulation: New clinical applications. *Ann N.Y.Acad.Sci*. 1022:90-99.
- Watkins ML, Rasmussen SA, Honein MA, Botto LD, Moore CA(2003)**. Maternal obesity and risk for birth defects. *Pediatrics* 111:1152-8.
- Wegmann TG, Lin H, Guilbert L and Mosmann TR (1993)**. "Bidirectional cytokine interactions in the maternal-fetal relationship: is successful pregnancy a TH2 phenomenon?" *Immunol Today* 14(7): 353-6
- Weiser S, Miu J, Ball HJ, Hunt NH(2007)**. Interferon-gamma synergises with tumour necrosis factor and lymphotoxin-alpha to enhance the mRNA and protein expression of adhesion molecules in mouse brain endothelial cells. *Cytokine*; 37: 84–91.
- Williams MA, Mahomed K, Farrand A, Woelk GB, Mudzamiri S, Madzime S, King IB, McDonald GB.(1998)** Plasma tumor necrosis factor-alpha soluble receptor p55 (sTNFp55) concentrations in eclamptic, preeclamptic and normotensive pregnant Zimbabwean women. *J Reprod Immunol.*; 40: 159–173.

- Williams M, Farrand A, Mittendorf R, Sorensen T, Zingheim R, Reily CO, King I, Zebelman A & Luthy D (1999).** Maternal second trimester serum tumor necrosis factor- α -soluble receptor p55 (sTNFp55) and subsequent risk of preeclampsia. *American Journal of Epidemiology*, 149: 323-329.
- Winkler G, Cseh K, Barany E, Melczer Z, Speer G, Hajós P, et al.(2002)** Tumor necrosis factor system in insulin resistance in gestational diabetes. *Diabetes Res Clin Pract*;56(2):93–9.
- Wira CR, Fahey JV, Sentman CL, Pioli PA, Shen L(2005).** Innate and adaptive immunity in female genital tract: cellular responses and interactions. *Immunol Rev*; 206: 306–335.
- World Health Organization.(1977)** Recommended definitions; terminology and format for statistical tables related to the perinatal period. *Acta Obstet Gynecol Scand*; 56:247–53
- Wright A, Zhou Y, Weier JF, Caceres E, Kapidzic M, Tabata T, Kahn M, Nash C, Fisher SJ(2004):** Trisomy 21 is associated with variable defects in cytotrophoblast differentiation along the invasive pathway. *Am J Med Genet*, 130:354-364.
- Wulf Schneider-Brachert¹, Vladimir Tchikov², Oliver Merkel¹, Marten Jakob², Cora Hallas², Marie-Luise Kruse³, Peter Groitl¹, Alexander Lehn¹, Eberhard Hildt⁴, Janka Held-Feindt⁵, Thomas Dobner¹, Dieter Kabelitz², Martin Krönke⁶ and Stefan Schütze²(2006)** Inhibition of TNF receptor 1 internalization by adenovirus 14.7K as a novel immune escape mechanism *J Clin Invest.*;116(11):2901–2913.
- Xu G, Makris A, Thornton C, Hennessy A(2005):** Glucocorticoids inhibit placental cytokines from cultured normal and preeclamptic placental explants. *Placenta*, 26:654-660.

- Yoneyama Y, Suzuki S, Sawa R, Yoneyama K, Power GG, Araki T(2002):** Relation between adenosine and T-helper 1/T-helper 2 imbalance in women with preeclampsia. *Obstet Gynaecol*, 99:641-646.
- Yoshiyuki Adachi, Andrei L. Kindzelskii, Naohito Ohno, Toshiro Yadomae and Howard R. Petty(1999)** Amplitude and Frequency Modulation of Metabolic Signals in Leukocytes: Synergistic Role of IFN- γ in IL-6- and IL-2-Mediated Cell Activation *J. Immunol.*;163;4367-4374
- Zaretsky MV, Alexander JM, Byrd W, Bawdon RE(2004).** Transfer of inflammatory cytokines across the placenta. *Obstet Gynecol*. 103(3):546-550.
- Zavalza-Gómez, A.B., R. Anaya-Prado, A.R. Rincón-Sánchez and J.M. Mora-Martínez, 2008.** Adipokines and insulin resistance during pregnancy. *Diabetes Res.Clin. Pract.*, 80: 8-15. PMID: 18291552
- Zhang, A Harada, H Bluethmann, JB Wang, S Nakao, N Mukaida and K Matsushima(1995)** Tumor necrosis factor (TNF) is a physiologic regulator of hematopoietic progenitor cells: increase of early hematopoietic progenitor cells in TNF receptor p55-deficient mice in vivo and potent inhibition of progenitor cell proliferation by TNF alpha in vitro, Volume 86, Issue 8, pp. 2930-2937, Copyright © 1995 by The American Society of Hematology
- Zhang H & Zhang C 2009** Regulation of microvascular function by adipose tissue in obesity and type 2 diabetes: evidence of an adipose-vascular loop. *American Journal of Biomedical Sciences* **1** 133–142.