

# References

## **References**

**Abboud TK, Sarkis F, and Hung TT., (1983):** Effects of epidural anesthesia during labour on maternal plasma beta-endorphin level. *Anesthesiology* 59: 1–5.

**ACOG. American College of Obstetricians and Gynecologists. , (2003):** Practice Bulletin. Dystocia and augmentation of labour .Clinical management guidelines for obstetricians-gynecologists. No 49.

**Apari P and Rózsa L .,(2006):** Deal in the womb: fetal opiates, parent-offspring conflict, and the future of midwifery. *Medical Hypotheses* 67: 1189–1194.

**Augustinson LE, Bohilin P and Bundsen P.,(1977):** Pain Relief During Delivery By TENS. *Pain.* 4:59-65.

**Ayman A and Olah K. (2002):** The sharp end of medical practice: the use of acupuncture in obstetrics: Acupuncture and gynaecology. *Br J Obstet Gynaecol.* 109:1–4.

**Bars D, Villanueva L, Willer J and Bouhassira D., (1991):** Diffuse Noxious Inhibitory Controls in Animals and Man. *Acupunct Med.*9:47–56.

**Bartz JA and Hollander E., (2008):** Oxytocin and experimental therapeutics in autism spectrum disorders. *Progress in Brain Research* 170: 451–62.

**Beilin Y, Bodian CA, Weiser J, Hossain S, Arnold I, Feierman DE, Martin G and Holzman I.,(2005):** Effect of labour epidural analgesia with and without fentanyl on infant breast-feeding: A prospective, randomized, double-blind study. *Anesthesiology.* 103:1211-1217.

**Bennett VR, (1999):** Myles textbook for midwives. 13th Edition. London: Churchill Livingstone. Bennett VR, Brown LK, eds. P 478.

**Bergal S, Dalayeun JF and Norès JM., (1993):** Physiology of endorphins. A close-up view and a review of the literature. *Biomedicine and Pharmacotherapy* .47: 311–320.

**Bibelkommissionen., (New International Version )Bibel 2000:** (In Swedish). Stockholm: Verum :Genesis 3:16.

**Bonica JJ.,(1990):**The pain of childbirth. The management of pain. 2:1313-1343.

**Bonica JJ., (1996):** The nature of pain in parturition. Pain relief and anaesthesia in obstetrics. In Vanzundert A and Ostheimer GW (eds.) New York: Churchill Livingstone: 19-52.

**Borup L, Wurlitzer W and Hedegaard M., (2009):** Acupuncture as pain relief during delivery: a randomized controlled trial. Birth. 36(1):5-12.

**Brinsmead, M., R. B Singh, T Lewin, P Owens and Smith R., (1985):** Peripartum concentrations of beta endorphin and cortisol and maternal mood states. Aust N Z J Obstet Gynaecol 25(3): 194-197.

**Buckley, S. J., (2003):** Undisturbed Birth: Mother Nature's blueprint for safety, ease and ecstasy. J Prenatal Psychol Health.17 (4):261-288.

**Budd S. ,(2000):**Acupuncture. In Tiran D, Mack S. Complementary therapies for pregnancy and childbirth, 2nd ed. New York: Balliere Tindall. 79-104.

**Bundsen P, Peterson L and Seltam U., (1981):**Pain Releif During LabourBy TENS.AProspective Matched Study.Acta Obstet Gynecol Scand.60 : 459-468.

**Bystrova, K. and A. M. Widstrom., (2003):** Skin-to-skin contact may reduce negative consequences of the stress of being born: a study on temperature in newborn infants, subjected to different ward routines in St. Petersburg." Acta Paediatr 92(3): 320-326.

**Bystrova, K., A. and M. Widstrom., (2007):** Early lactation performance in primiparous and multiparous women in relation to different maternity home practices. A randomised trial in St. Petersburg. Int Breastfeed J 2: 9.

**Cabyoglu MT, Ergene N andTan U., (2006):**The mechanism of acupuncture and clinical applications. Int J Neurosci;116:115–125.

**Cai Wuying.** (1992): Acupuncture and the Nervous System. American Journal of Chinese Medicine .20(3-4): 331-337.

**Cammu H, Clasen K, Van Wettere L and Derde MP. , (1994):** To bathe or not to bathe during the first stage of labour. Acta Obstet Gynecol Scand 73:468-472.

**Carroll D, Moore RA, Tramer MR and McQuay HJ. , (1997):** Transcutaneous electrical nerve stimulation does not relieve labour pain: updated systematic review. Contemporary Reviews in Obstetrics and Gynaecology 9(3):195-205.

**Carter, C. S., (2003):** Developmental consequences of oxytocin. Physiol Behav 79(3): 383-397.

**Carter, C. S., (2007):** Sex differences in oxytocin and vasopressin: Implications for autism spectrum disorders. Behav Brain Res 176(1): 170-186.

**Chan EC, Smith R, Lewin T, Brinsmead MW, Zhang HP and Cubis J., (1993):** Plasma corticotropin-releasing hormone, beta-endorphin and cortisol interrelationships during human pregnancy. Acta Endocrinol. 128(4):339-344.

**Chang, Duck Hee Kang and Mi Kyeong Lee., ( 2004):** Effects of SP6 Acupressure on Labour Pain and Length of Delivery Time in Women During Labour. The Journal of Alternative and Complementary Medicine. 10(6): 959-965.

**Chang, Z. M. and Heaman M. I. (2005):** Epidural analgesia during labour and delivery: effects on the initiation and continuation of effective breastfeeding. J Hum Lact 21(3): 305-314.

**Chard, T., (1989):** Fetal and maternal oxytocin in human parturition. Am J Perinatol 6(2): 145-152 .

**Charlton E., (2005):** Pain and Pregnancy and Labour. Core Curriculum for Professional Education in Pain. 3<sup>rd</sup> Edition.IASP Press, Seattle.Ch. 36:2-3.

**Chavkin LF and James A., (1982):** Goldstein, Dynorphin is a specific endogenous ligand of the opioid receptor. Science. 215: 413–415.

**Cheek TG, Gutsche BB and Gaiser RR.,(1999):** The pain of childbirth and its effect on the mother and fetus. In Chestnut DH [ed]. Obstetric Anesthesia. Principles and Practice. 2nd ed. St Louis, Mosby :p 323.

**Chen XH and Han JS. ,(1992):** All three types of opioid receptors in the spinal cord are important for 2/15 Hz electroacupuncture analgesia, Eur. J Pharmacol 211: 203–210.

**Christensson, K.and C. Siles., (1992):** Temperature, metabolic adaptation and crying in healthy full-term newborns cared for skin-to-skin or in a cot. Acta Paediatr 81(6-7): 488-493.

**Chung UL, Hung LC, Kuo SC and Huang CL., (2003):** Effects of L14 and BL67 acupressure on labour pain and uterine contractions in the first stage of labour. J Nurs Res 4: 251–260.

**Csaba, G., (2008):** Hormonal imprinting: phylogeny, ontogeny, diseases and possible role in present-day human evolution. Cell Biochemistry function .26(1):1-10.

**Csaba, G. and K. Tekes., (2005):** Is the brain hormonally imprintable? Brain Development. 27(7): 465-471.

**Deadman .P ,and Al Khfaji .M, (2007):A Manual Of Acupuncture. 2<sup>nd</sup> ed .** published by Journal of Chinese Medicine. California .USA.p 17-21.



**Deneux-Tharaux, C. and E. Carmona., (2006):** Postpartum maternal mortality and cesarean delivery. *Obstet Gynecol* 108(3): 541-548.

**Douglas, A. J. Russell JA, and G. Leng., (2002):** The importance of oxytocin mechanisms in the control of mouse parturition. *Reproduction* 123(4): 543-552.

**Eckert K, Turnbull D and MacLennan A., (2001):** Immersion in water in the first stage of labour. *Birth*.28:84-93.

**Enkin, M., Keirse, M.J.N.C., Neilson, J., Crowther, C., Duley, L., Hodnett, E. and Hofmeyr, J. (2000):** A guide to effective care in pregnancy and childbirth. Oxford University Press, 3rd Edition. p 255-266.

**Eriksson SV, Lundeberg T and Lundeberg S., (1991):** Interaction of diazepam and naloxone on acupuncture induced pain relief. *Am. J. Chin. Med.* 19 (1): 1–7.

**Ernst, Lee H and Edzard., (2006):** Acupuncture - a critical analysis. Journal of Internal Medicine 259 (2): 125–137.

**Ernst E, Pittler MH, Wider B and Boddy K., (2007):**Acupuncture: its evidence-base is changing. Am J Chin Med. 35 (1): 21–25.

**Ernst E , Cho S-H and Lee H., (2010):** Acupuncture for pain relief in labour: a systematic review and meta-analysis. British Journal Of Obstetrics And Gynecology. 117(8):907-920.

**Fei H, Xie GX and Han JS., (1987):** Low, high frequency electroacupuncture stimulation release met-enkephalin and dynorphin A in rat spinal cord, Sci. Bull. China 32: 1496–1501.

**Ferber, S. G. and I. R. Makhoul., (2004):** The effect of skin-to-skin contact shortly after birth on the neurobehavioral responses of the term newborn: a randomized, controlled trial. Pediatrics 113(4): 858-865.

**Ferber, S.G., Granot, M. and Zimmer, E.Z., (2005):**Catastrophizing labour pain compromises later maternity adjustments. Am J Obstet and Gynec 192: 826-831.

**Fernando, R. and E. Bonello., (1997):** Neonatal welfare and placental transfer of fentanyl and bupivacaine during ambulatory combined spinal epidural analgesia for labour. *Anaesthesia* 52(6): 517-524.

**Ferraz MB, Quaresma MR and Aquino LR., (1990):** Reliability of pain scales in the assessment of literate and illiterate patients with rheumatoid arthritis. *J Rheumatol* . 17: 1022–1024.

**Farrara J, Young Jr.b, Linda LaMoreauxb, L Werthb,J.and R.Michael Poole. , (2001):** Clinical importance of changes in chronic pain intensity measured on an 11-point numerical pain rating scale.*Pain*. 94(2):149-158.

**Fields H, Basbaum A and Heinricher M., (2006):** Central nervous system mechanisms of pain modulation. In: McMahon S, Koltzenburg M, editors. *Wall and Melzack's Textbook of Pain*. 5th. London: Elsevier; p: 125–142.

**Fisher, J. and J. Astbury., (1997):** Adverse psychological impact of operative obstetric interventions. A prospective longitudinal study. Aust N Z J Psychiatry 31(5): 728-738.

**Furlan AD, van Tulder MW and Cherkin DC. , (2005).** Acupuncture and dry-needling for low back pain. Cochrane database of systematic reviews (Online) (1): CD001351

**Gibb DMF, Arulkumaran E, Lun KC and Rathan SS., (1984):** Characteristics of uterine activity in nulliparous labour. British Journal Obstetric and Gynaecology. 91:220-227.

**Grant GZ, Yu CS, Lee WL, RLX. And Brian M. (2005):** Involvement of peripheral opioid mechanisms in electroacupuncture analgesia. Explore Journal. 1(5):366–371.

**Gross, J.B., Cohen, A.P. and Lang, J.M., (2002):** Differences in systemic opioid use do not explain increased fever incidence in parturients receiving epidural analgesia. Anesthesiology, 97: 157-161.

**Goland, R. S., S. L. Wardlaw, Blum M, Tropper PJ and Stark RI., (1988):** Biologically active corticotropin-releasing hormone in maternal and fetal plasma during pregnancy. *Am J Obstet Gynecol* 159(4): 884-890 .

**Guastella AJ, Einfeld EL, Gray, K, Rinehart N, Tonge B, Lambert TJ and Hickie IB., (2010):** Intranasal oxytocin improves emotion recognition for youth with autism spectrum disorders. *Biological Psychiatry* 67 (7): 692-694.

**Guyton A., (1991):** The cerebellum, the basal ganglia, and overall motor control. in *Basic Neuroscience-Anatomy & Physiology*, ed Guyton AC (W.B. Saunders Company, Philadelphia), 2nd ed.268-269.

**Habek D, Cerkez Habek J and Jagust M.,(2003):** Acupuncture Conversion of Fetal Breech Presentation.*Fetal Diagn Ther* , 18:418-421.

**Habek, D. and R. Becarevic., (2007):** Emergency peripartum hysterectomy in a tertiary obstetric center: 8-year evaluation. *Fetal Diagn Ther* 22(2): 139-142.

**Hamilton, B. E., A. M. Minino, A. M., Martin, J. A., Kochanek, K. D., Strobino, D. M. and Guyer, B., (2007):** Annual summary of vital statistics. *Pediatrics* 119(2): 345-360.

**Hammerschlag R., (1998):** Methodological and ethical issues in clinical trials of acupuncture. *J Altern Complement Med.* 4:159–71.

**Han JS. , (2004):** Acupuncture and endorphins. *Neuroscience Letters* 361 :258–261.

**Han JS and Xie CW., (1982):** Dynorphin: potent analgesic effect in spinal cord of the rat. *Life Sci* 31: 1781–1784.

**Han JS and Xie GX., (1984):** Dynorphin: important mediator for electroacupuncture analgesia in the spinal cord of the rabbit, *Pain.* 18: 367–376.

**Han JS, Ding XZ and Fan SG. , (1986):** The frequency as the cardinal determinant for electroacupuncture analgesia to be reversed by opioid antagonists, *Acta Physiol. Sin* 38: 475–482.

**Han JS., (2004) :** Acupuncture and endorphins. *Neurosci Lett.* 1-3:258–261.

**Han Z, Jiang YH, Wan Y, Wang Y, Chang JK and Han JS., (1999):** Endomorphin-1 mediates 2 Hz but not 100 Hz electroacupuncture analgesia in the rat, *Neurosci. Lett* 274:75–78.

**Hantoushzadeh S, Alhusseini N and Lebaschi AH.,(2007):** The effects of acupuncture during labour on nulliparous women: A randomised controlled trial. *Aust J Obstet Gynaecol.* 47(1):26-30.

**Harriet Hall .(2008):** Puncturing the Acupuncture Myth.Science Based Medicine.p 252.

**He CM and Han JS. , (1990):** Attenuation of low rather than high frequency electroacupuncture analgesia following microinjection of beta-endorphin antiserum into the periaqueductal gray in rats. *Acupunct. Sci Int J* .1: 19–27.

**Heasman, L., J. A. Spencer and Symonds, M.E., (1997):** Plasma prolactin concentrations after caesarean section or vaginal delivery. *Arch Dis Child Fetal Neonatal Ed* 77(3): 237-238.

**Hemmings GT, Carli F, Weeks SK, Mazza L, Zingg HH.and Scull TJ.,(1998):** Epidural analgesia in early labour blocks the stress

response but uterine contractions remain unchanged. Can J Anesth. 45(7):626–630.

**Hemminki, E. and J. Merilainen., (1996):** Long-term effects of cesarean sections: ectopic pregnancies and placental problems. Am J Obstet Gynecol 174(5): 1569-1574.

**Herman A, Zimmerman A, Arieli S, Tovbin Y, Bezer M, Bukovsky I and Panski M. , (2002.) :** Down-up sequential separation of the placenta. Ultrasound Obstet Gynecol.; 19(3):278-281.

**Huang LX. , (1997):** Series Hua-xia press, Beijing, China,,of ancient classics on Acu-moxibustion Science. Fertil Steril. 86: 700–710.

**IASP., (1996):** Classification of chronic pain: Descriptions of chronic pain syndromes and definitions of pain terms. Pain. 6 (3): 247–252.

**Irestedt L, Lagercrantz H and Belfrage P., (1995):** Causes and consequences of maternal and fetal sympatho-adrenal activation during parturition. Acta Obstetrica ET Gynecologica Scandinavica. 118: 111-115.



**Jacob S, Brune CW, Carter CS, Leventhal BL, Lord C and Cook EH., (2007).** Association of the oxytocin receptor gene in Caucasian children and adolescents with autism. *Neuroscience Letters* 417 (1): 6–9.

**Kaptchuk TJ, Goldman P, Stone DA and Stason WB. ,(2000):**Do medical devices have enhanced placebo effects? *J Clin Epidemiol.*53: 786–92.

**Kannan, S., Jamison, R.N.and Datta, S., (2001):** Maternal satisfaction and pain control in women electing natural childbirth. *Regional Anesthesia and Pain Medicine.* 26: 468-472.

**Kirsch, P., C. Esslinger, Chen Q, Mier D, Lis S, Siddhanti S and Gruppe H., (2005):** Oxytocin modulates neural circuitry for social cognition and fear in humans. *J Neurosci* 25(49): 11489-11493.

**Kloosterman, G., (1982):** The universal aspects of childbirth: Human birth as a socio-psychosomatic paradigm. *J Psychosom Obstet Gynaecol* 1(1): 35-41, p 40.

**Kvorning N, Holmberg C, Grennert L, Aberg A and Akeson J (2004):** severe pregnancy-related low back and pelvic pain. *Spine* 29: 478-484.

**Laatikainen, T. J. (1991):**Corticotropin-releasing hormone and opioid peptides in reproduction and stress. *Ann Med* 23(5): 489-496.

**Lagercrantz, H. and T. A. Slotkin., (1986):** The stress of being born. *Sci Am.* 254(4): 100-107.

**Lao L, Hamilton GR, Fu J and Berman BM., (2003):** Is acupuncture safe? A systematic review of case reports. *Altern Ther Health Med.* 9 (1): 72–83.

**Lao L, Zhang G, Wei F, Berman BM and Ren K., (2001):** Electroacupuncture attenuates hyperalgesia and modulates FOS protein expression in rats with unilateral persistent inflammation. *J Pain.* 2: 111–117.

**Lawrence ,L. , Fontaine,p.,King,V.,Klein,M. and Ratcliffe,S.,(2003):**  
The Nature and Management of Labour Pain PartI.Nonpharmacologic  
Pain Relief. Am Fam Physician. 68(6):1109-1113.

**Lederman.RP., (1985):** Anxiety and epinephrine in multiparous  
women in labour: relationship to duration of labour and fetal heart  
rate pattern. Am J Obstet Gynecol .153(8):870-877.

**Lee MK, Chang SB and Kang DH., (2004) :**Effects of SP6  
acupressure on labour pain and length of delivery time in women  
during labour. J Altern Complement Med .10: 959–965.

**Lee H., & Ernst E.,(2004):** Acupuncture for labour pain  
management. Am J Obst Gynecol.191: 1573–1579.

**Lee HJ, Macbeth AH, Pagani JH and Young WS., (2009).**  
Oxytocin: the great facilitator of life. Progress in Neurobiology 88 2:  
127–151.

**Levine, E. M.and V. Ghai., (2001):** Mode of delivery and risk of  
respiratory diseases in newborns. Obstet Gynecol .97(3): 439-442.

**Lett.,(2004):**Acupuncture and endorphin.Neuroscience J.361:258-261.

**Lewith GT, Machin D. ,(1983):** On the evaluation of the clinical effects of acupuncture. Pain. 16:111–127.

**Lieberman, E. and C. O'Donoghue (2002):** Unintended effects of epidural analgesia during labour: a systematic review. Am J Obstet Gynecol 186(5):31-68.

**Linde K, Streng A, Jürgens S, Hoppe A, Brinkhaus B, Witt C, Wagenpfeil S, Pfaffenrath V, Hammes MG, Weidenhammer W, Willich SN and Melchart D., (2005):** Acupuncture for patients with migraine: a randomized controlled trial. JAMA.293 (17): 2118–2125.

**Liu JL, Han XW and Su SN., (1990):** The role of frontal neurons in pain and acupuncture analgesia. Sci. China, Ser. B, Chem. Life Sci. Earth Sci. 33 (8): 938–945.

**Liu JY, Bian DL and Han Y., (2005):** Progress of acupuncturing Hegu (LI 4) during labour Chinese Journal of Information on TCM. 12(5): 108–110.

**Lo SY. , (2002):** Meridians in acupuncture and infrared imaging. *Medical Hypotheses* 58:72-76.

**Lowe N.K.,(2002):** The nature of labour pain. *Am J Obstet Gynecol* 186:S16-24.

**MacDorman, M. F., E. Declercq, Menacker, F., and Malloy, M. H., (2006):** Infant and neonatal mortality for primary cesarean and vaginal births to women with no indicated risk, United States, 1998-2001 birth cohorts. *Birth*. 33(3): 175-82.

**Matthiesen, A. S., A. B. Ransjo-Arvidson, Lilja G, Nissen E and Widstrom A. M., (2001):** Postpartum maternal oxytocin release by newborns: effects of infant hand massage and sucking. *Birth* .28(1): 13-19.

**Melzack, R and Wall. PD.,(1965):** Pain mechanisms: a new theory. *Science*. 150 (699): 971–979.

**Melzack R, Taenzer P, Feldman P and Finch RA. , (1981):** Labour is still painful after prepared childbirth training. Canadian Medical Association Journal .125:357-363.

**Melzack, R., (2003):** The Gate Control Theory: Reaching for the Brain. *Pain*: 104(3):597-608.

**Mendelson, C. R. and V. Boggaram., (1990):** Hormonal and developmental regulation of pulmonary surfactant synthesis in fetal lung. *Baillieres Clin Endocrinol Metab*.4 (2): 351-378.

**Mostyn, A., S. Pearce and Alves-Guerra MC ., (2004):** Hormonal and nutritional regulation of adipose tissue mitochondrial development and function in the newborn. *Exp Clin Endocrinol Diabetes*. 112(1): 2-9.

**Napadow V, Ahn A, Longhurst J, Lao L, Stener-Victorin E, Harris R and Langevin HM .,(2008):** The status and future of acupuncture clinical research. *Journal of alternative and complementary medicine* 14 (7): 861–9.

**Nesheim, , Ragnhild Kinge, , Bertha Berg, Birgitta Alfredsson, ,Eibjörg Allgot, , Gry Hove, , Wenche Johnsen, Ingunn Jorsett,**

**Sigrun Skei and Stani Solberg .,(2003)** Acupuncture during labor can rReduce the use of meperidine:a controlled clinical study .The Clinical Journal of Pain 19:187–191.

**Nissen, E., K. Uvnas-Moberg, Svensson K; Stock S; Widström A M and Winberg J., (1996):** Different patterns of oxytocin, prolactin but not cortisol release during breastfeeding in women delivered by caesarean section or by the vaginal route. Early Hum Dev. 45(1-2): 103-118.

**Nissen, E., P. Gustavsson, Widström AM and Uvnäs-Moberg K., (1998):** Oxytocin, prolactin, milk production and their relationship with personality traits in women after vaginal delivery or Cesarean section. J Psychosom Obstet Gynaecol .19(1): 49-58.

**Nissen, E.A. Ransjö-Arvidson<sup>1</sup>, Matthiesen,A. and Uvnäs-Moberg,K.,(2008):** Influence of oxytocin or epidural analgesia on personality profile in breastfeeding women: a comparative study.Women's Mental Health Journal.11:335-345.

**Odibo, A.O., A.G. Cahill, Stamilio DM, Stevens EJ, Peipert JF and Macones GA., (2007):** Predicting placental abruption and previa

in women with a previous cesarean delivery. Am J Perinatol. 24(5): 299-305.

**Odent, M., (1998):** Don't manage the third stage of labour. Practising Midwife. 1:9, p31-33

**Olofsson C, Ekblom A, Ekman-Ordeberg G, Hjelm A and Irestedt L. (1996):** Lack of analgesic effect of systemically administered morphine or pethidine on labour pain. Br J Obstet Gynaecol. 103:968–972.

**Oschman JL., (1993):** A biophysical basis for acupuncture. Proceedings of the first symposium of the society for acupuncture research. Rockville, : 23-24

**Phaneuf, S., B. Rodriguez Linares, R.L. TambyRaja, I.Z. MacKenzie and A. López Bernal. , (2000):** Loss of myometrial oxytocin receptors during oxytocin-induced and oxytocin-augmented labour. J Reprod Fertil .120(1): 91-97.



**Poggi, S.H., Allen, R.H. and Patel, C.,(2004):** Effect of epidural anesthesia on clinician-applied force during vaginal delivery. American Journal of Obstetrics & Gynecology. 191: 903-906.

**Pomerans,G.S.,(1999):**Basics Of Acupuncture Textbook.Springer-Verlag Berlin Heidelberg New York 3<sup>rd</sup> edition : 14

**Purves, Dale, George J. Augustine, David Fitzpatrick, William C. Hall, Anthony-Samuel LaMantia, James O. McNamara, and Leonard E. White.,(2008).** Neuroscience. 4th ed. Sinauer Associates. 137–138.

**Quiton, R.L. and Greenspan, J.D.,(2007):** Sex Differences in Endogenous Pain Modulation by Distracting and Painful Conditioning Stimulation. **Pain**, 132 :26-45.

**Radzyminski, S., (2005):** Neurobehavioral functioning and breastfeeding behavior in the newborn. J Obstet Gynecol Neonatal Nurs 34(3): 335-341.

**Rahm V-A, Hallgren A and Hogberg H., (2002):** Plasma oxytocin levels in women during labour with or without epidural analgesia a prospective study. Acta Obstet Gynecol Scand, 81: 1033-1039.

**Ramnero A, Hanson U and Kihlgren M., (2002):** Acupuncture treatment during labour, a randomized controlled trial. Br J Obstet Gynaecol. 109: 637–644.

**Ringler, N. M., J. H. Kennell, arvella, R., Navajosky, B.J. and Klaus, M.H. , (1975):** Mother-to-child speech at 2 years effects of early postnatal contact. J Pediatr .86(1): 141-144.

**Riordan, J., A. Gross, A, Angeron J, Krumwiede B and Melin J. (2000):** The effect of labour pain relief medication on neonatal suckling and breastfeeding duration. J Hum Lact 16(1): 7-12.

**Rivier, C., W. Vale, N. Ling, M. Brown and R. Guillemin. , (1977):** Stimulation in vivo of the secretion of prolactin and growth hormone by beta-endorphin. Endocrinology 100(1): 238-41.

**Rosen .,(2002):** Nitrous oxide for relief of labour pain. Am J Obstet Gynecol.186:S110-26.

**Rubik.,(1995):** Can Western Science Provide A Foundation For Acupuncture .Alternative Therapies Magazine .1:4.

**Rudra, A.(2004):** Pain Relief In Labour.Updates In Anasthesia.18:1-2.

**Segal, S., A.N. Csavoy, and Datta S., (1998):** The tocolytic effect of catecholamines in the gravid rat uterus. Anesth Analg. 87(4): 864-869.

**Sharma, S.K., McIntyre, D.D.and Wiley, J., (2004):** Labour analgesia and cesarean delivery. Anesthesiology, 100: 142-148.

**Shen J., (2001):** Research on the neurophysiological mechanisms of acupuncture: review of selected studies and methodological issues. Journal of alternative and complementary medicine .7 (1):121–127.

**Sim J., (1997):** The mechanism of acupuncture analgesia: a review. Com Ther Med. 5: 102–111.

**Simkin PP. ,(2002):** Nonpharmacologic relief of pain during labour: systematic reviews of five methods. *Am J Obstet Gynecol* .186(5): 31-59.

**Sakala C., (1988):** Content of care by independent midwives: assistance with pain in labor and birth. *Soc Sci Med* 26: 1141–1158.

**Skilnand E, Fossen D and Heiberg E., (2002):** Acupuncture in the management of pain in labour. *Acta Obstet Gynecol Scand*. 81:943–948.

**Smith CA, Collins CT, Cyna AM and Crowth CA.,(2003):**Complementary and alternative therapies for pain management in labour. *Cochrane Database Syst*; 2: 637–644.

**Smith, Caroline A. ,Crowther, Caroline A. ,Collins, Carmel T. Coyle and Meaghan E.,(2008):** Acupuncture to Induce Labour.*Obstetrics&Gynecology* .112:1067-1074.

**Sodipo JO, Gilly H and Pauser G., (1981):** Endorphins: Mechanism of acupuncture analgesia. *Am J Chin Med*. 9: 249–258.

**Steer P., (1993):** The methods of pain relief used. In: Pain and its relief in childbirth: the result of a National Survey conducted by the National Birthday: 49–67.

**Stefos, T., A. Sotiriadis, Tsirkas P; Messinis I and Lolis D., (2001):** Maternal prolactin secretion during labour. The role of dopamine. Acta Obstet Gynecol Scand. 80(1): 34-38.

**Steinbrook, R.A., Carr, D.B., Datta, S., Naulty, J.S., Lee, C. and Fisher, J. ,(1982):** Dissociation of plasma and cerebrospinal fluid beta-endorphin-like immuno-reactivity levels during pregnancy and parturition. Anesth. Analg. 61: 893-897.

**Subhuti Dharmananda., (1996):** An Introduction to Acupuncture and how it work. Journal of Chinese Medicine. 52:(3)5-6.

**Takayanagi Y, Yoshida M and Bielsky IF., (2005):** Pervasive social deficits, but normal parturition, in oxytocin receptor-deficient mice. Proceedings of the National Academy of Sciences of the United States of America 102 (44): 16096–16101.

**Takeshige C, Sato T, Mera T, Hisamitru T and Fang JQ., (1992):** Descending pain inhibitory system involved in acupuncture analgesia. Brain Res Bull. 29: 617–634.

**Thomas, T. A. and Fletcher, J. E. (1982):** Influence of medication of pain and progress in labour on plasma beta-endorphin-like immunoreactivity. Br J Anaesth 54(4): 401-408.

**Thomas, S. A., Fletcher, and R. D. Palmiter., (1997):** Impaired maternal behavior in mice lacking norepinephrine and epinephrine. Cell 91(5): 583-592.

**Tiran D (2006):** Midwives responsibilities when caring for women using complementary therapies during labour. Midirs 16(1):77-80.

**Tsuchiya M, Sato EF, Inoue M and Asada A., (2007):** Acupuncture enhances generation of nitric oxide and increases local circulation. Anesth. Analg. 104 (2): 301–307.

**Tsui, M.H., W.D. Ngan Kee, Ng FF and Lau TK., (2004):** A double blinded randomised placebo-controlled study of intramuscular

pethidine for pain relief in the first stage of labour. Br J Obstet Gynaecol .111(7): 648-655.

**Tyzio, R., R. Cossart, Khalilov I, Minlebaev M, Hubner CA, Represa A, Ben Ari Y and Khazipov R., (2006):** Maternal oxytocin triggers a transient inhibitory switch in GABA signaling in the fetal brain during delivery. Science. 314(5806): 1788-1792.

**Ulett GA., (1984):** Acupuncture update. Southern Medical Journal 78:233-234.

**Ulett GA, Han SP and Han JS.(1998):** Electroacupuncture: mechanisms and clinical application. Biol Psychiatry .44:129–138.

**Umeh B., (1986):** Sacral acupuncture for pain relief in labour: initial clinical experience in Nigerian women. Acupunct Electrother Res. 11:147–151.

**Uvnas-Moberg., (2003):** Role of efferent and afferent vagal nerve activity during reproduction: Integrating function of oxytocin on metabolism and behaviour. Psycho neuroendocrinology .19 :687-695.

**Varrassi G, Bazzano C and Edwards WT.,(1989):** Effects of physical activity on maternal plasma beta-endorphin levels and perception of labour pain. *Am J Obstet Gynecol.* 160(3):707-712.

**Wagner, T.D., Rilling, J.K., Smith, E.E., Sokolik, A.and Casey, K.L., (2007):** Placebo-Induced Changes in fMRI in the Anticipation and Experience of Pain. *Science.* 303 (5661): 1162–1167.

**Wallis L, Shnider SM, Palahniuk RJ and Spivey HT., (1974):** An evaluation of acupuncture analgesia in obstetrics. *Anesthesiology.*41 (6):596–601.

**Wang MQ, Zhu J, Zhang LF, Downs, T.and Fu X. D., (2005):** Experimental study on the mechanism of electroacupuncture at LI 4 and SP 6 points for promoting labour. *China Acupuncture.* 23:10 593–596.

**Ward ME., (1997):** Acute pain and the obstetric patient: recent developments in analgesia for labour and delivery. *International Anaesthesiology Clinics.*35:83-103.

**Wu, S., M. Kocherginsky and J.U. Hibbard.; (2005):** Abnormal placentation: twenty-year analysis. *Am J Obstet Gynecol .*192(5): 1458-61.



**Yanai N, Shalev E, Yagudin E and Zuckerman H., (1987):** The use of acupuncture during labour. *Am J Acupunct.*15:311–312 .

**Yang ZX., (2002):** Clinical experience in treating diseases with acupuncturing at Hegu and Sanyinjiao *Zejiang Journal of Sichuan of Traditional Chinese Medicine.*20(7):75–76.

**Yang ZX. , (2008):** Treatment with acupuncturing at Hegu and Sanyinjiao, *Zejiang Journal of Chinese Medicine.* 10:445–446.

**Yentis SM., (2001):** The use of Entonox for labour pain should be abandoned. *Int J Obstet Anesth.* 10:25–27.

**Zhao,CH.,Kung,J. and Adachi,N., (2003):** Effect Of Local Injection Of Steroid And Anesthetics On Electroacupuncture:Prevention Of Immediate AnalgesiaAnd Induction Of Hyperalgesia.*Medical Acupuncture Journal.*14(3):41-44.

**Zak, P.J., R. Kurzban, Borja, K., Matzner and W.T., (2005):** Oxytocin is associated with human trustworthiness.*Horm Behav.* 48(5): 522-527.

**Zanardo, S. Nicolussi, F. Favaro, D. Faggian, M. Plebani, F. Marzari and F. Freato., (2001):** Effect of postpartum anxiety on the colostrum milk  $\beta$ -endorphin concentrations of breastfeeding mothers. J Obstet Gynecol.21:130-134.

**Zhong JY and Cui HS., (2005):** Review of clinical researches on acupuncture Hegu (LI 4) and Sanyinjiao (SP 6) during labour Journal of Yunnan College of Traditional Chinese Medicine.28 (2):64–66.

**Zhu J, Wang MQ, Zhang., (2005):** On the prohibition of acupuncture of Hegu (LI 4) and Sanyinjiao (SP 6) during pregnancy, Acupuncture Research.30 (3):187–190.