

References

- 1-AbdRabbo, (1994):** SA. Stepwise uterine devascularization: a novel technique for management of uncontrolled postpartum hemorrhage with preservation of the uterus. ; 171(3):694-700.
- 2-AbouZahr C, (1998):** Ante partum and postpartum hemorrhage. Ln: Murray CJ, Lopez AD, eds. Health Dimensions of Sex and Reproduction. Boston, Mass: Harvard University Press; pp172-81.
- 3-Accurdy et al, (1993):** Activity of circular muscle of rat uterus at different times in pregnancy .Am J Physiol, 246:c21623.
- 4- Adekamni OA, Purmessur S, Edwards G, (2001):** Intrauterine misoprostol for the treatment of severe recurrent atonic secondary postpartum hemorrhage. Br J Obstet Gynecol; 108: 541-2.
- 5- Akhter S, Begum MR, Kabir Z, Rashid M, Laila TR, Zabeen F,(2003):** Use of a condom to control massive postpartum hemorrhage.MedGenMed Ob/Gyn Women health;(48):38.
- 6- Akhter S, Serum MR, Kabir J, (2005):** Condom hydrostatic tamponade for massive postpartum hemorrhage. Int J Gynecol Obstet; 90: 134-5.
- 7-Alderson et al, (2000):** The "tamponade test" in the management of massive postpartum hemorrhage. Obstet Gynecol; 101: 767-72.
- 8-Allam and B-Lynch, (2005):** Haemorrhagic shock B-Lynch and other uterine compression suture techniques.Int J Gynecol Obstet; 89:236.
- 9-American College of Obstetricians and Gynecologists, (1994):** ACOG Technical Bulletin No.196.Washington, DC, American College of Obstetricians and Gynaecologists.
- 10-American College of Obstetricians and Gynecologists ACOG Educational Bulletin, (1997):** Haemorrhagic shock Number 235. Washington, DC: ACOG, Int j Gynaecol Obstet; 57:219-26.

- 11- Arias F, (1993):** Postpartum complications in practical guide to high risk pregnancy and delivery. Mosby year Book Inc. 2nd ed. Eng. pp 433-51.
- 12-Bakri et al, (2001):** Tamponade-balloon for obstetrical bleeding. Int. J. Gynecol. Obstet. 2001; 74: 139-42.
- 13-Bang et al, (2004):** Maternal morbidity during labour and the puerperium in rural homes and the need for medical attention: a prospective observational study in Gadchirli, India.Br J Obstet Gynecol; 111:231-8.
- 14-Baskett TF, (1999):** Complications of the third stage of labour. In: Essential Management of Obstetrical Emergencies. 3rd ed. Bristol, England: Clinical Press; 1999:196-201.
- 15-Beisher and Mackay, (1996):** Complications in practical guide to high risk pregnancy and delivery. Mosby year Book Inc. 2nd ed. Eng. pp 433-51.
- 16-Bias et al, (2004):** Postpartum haemorrhage: definitions, medical and surgical management. A time for change. Br Med Bull; 67: 205-17.
- 17-Bjornerem A, Acharya A, Oian P. Maltau JM, (2002):** Postpartum hemorrhage prophylaxis and treatment in Norway. Tidsskr Nor Laegeforen; 122: 2536-7 (English abstr).
- 18-Blanchette, (1977):** Arterial embolisation for persistent primal postpartum hemorrhage: Before or after hysteroscopy? Br J. Obstet Gynecol; 111: 880-4.
- 19-B-Lynch C, Coker A, Lawal AH, Abu J, Cowen MJ, (1997):** The B-Lynch surgical technique for the control of massive postpartum haemorrhage: an alternative to hysterectomy? Five cases reported. Br J Obstet Gynecol. Mar 1997; 104(3):372
- 20-Bloom Al, Verstandig A, Gielchinsky Y, Nadjari M, Elchalal U , (2004):** Arterial embolisation for persistent primal postpartum hemorrhage: Before or after hysteroscopy? Br J. Obstet Gynecol; 111: 880-4.
- 21-Bollert et al, (2004):** Obstetric emergencies. BMJ; 318:1342-5.
- 22-Brandt M, (1933):** Mechanism and management of the third stage of labor. Coted from

- Best Pract Am J Obstet Gynecol 1933, 25:622.
- 23-Bukowski R. Silver H, (2000):** Postpartum hemorrhage. In: Evans AT, Niswander KR, eds. Manual of Obstetrics. 3rd ed. Philadelphia: Lippincott Williams and Wilkins; pp 482-90.
- 24-Caliskan E. Meydanli M, Dilbaz B, Aykan B, Sonmezer M, Hoberal A,(2002):**
Is rectal misoprostol really effective in the treatment of third stage of labor?
A randomized controlled trial .Am J Obstet Gynecol; 187:1038-45.
- 25-Chamberlain G, Steer P, (1999):** ABC of labour care: Obstetric emergencies. BMJ; 318:1342-5.
- 26-Chong YS. Su, LL Arulkumaran S, (2004):** Current strategies for the prevention of postpartum hemorrhage in the third stage of labor. Curr Opin Obstet Gynecol; 16: 143-50.
- 27-Choo WL, Chua S. Chong YS, (2000):** Correlation of change in uterine activity to blood loss in the third stage of labor. Gynaecol Obstet Invest; 46: 178-80.
- 28- Choo et al, (1998):** Correlation of change in uterine activity to blood loss in the third stage of labour. Gynaecol Obstet Invest; 46; 178-80.
- 29-Clark SL, Hankins GDV, Dudley DA, Diddy GA, Porter TF, (1995):** Amniotic fluid embolism: Analysis of the national registry. Am J Obstet Gynecol; 172: 1159-63.
- 30-Combs CA, Murphy EL, Laros RK, (1991):** Factors associated with postpartum haemorrhage with vaginal birth; 77;69-74.
- 31-Condous GS, Arulkumaran S, Symonds I, Chapman R. Sinha A, Razvi K ,(2003):**
The "tamponade test" in the management of massive postpartum hemorrhage. Obstet Gynecol; 101: 767-72.
- 32-Condous GS, Arulkumaran S, (2003):** Medical and conservative surgical management of postpartum hemorrhage. J Obstet Gynecol Can; 25: 931-6.

- 33-Cunningham FG, Gant NF, Leveno KJ, Glistrap LC, Hauth JC, Wenstrom KD, (2001):** Maternal adaptation to pregnancy. In: Williams Obstetrics 21st ed. New York, MY: McGraw-Hill; pp167-75.
- 34-Cunningham FG, Gant NF, Leveno KJ, Glistrap LC, Hauth JC, Wenstrom KD, (2005):** Obstetrical hemorrhage. In: Williams Obstetrics. 22nd ed. New York, NY: McGraw-Hill; i pp 823-35.
- 35-Dabelea V et al, (2007):** Management recommendations for postpartum haemorrhage. Arch Postnatal Med; 10:1-4.
- 36-Darryl, (2003):** MIRANZCOG consultant obstetrician and gynecologist surgical techniques Obstet Gynaecol; 391-395
- 37-Dildy GA, (2002):** Postpartum hemorrhage: New management options. Clin Obstet Gynecol 45: 330-44.
- 38-Donald I, (1979):** Practical Obstetric Problem 4th ed., London Lioyde-Lude. Obstet Gynecol; 187:1038-45.
- 39-Doumouchtsis SK, Papageorghiou AT, Arulkumaran S Aug, (2009):** Systematic review of conservative management of postpartum hemorrhage: what to do when medical treatment fails. Obstet Gynecol Surv. ; 62(8):540-7.
- 40-Duff et al, (1984):** Management of primary postpartum haemorrhage. British Journal of Obstetrics and Gynaecology; 104:275–7.
- 41-Egypt MOHP, (2003):** National Maternal Mortality Survey: 2001-2003. Preliminary report: Findings and conclusions. Ministry of Health and Population, Cairo.49:267-75.
- 42-Elbourne DR, Prendiville WJ, Carroli G, Wood J, McDonald S, (2002):** Prophylactic use of oxytocin in the third stage of labour. Cochrane Database Syst Rev.; CD001808
- 43-Elbourne et al, (1988):** Choice of oxytocic preparation for routine use in the management of third stage of labor. Br J Obstet Gynecol; 95: 17-30.
- 44-El Refaey H, Templeton A, (1995):** Induction of abortion in the second trimester by a

combination of misoprostol and mifepristone: A randomized comparison between two misoprostol regimens. Hum Reprod; 10: 475-8:

45-El-Refaey H, O'Brien P, Morafa W, Walder J, Rodeck C, (1996): Use of oral misoprostol in the prevention of postpartum haemorrhage. BJOG 1997; 104: 336-39.

46-El-Refaey H, Rodeck C, (2003): Postpartum hemorrhage: definitions, medical and surgical management. A time for change. Br Med Bull; 67: 205-17.

47-Figo, (2004): Management of third stage of labour. BMJ; 297:1259-300.

48-Fliegner, (1966): Postpartum haemorrhage: a continuing problem. British journal of Obstetrics and Gynaecology; 94: 67-71.

49-Frenzel D et al, (2005): The use of the tamponade test to stop massive obstetric haemorrhage in placenta accrete. Br J Obstet Gynecol; 112:676-7.

50-Friedman EA. Primigravid labo, (1955): A graphic statistical analysis. Cited from International Journal Obstet Gynaecol. Dec; 6(6):567-89.

51-Gartner, (1998): Management of primary postpartum haemorrhage. British Journal of Obstetrics and Gynaecology; 104: 275-7.

52-Glistrap LC, Hauth JC, Hankins GDV, Patterson AR, (1987): Effect of type of anesthesia on blood loss at caesarean section Obstet Gynecol; 69: 328.

53-Guideline for the management of massive obstetric haemorrhage CJR september (2008) review September 2010.version 1.0.

54-Gulmezoglu AM, (2000): Prostaglandins for prevention of postpartum hemorrhage. Cochrane Database Syst Rev; (2): CD000494.

55-Hayman RG, Arulkumaran S, Steer PJ, (2002): Uterine compression sutures: Surgical management of postpartum hemorrhage Obstet Gynecol; 99: 502-6.

56-Hill K, AbouZahr C, Wardlaw T, (2001): Estimates of maternal mortality. Bull World Health Organ; 79: 182-93.

57-Holtsema et al, (2004): The b-lynch technique for postpartum haemorrhage: an option

for every gynaecologist. Report Biol; 115:39-42

58-Hong et al, (2004): Preventing postpartum hemorrhage in low-resource setting Ink J Gynecol Obstet; 77: 267-75.

59-Hussain M, Jabeen T, Liaquat N, Nooranik, Bhuttas, (2004): Acute puerperal uterine inversion Call physicians Surg Pak; 14: 215-7.

60-Hytten and chamberlain, (1991): Clinical physiology in obstetric. Blackwell scientific publication oxford. 12:303-344.

61-Ijaija et al, (2003): Change in uterine activity to blood loss in the third stage of labour. Gynecol Obstet Invest; 46: 1 78-80.

62-Jackson KW Jr, Albert JR, Schemmer GK, Elliot M, Humphrey A, Taylor J,(2001):A randomized controlled trial comparing oxygen administration before and after placental delivery in the prevention postpartum hemorrhage Am J Obstet Gynecol; 185:873-7.

63-Jacobsen, (1990): Disseminated intravascular Coagulation. Best Pract Res Clin Obstet Gynecol; 15: 623-44.

64-Johnson et al, (2003): Peripartum hysterectomy Obstet Gynecol; 65: 3 65-70.

65-Katesmark et al, (1994): Estimates of maternal mortality for. Bull World Health Organ; 79:182-93.

66-Kartmark, (1994): Uterine packing in management of postpartum haemorrhage lancet; 349; 698.

67-keirse et al, (1995): The chocrane collaboration issue 2. Oxford update software: review no.2974.

68-kellner et al, (1984): Parental behaviour after perinatal death. Obstet Gynecol; 63:809-14.

69-Kundodyiwa Toy, Majoko F, Rusakaniko S, (2001): Misoprostol versus oxytocin in the third stage of labor. Int J Gynecol Obstet; 75: 235-41.

- 70-Lalonde A, Daviss BA, Acosta A, Herschderfer K, (2006):** Postpartum hemorrhage today. *Int J Gynecol Obstet*; 94: 243-53.
- 71-Ledee N, Ville Y, Musset D, Mercier F, Frydman A, Fernandez H, (2001):** Management intractable obstetric hemorrhage: An audit study on 61 cases. *Eur J Obs Gynecol Reprod Biol*; 94: 189-96.
- 72-Letsky EA, (2001):** Disseminated intravascular Coagulation. *Best Pract Res Clin Obstet Gynecol*; 15: 623-44.
- 73-Lokugamage AU, Sullivan KR, Niculescu I, et al, (2001):** A randomized study comparing rectally administered misoprostol versus syntometrine combined with an oxytocin infusion for the cessation of primary post partum hemorrhage. *Acta Obstet Gynecol Scand. Sep*; 80(9):835-9
- 74-Lopez et al, (1995):** Activation of stimulatory pathway. *Adv Med Biol*; 395-435-9.
- 75-Lucas W, (1980):** Post partum hemorrhage. *Clin Obstet Gynecol*; 23:637.
- 76-MacDonald, (1985):** Third stage of labor. Massive obstetric haemorrhage *Obstet Gynecol Clin North Am*; 32:323-32.
- 77-MacDonald S, Premidivill W, Blair E, (1993):** Randomized controlled trial of oxytocin alone versus oxytocin and ergometrine in acute management of third stage of labor. *BMJ*; 307: 1 167-71.
- 78-Magann EF, Evans S, Hutchinson M et al, (2005):** Postpartum hemorrhage after vaginal birth: an analysis of risk factors. *South Med J. Apr*; 98(4):419-22.
- 79-Maier RC, (1993):** Control of postpartum hemorrhage with uterine packing. *Am J Obstet Gynecol*; 169: 317-21; discussion 321-3.
- 80-Mategrano VA, Gabay MP, (2001):** Misoprostol in the prevention of postpartum haemorrhage. *Ann Pharmacother Dec*; 35: 1648- 52.
- 81-McCormick ML, Sanghvi HCG, Kinzie B, Melntosh N, (2002):** Preventing postpartum hemorrhage in low-resource setting *Ink J Gynecol Obstet*; 77: 267-75.

- 82-Miller S, Lester F, Hensleigh P, (2004):** Prevention and treatment of postpartum haemorrhage: New advances for low-resource settings. *J. Midwifery Women's Health*; 49: 283-92.
- 83-Myerscough P, (1982):** Uterine action in Labor. In: Munro Kerr's Operative Obstetrics. 10th ed. Baillière Tindall (pub), London, pp 36-46.
- 84-Naqvi S, Makhdoom T, (2004):** Conservative management of primary postpartum hemorrhage. *J Coll Physicians Surg Pak* 14: 296-7.
- 85-Narsal et al, (1999):** Management of primary postpartum hemorrhage. *J Coll Physicians Surg Pak* 14: 296-7.
- 86-Nordstrom, Fogelstan K, Friedman G, Larsson A, Rydhstroem H, (1997):** Routine oxytocin in the third labour. *Br J Obstet Gynecol*; 104: 781-786.
- 87-Nothnagle M, Taylor SA, Smith K, (2003):** Should active management of the third stage of labor be routine? *Am Fam Physician*; 67: 2119-20.
- 88-Nwagha UI, Okaro JM, Nwagha TU, (2005):** Intraoperative uterine packing with mops: an effective, but under utilized method of controlling postpartum hemorrhage experience from South Eastern Nigeria. *Niger J. Med*; 14:279-82.
- 89-Oboro V, Tabowei T, (2003):** A randomized controlled trial of misoprostol versus oxytocin in the active management of the third stage of labour. *Obstet Gynecol*; 23: 13-6.
- 90-O'Brien P, El-Refaey H, Gordon A, Geary M, Rodeck CH, (1998):** Rectally administered misoprostol for the treatment of postpartum hemorrhage unresponsive to oxytocin and ergometrine: A descriptive study. *Obstet Gynecol*; 92:212-4.
- 91-Odent, Michel, (1992):** The Nature of Birth and Breastfeeding. Sydney: ACE Graphics.
- 92-Ojala K, Perala J, Kariniemi J, Ranta P, Raudaskoski T, Tekay A, (2005):** Arterial embolization and prophylactic catheterization for the treatment for severe obstetric hemorrhage. *Acta Obstet Gynecol Scand*; 84:1075-80.
- 93-Onds I et al, (2006):** Treatment for primary postpartum haemorrhage. *Cochrane Database*

Syst Rev; 1:CD003249.

- 94-Pal et al, (2003):** Active management of the third stage of labour is routine? Am Fam Physician; 67: 2119-20
- 95-Percival, (1980):** The placenta. In: Percival ed. Holland and brews manual of obstetrics 14ed. London. J and achurill ltd pp 21-4
- 96-Porcu et al, (2005):** Uterus and bladder necrosis after uterine artery embolisation for postpartum hemorrhage. Br J Obstet Gynecol; 112:122-3.
- 97-Philip et al, (2004):** Increasing the flow rate of i.v. fluid administration. Br Anaesth; 75: 361
- 98-Prata N, Hamza S, Gypson R, Nada K, Vahidnia F, Potts M, (2005):** Misoprostol and active management of the third stage of labour. International Journal of Gynaecology and Obstetrics; 94: 149-55.
- 99-Prendiville et al, (2003):** Active versus expectant management in the third stage of labour .Cochrane review issue 4. Ed John Wiley 84:1271-82.
- 100-Prendiville et al, (1998):** Active versus expectant management third stage of labour. Pregnancy and child birth module. Cochrane Database Syst Rev. 111:495-8
- 101-Prendiville WJ, Harding JE, Elbourne DR, Stirrat GM, (1989):** The Bristol third stage trial: active versus physiological management of third stage of labour. BMJ. Nov 19 1988; 297(6659):1295-300
- 102-Prendiville WJ, Elbourne D, McDonald S, (2001):** Active versus expectant management in the third stage of labour. Cochrane Database Syst Rev. CD000007.
- 103-Prendiville WJ, Elbourne D, McDonald S, (2002):** Active versus expectant management in the third stage of labour (Cochrane Review). In: The Cochrane Library, issue 2. Oxford, UK: Update Software.
- 104-Ramanathan G. Arulkumaran S, (2006):** Postpartum haemorrhage. J Obstet Gynecol Can; 28: 967-73.

- 105-Rizvi F, Mackey R, Barrett T, Mckenna P, Geary M, (2004):** Successful reduction of massive: postpartum hemorrhage by use of guidelines and staff education. Br J Obstet Gynecol; 111:495-8.
- 106-Rogers J, Wood J, McCandlish R, (1998):** Active versus expectant management of third stage of labour: The Hinchings brooke randomized controlled trial lancet; 351: 693-9.
- 107-Romero et al, (1988):** Scottish programme for clinical effectiveness in reproductive health.
- 108-Royal College of Obstetricians and Gynaecologists, (2004):** Placenta Praevia: Diagnosis and management. Available at: <http://www.rcog.org.uk/guidelines.asp?PageID=106&GuidelineID=17>. London, England.
- 109-Scottish programme, (2005):** For clinical effectiveness in reproductive health ,Obstet and Gynecol,256-62.
- 110-Sergent F, Resch B, Verspyck E, Rachet B, Curvier E, Marpeau L, (2004):** Intractable postpartum hemorrhage: Where is the place of vascular ligations, emergency peripartum hysterectomy or arterial embolization? Gynecol Obstet Fertile; 32:320-9.
- 111-Seror J, Allouche C, Elhaik S, (2005):** Use of Sengstaken Blakemore tube in massive postpartum hemorrhage: A series of 17 cases Acta Obstet Gynecol Scand; 84:660-4.
- 112-Shriner et al, (2005):** Changes in transcripts encoding calcium channel subunits of rat myometrium during pregnancy Am J Physiol, 269 Cl008 17.
- 113-Sideman DS, Armon Y, Roll D, Stevenson DK, Gale R, (1988):** Grand multiparty: An Obstetric or neonatal risk factor. Am J Obstet Gynecol; 158:1034-39.
- 114-Sleep, (1993):** Physiology and management of the third stage of labour. In: Bennett VR, Brown LK, eds. Myles' Textbook for Midwives. 12th ed. London, UK: Churchill Livingstone; pp 216-29

- 115-Society of Obstetricians and Gynaecologists of Canada, (2000):** Advances in labour and risk management (ALARM) course manual 5th edition volume 1.
- 116-Society of Obstetricians and Gynaecologist of Canada, (2002):** Advances in labour and risk management (ALARM) course manual 9th ed.
- 117-Stainsby D, MacLennan S, Hamilton PJ, (2000):** Management of Missive blood loss: A template guideline. Br J Anaesth; 85: 487-91.
- 118-Stoneham MD, (1995):** An evaluation of methods of increasing the flow rate of i.v. fluid administration. Br Anaesth; 75: 361-5.
- 119-Strand RT, Da Silva F, Jangsten E, Bergstrom S ,(2005):** Postpartum hemorrhage: a prospective, comparative study in Angola using a new disposable device for oxytocin administration Acta Obstet Gynecol Scand; 84: 260-5.
- 120-Tamizian O, Arulkumaran S, (2002):** The surgical management of postpartum hemorrhage. Best Pract Res Clin Obstet Gynecol; 16: 81-98.
- 121-Tanaka YD, Shigemitsus M, Ichikawa Y, Sohda S, Yoshikawa H, Itai Y, (2004):** Postpartum MRI diagnosis of retard placenta accreta. Eur Radiol; 14: 945-52.
- 122-Vartner M, (1991):** Postpartum hemorrhage. Crit Care Clin; 7: 883-97.
- 123-Visentin L, Alemanno MG, Arduino S, Bellino R, Tessarolo M, Wierdis T, Lanza A, (1996):** Management of patients with intrauterine fetal death. Clin Exp Obstet Gynecol; 23:263-7.
- 124-WHO, (2004):** Maternal mortality in 2000: Estimates developed by WHO, UNICEF and UNFPA. Geneva.