
Table (1) Clinical character of patient

Age(y)	Gestational age(w)	Mode of o	deliv	ery
(SD±)	(SD±)			
30±7.3	38.9±1.2	NVD	٨	٦٦,٧
		%		
		Elective CD	3	Yo,. %
		Emergency CD	1	8.3 %
		Total	12	100

Chart (1) modes of dlivery among the study group




■NVD ■Elective CD □Emergency CD

Table (2) Success rate and failure rate among the study group

Study group	number	%
Success	11	91,7
Failure	١	۸,۳
Total	١٢	١٠٠,٠

Chart (3) sucess and failure rates among the study group



■Success ■Failed

Table (*) Difference between haemoglobin level before and 24 hr after procedure among successful patient.

НВ	± SD	± SD of	SE	Paired"t"	P
Time		difference			
Before	۹,۸±٠,٩				
		۰,٥ <u>+</u> ٠,٢	٠,٠٦	٨,٥٦	< 00001
After	۹,٣±٠,٩				

There was significant diffirence regarding haemoglobin level before and 24hr after procedure among successful patient.

Chart (6) means of HB % BP before and after treatment




Table (4) Difference between pulse before and 24 hr after procedure among successful patient.

Pulse Time	±SD	SD of the ±difference	SE	Paired "t"	Р
Before	±7,9 97,1	٦,٥±٩,٤	۲,٧	۲,۳۹	<0.05
ृfter़	(1,1	(,o± (, ē	1, 1	1,11	<0.05
A	±Υ, ξ Λο, ٦				

There was significant diffirence regarding difference between pulse before and 24hr after procedure among successful patient.

Chart (7) means ofpulse BP before and after treatment




Table (5) Systolic blood pressure before and after 24 hrs

Systolic B.P.	±SD	± SD of the difference	SE	Paired "t"	P
Before	۱۰۲,۹ <u>+</u> ۸,۹				
After	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	۱٠,٤±٨,٤	۲,٤	٤,٢٨	<0.001

There was significant diffirence regarding systolic blood pressure before and 24hr after procedure among successful patient.

Chart (4) means of systolic BP before and after treatment




Table (6) Diffirence between diastolic blood pressure before and 24hrs after procedure among successful patient.

Diastolic	±SD	± SD of the	SE	Paired	P
B.P.		difference		"t"	
Time					
After	٦٩,٢±٧,٩				
		$\Lambda, \Upsilon \pm \circ, \Lambda$	١,٦٨	٤,٩٥	< 0.001
Before	٦٠,٨±٦,٧				

 Table (7) Blood loss (L) among the study group.

Variable	±SD	
Blood loss(liter)	۱,۸٣±٠,٩٤	