

REFERENCES

- Abell , M.R. (1985) :** Endometrial biopsy : Normal and abnormal diagnostic characteristics . In: Gold, J.J.(ed.) : Gynecologic endocrinology, 2nd ed. Harper & Row .Virginia, Maryland, London, NewYork. P.165-169.
- Aiman,J. and Smentek,C. (1985):** Premature ovarian failure Obstet. Gynecol.; 66(1):9-14.
- Ames, BN. ; Shigenaga, MK. And Gold, LS. (1993):** DNA lesions, inducible DNA repair, and cell division: three key factors in mutagenesis and carcinogenesis. Environ. Health Perspect. 101 Suppl; 5:35-44.
- Andolf ,E. ; Dahlander ,K. and Aspenberg, P. (1993):** Ultrasonic thickness of the endometrium correlated to body weight in asymptomatic postmenopausal women. Obstet. Gynecol. ; 82(6):936-40.
- Armstrong,BK. (1988):** Oestrogen therapy after the menopause-boon or bane? Med. J. Aust. ; 148(5):213-4.
- Baber, P. (1991):** Premature menopause. In: Frank ,Chervenak. ;Seang,T. and John, S. (eds.):Progress in Obstet. & Gynaecol, 2nd ed. Chirchil Living Stone,London .P. 209-220.

- Baber, P. and Salteor (1987):** Diagnosis of premature menopause In: Vanda, R.;Lops ZeRosenwaks; Fred Benjamin and Martin,L.Stone(eds.): Gynecology: Principles & practice, 4th eds., Macmillan Publishing Company, New York. p. 165-186.
- Berker, B.; Erkmen, M. and Sevim,D. (2005):** Relationship between body mass index and endometrial thickness in postmenopausal women. Clin. Obstet. Gynecol.;11(3): 34-44.
- Bjorntorp, P. (1994):** Fatty acids, hyperinsulinemia, and insulin resistance: which comes first? J. Curr. Opin. Lipidol. ;5(3):166-74.
- Bokhman, JV.(1983):** Two pathogenetic types of endometrial carcinoma. Gynecol. Oncol. ; 15(1):10-17.
- Bradley, LD. ; Falcone, T. and Magen, AB. (2000):** Radiographic imaging techniques for the diagnosis of abnormal uterine bleeding. Clin. Obstet. Gynecol. North Am. ;27(2):245-76.
- Bray GA(2005):** Epidemiology, risk and pathogenesis of obesity. J. Science :71, 2-7.
- Campbell, S. and Thoms, A. (1977):** Ultrasound measurement of the fetal head to abdomen circumference ratio in the assessment of growth retardation. Br. J. Obstet. Gynecol. ;84(3):165-74.

- Chambers, CB. and Unis, JS. (1986):** Ultrasonographic evidence of uterine malignancy in the postmenopausal uterus. *Am. J. Obstet. Gynecol.* ;154 (6):1194-9.
- Cohen, CJ. ; Gusberg, SB. and Koffler, D. (1974):** Histologic screening for endometrial cancer. *Gynecol. Oncol.*;2(2-3):279-86.
- Colleoni M, Gelber RD and Price KN (2005):** Relation between chemotherapy dose, estrogen receptor expression and body-mass index. *Lancet* , **366**:1108–10.
- Dallenbach-Hellweg, G. ; Dallenbach, C. and Lang, G.(1988):** Histogenesis and morphology of mucoepidermoid adenocarcinoma of the corpus endometrium. *Pathol.* ; 9(6):351-7.
- Dandolu,V.; Singh, R.; Lidicker,J. and et al.(2010):** BMI and uterine size: is there any relationship ? *Int. J. Gynecol. Pathol.*, 27(6): 568-71.
- Deichert, U. ; Hackelöer, BJ. and Daume, E. (1986):** The sonographic and endocrinologic evaluation of the endometrium in the luteal phase. *Hum. Reprod.* ;1(4):219-22.
- Deligdisch, L .(1993):** Effects of hormone therapy on the endometrium *Pathol.* ; 6:94-106.

- Deutinger, J.; Rudelstorfer, R. and Bernaschek, G. (1988):** Vaginosonographic velocimetry of both main uterine arteries by visual vessel recognition and pulsed Doppler method during pregnancy. Am. J. Obstet. Gynecol. ; 159(5):1072-6.
- Dodson, MG. and Deter, RL. (1990):** Definition of anatomical planes for use in transvaginal sonography. J. Clin. Ultrasound. ; 18(4):239-42.
- Donald, I. ; Macvicar, J. and Brown, TG. (1958):** Investigation of abdominal masses by pulsed ultrasound. Lancet. ; 1(7032):1188-95.
- Douchi, T. ; Yoshinaga, M. ; Katanozka, M. and et al. (1998):** relationship between body mass index and transvaginal ultrasonographic endometrial thickness in postmenopausal women . Acta obstet. Gynecol. Scand. ; 77 :905-908 .
- Eknoyan and Garabed (2008):** The average man and indices of obesity. J. Nephrol. 23 (1): 47–51.
- Emons, G. ; Fleckenstein, G. ; Hinney, B. and et al. (2000):** Hormonal interactions in endometrial cancer. Endocr. Relat. Cancer. ; 7(4):227-42.
- Fleischer, AC. ; Kalemeris, GC. ; Machin, JE. And et al. (1986):** Sonographic depiction of normal and abnormal endometrium with histopathologic correlation. J. Ultrasound Med. ; 5(8):445-52.

Fleischer, AC. ;Dudley, BS. ;Entman, SS. And et al. (1987): Myometrial invasion by endometrial carcinoma: sonographic assessment. Radiology. ; 162(2):307-10.

Fleischer, AC. ; Jordon AN. ; Entmans, S. and et al. (1990): Transvaginal sonography of the endometrium. In: Fleischer, A.C. Romero, R. ; Manning ,FA.; Jeanty, PH. and James, AE.(eds.):principle and practice of ultrasonography in Obstet. and Gynecol. , 4th Ed . Prince Hall International Inc. New York. P.145:149.

Fleischer, AC; Gordon ,NA. and Stephen (1991):Transvaginal sonography of the endometrial current and potential clinical application . In: Fleischer, A.C. Romero, R. ; Manning ,FA.; Jeanty, PH. and James, AE.(eds.):principle and practice of ultrasonography in Obstet. and Gynecol. , 4th Ed . Prince Hall International Inc. New York. p .583.

Franks, S. (2006): Genetic and environmental origins of obesity relevant to reproduction .J. Reprod. Biomed. ;12:526 -531.

Freeman,EW.; sammel, MD. ; Lin (2007) : Symptoms associated with menopausal transition and reproductive hormones in midlife. Am .J. Obstet . Gynecol . ; pp.30 – 40

Frystyk, J. ;Vestbo, E. ; Skjaerbaek, C. and et al. (1995): Free insulin-like growth factors in human obesity.Metabolism. ;44(10 Suppl 4):37-44.

Gadzik, J. (2006): "How much should I weight? Quetelet's equation, upper weight limits and BMI prime. Conn. Med. ;70(2):81-8.

Gallagher,D .(1996): How useful is body mass index for comparison of body fatness across age , sex and ethnic groups . Am . J . Epidemiology :143 – 239.

Goldstein, SR.(1990): Incorporating endovaginal ultrasonography into the overall gynecologic examination. Am. J. Obstet. Gynecol. ;162(3):625-32

Goldstein,SR. ;Nachtigall,M. ;Snyder,JR. and et al. (1990): Endometrial assessment by vaginal ultrasonography before endometrial sampling in patients with postmenopausal bleeding. Am. J. Obstet. Gynecol. ;163 (1):119-23.

Gratton, D. ;Harrington,C. ;Holt,SC. and et al. (1991): Normal pelvic anatomy using transvaginal scanning. Obstet. Gynecol. Clin. North Am. ;18(4):693-711.

Hendrickson, MR. and Kempson, RL. (1979): Surgical pathology of the uterine corpus. Major Probl. Pathol. ;12:1-580.

- Hill,HA. and Austin,H. (1996):** Nutrition and endometrial cancer. *Cancer Causes Control.* ;7(1):19-32.
- Hofmeister,FJ. (1974):** Endometrial biopsy: another look. *Am. J. Obstet. Gynecol.* ;15; 118(6):773-7.
- Huang Z, Hankinson SE and Colditz GA,(1997) :** Dual effects of weight and weight gain on breast cancer risk. *J.Am. Med.Ass* ; 278(17):1407-11.
- Jewelewicz R. and Schwartz M (1986):** Premature ovarian failure. *Bullny Acad. Med.* 62 (3):219-36.
- Jones, J. (1988):** Endometrial hyperplasia & carcinoma. In: Jonathan S. Jones JJ, HW, Wentz AC & Burorett LS.(eds): *Novak's Textbook of Gynecology*, 11th Ed. Williams and Wilkins. Baltimore. p.790-797.
- Judd (1984):** Complication of menstruation. In: Alan.H.Decherney, Lauren. N. (eds):*Current Obstet. & Gynecol. Diagnosis*,5thed. McGraw-Hill Medical. p. 570-589.
- Karlsson, B. ;Granberg, S. ;Wikland,M. and et al. (1995):** Transvaginal ultrasonography of the endometrium in women with postmenopausal bleeding - A Nordic multicenter study. *Am. J. Obstet. Gynecol.* ;172(5):1488-94.

- Kemp,G.M. (1987):** Etiology of menopause In:Zev.RosenWakes(ed): Gynecology principles & practice,4th ed. Macmillan Publishing Company, New York .p.484-518.
- Kenneth and Fortire, MD.(1986):** Postmenopausal bleeding and the endometrium. Clin. Obstet. Gynecol. ;29(2):440-5.
- Key,TJ. and Pike,MC. (1988):** The dose-effect relationship between unopposed oestrogens and endometrial mitotic rate: its central role in explaining and predicting endometrial cancer risk. Br. J. Cancer. ;57(2):205-12.
- Keys and Ancel (1972):** Indices of relative weight and obesity. J. Chronic Dis.; 25 (6): 329–43.
- Krotkiewski, M. ;Seidell,JC. and Bjorntorp, P. (1990):** Glucose tolerance and hyperinsulinemia in obese women: Role of adipose tissue distribution, muscle fibre characteristics and androgens. J. Intern. Med. ; 228 (4): 385–392.
- Kurjak,A. ;Zalud,I. ;Jurkovic,D. and et al. (1989):** Transvaginal color Doppler for the assessment of pelvic circulation. Acta Obstet. Gynecol. Scand. 68(2):131-5.
- Kurman ,RJ. and Mazur, MT. (1994):** Benign diseases of the endometrium. In: Kurman, RJ. (ed.): Blaustein's Pathology of the Female Genital Tract,4th ed. Springer- Verlag, New York.p. 421-499.

- Kurman, R.J. and Norris, H.J. (1994):** Endometrial Hyperplasia and Related Cellular Changes. In: Kurman,RJ. (ed.): Blaustein's Pathology of the Female Genital Tract, 4th Ed. New York: Springer- Verlag .p. 425: 435.
- Kurman, R.J. ;Zaino, R.J. and Norris, H.J. (1994):** Endometrial carcinoma. In: Kurman, R.J.(Ed.): Blaustein's Pathology of the Female Genital Tract,4th ed. New York: Springer- Verlag. P. 439:470 .
- Lenton,EA. ;de Kretser,DM. and Woodward,AJ. (1991):** Inhibin concentrations throughout the menstrual cycles of normal, infertile, and older women compared with those during spontaneous conception cycles. J. Clin. Endocrinol. Metab. ;73(6):1180-90.
- Longcope,C. ;Hunter,R. and Franz,C. (1998):** Steroid secretion by the postmenopausal ovary. Am. J.Obstet. Gynecol. ;138(5):564-8.
- Maxson,WS. and Wentz,AC. (1983):** The gonadotropin resistant ovary syndrome. J. Rev. Med. Interne. ; 4(4):353-535.
- McKinley, SM. ; Nancy, LB. and McKiplay, JB. (1985):** Smoking and age at postmenopause in women. Ann. Of Internal Medicine ;103: 350-356.
- McTiernan A. (2005) :** Obesity and cancer: the risks, science, and potential management strategies. Oncology ; 19:871–81.

Meeks,GR. ;Woodliff,JM. ;Box, ST. and and et al. (1993): Evaluation of endometrial biopsy using an aspiration curette without cervical tenaculum. South Med. J. ; 86(9):1011-6.

Mei,Z. ; Grummer-Strwan,IM. ;Pietrobelli,A and et al. (2002):

Validity of body mass index compression screening indices for assessment of body fatness . Am . J . Clin . Nutr .;7 : 597 -985.

Mendelson ,EB. and Bohm-Velez, M. (1992): Transvaginal ultrasonography of pelvic neoplasms. Radiol. Clin. North Am. ;30(4):703-34.

Merce,LT. ;Garcia,GL. and Fuente, F. (1991): Doppler ultrasound assessment of endometrial pathology. Acta Obstet. Gynecol. Scand.;70: 525-530.

Merrill, JA. (1981): Management of postmenopausal bleeding. Clin.Obstet. Gynecol. ;24(1):285-99.

Merrill, JA. (1991): The interpretation of endometrial biopsies. Clin. Obstet. Gynecol. ;34 (1): 211-221.

Meyer,WC. ;Malkasian,GD. ;Dockerty,MB. and et al.(1983): Postmenopausal bleeding from atrophic endometrium. Obstet. Gynecol. ;38(5):731-8.

Miller,WR. and Anderson,TJ. (1988): Estrogen, progesterone and the breast. Menopause. Lancet J. :900-902.

- Nakamura,S. ;Douchi,T. ;Oki,T. and et al. (1996):** Relationship between sonographic endometrial thickness and progestin-induced withdrawal bleeding. *Obstet. Gynecol.* ;87 (5 Pt 1):722-5.
- Nasri,MN. and Coast,GJ. (1989):** Correlation of ultrasound findings and endometrial histopathology in postmenopausal women. *Br. J.Obstet. Gynecol.*; 96(11):1333-8.
- Nasri, MN. ;Shepherd,JH. and Setehell,ME. (1991):** The role of vaginal scan in measurement of endometrial thickness in postmenopausal women. *Br. J. Obstet. Gynaecol.*; 98 (5): 470 - 475 .
- Neele,SJM. ;Marchien Van baal,W. ; Van Der Mooren, P. and et al. (2002):**Ultrasound assessment of endometrium in healthy , asymptomatic early postmenopausal women . *Ultrasound in obstet. Gynecol.* ;16 (3) :254- 259 .
- Nichols,DH. and Randall,CL. (1996):** Minor and ambulatory surgery. In: Nichols, DH. and Randall,CL. (Eds.): *vaginal surgery*.4thed. Williams and Wilkins, Baltimore, New York. p. 95-100.
- Novak,ER.(1956):** Postmenopausal endometrial hyperplasia. *Am. J. Obstet. Gynecol.* ;71(6):1312-21.

- Okaunofa, FE. ;Lawal, A. and Bamgbose,JK. (1990):** Features of menopause and menopausal age in Nigerian women. Int. J.Gynecol. Obstet. ;31(4):341-5.
- Otake S, Takeda H and Suzuki Y (2005):** Association of visceral fat accumulation and plasma adiponectin with colorectal adenoma: evidence for participation of insulin resistance. Clin. Cancer , 11: 3642–6.
- Parazzini,F. ;La Vecchia,C. ;Bocciolone,L. and et al. (1992):** The epidemiology of endometrial cancer. Gynecol. Oncol. ;41(1):1-16.
- Pasquali, R. ;Vicennati,V. ;Bertazzo,D. and et al. (1997):** Determinants of sex hormone-binding globulin blood concentrations in premenopausal and postmenopausal women with different estrogen status. Virgilio-Menopause-Health Group. Metabolism. ; 46: 5–9.
- Pasquali, R. ;Gambineri ,A. ; Biscotti ,D. and et al. (2000):** Effect of long-term treatment with metformin added to hypocaloric diet on body composition, fat distribution, and androgen and insulin levels in abdominally obese women with and without the polycystic ovary syndrome. J. Clin. Endocrinol. Metab.;85(8):2767-74.
- Patten,RM. ;Vincent,LM. ;Wolner-Hanssen,P. and et al. (1990):** Pelvic inflammatory disease: Endovaginal sonography with laparoscopic correlation. J. Ultrasound Med. ;9(12):681-9.

- Pratap,K. and Narenda,M. (1987):**Pathology of post menopausal endometrium In: Tindall,VR.(ed):Jeffcopates Principles of gynaecology.4th ed. Butterworths, London. p91-100.
- Romagnolo,C. ;Zaps,B. ;and Meagan,T. (1993):** Histopathological characterization in carcinoma of endometrium. Comparison between biopsy and pathological examination of specimen. Eur. J. Gynecol. Oncol. ; 14 (2): 106-108.
- Romero-Corral, A. ;Somers, VK. ;Sierra-Johnson, J. and et al.(2008):** "Accuracy of body mass index in diagnosing obesity in the adult general population". Intern. J. Obesity ;32 (6): 959–956.
- Rubin,SC.(1987):** Postmenopausal bleeding: Etiology, Evaluation and Management. J. Med. Clin. North Am. ; 71(1): 59-69.
- Sherer,DM. ;Liberto,L. ;Abramowicz,JS. and and et al. (1991):** Endovaginal sonographic features associated with isolated torsion of the fallopian tube. J. Ultras. Med.; 10(2): 107-9.
- Sherman,ME. ;Bitterman,P. ;Rosenshein,NB. And et al.(1992):**Uterine serous carcinoma. A morphologically diverse neoplasm with unifying clinicopathologic features. Am. J. Surg. Pathol.;16(6):600-10.
- Sheth, S. ; Hamper,LJVL. and Kunnan, RJ. (1993):** Thickened endometrium in the postmenopausal women. Sonographic pathologic correlation. Radiology.;187(1):135-9.

- Siiteri,PK.(1987):** Adipose tissue as a source of hormones. Am. J. Clin. Nutr.; 45(1 Suppl):277-82.
- Siladitya,B. (1999):** Infertility, In: Keith Edmonds(ed):Dewhurst's textbook of Obstet. and Gynecol . 7th ed. BlackWeel, London. p.441-461.
- Silva,EG. and Jenkins,R.(1990) :** Serous carcinoma in endometrial polyps. Pathol. ;3(2):120-8.
- Smith,KE. and Judd,H.L. (1994):** Menopause and postmenopause. In: DeCherney,AH.;Pernoll,ML.(eds):Current Obstetric and Gynecologic Diagnosis and Treatment.8th ed. Appleton & Lange, USA. p.1030-1050.
- Speroff,L.(1999):** Postmenopausal hormone therapy and the risk of breast cancer, Maturitas. 16;32(3):123-9.
- Stovall,TG. ;Solomon ,SK. and Ling, FW. (1989):** Endometrial sampling prior to hysterectomy. Obstet. Gynecol.; 73(3 Pt 1):405-9.
- Sullivan, J. ;Kopher, RA. and Nagel ,TC. (1994):** Advanced in treatment of menopause and osteoprosis. Br. J. Obstet. & Gynecol.; 101: 170-174.

Swartz,D. and Buttler, W. (1992): Normal and abnormal uterine bleeding.
In: Tompson, J.and Rock,J. (eds):TeLinde's operative Gynecology.7th
ed. JB Lippincott, Philadelphia. p. 297-316.

Timmermans,A. ;Opmeer,BC. ;Ersema,SV. And et al. (2007) : Patient`s
references in the evaluation of postmenopausal bleeding. An. Intern.
J.Obestet. Gynecol . ;114 (9):1146 – 1149 .

Timor-Tritsch,IE. ;Bar-Yam,Y. ;Elgali,S. and et al. (1988): The technique of
transvaginal sonography with the use of a 6.5 MHz probe. Am. J.
Obstet. Gynecol.; 158(5):1019-24.

Tsuda,H. ;Kawabata,M. ;Kawabata,K. and et al. (1997): Improvement of
diagnostic accuracy of transvaginal ultrasound for identification of
endometrial malignancies by using cut off level of endometrial
thickness based on length of time since menopause. Gynecol. Oncol.;
64(1):35-7.

Van Baak, MA. and Borghouts, LB. (2000): Relationships with physical
activity. Nutr. Rev. ;58(3 Pt 2):S16-8.

Van den Bosch,T. ;Vandendael,A. ;Van Schoubroeck,D. and et al. (1996):
Age, weight, body mass index and endometrial thickness in
postmenopausal women. Acta Obstet. Gynecol. Scand.; 75(2):181-2.

- Verkasalo, PK. ;Thomas, HV. ;Appleby, PN. And et al. (2001):** Circulating levels of sex hormones and their relation to risk factors for breast cancer: a cross-sectional study in 1092 pre- and postmenopausal women (United Kingdom). *Cancer Causes Control*; 12: 47–59.
- Vilaro,MM. ;Rifkin,MD. ;Pennell,RG. And et al.(1987):** Endovaginal ultrasound. A technique for evaluation of nonfollicular pelvic masses. *J.Ultrasound Med.* ;6(12):697-701.
- Wageman,HL. (1988):** Postmenopausal bleeding. *Bailliere's Clin. Obstet. Gynecol.*; 2(2): 355-362 .
- Walker,ARF. ;Walker,PE. ;Ncongware,J. and et al. (1984):**Age at menopause in black women in South Africa. *Br. J. Of Obstet. Gynecol.*; 91: 797-801.
- Warming,I. ;Ravn,P. ;Skouby,S. and et al. (2002):** Measument precision and normal range of endometrial thickness in a postmenopausal by transvaginal ultrasound .*Ultrasound in Obestet. Gynecol .* ;20(5): 492 – 495 .
- Whitehead,MI. and Marsh,MS.(1992):** Management of the menopause. *Br. Med. Bull.*;48(2):426-57.
- World Health Organization (2005):**Obesity :preventing and managing the global Epidemic .Report of a WHO consultation on obesity .Geneva,World Health Organization ;894 :1-253.