REFERENCES

- Abdelwahed HM, Saaban HA, and Ahmed MM (1990): Statistical study of cases of abortion in AL-Kasr- Al-Ainin Hospital in the last five years (985-1989). Ciro faculty of Medicine, Cairo university, Egypt Cairo, 3: 98-110.
- Adanu R M, Ntumy MN and Twenboah E (2005): Profile of women with abortion complications in Ghana. Trop Doct 35:139-42.
- Adelberg, A M and Kuller JA (2002): Thrombophillias, and recurrent miscarriage. Obstet Gynecol surv. 57:703-743.
- Altay MM, Yaz H and Haberal AH (2009): The assessment of the gestational sac diameters, crown-rump lengths, progesterone and fetal heart rate measurements at 10th gestational week to predict the spontaneous abortion risk. J Obstet Gynecol Res. 35: 287-290.
- Althuisius SM, Dekker G A and Hummel P (2001): Final results of the cervical incompetence prevention randomized cerclage trial (CIPRCT): therapeutic cerclage with bed rest versus bed rest alone. Am J Obstet Gynecol 185: 1106-1145.
- ACOG; (2001): Management of recurrent early pregnancy loss. Practice Bulletin No. 24.
- ACOG; (2004): Abortion policy. ACOG Statement of policy issued by the ACOG Executive Board

- ACOG; (2005): Medical management of abortion. Practice Bulletin No. 67.
- Arnot D; Margaret L. and Cornelie U (2001): Gender and Crime in Modern Europe. Obstet Gynecol 231:657-665.
- Barreto T, Cambell OM, Daris JL, Favvio V and Fillippi VG (2000):

 Investigating induced abortion in developing countries:

 methods and problems. Stud fam plan 23: 159-170.
- Barrett, J.P. Whiteside JL and Boardman LA (2002): Fatal clostridia sepsis after spontaneous abortion. Obstet Gynecol 99:899-908.
- Bartlett LA, Berg CJ, Shulman HB, Zane SB and et al; (2004): Risk factors for legal induced abortion-related mortality in the United States. Obstet Gynecol. 103:729-737.
- Benthem BS, Wilcox AL, Zan SB and Green CA (2000): Association between viral infection and recurrent abortion befor 16 w: Prospective, community based cohort study. Br Med J, 28: 115-121.
- Blohm F, Friden B and Hay SB(2005): A randomized double blind trial comparing mesoprostol or placbo in the management of early miscarriage. British J Obstet Gynecol, 17:140-165.
- Branch DW and Khamashta MA and Watson F(2003): Anti Phosphoipid Syndrom: Obstetric diagnosis, management, and controversis. Obstet Gynecol 101: 1333-1341.
- **Braude**, **P and Peckering S (2002):** Prei-mplantation genetic diagnosis. Nat Rev Genet 3: 941-953.

- Brogman CE and Jones BS (2000): Legal issues in the provision of medical abortion. Am. J. Obstet Gynecol 31: 183-184.
- Car OE and Feldman B(2004): Parental karyotype and subsequent live birth in recurrent miscarriage Fertil Steril 81: 1296-1301.
- Carbonell L, Esteve JL, Varela L, Velasko A and Sanchez C (1999): 25 mg or 50 mg of oral methotrexate followed by vaginal misoprostol 7 days after, for early abortion: a randomized trial. Obstet Gynecol; 47:182-187.
- Carp, H Dolitzky M and Trakasa E (2004): Hereditary thrombophillias are not associated with a decreased live birth rate in women with recurrent miscarriage. Fertil Steril 78: 158-169.
- Carrell DT and Wilcox AL (2003): Male chromosomal factors of unexplained recurrent Pregnancy loss. Obstet Gynaecol 101:1229-1237.
- Caruso A, Trivellini C and Lowy L (2000): Emergency cerclage in the presence of protruding membranes: Is pregnancy outcome predictable? Obstet Gynecol 79:265-273.
- Castenada R, Lechuga D and Ramos RI (2002): Endemic goiter in pregnant women: utility of simplified classification of thyroid size by palpation and urinary iodide as screening tests. Br J Obstet Gynecol 109:1355-1363.
- Centers for Disease control and Prevention (2003): Abortion surveillance- United States, 2000. Am J Obstet Gynecol 52: 552-566.

- Chatterjee C, Joarder GK, Mukhergee G and Charkaborty M (2007):

 Complications of septic abortion. Indian J Public Health 51:
 193-214.
- *Clark*, *D.A.* (2003): Is there any evidence for immunologically mediated or immunologically modifiable early pregnancy failure? J. Assisted Reprod. Genet., 20: 63-72.
- Clifford, K.; Ria, R.; Watson, H, Fransik S and Regan L (2002): Does suppressing luteinising hormone secretion reduce the miscarriage rate? Results of a randomized controlled trial. Br Med J 13:95-103.
- Cnattingius, S.; Singorello, LB and Anneren G (2000): Caffeine intake and the risk of first-trimester Spontaneous abortion. Nat. Engl J Med. 143:1834-1849.
- Cowett AA, Cohew LS, and et al;(2004): Ultrasound evaluation of the endrometrium after medical termination of pregnancy.

 Obstet Gynecol 103: 871-879.
- Craig TB, Kutteh WH, Lechenberge ES and Stika S (2002): Increase prevalence of insulin resistance in women with a history of recurrent pregnancy loss. Fertil Steril 78: 487-495.
- Crenin, MD (2003): Current medical abortion care. Contraception; 3: 461-490.
- Crenin M and Aubeny E (2004): Medical abortion in early pregnancy, clinician's guide to medical and surgical abortion, Stubblefield, P., Editor, 1st ed. Churchill Livingstone: New York. 91:106-117.

- Cuningham FG, Leveno KJ Bloom SL Hauth JK, Glestrapll LC and Wenstrom KD (2005): Reproductive success and failure, Abortion- Williams Obstetrics, 22ed. McGrew. Hell, New York 232-247.
- **Daniel C and Sacred R (2003):** Family planing, Contraciption, and abortion in Maguire. Oxford university press, 18:80-95.
- *Dewhurst GJ; (2007):* Spontaneous miscarriage. Integrated Obstet and Gynecol. 7th ed., Blackwell scientific Pub., Oxford, London, 96:94-100.
- Diedrich J and Steinauer J(2009): Complications of surgical abortion.

 Clin Obstet Gynecol,52:205-212.
- Dighe M, Cuevas C Mosheri M Dubenky T andDogra VS (2008):Review sonography in the frist trimester bleeding. J Clin Ultrasound.36:352-66.
- Edwards A, Megens A and Peek M (2000): Sexual origins of placental dysfunction. Obstet Gynecol 355:203-204.
- **EL-Refaey H and Hinshaw R**(1992):Medical management of missed abortion and anemberionic pregnancy.Br Med J, 305:1399-6866.
- Elson, J. and Jurkovic; (2004): Biochemistry in diagnosis and management of abnormal early pregnancy. Obstet Gynecol 16, 339-344.
- Farquharson RG, Quenby S and Graves M (2002): Antiphospholipid syndrome in pregnancy: A randomized, controlled trial of treatment. Obstet Gynecol 100:408-416.

- Gamble, SB and Struss L(2008): Abortion surveillance- United states 2005.Am J Obstet Gynecol. 57;1-32
- Gary MM and Harrison DJ (2006): Analysis of severe adverse events related to the use of mifepristone as an abortifacient. Am. Pharmacotherapy 4:191-207.
- Giannolil L, Magli MC and Ferraretti A (2000): Gonadal activity and chromosomal constitution of in vitro generated embryos.J Clin Endocrinol Metab 61: 111-116.
- Glueck, C.J.; Wang PA and et al (2002): Pregnancy outcome among women with polycystic ovary syndrome treated with metformin. Hum Reprod, 17:858-864.
- Goldbergy AB, Dean G and et al; (2004): Manual versus electric vacuum aspiration for early first- trimester abortion: A controlled study of complications rates. Obstet Gynecol, 30:101-121.
- Goldstein SR, and Wolfson GF; (1999): Endovaginal ultrasound meassurment of early embryonic size as a means of assessing gestational age. J Ultrasound Med; 13: 27-31.
- Grimbizis GF, Camus MD; (2001): Clinical implications of uterine malformations and hystroscopic treatment results. Hum Reprod,7:161-174.
- Grudiznskas JG, Stabile ID and et al;(2001): Ultrusound assessment in complication of first trimester pregnancy. Lancet, 12: 123-142.

- Hassoun, DH and Perin ID (2006): Ambulatory medical abortion performed in a family planning centers. J Gynecol Obstet Biol Reprod, 35: 483-499.
- Hausknecht R (2000): Mifeprestone and misoprostal for early medical abortion: 18 months experience in the United Utates. Contra caption, 67: 463.
- **Hebpurn A (2000):** A review of techniques of induced abortion. Fam Plan, 1:9-16.
- Henshaw SK, Adewole I, Singh S, Bancole A, Oye E and et al;(2008):

 Sevrerity and cost of unsafe abortion complications treated in Nigerian hospitals. Fam Plan, 34: 4-50.
- Jain JK, Meckstroth KR, and et al;(2001): A comparison of tamoxifen and misoprostol to misoprostol alone for early pregnancy termination. Contraception, 60:353-56.
- Jones RK and Kost K (2009): Trends in abortion in the United States Clin. Obstet Gynecol, 52: 119-129.
- *Kahn, JG, Beeker BJ and Macissa L (2000):* The efficacy of medical abortion: A meta- analysis. Contraception, 16: 29-35.
- *Kambiss SM*, *Hibbert ML and Macidonia C (2000):* Uterine Perforation resulting in bowel infarction: sharp traumatic bowel and mesenteric injury at the time of pregnancy termination. Milit Med 81:165-172.
- Katy C and Lesley R; (2001): Sporadic and recurrent, miscarriage, in Turnbull's Obstetrics edited by Chamberlin G. and Steer P.,3red edition, Harcourt Publishers limited, 9: 117-127.

- Katzalezky G, Gracia CR Brline GA, and Sammel MD (2004):

 Evaluation of the association between hereditary thrombophillias and recurrent pregnancy loss: a meta analysis.

 N Engl J Med, 164: 558-563.
- **Kesmodel U and Wesborg K (2002):** Moderate alcohol intake and risk of spontaneous abortion. Alcohol, 37:87-91.
- Kigub JH, Daru PH and Ugah IA (2009): Review of maternal deaths from unsafe abortion in Jos, Nigeria. Niger J Med, 18: 103-116.
- Kulier R, Gulmezoglu A, Hofmery G, and cheng L (2004): Medical methods for first trimester abortion. Obstet Gynecol, 25:267-268.
- Kupesics R and Aksomija A (2002): Severity and cost of unsafe abortion complications treated in Nigerian hospital. Niger J Med, 56: 171-180.
- Lancet (2006): Hospital admission resulting from unsafe abortion estimated from 13 developing countries. Obstet Gynecol, 25 368:84.
- Lassey, A.T. (2001): Complications of induced obortions and their prevention in Ghana. East Afr Med J, 72: 774-747.
- Lefkowitz A, Mary R and Fant B (2008): Women's life in Greece and Rome. Baltiomre: Johns Hopkins, university press, 818-447-456.

- Lichtenberg ES, Grimes DA and Brrgatta L (2000): A Clinician's guide to medical and surgical abortion. Stubblefield P Editor, 1st ed. Churchill Livingstone, New York, 197-210-225.
- Lindsay DJ, Lovett IS and Lyons EA (2000): Yolk sac at endovaginal ultrasound. J Ultrasound Med, 183:115-118.
- Luesley DM and Baker PN (2004): Problems in early pregnancy. Evidence-based text, Obstet Gynecol for MRCOG, 1st ed, Arnold, Great Britain 57: 606-615.
- Luise C, Jermy Ket AL and May C (2002): Outcome of expectant management of spontaneous first-trimester miscarriage.

 Observational Study, Britain Med J 329:873-879.
- MacIsaac L and Darney P; (2000): Early surgical abortion an alternative to and backup for medical abortion. Am J Obstet Gynecol. 82:183-189.
- Morikaw M, Yamada H, Kato EH, Shimada S and Minakami H (2004): Embryo loss pattern is predominant in miscarriages with normal chromosomal karyotype among women with repeated miscarriage. Hum Reprod 19:264-274.
- Muffley PE, Stitely ML and Gherman RB (2002): Early intrauterine pregnancy failure; A randomized trial of medical versus surgical treatment. Am J Obstet Gynceol, 187:321-332.
- MurAd AW (2001): Study of obstetric emergencies in Benha University Hospital during the period from 1996 to 2000. Benha Faculty of Medicine, Benha University, Egypt, Benha, 3:75-76

- Nybo AM, Wohlfohrt J and Chrestin B (2000): Maternal age and fetal loss: population based register linkage study. Br. Med J, 320: 708-717.
- Oakeshott P, Hay P and Hay S (2002): Association between bacterial vaginosis and chlamedial infection and miscarriage before 16 weeks gestation: Prospective, community based cohort study. Br Med J, 325: 334-341.
- Otes-Whitehead RM, Hass DM and et al; (2003): Progesteron for preventing miscarriage. Hum Reprod Med, 17: 135-141.
- Owen Jand Iams JD (2003): Vaginal sonography and cervical incompetence. Am J Obstet Gynecol 188: 586-591.
- **Paul ME, Mitchell CM and et al; (2002):** Early surgical abortion: efficacy and safety. Am J Obstet Gynecol 187: 407-416.
- Perriera L, Reeves MF and Rogers AJ (2008): Ultrasound criteria for diagnosis of early pregnancy failure and ectopic pregnancy. Semin Reprod Med, 26:373-382.
- Philipp T, Philipp K Reiner A, and Beer F (2003): Embryo-scopic and cytogenic analysis of 233 missed abortions: factors involved in the pathogenesis of developmental defects of early failed pregnancies. Hum Rerprod Med, 18:724-732.
- **Pratola DC and Poland BJ (2001):** Vaginal bleeding in early pregnancy. Am J Obstet and Gynecol 35: 937-941.
- Provansal M, Miamari R and Grgoire B (2009): Medical abortion at home and at hospital: A trial of efficacy and acceptability.

 Obstet Gynecol, 41: 170-181.

- Raghupathy R, Makhseed M Azizich F and Omar A (2000): Cytokin production by maternal lymphocytes during normal human pregnancy and in unexplained recurrent spontaneous abortion. Hum Reprod Med, 15:713-721.
- Rey E, Kshn SH and Senerier I(2003): Thrombophillic disorders and fetal loss. A meta analysis. Obstet Gynecol, 35: 361 -367.
- **Riddle JM**;(2001)): Contraception and abortion from the ancient world to the Renaissance. Cambridge, Harvard University press, 11: 46-52.
- Rushowrth FH and Backos M(2003): Prospective pregnancy outcome in untreated recurrent miscarriages with thyroid autoantibodies. Hum Repord 15: 1637-1745.
- Rust OA, Atlas RO and Reed G (2001): Revisiting the short cervix detected by transvaginal ultrasound in the second trimester: Why cerclage may not help. Am J Obstet Gynecol 185:1098-2012.
- Sairam S, Khare M and Mecheilides A (2001): The role of ultrasound in the expectant management of early pregnancy loss.

 Ultrasound Obstet Gynecol 17:506-509.
- Salin R, Raganl W and Wolfer B and Backos M (2003): A comparative study of morphology of congenital uterine anomalies in women with and without history of recurrent first trimester miscarriage. Hum Reprod, 19 162-166.

- Sawyer E and Jurkovic D(2007): Ultrasonography in the diagnosis and management of abnormal early pregnancy. Clin Obstet Gynecol, 50:31-54.
- Schauberger CW, Mathiason MA and Rony BL (2005): Ultrasound assessment of frist-trimester bleeding. Obstet Gynecol 105: 333-338.
- Scott JR (2003): Immunotherapy for recurrent miscarriage .Fertil Steril,17:107-125.
- Sedgh G, Stanley KH, Suchila S, Akinrinola B and Gona D (2007):

 legal abortion worldwide: Incidence and recent trends.

 Internat. family planing perspectives 33:117-138.
- Sheikh M and Sadiyya N (2003): Family planning, contraception, and abortion in Islam. Oxford University Press,27: 105-128.
- Signorello LB and Mclaughlin JK (2004): Maternal caffeine consumption and spontaneous abortion: A review of the epidemiologic evidence. Epidemiology 15:229-241.
- Simon NM, Rothmon D; (1998): Psychological factors related to spontaneous and theraputic abortion. Am J Obstet Gynecol, 104:799-815.
- Sophie CM, Philippe B and Groff JT (2000): Medical termination of pregnancy. New Eng J Med, 342:946-956.
- Sotiriadis A, Makrydimas G (2005): Expectant, medical, or surgical management of first-trimester miscarriage: a meta-analysis.

 Obstet Gynecol. 105:1104-1113.

- Soulat C and Gelly M (2006): Immediate complications of surgical abortion. Am J Obstet Gynecol 35: 157-162.
- Stephenson MD, Awartani KA and Robenson WB (2002): Cytogenic analysis of miscarriage from couples with recurrent miscarriage: a case control study. Hum Reprod, 17: 446-451.
- Stirrat GM (2000): Recurrent miscarriage: Definition and epidemiology. Lanct, 366: 673-678.
- Stricker RB Steinleitner A Bokoff CN, Weekestin L and Wenger EE (2000): Successful treatment of immunologic abortion with low dose intravenous immunoglobulin. Fertil steril 73:563-569.
- Stubblefield PG, Naftrolin F and et al; (2004): Lamenaria augmentation of intra-amniotic PGF mid-trimester pregnancy termination. Prostaglandins, 10:1413-1975.
- Terkildson MF, Parilia BV and Kumar B (2003): Factors associated with success of emergent second-trimester cerclage. Obst Gynecol 101:565-571.
- Tristan SB and Gilliam M;(2009): First trimester surgical abortion.

 Obstet Gynecol. 52:151-160.
- **Tulandi T and Sammour A (2000):** Evidence based management of ectopic pregnancy. Curr Opin Obstet Gynecol, 12:289-292.
- Ventura SJ, Abona JC Musher WD and Henshaw S (2003): Revised pregnancy rates, 1990-97, and new rates for 1998-99: United States. Natal Vital Stat Rep 11: 52-61.

- Wegmann TG and Lin H(2000): Bidirectional cytokine interaction in the maternal fetal relationship: is successful pregnancy a TH₂ phensmenon? Immunol Today, 14:353-356.
- Weisman AM, Dawson JD and Rijhsinghani A (2002): Non- evidence-based use of Rho (D) immunoglobulin for threatened abortion by family practice and obstetric faculty physicians. J Reprod Med 47: 909-1006.
- Weiss JL, Malone FD and Vedaver J (2004): Threatened abortion: A screening study. Am J Obstet Gynecol 190: 745-753.
- Wisborg K, Kesmodel U and Henreksen TB (2003): A prospective study of maternal smoking and spontaneous abortion. Obstet Gynecol, 82: 936-943.
- Younis JS, Brenner B Ohel J, Tale J, Lanir N and Benami M (2000):

 Activated protein C resistance and factor V leaden mutation can be associated with first as well as second trimester recurrent pregnancy loss. Am J Repord Immunol 43: 131-137.