

REFERENCES

- Abdel-Aleem, H.; Villar, J.; Guimezoglu, A.M. ., (2003):*** The pharmacokinetics of the prostaglandin E1 analogue misoprostol in plasma and colostrums after postpartum oral administration, Eur J Obstet Gynecol Reprot Biol; 108:25-8.
- Acalovschi, M.; Durnitrascu, D.L.; Suteu, T.; Veracs, A.; Alii, and Badea, R.L. (2002):*** acute intravaginal misoprostol toxicity with fetal demise, Acta Gastroenicol Mol[^]-65:191-2.
- Adair, C.D. (2000):*** Nopharmacologic approaches to cervical priming and labor induction. Clinical Obstetrics and Gynecology; 43: 447-54.
- Agarwal, A.; Gupta, S. and Sharma, R.K. (2005):*** Role of oxidative stress in female reproduction. Reproductive Biological Endocrinology; 3: 28.
- Aktan, F. (2004):*** Inducible nitric oxide synthase - mediated nitric oxide production and its regulation. Life Science; 75: 639-653.
- Altabef, K.; Spencer, J.T. and Zinberg, S. (1992):*** Intravenous nitroglycerin for uterine relaxation of an inverted uterus. American Journal of Obstetrics and Gynecology; 166:1237.
- American College of Obstetricians and Gynecologists, (2003):*** New U.S. food and drug administration labeling on cytotec (misoprostol) use and pregnancy: ACOG committee opinion. No. 283. Washington, D.C.

American College of Obstetricians and Gynecologists, (2000): response to searle's drug warning on misoprostol: ACOG committee opinion. No. 248. Washington, D.C.

American College of Obstetricians and Gynecologists (ACOG 1999): Induction of labor with misoprostol. AGOG committee opining 228 Washington DC.

American College of Obstetricians and Gynecologists (1999): Induction of labor. ACOG Practice Bulletin. No. 10. Washington, DC: ACOG.

American College of Obstetricians and Gynecologists (1999): Induction of labor with misoprostol. ACOG committee opinion 228. Washington, D.C.: ACOG; P. 2.

American Journal of Obstetric and Gynecology March vol. 112, p. 26,72.

Anderson, K.E. Uif Ulmsten, and Kent UeIand, (1983): The forces of labor, uterine contractions and the resistance of the cervix. Clin Obstet Gynecol.26:100-105.

Armstrong, C.P. and Blower, A.L. (1987): gastrointestinal tract complications with NSAIDs. GUT; 28:527.

Aslan, H.; Unlu, E.; Agar, M. and Ceylan, Y. (2004): Uterine rupture associated with misoprostol labor induction in women with previous Cesarean delivery. Eur J Obstet Gynecol Reprod Biol; 113:45-48.

- Austin, J.F.; Rouse, A. and Hanna, E. (1997):** Acute intravaginal misoprostol toxicity with fetal demise J Emergency Med; 15:61-2.
- Beales, L.L.; Demons, M. and Kong, W.M. (1997):** Misoprostol associated platelet aggregation dysfunction. European J of gastroenterology and hepatology.; 1:91 -92.
- Bergstrom, S.; Danielsson, H. and Samuelsson, B.(1964):** The enzymatic formation of prostaglandin E₂ from arachidonic acid Biochemica et Biophysica. 1964a;90:207-210.
- Bishop, E.H. (1964):** Pelvic scoring for elective induction. Obstetrics and Gynecology; 24:266.
- Bootstaylor, B.; Roman, C.; Heymann, M.A. and Parer, J.T. (1994):** fetal cardiorespiratory effects of nitroglycerin in the near term pregnant sheep. American Journal of Obstetric Gynecology; 170 (suppl): 281.
- Boulvain, M.; Irion, O.; Marcoux, S. and Fraser, W. (1999):** sweeping of the membranes to prevent post-term pregnancy and to induce labour: a systematic review. Br J Obstet Gynaecol 1999; 106:481-5.
- Boulvain, M.; Kelly, A. and Lohse, C. (2001):** Mechanical methods for induction of labour. Cochrane Database Syst Rev; CD001233.
- Boulvain, M.; Stan, C. and Irion, O. (2001):** Elective delivery in diabetic pregnant women. Cochrane Database Syst Rev; Issue 2.

- Brune, B.; Von Knethen, A. and Sandau, K.B. (1998):** Nitric oxide and its role in apoptosis. *European Journal of Pharmacology*; 351: 261-72.
- Bugalho, A.; Faundes, A. and Jamisse, L. (1996):** Evaluation of the effectiveness of vaginal misoprostol to induce first trimester abortion. *Contraception*; 53:244- 6.
- Buhimschi, C.; Buhimschi, L.A. and Malinow, A.M. (2003):** The forces of labour. *Fetal and Maternal Medicine Review*; 14(4):273- 307.
- Buhimschi, C.S.; Buhimschi, L.A. and Maiinow, A.M. (2002):** Pushing in labor: performance and notendurance. *Am J Obstet .Gynecol*; 186(6):1339-44.
- Buhimschi, Yallampalli C.; Dong, Y.L. and Garfield, R.E. (1995):** Involvement of a nitric oxide-cyclic guanosine monophosphate pathway in control of human uterine contractility during pregnancy. *American Journal of Obstetrics and Gynecology*; 172: 1577-84.
- Caldwell, W.E. and Moloy, H.C. (2005):** Anatomical variations in the female pelvis and their effect on labor with a suggested classification. *Am J Obstet Gynecol* 1933; 26:479- 505. (Quoted from: **Liao JB, 1 Buhimschi CS, Norwitz ER**, Normal Labor: Mechanism and Duration. *Obstet Gynecol Clin N Am*; (32): 145-164).
- Cammu, H.; Martens, G.; Ruyssinck, G. and Amy, J.J. (2002):** Outcome after elective labor induction in nulliparous women: a matched control study. *Am J Obstet Gynecol*; 186:240-4.

- Casey, M.L. and MacDonald, P.C. (1998a):** Biomolecular processes in the initiation of parturition: decidual activation. *Clinical Obstetrics and Gynecology*; 31:533-52.
- Castleman, L.D.; Oanh, K.T.; Hyman, A.G.; Thuy, le T. and Blumenthal, B.D. (2006):** Introduction of the dilation and evacuation procedure for second-trimester abortion in Vietnam using manual vacuum aspiration and buccal misoprostol, *Contraception*; 74: 272-6.
- Chan, C.C.; Tang, O.S.; Ng, E.H.; Li, C.F. and Ho, P.C. (2005):** Intracervical sodium nitroprusside versus vaginal misoprostol in First trimester surgical termination of pregnancy: a randomized double-blinded controlled trial. *Hum Reprod.*, 20: 829-833.
- Chandra, S.; Crane, J.M.G. and Hutchens, D.:** Trans-vaginal ultrasound and digital examination in predicting successful labor induction. *Obstet Gynecol* 2001; 98:2-6.
- Chanrachakul, B.; Herabutya, Y. and Punyavachira, P. (2002):** A Randomized trial of isosorbide mononitrate versus misoprostol for cervical ripening at term. *International Journal of Gynaecology and Obstetrics*; 78: 139-45.
- Chanrachakul, B.; Herabutya, Y. and Punyavachira, P. (2000):** Randomized comparison of glyceryl trinitrate and prostaglandin E2 for cervical ripening at term. *Obstet Gynecol.*, 96: 549-553.
- Cocks, D.P. (2006):** Significance of initial condition of cervix uteri to subsequent course of labour. *Br Med J* 1955; 1: 327-8. (Quoted , from: **Crane JMG.** Factors Predicting Labor

Induction Success: A Critical Analysis. Clin Obstet Gynecol.; 49(3): 573-84).

CRAIG T.B. (1988): Human cervical connective tissue contribution in labour 77:488.

Crane, J.M.G. (2006): Factors Predicting Labor Induction Success: A Critical Analysis. Clin Obstet Gynecol; 49(3): 573-84.

Crane, J.M.G.; Delaney, T. and Butt, K.D. (2004): Predictors of successful labor induction with oral or vaginal misoprostol. J Maternol Fetal NeonatMed; 15:319-23.

Crowley, P. (2004): Interventions for preventing or improving the outcome of delivery at or beyond term. Cochrane Database Syst Rev;(3):CD000170.

Cunnninham, F.G.; Levano, K.J.; Bloom, S.L. and Hawth, J.C.F.G.K.D. (2005): Gilstrap L and Williams Obstetrics; 22 edition. Hill New York. CH9 (Abortion): 231-51.

DA, Rahall A.; Jones, M.M.; Goodwin, T.M. and Paul, R.H. (1995b): Misoprostol :an effective agent for cervical ripening and labor induction. Am J Obstet _ Gynecol; 172(6): 181-6.

Danforth, D.N. (1947): The fibrous nature of the human cervix, and its relation to isthmic segment in gravid and non gravid uteri. Am J Obstet Gynecol; 53: 541-60.

Danforth, D.N. (1983): The morphology of the human cervix. Clin Obstet Gynecol 1983; 26 (1): 7-13.

- Deborah, A.; Wing, M.D.; Cecilia, A. and Lyons Gaffaney, M.D. (2006):** Vaginal Misoprostol Administration for Cervical Ripening and Labor Induction. Clinical obstetrics and gynecology; 49(3): 627-41.
- Den Hertog, D.F.; Michael, C.; Samouel, D. and Tiffany, H. (2001):** octa peptide biosynthesis and metabolism Am J Physiol, cell physiol; 130: 17-9.
- DMKG, (2003):** Misoprostol in the treatment of trigeminal neuralgia associated with multiple sclerosis. J Neurol; - 250:542-3.
- Edwards, R.K. and Richards, D.S. (2000):** Preinduction cervical assessment. Clin Obstet Gynecol; 43:440-6.
- Ekerhovd, E.; Weijdegard, B.; Brannstrom, M. and Norstrom, A. (1999):** Nitric oxide-mediated effects on myometrial contractility at term during prelabor and labor. Obstet Gynecol, 93: 987-994.
- Ekerhovd, E.; Weijdegard, B.; Brannstrom, M.; Mattsby-Baltzer, I. and Norstrom, A. (2002):** Nitric oxide induced cervical ripening in the human: involvement of cyclic guanosine monophosphate, prostaglandin F (2 alpha), and prostaglandin E (2). Am J Obstet Gynecol, 186:745-50.
- Elder, M.G. (1992):** Foreword. In: Elder MG (ed). Baillieres Clinical Obstetrics and Gynecology, international practice and research vol.6, No.6: Prostaglandins ppix. Pallierre^fs Tindall London, UK.
- Eppel, W.; Facchinetti, F.; Schleussner, E.; Piccinini, F.; Pizzi, C.; Gruber, D.M.; Schneider, B. and Tschugguel, W. (2005):** Second trimester abortion using isosorbide mononitrate in

addition to gemeprost compared with gemeprost alone: a double-blind randomized, placebo-controlled multicenter trial. *American Journal of Obstetrics and Gynecology*; 192, 856-61.

Facchinetti, F.; Piccinini, F. and Volpe, A. (2000): Chemical ripening of the cervix with intracervical application of sodium nitroprusside: a randomized controlled trial. *Hum Reprod.*, 15: 2224-2227.

Fich, A.; Arber, T.V.; Sestieri, M.L. and Zajieek, G. (1985): misoprostol adverse reactions, *gastroenterology*; 89:57-8.

Fonseca, W.V.; Alencer, A.J.; Morta, F.S. and Coelho, H.L. (1991): Misoprostol and congenital fetal malformations. , *Lancet*; 338:56.

Forman, A.; Ulmsten U, Banyai J, Wingerup L and Uldbjerg N (1992): Evidence for a local effect of the intracervical prostaglandin E2 gel. *American Journal of Obstetrics , and Gynecology*; 143 756-60.

Forrl, A.; Sheppard, B.L.; Burke, G. and Bonnar, J. (1994): Platelet activation in normotensive and hypertensive pregnancies complicated by intrauterine growth retardation. *British Journal of Obstetrics and Gynecology*; 101:209-14.

Fraser, W.D.; Marcoux, S. and Krauss, I. (2000): Multicenter, randomized, controlled trial of delayed pushing for nulliparous women in the second stage of labor with continuous epidural analgesia: the PEOPLE (Pushing Early or Pushing Late with Epidural) Study Group. *Am J Obstet Gynecol*; 182(5): 1165- 72.

- Gary, F.; Cunningham, Kenneth J. AND Leveno, Steven, L. (2005):** Bloom, Williams Obstetrics 22nd edition, chapter 5, section 2 maternal physiology p. 123-24 and, Chapter 6 section 2 Parturition, p. 152, 153,154, 157.
- Gelber, S. and Sciscion, A. (2006):** Mechanical Methods of Cervical Ripening and Labor Induction. Clin Obstet Gynecol; 49(3): 642-57.
- Gifford, D.; Morton, S. and Fiske, M. (2000):** Lack of progress in labor as a reason for cesarean. Obstet Gynecol; 95:589- 95.
- Gimple and Farenholz (2001):** oxytocin biophysiology and metabolism. Am J physiol; 16: 98-101.
- Goldman, (2000):** Cecil Textbook of Medicine, 21st ed. Published by W.B Saunders company :233,1 189-1195.
- Gonzalez, C.H.; Marques, M.J.; Kirn , C.A. and Sugagama, S.M. (1998):** Congential abnormalities with misoprostol misuse in first trimester of pregnancy. Lancet; 351:1624-5.
- Goszez, A.; Crodzinska, L.; Kostka, Trebka, E. and Bleron, K. (1996):** Misoprostol in the treatment of peripheral ischemic disease Acta physiologic Hungarica; 84:415-6.
- Graham, D.Y.; Agrawal, N.M. and Roth, S.H. (1988):** A study on prevention of NSAIDs induced gastric ulcers, Lancet; 2:1277-8.
- Graham, D.Y.; Agrawal, N.M. and Roth, S.J.L. (1988):** A study on prevention of NSAIDs induced gastric ulcers with misoprostol The lancet 2:1277-1280.

- Granstrom L, Ekman G, Ulmsten U and Malmstrom A:** Changes in the connective tissue of corpus and cervix uteri during ripening and labour in term pregnancy. *Br J Obstet Gynaecol* 1989; 96 (10): 1198-202.
- Guinn, D.A.; Davies, J.K. and Jones, R.O. (2004):** Labor induction in women with an unfavorable Bishop score: randomized controlled trial of intrauterine Foley catheter with concurrent oxytocin infusion versus Foley catheter with extraamniotic saline infusion with concurrent oxytocin infusion. *Am J Obstet Gynecol* 191:225-9.
- Hadi, H. (2000):** Cervical ripening and labor induction: clinical guidelines. *Clin " Obstet Gynecol*; 43: 524-36.
- Hannah ME, Hannah WJ, Hellmann J, Hewson S, Milner R, Willan A, and the Canadian multicenter post-term pregnancy trial group (1992):** Induction of labour as compared with serial antenatal monitoring in post-term pregnancy. A randomized controlled trial. *N Engl J Med*; 326:1587-92.
- Hatfield, A.S.; Sanchez-Ramos, L. and Kaunitz, A.M. (2001):** Sonographic cervical assessment to predict the success of labor induction: a systematic review with metaanalysis. *AJOG*; 186-192.
- Henry, Lepoivre, A.; Drapier, J.C.; Ducrocq, C.; Boucher, J.L. and Guissani, A. (1993):** EPR characterization of molecular targets for NO in mammalian cells and organelles. *FASEB J*; 7: 1124-34.

Hofmeyr, G.J. and Gulmezoglu, A.M. (2003): Vaginal misoprostol for cervical ripening and induction of labour. Cochrane Database of Sys Rev(1):CD000941.

Hughesdon, P.E. (1952): The fibromuscular structure of the cervix and its changes during pregnancy and labour. In: Studd J (ed.): The cervix in pregnancy and labour. Progress in Obstet and Gynecol. Churchill Livingstone; 12: 108.

Ingemarsson, I.; Arulkumaran, S. and Ratnam, S.S. (1985): Single injection of butaline in term labor, Effect on uterine activity. Am J Obstet Gynecol ; 153:865-9.

James, E.F. and Rynolds. Prostaglandins Martindale Librairie du liban.199j;1367-1376 Quoted in internet midline.

Josie Tenore (2003): Methods for Cervical Ripening and Induction of Labor, American Family Physician May 15; 67: 2123-8.

Junqueira, L.C.U.; Zugaib, M.; Montes, G.S.; Toledo, O.M.S.; Krisztan, R.M. and Shigihara, K.M. (1980): Morphological and histochemical evidence of the occurrence of collagenolysis and for the role of neutrophilic polymorphonuclear leukocytes during cervical dilatation. Am. J. Obstet. Gynecol., 138:273-281.
A: Pharmacokinetic profile of anti-ulcer prostaglandin, misoprostol, in human Postgraduate Med J.1988;64(I): 80.

Karim, S. and Sharma, S. (1971): Oral administration of prostaglandin for the induction of labor BMJ.1971; 1:260-267.

- Kavanagh, J.; Kelly, A.J. and Thomas, J. (2001b):** Breast stimulation for cervical ripening and induction of labour. Cochrane Database of Systematic Reviews. CD003392.
- Kavanagh, J.; Kelly, A.J. and Thomas, J. (2001a):** Sexual intercourse for cervical ripening and induction of labour. Cochrane Database of Systematic Reviews. CD003093.
- Keirse, M. (1979):** Prostaglandins in parturition. In Human Parturition: Kerise M, Anderson A and Gravenhorst. J (ed); The Hague, Netherlands. Martinus Nijhoff 1979; 101-120. **Keirse M.::** Enkin MW, Ren Freur MJ, Neilson IP (eds).: Vaginal PG versus ocytocin for induction of labor. In Pregnancy and childbirth, module. Cochrane database of systemic reviews No. 04538. 1992; Disc Issue 2 Oxford Update Software. **1993a.**
- Keirse, M.N. (1992):** Therapeutic use of prostaglandins. Bailliere's Clin Obstet Gynecol; 6:787-809.
- Kirby, R.S. (2004):** Trends in labor induction in the United States: is it true that what goes up must come down? Birth; 31: 148-51.
- Kurup, A.; Arulkumaran, S.; Tay, D.; Ingemarsson, I.; Ratnam, S.S. (1991):** Can terbutaline be used as a nebuliser instead of intravenous injection for inhibition of uterine activity? Gynecol Obstet Invest 1991 ;32:84-7.
- Lang, W. (1955):** Vaginal acidity and pH: a review obstet gynecol surv. 1955, Aug; 10(4):546-60.
- Langlios, P.L. (1970):** The size of the normal uterus. J Reprod Med; 4: 220.

- Lapiere, C.M.; Lenaers, A. and Kohn, L.D. (2005):** Procollagen peptidase: An enzyme excising the coordination peptides of procollagen. Proc Natl Acad Sci USA 1971; 68(12):3054-8. (Quoted from: **Richard I.** The genetic and molecular bases of monogenic disorders affecting proteolytic systems. J Med Genet. July; 42(7): 529-39).
- Larsen, B. (1993):** Vaginal flora in health and disease. Clin Obstet Gynecol; 36:107-21.
- Lawrie, A.; Penney, G. and Templeton, A. (1996):** A randomized comparison of oral and vaginal misoprostol for cervical priming before suction termination of pregnancy. Br J Obstet Gynaecol; 103:1117-9.
- Lazanakis, M.; Marsh, M.; Brockbank, E. and Economides, D. (2002):** Assessment of the cervix in the third trimester of pregnancy using transvaginal ultrasound scanning. Eur J Obstet Gynecol Reprod Biol; 105:31-5.
- Ledingham, M.A.; Thomson, A.J.; Young, A.; Macara, L.M.; Greer, I.A. and Norman, J.E (2000):** Changes in the expression of nitric oxide synthase in the human uterine cervix during pregnancy and parturition. Mol Hum Reprod., 6: 1041-1048.
- Leppert, P.C. and Yu, S.Y. (1991):** Three dimensional structures of uterine elastic fibers: Scanning ' electron microscopic studies. Connect Tissue Res., 26:15-31.
- Leppert, P.C. (1992):** Cervical softening, effacement and dilatation. A complex biochemical cascade journal of Maternal and Fetal Medicine, 1: 213-23.

- Leppert, P.C. (1995):** Anatomy and physiology of cervical ripening. Clin Obstet Gynecol., 38(2):267-79.
- Leppert, P.C.; Yu, S.Y.; Keller, S.; Cerreta, J. and Mandl, I. (1987):** Decreased elastic fibers and desmosine content in incompetent cervix. Am J Obstet Gynecol; 157 (5): 1134-39.
- Leung, A.; Kwok, P. and Chang, A. (1987):** Association between prostaglandin E2 and placental abruption. Br J Obstet Gynaecol; 94:1001- 2.
- Li, C.; Chan, C. and Ho, P. (2003b):** A study of the efficacy of cervical ripening with nitric oxide donor versus placebo for cervical priming before second- trimester termination of pregnancy. Contraception, 68: 269-272.
- Liao, J.B.; Buhimschi, C.S. and Norwitz, E.R. (2005):** Normal Labor: Mechanism and Duration. Obstet Gynecol Clin N Am 2005; 32: 145- 64.
- Lichtenberg, E.S. (2004):** Complications of osmotic dilators. Obstet Gynecol Surv; 59:528-36.
- Liggins, G.C.; Fairclough, R.J. and Grieves, S.A. (1973):** The mechanism of initiation of parturition in the ewe. Recent Prog Horm Res 1973; 29:111 -59. (Quoted from: **Liao JB, Buhimschi CS, Norwitz ER. Normal Labor: Mechanism and Duration. Obstet Gynecol Clin N Am 2005; (32): 145- 164).**
- Lin, M.G. and Rouse, D.J. (2006):** What is a failed labor induction? Clin Obstet Gynecol.; 49(3):585-93.

- MacKenzie, I.Z.** : Clinical studies on cervical ripening. In Ellwood DA person ABM, eds. *The Cervix in Pregnancy and Labour* 1981. London: Churchill Livingstone: 161-86.
- MacKenzie, I.Z. and Embery, M.P. (1979):** A comparison of PGE₂ and PGF₂ vaginal gel for ripening the cervix before induction of labor. *Br J Obstet Gynaecol*; 85:657.
- MacKenzie, I.Z. and McKinlay, E. (1995):** Paediatric follow-up study: follow-up of infants delivered by spontaneous vaginal delivery, following PGE₂ or oxytocin induction, or by cesarean section. In Calder AA, Keirse MJNC, MacKenzie IZ, eds. *Propan RS. A Clinical Review*, Hastenove: Euromed:37-9.
- Marletta, M. (1994):** Nitric oxide synthase: Aspects concerning structure and catalysis cell; 78: 927-30.
- Martin, J.A.; Hamilton, B.E. and Sutton, P.D.E.T. (2005):** Al. Births: Final data for 2003. *National Vital Statistics Reports*: 54(2). Hyattsville, MD: National Center for Health Statistics.
- Maul, H.; Longo, M.; Saade, G.R. and Garfield, R.E. (2003a):** Nitric oxide and its role during pregnancy: from ovulation to delivery. *Current Pharmacology* ; 9: 359.
- Maul, H.; Mackay, L. and Robert, E. Garfield (2006):** Biochemical, molecular and clinical considerations. *Clinical Obstetrics and Gynecology*: Lippincott Williams & Wilkins; 49 (3) 551-63.
- McCarthy, A.L.; Woolfson, R.G.; Raju, S.K. and Poston, L. (1993):** Abnormal endothelial cell function of resistance arteries

from women with pre-eclampsia. *American Journal of Obstetrics and Gynecology*; 168: 1323-30.

Meckstroth, K.R.; Whitaker, A.K.; Bertisch, S.; Goldberg, A.B. and Darney, P.D. (2006): Misoprostol administered by epithelial routes: Drug absorption and uterine response *Obs.&Gyn.*;108 :582-590.

Mervi Vaisanen-Tommiska, (2006): Nitric Oxide and uterine cervix: role in cervical ripening. Helsinki University Central Hospital University of Helsinki, Finland. *nmqtaelandin in drugs*.

Moncada, S. and Higgs, A. (1997): The L-arginine-nitric oxide pathway. *N Engl J Med*, 329: 2002- 20127.

Murray, H.M.; Greer, I.A. and Norman, J.E. (2006): The prim study; a randomized comparison of prostaglandin E2 with the nitric oxide donor isosorbide mononitrate for cervical ripening before the induction of labor at term. *American Journal of Obstetrics and Gynecology*; 194: 1012-21.

Nakane, M. Schmidt, H.H.; Pollock, J.S.; Forstermann, U. and Murad, F. (1993): Cloned human brain nitric oxide synthase is highly expressed in skeletal muscle. *FEES Lett.*, 316: 175-180.

Neilson, D.R.; Prins, R.P. and Bolton, R.N. (1983): A comparison of prostaglandin E2 gel and prostaglandin F2 gel for pre-induction cervical ripening. *Am J Obstet Gynecol*; 146:526.

Neilson, J.P. (2002): Mifepristone for induction of labour. *Cochrane Database Syst Rev*; 2:CD002865.

- Ness, A.; Goldberg, J. and Berghella, V. (2005):** Abnormalities of the First and Second stages of Labor. *Obstet Gynecol Clin N Am* 32: 201- 20.
- Ngai, S.W.; Chan, Y.M. and Ho, P.C. (2001):** The use of misoprostol prior to hysteroscopy in postmenopausal women *hum. Reprod*; 16: 186-9.
- Normal, J.E.; Thong, K.J. and Baird, D.T. (1991):** Uterine contractility and induction of abortion in early pregnancy by misoprostol and mifepristone. *V Lancet*; 338:1233-6.
- Norris, M.C. and Leighton, B.L. (1990):** Intravenous nitroglycerin aids manual removal of retained placenta. *Anesthesiol (letter)*; 73:787.
- Norwits, E.R.; Snegovskikh, V.Y. and Caughet, A.B. (2007):** Prolonged Pregnancy: x When Should We Intervene? *Clin Obstet Gynecol*; 50(2): 547-57.
- Norwitz, E.; Robinson, J. and Repke, J. (2002):** Labor and delivery. In: Gabbe SG, Niebyl JR, Simpson JL, eds. *Obstetrics: normal and problem pregnancies*. 4th ed 2002. New York: Churchill Livingstone,:353-94.
- O'Brien, P.; El-Rafaey, H.; Gordon, A.; Geory, M. and Rodeck, C.H. (1998):** Rectal administered misoprostol for the treatment of postpartum hemorrhage unresponsive to oxytocin and ergometrine: a descriptive study. *Obstet Gynecol.*, 92(2): 212-4.
- Odendaal, H.J.; Pattinson, R.C.; Bam, R.; Grove, D. and Kotze, T.J. (1990):** Aggressive or expectant management for patients with

severe pre-eclampsia between 28-34 weeks gestation: A randomized controlled trial. *Obstet Gynecol*; 76:1070-5.

Olesen, A.W.; Westergaard, J.G. and Olesen, J. (2003): Perinatal and maternal complications related to postterm delivery: a national register-based study, 1978-1993. *Am J Obstet Gynecol* 189:222-7.

Palmer, R.M.J. and Moncada, S. (1987): Endogenous Nitric Oxide Inhibits Human Platelet Adhesion to Vascular Endothelium. *Lancet*; 2: 1057-82

Quiroga, Diaz R.; Esperaza Arechiga, M.; Batiza Resendiz, V.; Coronado Lopez, Hernandez Ayup, S.; Martinet, X. and Cuervo, J. (2002): Vaginal misoprostol in the prevention of post partum hemorrhage. *Gynecol ' Obstet*; 70: 52-3.

Rechberger, T. and Woessner, J.F. (1994): Its inhibitors, and decorin in the lower uterine segment in pregnant women. *Am. J. Obstet. Gynecol.*, 168: 1598-1603.

Ringel, R.E.; Haney, P.J. and Brenner, J.I. (1983): Fetal abnormalities with misoprostol misuse in abortion. *J I pediat*; 103:251-5.

Romero, R. (1998): Clinical application of nitric oxide donors and blockers. *Hum Reprod.*; 13:248-250.

Romero, R.; Munoz H. and Gomez R. (1996): Increase in prostaglandin bioavailability precedes the onset of human parturition. *Prostaglandins Leukot Essent Fatty Acids* 1996; 54:187.

Rosselli, M (1997): Nitric oxide and reproduction. *Molecular Human Reproduction*; 3: 639-41.

- Sanchez-Ramo, L.; Kaunitz, A.M. and Delke, I. (2002):** Labor induction with 25 microg versus 50 microg intravaginal misoprostol: a systematic review. *Obstet Gynecol*; 93:145-51.
- Sanchez-Ramos, L.; Kaunitz, A.M. and Wears, R.L. (1997):** Misoprostol for cervical ripening and labor induction: a meta-analysis. *Obstet Gynecol*; 89:633-42.
- Sanchez-Ramos, L.; Olivier, F.; Delke, I. and Kaunitz, A.M. (2003):** Labor induction versus expectant management for postterm pregnancies: a systematic review with meta-analysis. *Obstet Gynecol*; 101(6): 1312-8.
- Sanchez-RamosL, Gaudier, F.L. and Kaunitz, A.M. (2000):** cervical pipening and labour induction after previous cesarean delivery, *clin Obestet Gynecol*; 43: 51.3-23.
- Schoenhard, G.; Fitzpatrick, R.J. and Dixon, W.C.W. (1985):** Metabolism and pharmacokinetic studies of misoprostol. *Drug Dis Sci*;30: 126-128.
- Schonfofer, S. and Fonesca, (1992):** Teratogenicity of misoprostol. *Lancet*; 139: 437-40.
- Sennstrom, M.B.; Eknian, G.; Westergren-Thorsson, G.; Malmstrom, A.; Bysfrom, B.; Endresen, U.; Mlambo, N.; Norman, M.; Stabi, B. and Brauner, A. (2000):** Human cervical ripening, an inflammatory process mediated by cytokines. *Mol Hum Reprod.*, 6: 375-381.
- Sherman DJ, Frenkel E, Pansky M, Caspi E, Bukovsky I and Langer R (2001):** Balloon cervical ripening with extra-amniotic infusion of

saline or prostaglandin E2: a double-blind, randomized controlled study. *Obstetrics and Gynecology*; 97:375-80.

Singhal, K.K. and La Gamma, E.F. (2000): Management of 168 neonates weighting more than receiving intrapartum chemoprophylaxis for chorioamnionitis. Evaluation of an early discharge strategy. *Arch Pediatr Adolesc Med* 1996; 150:158-63.

Skiba, M.Y.; Maciejewska-Paszek, Pawlowskagoral K.Y.; Aleksiewicz, R. and Wardas, M. (2003): Hepatoprotective action of PGE1 analogue estimated by measuring the concentrations of acetoacetate and beta-hydroxybutyrate. *J N Engl Med*; 36:447-50.

Smith, C.A. and Crowther, C.A. (2002): Acupuncture for induction of labour. *Cochrane Database Syst Rev*; 2:CD002962.

Society of Obstetricians and Gynecologists of Canada (2001): SOGC clinical practice guidelines. Induction of labour at Term. Number 107.

Soder, H.; Bauschim, S. and Miestriner, L.A. (1990): Arterial abnormalities in talipes eqinovarus as assessed by angiography and Doppler. *J ped*; 10: 101-6.

Spanhaka, E.W.; Hyman, A.L. and Kandomtz, P.J. (1981): Bronchoactive metabolites of arachidonic acid and their role in airway function. *Prostaglandins*; 22: 113-9.

Speroff, L.; Galss, R. and Kase, N. (1999): (eds) *Clinical Gynecologic Endocrinology and Fertility. Williams & Wilkins, USA*,(4): 317-350.

- Speroff, L.; Glass, R. and Kase, N. (1994):** Prostaglandin in clinical Gynecologic Endocrinology and 5th ed. Williams and Wilkins. Maryland USA, p 291.
- Stjernholm, Y.; Sahlin, L. and Akerberg, S. (1996):** Cervical ripening in humans: potential roles of estrogen, progesterone, and insulin-like growth factor-1. Am J Obstet Gynecol; 174:1065-1071.
- Tan, B.P. and Hannah, M.E. (2001):** Prostaglandins versus oxytocin for prelabour rupture of membranes at or near term. Cochrane Database Syst Rev; Issue.
- Tang, O.S.; Gemzell-Danilsson, K. and Ho, P.C. (2007):** Misoprostol Pharmacokinetic profiles, effects on the uterus and side effects. Int. Jour, of Obs.& Gyn. 99: 160-167.
- Taylor, A.V.; Boland, J.C. and MacKenzie, I.Z. (1990):** Prostaglandin induced cervical ripening under tocolytic cover in primigravida: results of a double-blind placebo-controlled trial. Br J Obstet Gynaecol; 97:827-31.
- Thomson, A.; Telfer, J.; Kohnen, G.; Young, A.; Cameron, I.; Greer, I. and Norman, J. (1997):** Nitric oxide synthase activity and localization do not change in uterus and placenta during human parturition. Human Reproduction; 12: 2546-552.
- Tschugguel W, Schneeberger C, Lass H, Stonek F, Zaghlula MB and Czerwenka K (1999):** Human cervical ripening with an increase in cervical inducible nitric oxide synthase expression. Biological Reproduction; 60:1367-72.

Tschugguel, W.; Schneeberger, C.; Lass, H.; Stonek, F.; Zaghlula, M.B.; Czerwenka, K.; Schatten, C.; Kaider, A.; Husslein, P. and Huber, JC (1999): Human cervical ripening is associated with an increase in cervical inducible nitric oxide synthase expression. Biol Reprod., 60: 1367-1372.

Uldbjerg, N.; Ekman, G.; Herltoft, P.; Malmstrom, A.; Ulmsten, U. and Wingerup, L. (1983): Human cervical connective tissue and its reaction to prostaglandin E2. Acta Obstet Gynecol Scand; 113 (Suppl.) 16366.

Vrouenraets, F.P.; Roumen, F.J.; Dehing, C.J.; Van deli Akker, E.S.; Aarts, M.J. and Scheve, E.J. (2005): Bishop score and risk of cesarean delivery after induction of labor in nulliparous women. Obstet Gynecol; 105:690-7.

Weeks, A. and Alfirevic, Z. (2006): Oral Misoprostol Administration for Labor Induction. Clin Obstet Gynecol;49(3): 658-71.

Wikland, M.; Lindblom, B.; Wilhelmsson, L. and Wiqvist, N. (1982): Oxytoxin, prostaglandins and contractility of the human uterus at term pregnancy. Acta Obs.& Gyn.Scan 1982; 61: 4467- 4472.

William, F.; Rayburn Fredrick, P. and Zuspan, (1991): Uterine stimulation in drug therapy in Obs.& Gyn.; 3rd edition :197.

Wing, D. and Paul, R.H. (1999): Misoprostol for cervical ripening and labor induction: the clinician's perspective and guide to success. Contemplation Obstetrics and Gynecology; 44: 46.

Wing, D.A. and Paul, R.H. (1996): A comparison of differing dosing regimens of vaginally administered misoprostol for pre-induction

cervical ripening and labor induction. Am J Obstet Gynecol; 75: 158.

Wing, D.A.; Fassett, M.J. and Mishell, D.R. (2000): Mifepristone for pre-induction cervical ripening beyond 41 weeks' gestation : a randomized controlled trial. Obstet Gynecol; 96: 543.

Wing, D.A.; Jones, M.M.; Rahall, A.; Goodwin, T.M. and Paul, R.H. (1995a): A comparison of misoprostol and prostaglandin E2 gel for pre-induction cervical ripening and labor induction. Am J Obstet Gynecol ; 172: 184-6.

Wolfler, M.M.; Facchinetti, F.; Venturini, P.; Huber, A.; Helmer, H.; Husslein, P. and Tschugguel, W. (2006): Induction of labor at term using isosorbide mononitrate simultaneously with dinoprostone compared to dinoprostone treatment alone. Am J Obstet Gynecol., 195: 1617-22.

Xiao, (2003): Misoprostol and mifepristone combine to open . emergency contraception window. Contraception; 68:477-450.

Xie, Q.W.; Cho, H.J.; Calaycay, J.; Mumford, R.A.; Swiderek, K.M. Lrc, T.D.; Ding, A.; Troso, T. and Nathan, C. (1992): Cloning and characterization of inducible nitric oxide synthase from mouse macrophages. Science, 256: 225-228.

Yallampalli, C. and Garfield, R.E. (1993): Inhibition of nitric oxide synthesis in rats during pregnancy produces signs similar to those of pre-eclampsia .American Journal of Obstetrics and Gynecology; 169: 1316-20.

- Yaung, C.W.; Lee, S.H.; Kirn, Y.S. and Chang, Y.S. (1996):** Prevention of post transplant peptic ulcer by misoprostol. *Nephron*; 74: 131-5.
- Yoshida, M.; Sagawa, N.; Itoh, H.; Yura, S.; Korita, D.; Kakui, K.; Hirota, N.; Sato, T.; Ito, A. and Fujii, S. (2001):** Nitric oxide increases matrix metalloproteinase-1 production in human uterine cervical fibroblast cells. *Molecular Human Reproduction*; 7: 979-85.
- Zhang, X.; Lin, H.Y.; Liu, G.Y.; Wang, H.M.; Li, Q.L. and Zhu, C. (2005):** Expressions and regulation of endothelial and inducible nitric oxide synthases in mouse uterus during the estrous cycle and early pregnancy. *Front Bioscience*; 10:3172-182.
- Zieman, M.; Fong, S.K.; Benowitz, N.L.; Banskter, D. and Darney, P.D. (1997):** Absorption kinetics of misoprostol with oral or vaginal administration, *Obstet Gynecol*; 90: 88-92.
- Norris, L.A.; Sheppard, B.L.; Burke, G. and Bonnar, J. (1994):** Platelet activation in normotensive and hypertensive pregnancies complicated by intrauterine growth retardation. *British Journal of Obstetrics and Gynecology*; 101:209-14.
- Osman, I.; Mackenzie, F.; Norrie, J.; Murray, H.M.; Greer, I.A. and Norman, J.E. (2006):** The prim study; a randomized comparisons of prostaglandin E2 with the nitric oxide donor isosorbide mononitrate for cervical ripening before the induction of labor at term. *American Journal of Obstetrics and Gynecology*; 194: 1012-21.

Radomski, M.W.; Palmer, R.M.J. and Moncada, S. (1987): Endogenous Nitric Oxide Inhibits Human Platelet Adhesion to Vascular Endothelium. *Lancet*; 2:1057-58.