

References

References

- Aalders J, Abeler V, Kolstad P, et al: Postoperative external irradiation and prognostic parameters in stage I endometrial carcinoma: Clinical and histopathologic study of 540 patients. *Obstet Gynecol* .1980; 56:419,.
- Abeler VM, Kjorstad KE: Clear cell carcinoma of the endometrium: A histopathological and clinical study of 97 cases. *Gynecol Oncol* .1991;40:207,.
- Altgassen C, pagenstecher J, Hornung D, Diedrich K, Hornemann A. Anew approach to label sentinel nodes in endometrial cancer. *Gynecol Onco* .2007;105:457-61.
- Awtrey CS, Cadungog MG, Leitao MM, et al: Surgical resection of recurrent endometrial carcinoma. *Gynecol Oncol* . 2006;102:480,.
- Bafna UD, Umadevi K, Savitha M: Closed suction drainage versus no drainage following pelvic lymphadenectomy for gynecological malignancies. *Int J Gynaecol Carcinoma* .2001;11:143,.
- Bakkum-Gamez JN, Gonzalez-Bosquet J, Laack NN, Mariani A, Dowdy SC. Current issues in the management of endometrial cancer. *Mayo Clin Proc* 2008;83:97–112.
- Barakat RR, Goldman NA, Patel DA, et al: Pelvic exenteration for recurrent endometrial carcinoma. *Gynecol Oncol* .1999;75:99,.
- Bats AS, Clement D, Larousserie F, Le Frere-Belda MA, Pier-quet-Ghazzar N, Hignette C, et al. Does sentinel node biopsy improve the

- management of endometrial cancer? Data from 43 patients. *J Surg Oncol* .2001;97:141-5.
- Bats AS, Clement D, Larousserie F, Lefrère-Belda MA, Faraggi M, Froissart M, et al. La biopsie du ganglion sentinelle dans les cancers de l'endometre est-elle faisable? *J Gynecol Obstet Biol Reprod* 2005;34:768–74.
 - Benedet JL, Bender H, Jones H, : FIGO staging classifications and clinical practice guidelines in the management of gynecologic cancers. FIGO Committee on Gynecologic Oncology. *Int J Gynaecol Obstet* .2000;70(2):209,.
 - Benedetti, Panici, P.; Scambia, G.; Baiocchi, G.: Radical hysterectomy: a randomized study comparing two techniques for resection of the cardinal ligament *Gynecol oncol* .1993;50:226,.
 - Benedetti, Panici, P.; Maneschi, F.; Scambia, G.: A new transabdominal approach to the left retroperitoneum for systematic removal of lymph nodes left of the aorta in gynecologic malignancies. *Obstet Gynecol* .1994;83:1060,.
 - Benedetti-Panici, P.; Maneschi, F.; D'Andrea, G.: Anatomic abnormalities of the retroperitoneum encountered during aortic and pelvic lymphadenectomy. *Am. J. Obstet Gynecol*. 1994;170: 111.
 - Benedetti, Panici, P.; Maneschi, F.; Cutillo, G.: Lymphadenectomy in early epithelial ovarian carcinoma. When and how (abstract) presented at the 6th biannual meeting of the international Gynecologic carcinoma society. *Int. J. gynecol. Carcinoma* 7 (suppl 2). 1997; 17,.
 - Benedetti, Panici, P.; Maneschi, F.; Cutillo, G.: Anatomical and pathological study of retroperitoneal nodes in endometrial carcinoma. *Int. J. gynaecol. Carcinoma* .1998;8:322,.

- Benedetti, Panici, P.; maneschi, F.; Cuttillo, G.: A randomized study comparing retroperitoneal drainage with no drainage after lymphadenectomy in gynecologic malignancies. *Gynecol oncol.* .1998;65: 478.
- Benedetti-Panici, P.; Maneschi, F.; Cuttilo, G.: Staging surgery for early ovarian and tubal carcinomas. In Greer BE, Montz FJ (eds): contemporary clinical management of gynecologic malignancies (atlas of clinical gynaecology, vol 4). Philadelphia, Appleton & Lange, .1999; P. 12.1.
- Benedetti-Panici, P.; Maneschi, F; Cuttillo, G.: Tailored surgery in stage IA2-IB1 cervical carcinoma based on lymphadenectomy of primary pelvic groups: Preliminary results of a pilot study. *Int. J. Gynaecol. Obstet.* .2000;12:1,.
- Bokhman JV. Two pathogenetic types of endometrial carcinoma. *Gynecol Oncol* .1983;15:10-17.
- Boronow RC. Morrow CP. Creasman WT, et al: Surgical staging in endometrial carcinoma: Clinical-pathologic findings of a prospective study. *Obste Gynecol* .1984;63:825-832,.
- Bostick PJ, Giuliano AE Vital dyes in sentinel node localization. *Semin Nucl Med.* 2000; 30:18–2,.
- Brinton LA, Lacey JV Jr, Devesa SS, et al: Epidemiology of uterine corpus cancer. In Gershenson DM, McGuire WP, Gore M, et al (eds): *Gynecologic Cancer: Controversies in Management*. New York, Churchill Livingstone .2004; p 190.
- Bristow RE, Zerbe MJ, Rosenshein NB, et al: Stage IVB endometrial carcinoma: The role of cytoreductive surgery and determinants of survival. *Gynecol Oncol* .2000;78:85,.

- Bristow RE, Asrari F, Trimble EL, et al: Extended surgical staging for uterine papillary serous carcinoma: Survival outcome of locoregional (stage I–III) disease. *Gynecol Oncol*. 2001;81:279,.
- Burghardt, E.: Surgical treatment of cervical carcinoma. In burghardt. E. (Ed): surgical gynecologic oncology. Stuttgart, 1993.
- Burke TW, Levenback C, Tornos C, et al: Intra-abdominal lymphatic mapping to direct selective pelvic and para-aortic lymphadenectomy in women with high-risk endometrial carcinoma: Results of a pilot study. *Gynecol Oncol* .1996;62:16,.
- Chao CK, Grigsby PW, Perez CA, et al: Medically inoperable stage I endometrial carcinoma: A few dilemmas in radiotherapeutic management. *Int J Radiat Oncol Biol Phys*.1996; 34:2,.
- Chuang L, Burke TW, Tornos C, et al: Staging laparotomy for endometrial carcinoma: Assessment of retroperitoneal lymph nodes. *Gynecol Oncol* .1995;58:189-193..
- Cianciulli AM, Guadagni F, Marzano R, et al. HER-2/neu oncogene amplification and chromosome 17 aneusomy in endometrial carcinoma: correlation with oncoprotein expression and conventional pathological parameters. *J Exp Clin Cancer Res*.2003;22:265-271.
- Clagett, G.; Anderson, F.A.; Heit, J.: Prevention of venous thromboembolism. *Chest*. 1995;108(suppl) 312.
- Cochran AJ: Surgical pathology remains pivotal in the evaluation of sentinel lymph nodes. *Am J Surg Pathol*. 1999; 23:1169-1172.
- Cody HS, Hill AD, Tran KN, Borgen PI. Credentialing for breast lymphatic mapping: how many cases are enough? *Ann Surg* 1999;229:723–8

- Coleman RL, Whitten CW, O'Boyle J, et al. Unexplained decrease in measured oxygen saturation by pulse oximetry following injection of Lymphazurin 1% (isosulfan blue) during a lymphatic mapping procedure. *J Surg Oncol*. 1999;70:126–9.
- Conte, M.; Beedett-Panici, P.; Guariglia, L.: pelvic lymphocele following radical para-aortic and pelvic lymphadenectomy for cervical carcinoma: Incidence rate and percutaneous management *obstet gynecol*. 1990;76:268.
- Cosin, J.A.; Fowler, J.M.; Chen, M.D.: Pretreatment surgical staging of patients with cervical carcinoma. The case lymph node debulking. *Carcinoma*. 1998;82:2241.
- Cowles TA, Magrina JF, Materson BJ, et al. Comparison of clinical and surgical staging in patients with endometrial carcinoma. *Obstet Gynecol* .1985;66:413-416.
- Creasman WT. Morrow CP. Bundy BN, et al: Surgical pathologic spread patterns of endometrial carcinoma: A gynecologic oncology group study. *Carcinoma*. .1987;60:2035-2041.
- Creasman WT (2000): New gynecologic carcinoma staging: *Obstetric Gynecology* .2000;75:287-288.
- Creutzberg CL, van Putten WL, Koper PC, et al: The morbidity of treatment for patients with stage I endometrial carcinoma: Results from a randomized trial. *Int J Radiat Oncol Biol Phys*. 2001;51:1246.
- Creutzberg CL, van Putten WL, Warlam-Rodenhuis CC, et al: Outcome of high-risk stage IC, grade 3, compared with stage I endometrial carcinoma patients: The Postoperative Radiation Therapy in Endometrial Carcinoma Trial. *J Clin Oncol*.2004; 22:123,.

- Creutzberg CL. GOG-99: ending the controversy regarding pelvic radiotherapy for endometrial carcinoma? *Gynecol Oncol* 2004;92:740–3.doi:10.1016/j.ygyno.2004.01.009
- Dargent, D.: Laparoscopic lymphadenectomy in the management of cervical carcinoma. In benedetti-Panici, P.; Scambia, G.; Maneschi, F. et al. (eds): *Wertheim's Radical hysterectomy*. Rome, SEU Edizioni, .1996; P. 175.
- De Cicco C, Cremonesi M, Luini A, et al. Lymphoscintigraphy and radioguided biopsy of the sentinel axillary node in breast cancer. *J Nucl Med*. 1998;39:2080–4.
- de Hullu JA, Doting E, Piers DA, et al.: Sentinel lymph node identification with technetium-99m-labeled nanocolloid in squamous cell carcinoma of the vulva. *J Nucl Med* .1998;39:1381–5.
- Delaloye JF, Pampallona S, Chardonnens E, Fiche M, Lehr HA, De Grandi P, et al. Intraoperative lymphatic mapping and sentinel node biopsy using hysteroscopy in patients with endometrial cancer. *Gynecol Oncol*. 2007; 106:89-93.
- Delpech Y, Cortez A, Coutant C, et al. The sentinel node concept in endometrial cancer: histopathologic validation by serial section and immunohistochemistry. *Ann Oncol*. 2007;18:1799–1803.
- Dhar KK, NeedhiRajan T, Koslowski M, et al: Is levonorgestrel intrauterine system effective for treatment of early endometrial carcinoma? Report of four cases and review of the literature. *Gynecol Oncol* .2005;97:924.
- Di Re F, Fontanelli, R.; Raspagliesi, F.: Pelvic and para-aortic lymphadenectomy in carcinoma of the ovary. *Baillieres Clin. Obstet gynaecol*. 1989;3:311.

- DiSaia PJ, Creasman WT, Boronow RC, et al. Risk factors and recurrent patterns in stage I endometrial carcinoma. *Am J Obstet Gynecol* .1985;151:1009-101,.
- Echt ML, Finan MA, Hoffman MS, et al.: Detection of sentinel lymph nodes with lymphazurin in cervical, uterine, and vulvar malignancies. *South Med. J* 1999; 92:204–8.
- Ege GN: Internal mammary lymphoscintigraphy. The rationale, technique, interpretation and clinical application: a review based on 848 cases. *Radiology* .1976;118:101–7.
- Eltabbakh GH, Shamonki J, Mount SL: Surgical stage, final grade, and survival of women with endometrial carcinoma whose preoperative endometrial biopsy shows well-differentiated tumors. *Gynecol Oncol* .2005;99:309.
- Eric Eisenhaue and Yukio Sonoda. Surgical staging of gynecologic malignancies 1. In: Douglas A Levine, Richard R Barakat and Nadeem R Abu-Rustum editors. *Atlas of Procedures in gynecologic oncology*. second edition UK. Informa healthcare 2008.
- Eshima D, Fauconnier T, Eshima L, et al.: Radiopharmaceuticals for lymphoscintigraphy:including dosimetry and radiation considerations. *Semin Nucl Med* .2000;30:25–32.
- Eudy GE, Carlson GW, Murray DR, et al.: Rapid immunohistochemistry of sentinel lymph nodes for metastatic melanoma. *Hum Pathol* .2003;34:797–802,.
- Fanning J, Evans MC, Peters AJ, et al. Endometrial adenocarcinoma histologic subtypes: clinical and pathologic profile. *Gynecol Oncol*.1989; 32:288-291.

- Feltmate CM, Duska LR, Chang Y, et al: Predictors of recurrence in surgical stage II endometrial adenocarcinoma. *Gynecol Oncol* .1999;73:407.
- Ferenczy A, Gelfand M. The biologic significance of cytologic atypia in progestin-treated endometrial hyperplasia. *Am J Obstet Gynecol* .1989;160:126-131.
- Ferrandina G, Zannoni GF, Gallotta V, et al: Progression of conservatively treated endometrial carcinoma after full term pregnancy: A case report. *Gynecol Oncol* .2005;99:215.
- FIGO. Annual report on the results of treatment in gynecologic carcinoma. *Int J Gynecol Obstet* .1989;28: 189-193,.
- Fiorica JV, Brunetto VL, Hanjani P, et al: Phase II trial of alternating courses of megestrol acetate and tamoxifen in advanced endometrial carcinoma: A Gynecologic Oncology Group study. *Gynecol Oncol* .2004;92:10.,
- Fishman DA, Roberts KB, Chambers JT, et al: Radiation therapy as exclusive treatment for medically inoperable patients with stage I and II endometrioid carcinoma with endometrium. *Gynecol Oncol*. 1996; 61:189,.
- Fleming GF, Brunetto VL, Cella D, et al: Phase III trial of doxorubicin plus cisplatin with or without paclitaxel plus filgrastim in advanced endometrial carcinoma: A Gynecologic Oncology Group study. *J Clin Oncol*. 2004; 22:2159
- Fortier KJ. Postmenopausal bleeding and the endometrium. *Clin Obstet Gynecol*. 1986;29:440-445

- Franchi M, Chezzi F, Melpignono M, et al. Clinical value of intraoperative gross examination in endometrial carcinoma. *Gynecol Oncol.*2000; 76:357-361
- Frumovitz M, Slomovitz BM, Singh DK, Broaddus RR, Abrams J, Sun CC, et al. Frozen section analyses as predictors of lymphatic spread in patients with early-stage uterine cancer. *J Am Coll Surg* 2004a;199:388–93.
- Frumovitz M, Singh DK, Meyer L, et al: Predictors of final histology in patients with endometrial carcinoma. *Gynecol Oncol.* 2004b; 95:463.
- Frumovitz M, Bodurka DC, Broaddus RR, Coleman RL, Sood AK, Gershenson DM, et al. Lymphatic mapping and sentinel node biopsy in women with high-risk endometrial cancer. *Gynecol Oncol.* 2007; 104:100-3..
- Gal D. Hormone therapy for lesions of the endometrium. *Semin Oncol.*1986; 13:33-36
- Gargiulo T, Giusti M, Bottero A, et al.: Sentinel Lymph Node (SLN) laparoscopic assessment early stage in endometrial carcinoma. *Minerva Gyneco.*,2003; 55:259–62.
- Gerber B, Krause A, Heiner M, et al. Effects of adjuvant tamoxifen in postmenopausal women with breast carcinoma: a prospective long-term study using transvaginal ultrasound. *J Clin Onco.*,2000; 18: 3464-3470.
- Giradi, F.; Petru, E.; Heydarfadaei, M: pelvic lymphadenectomy in the surgical treatment of endometrial carcinoma. *Gynecol oncol.*1993; 49: 177.

- Giradiy F.; Lichtenegger, W.; Tamussino, K.: The importance of parametrial lymph nodes in the treatment of cervical carcinoma. *Gynecol. Oncol.* 1989; 34:206,.
- Goff BA, Goodman A, Muntz HG, et al: Surgical stage IV endometrial carcinoma: A study of 47 cases. *Gynecol Oncol.* 1999;. 52:237.
- Gordon AN, Fleischer AC, Dudley BS, et al. Preoperative assessment of myometrial invasion of endometrial adenocarcinoma by sonography (US) and magnetic resonance imaging (MRI). *Gynecol Oncol.* 1989;34:175-179..
- Gordon MD, Ireland K. Pathology of hyperplasia and carcinoma of the endometrium. *Semin Oncol.*1994;21:64-70.
- Gotlieb WH, Beiner ME, Shalmon B, et al: Outcome of fertility-sparing treatment with progestins in young patients with endometrial carcinoma. *Obstet Gynecol.*, 2003; 102:718.
- Gould EA, Philbin WT, Hyland PH, et al.: Observations on a 'sentinel node' in carcinoma of the parotid 1960.
- Grady D, Gebretsadik T, Kerlikowske K, et al. Hormone replacement therapy and carcinoma risk: a meta analysis. *Obstet Gynecol.*1995; 85:304-313.
- Hacker N.: Uterine cancer, In: Berek J, Hacker N, Editors. *Practical gynecologic oncology*. Philadelphia: Lippincott Williams & Wilkins; 2005; P. 397-442.
- Hacker, N.F.; Wain, G.V.; Nicklin, J.L.: Resection of bulky positive lymph nodes in patients with cervical carcinoma. *Int. J. Gynaecol. Carcinoma.*,1995; 5:250.

- Halsted W: *The results of radical operations for the cure of carcinoma of the breast.* Ann Surg.1997; 46:1
- Hanson MB, Van Nagell JR, Powell DE, et al. *The prognostic significance of lymph-vascular space invasion in stage I endometrial carcinoma.* Carcinoma.1985; 55:1753-1757.
- Hardesty LA, Sumkin JTH, Nath ME. et al: *Use of preoperative MR imaging in the management of endometrial carcinoma: Cost analysis.* Radiolog.2000; 215:45-49.
- Havrilesky LJ, Cragun JM, Calingaert B, et al: *Resection of lymph node metastases influences survival in stage IIIC endometrial carcinoma.* Gynecol Oncol. 2005; 99:689.
- Hauspy J, Beiner M, Harley I, Ehrlich L, Rasty G, Covens A. *Sentinel lymph nodes in early stage cervical cancer.* Gynecol Oncol 2007;105:285e90.
- Henricksen E: *The lymphatic dissemination in endometrial carcinoma: A study of 188 necropsies.* Am J Obstet Gynecol.1975;. 123:570-576.
- Henze E, Schelbert HR, Collins JD, et al.: *Lymphoscintigraphy with Tc-99m-labeled dextran.* J Nucl Med.1982; 23:923–9,
- Holub Z, Jabor A, Kliment L *Comparison of two procedures for sentinel lymph node detection in patients with endometrial cancer: a pilot study.* Eur J Gynaecol Oncol.2002; 23:53–7..
- Holub Z, Jabor A, Lukac J, Kliment L. *Laparoscopic detection of sentinel lymph nodes using blue dye in women with cervical and endometrial cancer.* Med Sci Monit. 2004;10:CR587-91.

- Holub Z, Kliment L, Lukac J, et al. Laparoscopically-assisted intraoperative lymphatic mapping in endometrial carcinoma: preliminary results. *Eur J Gynaecol Oncol.*2001; 22:118–21,
- Homesley HD, Kadar N, Barrett RJ, et al. Selective pelvic and peri-aortic lymph-adenectomy does not increase morbidity in surgical staging of endometrial carcinoma. *Am J Obstet Gynecol.* 1992; 167:1225-1230,
- Horowitz NS, Powell MA, Drescher CW, et al: Adequate staging for uterine carcinoma can be performed through Pfannenstiel incisions. *Gynecol Oncol.*2003; 88:404,.
- Hoskins PJ, Swenerton KD, Pike JA, et al: Paclitaxel and carboplatin, alone or with irradiation, in advanced or recurrent endometrial carcinoma: A phase II study. *J Clin Oncol.* 2001; 19:4048,.
- Inter Federation of Obstets & Gynecol FIGO. Classification and staging of malignant tumors in the female pelvis. *Int J Gynecol Obstet.*1971;9:172-180..
- Iversen T, Aas M: Lymph drainage from the vulva. *Gynecol Oncol* .1983; 16:179–89.
- Jemal A, Siegel R, Ward E, Murray T, Xu J, Thun MJ. Cancer statistics, 2007. *CA Cancer J Clin* 2007;57(1):43–66.
- Jobo T, Sato R, Arai T, Tamura T, Watanabe J, Kuranoto H.: Lymph node pathway in the spread of endometriqual carcinoma. *Eur J Gynaecol Oncol.* 2005; 26(2): 167-9,.
- Kadar N, Malfetano JH, Homesley HD. Determinants of survival of surgically staged patients with endometrial carcinoma histologically

- confined to the uterus: implications for therapy. Obstet Gynecol.*1992; 80:655-659,.
- Kaku T, Tsuruchi N, Tsukamoto N, et al. Reassessment of myometrial invasion in endometrial carcinoma. *Obstet Gynecol.*1994; 84:979-982
 - Kaunitz AM, Masciello A, Ostrowski M, et al. Comparison of endometrial biopsy with the endometrial Pipelle and Vabra aspirator. *J Reprod Med.* 1988; 33:427-43
 - Khoury-Collado F, Abu-Rustum NR. Lymphatic mapping in endometrial cancer: a literature review of current techniques and results. *Int J Gynecol Cancer* 2008;18 (6):1163–8.
 - Kilgore LC, Partridge EE, Alvarez RD, et al: Adenocarcinoma of the endometrium: survival comparisons of patients with and without pelvic node sampling. *Gynecol Oncol.*1995; 56:29.
 - Kodama J, Seki N, Ojima Y, Nakamura K, Hongo A, Hiramatsu Y. Risk factors for early and late postoperative complications of patients with endometrial cancer. *Eur J Obstet Gynecol Reprod Biol* 2006;124(2):222–6.
 - Kurman RJ, Kaminski PF, Norris HJ. The behavior of endometrial hyperplasia: a long term study of endometrial hyperplasia in 170 patients. *Cancer* .1985;56:403-412.
 - Larson DM, Johnson K, Olson K : Pelvic and para-aortic lymphadenectomy for surgical staging of endometrial carcinoma: Morbidity and mortality. *Obstet Gynecol.* 1992.;9:998-1001,
 - Lentz SS, Brady MF, Major FJ, et al: High-dose megestrol acetate in advanced or recurrent endometrial carcinoma: A Gynecologic Oncology Group study. *J Clin Oncol.* 1996; 14:357

- Leong SP, Donegan E, Heffernon W, et al.: Adverse reactions to isosulfan blue during selective sentinel ,2000.
- Levenback C, Coleman RL, Burke TW, et al. Intraoperative lymphatic mapping and sentinel node identification with blue dye in patients with vulvar cancer. *Gynecol Oncol.* 2001; 83:276–81.
- Lewandowski G, Torrisi J, Potkul RK, Holloway RW, Popescu G, Whitfield G, et al. Hysterectomy with extended surgical staging and radiotherapy versus hysterectomy alone and radiotherapy in stage I endometrial cancer: a comparison of complication rates. *Gynecol Oncol* 1990;36:401e4.
- Li B, Li XG, Wu LY, Zhang WH, Li SM, Min C, et al. A pilot study of sentinel lymph nodes identification in patients with endometrial cancer. *Bull Cancer.* 2007.;94:E1-4
- Liao BS, Twiggs LB, Leung BS, et al. Cytoplasmic estrogen and progesterone receptors as prognostic parameters in primary endometrial carcinoma. *Obstet Gynecol.* 1986;67:463-467..
- Lidor A, Ismajovich B, Confino E, et al. Histopathological findings in 226 women with postmenopausal uterine bleeding. *Acta Obstet Gynecol Scand.* 1986; 65:41-43,
- Lopes AD, Hall JR, Monaghan JM: Drainage following radical hysterectomy and pelvic lymphadenectomy: Dogma or need? *Obstet Gynecol.* 1995; 86:960,.
- Lundgren C, Auer G, Frankendal B, et al. Nuclear DNA content, proliferative activity, and p53 expression related to clinical and histopathologic features in endometrial carcinoma. *Int J Gynecol Cancer.*2002;12:110-118..

- Lurain JR, Rice BL, Rademaker AW, et al. Prognostic factors associated with recurrence in clinical stage I adenocarcinoma of the endometrium. *Obstet Gynecol*. 1991;78:63-69.
- Lurain JR. The significance of positive peritoneal cytology in endometrial carcinoma. *Gynecol Oncol* . 1991;46:143-144,.
- Lutman CV, Havrilesky LJ, Cragun JM, et al: Pelvic lymph node count is an important prognostic variable for FIGO stage I and II endometrial carcinoma with high-risk histology. *Gynecol Oncol*. 2006 ; 102:92
- Maccauro M, Lucignani G, Aliberti G, Villano C, Castellani MR, Solima E, et al. Sentinel lymph node detection following the hysteroscopic peritumoral injection of 99mTc-labelled albumin nanocolloid in endometrial cancer. *Eur J Nucl Med Mol Imaging* 2005;32:569–74.
- Maggino T, Romagnolo C, Landoni F, Sartori E, Zola P, Gadducci A.: An analysis of approaches to the management of endometrial cancer in North America: a CTF study. *Gynecol Oncol*, . 1998; 68:274-9.
- Manfredi R, Mirk P, Maresca G et al.,: Local-regional staging of endometrial carcinoma: role of MR imaging in surgical planning. *Radiology*. 2004; 231:372-378.
- Mariani A, Webb MJ, Galli L, et al. Potential therapeutic role of para-aortic lymphadenectomy in node-positive endometrial carcinoma. *Gynecol Oncol* . 2000a;76:348-356.
- Mariani A, Webb MJ, Keeney GL, Podratz KC. Routes of lymphatic spread: a study of 112 consecutive patients with endometrial cancer. *Gynecol Oncol* 2001;81:100e4.

- Mariani A, Webb MJ, Keeney GL, et al: Hematogenous dissemination in corpus carcinoma. *Gynecol Oncol.* 2001; 80:233.
- Mariani A, Webb MJ, Keeney GL, Haddock MG, Calori G, Podratz KC. Low-risk corpus cancer: is lymphadenectomy or radiotherapy necessary? *Am J Obstet Gynecol.* 2000b;182-1506-19.
- Marisa R. Nucci, Esther Oliva: *Gynecologic pathology: Endometrial Neoplasia Elsevier Churchill Livingstone.* 2009;233-259;.
- Martimbeau, P.W.; Kjorstad, K.E.; Kolstad, P.: Stage I carcinoma of the cervix, the Norwegian radium hospital, 1986-2970: Results of treatment and major complications: Lymphedema. *Am. J. Obstet Gynecol.* 1978; 131:389,.
- Marziale P, Atlante G, Pozzi M, et al. 426 Cases of stage I endometrial carcinoma: a clinicopathologic analysis. *Gynecol Oncol.* 1989; 32:278-281.
- Matsumoto K, Yoshikawa H, Yasugi T, Onda T, Nakagawa S Yamada M, et al. Distinct lymphatic spread of endometrial Carcinoma in comparison with cervical and ovarian carcinomas *Cancer Lett* 2002;180:83e9.
- Maxwell GL, Risinger JI, Alvarez AA, et al. Favorable survival associated with microsatellite instability in endometrioid endometrial cancers. *Obstet Gynecol.* 2001;97:417-422..
- Melhem MF, Tobon H: Mucinous adenocarcinoma of the endometrium: A clinico-pathological review of 18 cases. *Int J Gynecol Pathol.*1987 ; 6:347.
- Mettlin C, Jones G, Averette H, et al. Defining and updating the American Carcinoma Society guidelines for the carcinoma-related

- checkup: prostate and endometrial carcinomas. *CA Carcinoma J Clin*. 1993;43:42-46,.
- Molpus KL, Kelley MC, Johnson JE, et al. Sentinel lymph node detection and microstaging in vulvar carcinoma. *J Reprod Med*. 2001; 46:863–9..
 - Montz FJ, Bristow RE, Bovicelli A, et al: Intrauterine progesterone treatment of early endometrial carcinoma. *Am J Obstet Gynecol*,. 2002; 186:651.
 - Moore DH, Fowler WC, Jr, Walton LA, Droegenmueller W: Morbidity of lymph node sampling in carcinomas of the uterine corpus and cervix. *Obstet Gynecol*.1990; 74:180-184,
 - Moore RG, DePasquale SE, Steinhoff MM, et al. Sentinel node identification and the ability to detect metastatic tumor to inguinal lymph nodes in squamous cell cancer of the vulva. *Gynecol Oncol*. 2003; 89:475–9
 - Moreno-Bueno G, Hardisson D, Sanchez C, et al. Abnormalities of E- and P-cadherin and catenin (beta-gamma-catenin, and p120 ctn) expression in endometrial cancer and endometrioid cancer and endometrial atypical hyperplasia. *J Pathol*.2003;199:471-478
 - Morice P, Lassau N, Pautier P, et al: Retroperitoneal drainage after complete para-aortic lymphadenectomy for gynecologic carcinoma: A randomized trial. *Obstet Gynecol* . 2001;97:243.
 - Morice P, Joulie F, Camatte S, et al: Lymph node involvement in epithelial ovarian carcinoma: Analysis of 276 pelvic and paraaortic lymphadenectomies and surgical implications. *J Am Coll Surg*. 2003 ; 197:198.

- Morice P, Fourchotte V, Sideris L, et al: A need for laparoscopic evaluation of patients with endometrial carcinoma selected for conservative treatment. *Gynecol Oncol.* 2005; 96:245.
- Morris M, Alvarez RD, Kinney WK, et al: Treatment of recurrent adenocarcinoma of the endometrium with pelvic exenteration. *Gynecol Oncol.*1996; 60:288,
- Morrow C P, Curtin JP : *Surgical Anatomy*, in Morrow CP (ed): *Gynecologic Carcinoma Surgery*. New York, NY, Churchill Livingstone Inc. 1996.
- Morrow C P.; Curtin, J. *Surgery for cervical neoplasia*. In Morrow, P. Curtin, J. (eds): *Gynecological carcinoma surgery*. New York, Churchill Livingstone, 1996.
- Morrow CP, Bundy BN, Kurman RJ, et al: Relationship between surgical-pathological risk factors and outcome in clinical stage I and II carcinoma of the endometrium: A Gynecologic Oncology Group study. *Gynecol Oncol.* 1991; 40:55.
- Munakata S, Aihara T, Morino H, et al.: Application of immunofluorescence for intraoperative evaluation of sentinel lymph nodes in patients with breast carcinoma. *Carcinoma* . 2003;98:1562–8,.
- Mutter GL, Zaino RJ, Baak JP, et al. Benign endometrial hyperplasia sequence and endometrial intraepithelial neoplasia. *Int J Gynecol Pathol.*2007.;26:103–114,
- Mutter GL, Kandeser J, Baak JPA, et al. Biopsy histomorphometry predicts uterine myoinvasion by endometrial carcinoma. A Gynecologic OncologyGroup Study. *Hum Pathol.*2008;39:866–874..

- Nahrig JM, Richter T, Kuhn W, et al.: Intraoperative examination of sentinel lymph nodes by ultrarapid immunohistochemistry. *Breast J.* 2003; 9:277–81,.
- Niikura H, Okamura C, Utsunomiya H, Yoshinaga K, Akahira J, Ito K, et al. Sentinel lymph node detection in patients with endometrial cancer. *Gynecol Oncol* 2004;92:669e74.
- Niikura H, Okamoto S, Yoshinaga K, Nagase S, Takano T, Ito K, et al. Detection of micrometastases in the sentinel lymph nodes of patients with endometrial cancer. *Gynecol Oncol.* 2007; 105:683-6.
- Niwa K, Tagami K, Lian Z, et al: Outcome of fertility-preserving treatment in young women with endometrial carcinomas. *Br J Obstet Gynaecol.*2005. 112:317.
- Obermair A, Geramou M, Gucer F, Denison U, Graf AH, Kapshammer E, et al. Does hysteroscopy facilitate tumor cell dissemination? Incidence of peritoneal cytology from patients with early stage endometrial carcinoma following dilatation and curettage (D & C) versus hysteroscopy and D& C. *Cancer* 2000;88:139e43.
- Onda, T.; Yoshikawa, H.; Mutuzani, K. : Treatment of node positive endometrial carcinoma with complete node dissection; chemotherapy and radiation therapy. *Br. J. Carcinoma.* .1997;75:1836.
- Orr J.W.; Holimon J.L.; Orr P.F. : Stage 1 ccrupus carcinoma. Is teletherapy necessary ? *Am. J. Obstet Gynecol.* 1997; 176:777.
- Orr J, Chamberlain D: ACOG practice bulletin, clinical management guidelines for obstetrician-gynecologists, no. 65, August 2005: Management of endometrial carcinoma. *Obstet Gynecol.*2005; 106:41.,

- Ozalp S, Yalcin OT, Polay S, et al: Diagnostic efficacy of th3 preoperative lymphoscintigraphy, Ga-67 scintigraphy and computed tomography for detection of lymph node metastasis in cases with ovarian or endometrial carcinoma. *Acta Obstet Gynecol Scand*. 1999; 78:155-159.
- Palmer DC, Muir IM, Alexander AI, et al. The prognostic importance of steroid receptors in endometrial carcinoma. *Obstet Gynecol*. 1988;72:388-393..
- Parazzini F, LaVecchia C, Negri E, et al. Reproductive factors and risk of endometrial cancer. *Am J Obstet Gynecol*. 1991;64:522-527.
- Parry-Jones E: Lymphatics of the vulva. *J Obstet Gynaecol Br Commonw*. 1963;70:751–65.
- Pelosi E, Arena V, Baudino B, et al.: Preliminary study of sentinel node identification with 99mTc colloid and blue dye in patients with endometrial carcinoma. *Tumori*.2002; 88:S9–S10,.
- Pelosi E, Arena V, Baudino B, Bellò M, Giusti M, Gargiulo T, et al. Pre-operative lymphatic mapping and intra-operative sentinel lymph node detection in early stage endometrial cancer. *Nucl Med Commun* 2003;24:971–5.
- Plenti AA, Friedman EA (eds) : *Lymphatics of the Vulva, Lymphatic System of the Female Genitalia:The Morphologic Basis of Oncologic Diagnosis and Therapy*. Philadelphia, PA, Saunders, 1971.
- Plentl A, Friedman E,: *Lymphatic system of the female genitalia*. Philadelphia: WB Saunders, 1971.

- Pothuri B, Ramondetta L, Martino M, et al: Development of endometrial carcinoma after radiation treatment for cervical carcinoma. *Obstet Gynecol* . 2003;101:94.
- Ramirez PT, Frumovitz M, Bodurka DC, et al. Hormonal therapy for the management of grade 1 endometrial adenocarcinoma: a literature review. *Gynecol Oncol*. 2004; 95:133-138,
- Randall TC, Kurman RJ. Progestin treatment of atypical hyperplasia and well-differentiated carcinoma of the endometrium in women under age 40. *Obstet Gynecol*.1997; 90:434-440.
- Raspagliesi F, Ditto A, Kusamura S, Fontanelli R, Vecchione F, Maccauro M, et al. Hysteroscopic injection of tracers in sentinel node detection of endometrial cancer: a feasibility study. *Am J Obstet Gynecol* 2004;191:435e9.
- Rittenberg PV, Lotocki RJ, Heywood MS, et al: Stage II endometrial carcinoma: Limiting post-operative radiotherapy to the vaginal vault in node-negative tumors. *Gynecol Oncol*. 2005; 98:434.
- Rob L, Charvat M, Robova H, et al. Sentinel lymph node mapping in early-stage cervical cancer. *Ceska Gynekol* .2004; 69:273–7.
- Sadiq TS, Burns WW, 3rd, Taber DJ, et al.: Blue urticaria: a previously unreported adverse event associated with isosulfan blue. *Arch Surg*. 2001;136:1433–5.
- Sanjuan A, Cobo T, Pahisa J, et al: Preoperative and intraoperative assessment of myometrial invasion and histologic grade in endometrial carcinoma: Role of magnetic resonance imaging and frozen section. *Int J Gynecol Carcinoma*.2006; 16:385.

- Schink JC, Lurain JR, Wallemark CB, et al. Tumor size in endometrial carcinoma: a prognostic factor for lymph node metastasis. *Obstet Gynecol.*1987; 70:216-219.
- Schink JC, Rademaker AW, Miller DS, et al. Tumor size in endometrial cancer. *Cancer.* 1991;67:2791-2794.
- Schorge JO, Molpus KL, Goodman A, et al: The effect of postsurgical therapy on stage III endometrial carcinoma. *Gynecol Oncol.*1996; 63:34,.
- Semczuk A, Berbec H, Kostuch M, et al. K-ras gene point mutations in human endometrial carcinomas: correlation with clinicopathological features and patients' outcome. *J Cancer Res Clin Oncol*1998.;124:695-700.
- Silverberg SG, Kurman RJ, Nogales F, et al: Tumors of the uterine corpus (epithelial tumors and related lesions). In Tavassoli FA, Devilee P (eds): *World Health Organization Classification of Tumours*, Lyon, IARC Press . 2003;,. p 221,
- Smith M, McCartney AJ. Occult, high-risk endometrial carcinoma. *Gynecol Oncol* 22:154-161,1985.
- Smith RA, Cokkinides V, von Eschenbach AC, et al. American Cancer Society guidelines for the early detection of cancer. *CA Cancer J Clin.* 2002.;52:8-22.
- Sovak MA, Hensley ML, Dupont J, et al: Paclitaxel and carboplatin in the adjuvant treatment of patients with high-risk stage III and IV endometrial carcinoma: A retrospective study. *Gynecol Oncol.* 2006; 103:451,

- Suzuki M, Ohwada M, Sato I: Pelvic lymphocysts following retroperitoneal lymphadenectomy: Retroperitoneal partial "no-closure" for ovarian and endometrial carcinomas. *J Surg Oncol.* 1998; 68:149.
- Tavares MG, Sapienza MT, Galeb NA, Jr, et al. The use of 99mTc-phytate for sentinel node mapping in melanoma, breast cancer and vulvar cancer: a study of 100 cases. *Eur J Nucl Med.*2001; 28:1597–604.
- Thigpen JT, Brady MF, Alvarez RD, et al: Oral medroxyprogesterone acetate in the treatment of advanced or recurrent endometrial carcinoma: A dose-response study by the Gynecologic Oncology Group. *J Clin Oncol* 1999; 17:1736,.
- Turner DA, Gershenson DM, Atkinson N, et al. The prognostic significance of peritoneal cytology for stage I endometrial carcinoma. *Obstet Gynecol.*1989; 74:775-780.
- Uren RF, Howman-Giles R: The role of nuclear medicine. In: Cody HS, ed. *Sentinel lymph node biopsy* London: Martin Dunitz: .2002;19–43.
- Wang CB, Wang CJ, Huang HJ, et al: Fertility-preserving treatment in young patients with endometrial adenocarcinoma. *Carcinoma.* 2002; 94:2192,
- Whitney CW, Stehman FB: The abandoned radical hysterectomy: A Gynecologic Oncology Group study. *Gynecol Oncol.* 2000; 79:350.
- Whitney CW, Brunetto VL, Zaino RJ, et al: Phase II study of medroxyprogesterone acetate plus tamoxifen in advanced endometrial carcinoma: A Gynecologic Oncology Group study. *Gynecol Oncol.*2004; 92:4,

- Whitney CW: *GOG Surgical Procedures Manual*. Gynecologic Oncology Group, 2005.
- Wong JH, Cagle LA, Morton DL: *Lymphatic drainage of skin to a sentinel lymph node in a feline model*,1991.
- Yang YC, Wu CC, Chen CP, et al: *Reevaluating the safety of fertility-sparing hormonal therapy for early endometrial carcinoma*. *Gynecol Oncol*. 2005; 99:287.
- Yokoyama Y, Maruyama H, Sato S, Saito Y : *Indispensability of pelvic and para-aortic lymphadenectomy in endometrial cancers*. *Gynecol Oncol*,. 1997; 64:411-7
- Zaino RJ, Kurman RJ, Diana KL, et al. *Pathologic models to predict outcome for women with endometrial adenocarcinoma*. *Carcinoma*.1996; 77:1115-1121.
- Zaino RJ, Davis ATL, Ohlsson-Wilhelm BM, et al. *DNA content is an independent prognostic indicator in endometrial adenocarcinoma*. *Int J Gynecol Pathol*. 1998;17:312-319.
- Zerbe MJ, Bristow R, Crumrine FC, et al. *Inability of preoperative computed tomography scans to accurately predict the extent of myometrial invasion and extracorporeal spread in endometrial carcinoma*. *Gynecol Oncol*.2000; 78:67-70.