

References

- **Abudu N., Levinson S. and Miller J. (2003):** Triglycerides/HDL-c levels versus ApoA-1 and apoB in Coronary Artery Disease (CAD) for recognition of Familial Combined Hyperlipidemia (FCH). Journal of the American Diabetes Association; 52(1): F67.
- **Adams-Chapman I and Stoll BJ. (2006):** Neonatal infection and long-term neurodevelopmental outcome in the preterm infant. Curr Opin Infect Dis;19(3):290-297.
- **Aitken C. and Jeffries DJ. (2001):** Nosocomial spread of viral disease. Clinical Microbiology Review. July 2001; 14(3): 528-546.
- **American Academy of Pediatrics (2003):** Risk factors for nosocomial infection in a neonatal Intensive Care Unit . J Hosp infecu 53 (1) :25-30.
- **Anderson B., Linda L. and Bryan L. (2006):** Neonatal sepsis, from e-medicine, webMD.
- **Assman G., (1988) :** Triglycerides and atherosclerosis. Lipid Rev; 2:89-96.

- **Assman G., Paoletti R., Austin MA. et al. (1991):** Hypertriglyceridemia risk and management. *Am J Cardiol*; 68(3): 1-42.
- **Assman G., Von Eckardstein A and Funke H. (1992):** HDL-C, reverse cholesterol transport and CHD. *Circ*; 85:132.
- **Austgulen R., Arntzen KJ., Haereid PE. et al. (1997):** Infections in neonates delivered at term are associated with increased serum levels of ICAM-1 and E-selectin. *Acta Paediatrica*. 86, (3), 274-280.
- **Baltimore RS. (1998):** Neonatal nosocomial infections. *Semin Perinatal* ; 22:25 -32.
- **Baquero Latorre H., Neira Safi F and Gonzalez Vargast (2006):** Endogenous endophthalmitis due to *Serratia marcescens* in the course of early neonatal sepsis. *An Pediat*; 64(3):291-292.
- **Barlage S., Gnewuch C., Liebisch et al. (2009):** Changes in HDL- associated apolipoproteins relate to mortality in human sepsis and correlate to monocyte and platelet activation. *Intensive Care Med*.;35(11):1877-85.
- **Baron S. (2003):** Bacteriology- *Hemophilus* species. *Medical microbiology*, 4th edition, Univ of Texas.

- **Barbara J. Stoll, (2003):** Infections of the neonatal infant, Nelson textbook of pediatrics, Behrman R., Kliegman R. and Jenson H. (eds) 17th Edition, WB Saunders Company ,Philadelphia., 623-640.
- **Belling L. (2004):** Neonatal sepsis. E-medicine. <http://www.emedicine.com/ped/topic2630.htm>.
- **Bhutta K and Yusu F. (1997):** Neonatal sepsis in Karachi factors determining outcome and mortality. J Trop Pediatr; (43):65-70.
- **Bilal Y., Birsan U., Arif A. et al. (2009):** Diagnostic values of lipid and lipoprotein levels in late onset neonatal sepsis. Scandinavian Journal of Infectious Diseases, 2009: 41: 263-267.
- **Boland JAV., Kuhu M. and Berch P. (2001):** Listeria pathogenesis and molecular virulence determinants. Clin Micro Rev; 14 (3): 584-640.
- **Boyce J. and Pittet D. (2002):** Guidelines for hand hygiene in health-care settings: Recommendations of the healthcare infection control practices advisory committee and the HICPAC/SHEA/APIC/IDSA hand hygiene task force. MMWR;51:1-44.
- **Breslow JL. (1995):** Familial disorders of high-density lipoprotein metabolism. In: The Metabolic and Molecular

Bases of Inherited Disease. 7th ed., Scriver CR, Beaud, Sly WS and Valle D (eds.). New York, McGraw-Hill; 2: 2031-52.

- **Brew HB., Gregg RE. , Hoeg JM. et al. (1988):** Apolipoprotein A-1 and lipoproteins in human plasma. An overview. Clin Chem; 34: 4.
- **Brewer, H.B. Jr., T. Fairwell et al. (1978):** The amino acid sequence of human Apo A-1, an apolipoprotein isolated from high density lipoproteins. Biochem. Biophys. Res. Commun 80:623-630.
- **Brown MS. and Goldstein JL. (1986):** A receptor mediated pathway for cholesterol homeostasis. Science; 232:4.
- **Burger D. and Dayer JM. (2002):** High-density lipoprotein-associated apolipoprotein A-1: the missing link between infection and chronic inflammation. 2002 Feb;1(1-2):111-7.
- **Byington CI., Enriquez FR .and Hoff C. (2004):** Serious bacterial infections in febrile infants 1 to 90 days old with and without viral infections. Pediatrics ;113 (6): 1662-6.
- **Calvert GD. and Abbey M. (1985):** Plasma Lipoproteins, apolipoproteins A-1 and proteins concerned with lipid metabolism. Adv Chem; 24:217-98.

- **Carl A., Burtis PhD and Edward R. et al. (1999):** Lipids, lipoproteins and apolipoproteins. In: Tietz Textbook of Clinical Chemistry. Carl A, Burtis PhD, Edward R, Ashweod MD (eds.). 3rd Edition, WB Saunders Company, Philadelphia; 809-61.
- **Carrier MP., Stolfi I. and Moro ML. (2003):** Intercenter variability and time of onset: two crucial issues in the analysis of risk factors for nosocomial sepsis. *Pediatr Infect Dis J*; 22 (7) : 599-609.
- **Casanova- Roman M., Rios J., Sanchez-Porto A. et al. (2002):** Early-onset of neonatal sepsis due to group A streptococcus. *Minerva – Pediatr*; 54 (2): 161-3.
- **Centers for Disease Control and Prevention (CDC). (2002):** Prevention of perinatal group B streptococcal disease: revised guidelines from CDC. *Morbidity and Mortality Weekly Report*, volume 51, no. RR-11, August 16.
- **Chacko B. and Sohi I. (2005):** Early onset neonatal sepsis. *Indian J Pediatr*; 72(1):23-26.
- **Chapman MJ., Caslake M. , Packard C. et al. (2003):** New dimension of statin action on apoB atherogenicity. *Clin Cardiol*: 1(1): 17-20.

- **Chenaud, Catherine MD., Merlani et al. (2004):** Low apolipoprotein A-1 level at intensive care unit admission and systemic inflammatory response syndrome exacerbation. *Critical Care Medicine*; 32- (3)- 632-637.
- **Chiesa C., Panero A., Rassi N. et al. (1998):** Reliability of procalcitonin concentrations for the diagnosis of sepsis in critically ill neonates. *Clin Infect Dis* 26: 664-72.
- **Claudio Chiesa, Alessandra Panero, John F. et al. (2004):** Diagnosis of neonatal sepsis: A clinical laboratory challenge. *Clinical chemistry*; 50: 279-287.
- **Dar-Shang L., Shing-Guey H., Chun-Chun L. et al. (2000):** Urinary Tract Infection in febrile infants younger than eight weeks of age *pediatrics*;105(2):20.
- **Davis RA. (1991):** Lipoprotein structure and secretion. In: *Biochemistry of Lipids, Lipoproteins and Membranes*. Vance DE. And Vance J. (eds). Elsevier Science Publishers, BV. Chapter 14.
- **Easterbrook PJ. (1999):** Lipids and metabolism. In: *Basic Medical Science*. 2nd ed, Easterbrook (ed.). Churchill Livingstone, Harcourt Brace and Company Limited; 243-54.

- **Edmond K. and Zaidi A. (2010):** New Approaches to Preventing, Diagnosis and Treating Neonatal Sepsis. Plos Med 7(3):e1000213.
- **Farmer JA. and Gotto AM .(1992):** Risk factors for coronary artery disease. In: Heart Disease: A Textbook of Cardiovascular Medicine., 4th ed. Farmer JA and Gotto AM (eds.). WB Saunders Company, Philadelphia, London; 1124.
- **Fergany AL. (2006):** Serum cortisol and thyroid hormone levels in neonates with sepsis. Master Thesis, Faculty of Medicine, Ain-Shams University.
- **Fjaertoft G., Harkanson L., Ewald U. et al. (1999):** Neutrophils from term and newborn infants express the high affinity Fc gamma-receptor 1 (CD64) during bacterial infection. Paediatr Res; 45:871-6.
- **Fowlie P. and Schmidt B. (1997):**diagnosis test for bacterial infection from birth to 90th days. Arch. Dis. Child. (Fetal Neonatal Edition); 78(2): 92-8.
- **Garges HP., Moody MA. and Cotton CM. (2006):** Neonatal meningitis what is the correlation among cerebrospinal fluid cultures. Blood cultures. And cerebrospinal fluid parameters?. Pediatrics Apr; 117(4): 1094-100 graham PL, Begg MD and Larson E (2006):

Risk factors for late onset gram negative sepsis in low birth weight infants hospitalized in the neonatal intensive care unit. *Pediatr infect Dis J* Feb; 25(2): 113-7.

- **Gerald L., Marie B. , Jean F. et al. (2002):** Value of HDL-C, Apolipoprotein A-1, lipoprotein A and lipoprotein A1/A11 in prediction of coronary heart disease. *Arterioscler Thromb Vasc Biol*; 22(7): 1155-61.
- **Gerber Y., Goldbourt U. and cohen H. et al. (2002):** Association between serum apolipoprotein C (II) concentration and coronary heart disease. *Prev Med*; 35(1): 42-7.
- **Gerdes JS. (2004):** Diagnosis and management of bacterial infections in neonate. *Pediatr Clin N Am*; 51:939-959.
- **Glew HR. (2002):** Lipid metabolism. II. Pathways of metabolism of special lipids. In: *Textbook of Biochemistry with Clinical Correlations*. 5th ed. Devlin (ed). Wiley-liss; 728.
- **Goldstein JL. And Brown MS. (1989):** The low density lipoprotein receptor pathway and its relation to atherosclerosis. *Ann Rev Biochem*; 46:897-930.
- **Golombek S., Rohan A., Parvez B. et al. (2002):** Proactive management of percutaneously inserted central

catheters results in decreased incidence of infection in the ELBW population. *J Perinatol*;22:209-213.

- **Gordon, BR., Parker TS., Levine DM., et al., (2001):** Relationship of hyperlipidemia to cytokine concentrations and outcomes in critically ill surgical patients. *Crit. Care Med.* 29:1563-1568.
- **Gottof SP. (2000):** Infections of the neonatal infant. In *Nelson Textbook of Pediatrics*. Behrman R.E kleigman R.M. and Jenson h. B. (eds.) 16th Edition, WB Saunders Company Pp. 538-552.
- **Gowenlock AH., McMurray JR. and McLauchlan DM. (2002):** Lipids and lipoproteins. In: *Varley's Practical Clinical Biochemistry*. 6th ed. Gowenlock AH, McMurray JR and McLauchlan DM (eds.). CBS Publishers and Distributors, India,;454-5.
- **Graham PL. and Begg MD. (2006):** Neonatal sepsis. *Pediatr Infect Dis J.* 117(4):664-72.
- **Guibourdenche J., Bedu A., Petzold L. et al. (2002):** Biochemical markers of neonatal sepsis: value of procalcitonin in the emergency setting. *Ann Clin Biochem.* 39(2):130-5.

- **Harvas JA., Ballesteros F .and Alomar A .(2001):** Increase of enterobacter in neonatal sepsis: A twenty two year study. *Ped Infec Dis J*:20(2): 134-40.
- **Haslett C., Edwards C., Chilvers E. et al. (2003):** Diseases due to infection. *Davidson's Principles and Practice of Medicine*, 19 th ed., 65-71.
- **Havel RJ., Goldstein JL. And Brown MS (1980):** Lipoproteins and lipid transport. In: *Metabolic Control and Disease*. Bondy and Rosenberg LK (eds). Philadelphia, WB Saunders; 393-494.
- **Heng CK., Low PS. and Saha N. (2001):** Variations in the promoter region of the Apolipoprotein A-1 gene influence plasma lipoprotein (a) levels in Asian Indian neonates from Singapore. *Pediatr Res*; 49: 514-8.
- **Hintz SR., Kendrick DE., Stoll BJ. et al. (2005):** Neurodevelopmental and growth outcomes of extremely low birth weight infants after necrotizing enterocolitis. *Pediatrics*. 2005;115(3):696-703.
- **Hoover-plow J., Hart E., Gong Y. et al. (2009):** A physiological Function for Apolipoprotein (a): A Natural Regulator of the Inflammatory Response. *Biol, Med*; 234:28-34.

- **Howard MR., and Smith RA. (1999):** Early diagnosis of neonatal sepsis using a hematological scoring system. *J Matern Fetal Med.* 1999 May-Jun;8(3):88-94.

- **Judith D., Nicki R., and Brenda D. (2004):** Impact on the Incidence of Nosocomial Infections in the Newborn Intensive Care Unit. Neonatal Intensive Care Unit, West Virginia University Children'S Hospital, Morgantown, WV, USA.

- **Kenner C. and Lott J. (2003):** Assessment and management of the immune system. In *Comprehensive Neonatal Nursing.* Gewolb I., Vice H., Frank L., et al (eds) Saunders, Philadelphia, 3rd Edition, 550-579.

- **Khovidhunkit W., Kim MS., Memon RA. et al. (2004):** Effects of infection and inflammation on lipid and lipoprotein metabolism: mechanisms and consequences to the host. *J. Lipid Res.* 45: 1169-1196.

- **Kilbride H., Wirtschafter D., Powers R. et al. (2003):** Implementation of evidence-based potentially better practices to decrease nosocomial infections. *Pediatrics;* 111:519-533.

- **Klein JO. and Marcy SM. (1995):** Bacterial sepsis and meningitis. In: Remington JS, Klein JO (eds). *Infectious*

disease of the fetus and the newborn infant. Philadelphia: WB Saundeers Company, 4th Edition, 835-90.

- **Kloos WE. and Bannerman TL. (1994):** Update on clinical significance of coagulase-negative staphylococci. Clin Micro Rev;7:117-140.
- **Krauss RM., Perlman JA. and Petitti D. (1988):** Effects of estrogen dose and smoking on lipid and lipoprotein levels in the postmenopausal women. Am J Obstet Gynecol; 158:1606-11.
- **Law YL. and Hey EN. (1991):** Sensitivity and specificity of daily tracheal aspirate culture in predicting organisms causing bacteremia in ventilated neonates. Pediatr infect Dis J; 10(4):290-294.
- **Liberopoulos E., Apostolou F., and Elisaf M. (2004):** Serum Lipid profile in patients with severe Leptospirosis. Nephrol. Dial. Transplant. 19:1328-1329.
- **Linda L. and Bryan L. (2004):** Track coordinator, instructor neonatal Nurse Practitioner Program, Medical university of South Carolina, College of nursing.
- **Litman G., Cannon J. and Dishaw L. (2005):** Reconstructing immune phylogeny: new perspectives. Nat Rev Immunol 5(11):866-79.

- **Lizewsky SE., Lundberg DS. and Schyrr UJ. (2002):** The transcriptional regular Alg R is essential for *Pseudomonas aeruginosa* pathogenesis. *Infect and Immun*; 70(11):6083-6093.
- **Lu S., Yao Y. , Meng S.et al. (2002):** Overexpression of Apolipoprotein A-1-IV enhances lipid transport in newborn swine intestinal epithelial cells. *J Biol Chem*; 277:31929-37.
- **Lucos JF., Brabley SM. , Eicher D. et al. (2005):** Fatal complete atrioventricular block as a complication of bacterial sepsis in a premature newborn. *Pediatr cardiol*; 26(2):677-9.
- **Martin M. meremikwu., Chukwuemcka E., et al. (2005):** Bacterial isolates from blood cultures of children with suspected septicemia in Calabar, Nigeria. *BMC Infect*;5:110.
- **Mathers NJ. and Pohlandt F. (1987):** Diagnostic audit of C- reactive protein in neonatal infection. *Eur J Pediatr*;146:147-51.
- **Mayes PA. (1993):** Lipid transport and storage. In: *Harper's Biochemistry*. 23rd ed. Murraroy RK, Mayes PA, Granner DK and Rodvell VW (eds). Appleton and Lange ;251-76.

- **Mc Intosh N., Peter H. and Rosalind S. (2003):** Infection and immunity in the newborn, Forfar& Arneil's Textbook of pediatrics, Peter J., Helms, Neil Mc Intosh, et al (eds), Elsevier Health Sciences, London, 6th edition, 336-353.
- **Mckenney W. (2001):** Neonatal nursing. Understanding the neonatal immune system: high risk for infection. Critical Care Nurse; 21(6),35-47.
- **McPherson RJ. and Juul S. (2005):** Patterns of thrombocytosis and thrombocytopenia in hospitalized neonates. Journal of perinatology; 2:166-172.
- **Merck (2005):** Neonatal infections. The Merk Manual of Diagnosis and Therapy website. Available at: www.merk.com/mrkshared/mmanual/section19/ch.
- **Navab M., Roger YU., Gharavi N. et al. (2007):** High-density lipoprotein: antioxidant and anti-inflammatory properties. Curr Atheroscler Rep 9(3):244-8.
- **Ng PC. and Lam HS. (2006):** Diagnostic marker for neonatal sepsis. Curr Opin pediatr; 18(2):125-31.
- **Ng PC., Leung TF., Wong RP. et al. (2006):** Early prediction of sepsis induced Disseminated intravascular coagulopathy with interleukin 10 interleukin-6 and RANTES in preterm infants. Clin Chem; 13.

- **O'Grady N., Alexander M., Dellinger E. et al. (2002):** Guidelines for the prevention of intravascular catheter-related infections. *MMWR*; 51:1-26.
- **Ottolini M., Lungren K., Cason S. et al. (2003):** Utility of complete blood count and blood culture screening in the diagnosis of neonatal sepsis. *Pediatr Infect Dis J*.17:22-235.
- **Petit J. (2002):** Assessment of infants with peripherally inserted central catheters : Part 1. *Adv Neonatal Care*; 2:204-314.
- **Podschun R. and Ullmann U. (1998):** *Klebsiella* as nosocomial pathogen: Epidemiology, taxonomy, method and pathogenicity factors. *Clin Micro Rev*; 11(4):589-603.
- **Prats JA., Cooper TR., Schneider VF. et al. (2000):** Rapid detection of microorganisms in blood cultures of newborn infants. *Pediatr*; 105(30):523-7.
- **Radunovic N., Kuczynski E. , Rosen et al. (2000):** Plasma Apolipoprotein A-1 and B concentrations in growth-retarded fetuses, a link between low birth weight and adult atherosclerosis. *J Clin Endocrinol Metab*; 85(1):85-8.
- **Remington S., Klein JO., Tenover JC. et al. (2005):** Infectious diseases of the fetus and newborn infants, 6th edition.

- **Rieger- Fackeldey E., Aumeier S .and Genzel-Boroviczeny O. (2000):** Disseminated adenovirus infection in two premature infants. *Infect*; 28(4):237-9.
- **Rifai N., Bachorik S. and Albers J. (1999):** Lipids, lipoproteins, and apolipoproteins. In: *Tietz Textbook of Clinical Biochemistry*, 3rd ed. Burlis AC and Ashwood RE (eds). WB Saunders Company, America, Chap. 25;809-60.
- **Roberts IA. and Murray NA. (2003):** Thrombocytopenia in the newborn. *Curr Opin pediatr*; 15(1):17-23.
- **Rowen JL., Judy M., Tate I. et al. (1999):** Candida isolate from neonates: Frequency of miss identification and reduced fluconazol susceptibility. *Clin Microb J*; 37(11):3735-3737.
- **Saez X. and McCracken GH. (2006):** Clinical pharmacology of Antibacterial Agents infectious diseases of the fetus and newborn infant, 4th ed. 1294-1327.
- **Sanker MJ., Agarwal R., Deorari AK. et al. (2008):** Sepsis in the newborn. *Indian J Pediatr*. 2005 Mar; 65(3):261-6.
- **Segrest JP. (2000):** Non-LDL lipid risk factors: HDL, triglycerides, small dense LDL-c and Lp(a). *Clin Rev Spring*; 23-8.

- **Sithembiso Velophi MB., Jane D., Siegel MD. et al. (2003):** Early-onset Group B Streptococcal Infection After a Chemoprophylaxis Strategy. *PEDIATRICS* VOL. 111:(3), 541-547.
- **Springer TA. (1990):** Adhesion receptors of the immune system . *Nature* 346:425-434.
- **Stein EA. and Gray LM. (1994):** Lipids, lipoproteins and apolipoproteins. In: Tietz N (ed). *Textbook of Clinical Chemistry*, 2nd ed. Burtis and Ashwood ER (eds). Philadelphia: WB Saunders Co; 1002.
- **Stein EA. and Myers GL. (1994):** Lipids, lipoproteins and apolipoproteins. In: Tietz *Textbook of Clinical Chemistry*. 5th ed. Edward R and Ashwood (eds). WB Saunders Company; 1019-32.
- **Stokowski LA. (2003):** JCAHO issues sentinel event alert about death from nosocomial infection. *Adv Neonatal Care*; 3:49.
- **Stoll BJ. and Hansen N. (2003):** Infections in VLBW infants: Studies from NICHD Neonatal Network Seminars in Perinatology; 27:293-301.
- **Szymankiewicz M., Oszkinis G. and Uchman J. (2005):** Abdominal aortic iliac thrombosis as a complication of newborn's bacterial sepsis. *I grinekol pol*; 76(1):61-6.

- **Tollner U. (1982):** Early diagnosis of septicemia in newborn. Clinical studies and sepsis score Eur. J pediatric. 138 (10): 331-7.
- **Vaciloto E., Richtmann R., Kusano EJ. et al . (2002):** A survey of the incidence of neonatal sepsis by group B Streptococcus during a decade in a Brazilian maternity hospital. Braz J Infect Dis. 2002 Apr.8(2):55-82.
- **Vartzelis G., Theodoridou M., Daikos GL. et al. (2005):** Brain abscesses complicating staphylococcus aureus sepsis in a premature infant. Infectio; 33(1):36-8.
- **Vjekoslava R. and Jeren S. (2002):** Dyslipoproteinemia and coronary disease. Angiology; 53(5):557-62.
- **Walmsley RN. and White GH. (1994):** Plasma lipids. In: A guide to diagnostic clinical chemistry, 3rd ed. Walmsley RN and White GH (eds). Oxford, Blackwell Scientific Publication, Chap; 13:348-67.
- **Warnick RG., McNamara RJ., Wu LL. et al. (1996):** Lipids and lipoproteins. In: Clinical Chemistry Principles, Procedures and Correlations, 3rd ed. Bishop ML, Duben, Engrelkirk JL and Fody EP (eds). Lippincott-Raven Publishers, Philadelphia and New York, Chap; 15:313-33.
- **Weinberg GA., and Powell KR. (2001):** Laboratory aids for diagnosis of neonatal sepsis. In: Remington JS and

Klein JO, editor. Infectious diseases of the fetus and newborn infant. fifth. Philadelphia, Saunders; 1327–1344.

- **Whicher J., Bienvenu J., and Monneret G. (2001):** Procalcitonin as an acute phase marker. *Ann Clin Biochem*; 38:483-93.
- **Wu A., Hinds CJ., and Thiemermann C. (2004):** High density lipoproteins in sepsis and septic shock: metabolism, actions, and therapeutic applications. *Shock* 21: 210-221, 2004.
- **Wu LH., Wu JT., and Hopkins PN. (1994):** Apolipoprotein E. Laboratory determination and clinical significance. In: *Methods for Clinical Laboratory Measurement of Lipid and Lipoprotein Risk factors*, 2nd ed. Rifia N, Warnick GR (eds). Washington, DC , AACC Press; 279.
- **Zhenghao Z., Geela D., Yun Z. et al. (2009):** Apolipoprotein A-I mimetic peptide treatment inhibits inflammatory responses and improves survival in septic rats. *Am J Physiol Heart Circ Physiol*. 2009 August ; 297(2):H866-H873.