

References

1. **Accardo J, Kammann H, Hoon AH Jr (2004):** Neuroimaging in cerebral palsy. *J Pediatr.* 2004 Aug;145(2 Suppl):S19-27.
2. **Alexander G.R. (1997):** The accurate measurement of gestational age a critical step towards improving fetal death. *Am. J. Public Heath*; 87:1278.
3. **Als H, Duffy FH, McAnulty GB, et al.** Early experience alters brain function and structure. *Pediatrics* 2004;113(4):846–57.
4. **Als H, Gilkerson L, Duffy FH, et al.(2003):** A three-center, randomized, controlled trial of individualized developmental care for very low birth weight preterm infants: medical, neurodevelopmental, parenting, and caregiving effects. *J Dev Behav Pediatr* 2003;24(6):399–408.
5. **Aly H, Milner JD, Patel K, El-Mohandes AA.(2004):** Does the experience with the use of nasal continuous positive airway pressure improve over time in extremely low birth weight infants? *Pediatrics.* 2004 Sep; 114(3):697-702.
6. **Amess, P., C. McFerran, et al. (2009).** "Early prediction of neurological outcome by term neurological examination and cranial ultrasound in very preterm infants." *Acta Paediatr* 98(3): 448-53.
7. **Andrew B. Onderdonk and Mei-Ling Lee.(2003):** Quantitative microbiologic models for preterm delivery .*J Clin Microbiol.* March; 41(3): 1073–1079., American Society for Microbiology.
8. **Ann R, Li-Yin C, Abraham P, Ranjita B, and Shoo K. (2001):** Variations in intraventricular haemorrhage incidence rates among Canadian neonatal intensive care units. *J Pediatr*; 138(4):1-9.
9. **Attia N, Taman K, Abdel Rahman Z, and Faris L (1997):**Role of SPECT and cranial sonography in the

- evaluation of brain insult in perinatal asphyxia . The Egyption Journal of pediatrics ;14(4)619-631.
10. **Aurora Sand Snyder Y Evan(2004):**perinatal asphyxia. In: Manual of Neonatal Care.Cloherty J . P. and Stark A. P. (eds);4th edition, Little Brown and Lippincott-Raven Co.27:536 -555.
 11. **Badiee, Z. (2007).** "Intraventricular hemorrhage in very low birth weight infants. Associated risk factors in Isfahan, Iran." Saudi Med J 28(9): 1362-6.
 12. **Bassan. H. (2009):** Intracranial Hemorrhage in the Preterm Infant: Understanding It, Preventing It. Clin Perinatol 36 (2009) 737–762.
 13. **Beherman RE, Kliegman RM, Arvin, Nelson WE. (2000):** Respiratory distress syndrome .In Behrman RE, Kliegman RM,Jenson HB Nelson textbook of Pediatrics,16th edition Philadiphia , London Toronto,WB Saunders company 478:-480.
 14. **Beverley DW, Pitts-Tucker TJ, Congdon PJ, et al.(1985):** Prevention of intraventricular haemorrhage by fresh frozen plasma. Arch Dis Child 1985;60(8):710–3.
 15. **Blankenberg FG, Loh NN, Bracci P, et al.(2000):** Sonography, CT, and MR imaging: a prospective comparison of neonates with suspected intracranial ischemia and hemorrhage. AJNR 2000;21(1):213-218.
 16. **Boal DK, Watterberg KL,Miles S ,Gifford KL.(1995):** Optimal cost effective timing of cranial ultrasound screening in low birth weigh infants. Pediatr. Radiol;25:425-8.
 17. **Born CM, Meisenzahl EM, Frodl T, Pfluger T, Reiser M, Moller HJ, Leinsinger GL. (2004):** The septum pellucidum and its variants. An MRI study. Eur. Arch Psychiatry Clin. Neurosci.2004 Oct; 254(5):295-302.
 18. **Brangenberg R, Bodensohn M, Burger U.** Antithrombin-III substitution in preterm infants: effect on intracranial hemorrhage and coagulation parameters. Biol Neonate 1997;72(2):76–83

19. **Brion LP, Bell EF, Raghuveer TS.(2003):** Vitamin E supplementation for prevention of morbidity and mortality in preterm infants. *Cochrane Database Syst Rev*2003;(4):CD003665.
20. **Burguet A, Kaminski M and Abraham-Lerat L.(2004):** The Complex Relationship between Smoking in Pregnancy and Very Preterm Delivery. Results of the Epipage Study. *Br J Obstet& Gynaecol*; 111:258-265.
21. **Caravale B, Allemand F, Libenson MH. (2003):** Factors predictive of seizures and neurologic outcome in perinatal depression. *Pediatr Neurol*. 2003 Jul; 29(1):18-25.
22. **Chang S C, O'Brien K O, Nathanson M S, Mancini J and Witter F R.(2003):** Characteristics and risk factors for adverse birth outcomes in pregnant black adolescents. *J Pediatr*; 143:250-257.
23. **Chang YC, Huang CC, Wang ST, Chio CC. (1999):** Risk factor of complications requiring neurosurgical intervention in infants with bacterial meningitis. *Pediatr Neurol*. 1999 Sep;17(2):144-9.
24. **Changlain, Z., X., Xiaoyang,W., (2003):** Different apoptotic mechanisms are activated in male and female brains after NHIE. *J. Neuro.Chem.*;96:1061:1091.
25. **Chen C, Hus J and Hsieh T.(1997):** Analysis of birth weight and gestational age. *Taiwn J for Mosan Med assoc*; 90:382-387.
26. **Chess P, Chess M, Manuli M, and Guillet R. (1997):** Screening head ultrasound to detect intraventricular haemorrhage in premature infants. *Pediatr Radiol*,27(4):305-8.
27. **Cohen HL. (1996):** Neurosonography of the infant:diagnosis of abnormalities. In Timor-tritsch IE, Monteagudo A, and Cohen HL, *ultrasonography of the prenatal and neonatal brain*. Stamford: Appleton &Lange, 259-85.

28. **Cools F, Offringa M.(2000):** Neuromuscular paralysis for newborn infants receiving mechanical ventilation. *Cochrane Database Syst Rev* 2000;(4):CD002773.
29. **Correa F, Enriquez G, Rossello J, Lucaya J, Piqueras J, Aso C, Vazquez E, Ortega A, Gallart A.(2004):** Posterior fontanelle sonography: an acoustic window into the neonatal brain. *AJNR Am. J. Neuroradiol.* 2004 Aug; 25(7):1274-82.
30. **Crowly P. (2000):** Prophylactic corticosteroids for preterm birth. In: *The Cochrane Library*, Issue 1, Oxford.
31. **Crowley P. WITHDRAWN:(2006):** prophylactic corticosteroids for preterm birth. *Cochrane Database Syst Rev* 2006;(3):CD000065.
32. **Curtis P.D., Matthews T.G., Clarke T.A., et al. (1998):** Neonatal seizures; the Dublin collaborative study. *Archives of Disease in Childhood*, 63; 1065-1068.
33. **Dammann O, Leviton A, Gappa M, Dammann CE.(2005):** Lung and brain damage in preterm newborns, and their association with gestational age, prematurity subgroup, infection/inflammation and long term outcome. *BJOG.* 2005 Mar; 112 Suppl 1:4-9.
34. **Developmental Epidemiology Network Investigators. (1998):** The correlation between placental pathology and intraventricular hemorrhage in the preterm infants. *Pediatr Res*,34(1):15-9.
35. **De Vries, L.S., Van Haastert. I.C., Rademaker, K.J., Koopman, C. and Groenendaal, F. (2004):** Ultrasound abnormalities preceding cerebral palsy in high-risk preterm infants. *J. pediatr.*;144:815-20.
36. **Di Renzo GC, Mignosa M, Gerli S, et al.(2005):** The combined maternal administration of magnesium sulfate and aminophylline reduces intraventricular hemorrhage in very preterm neonates. *Am J Obstet Gynecol* 2005;192(2):433–8.
37. **Di Salvo DN.(2001):** A new view of the neonatal brain: clinical utility of supplemental neurologic US imaging windows. *Radiographics*2001;21(4):943-955.

38. **Dixon, G., Psych, B., Badawi, N., Kurinczuk, J., Keogh, J. M., Siburn, S.R. Zubric, S.R. and Stanley, F.J.(2002):** Early developmental outcome after newborn encephalopathy. *Peditrics*;109(1):26-33.
39. **Doyle LW, Betheras FR, Ford GW, Davis NM, Callanan C. (2000):** Survival, cranial ultrasound and cerebral palsy in very low birth weight infants:1980s versus 1990s.*J Pediatr Child Health*;36:7-12.
40. **Du Plessis AJ.(1998):** Posthemorrhagic hydrocephalus and brain injury in the preterm infant: dilemmas in diagnosis and management. *Semin Pediatr Neurol* 1998; 5(3):161–79.
41. **Egarter C, Leitich H, Karas H,(1999):** Antibiotic treatment in preterm premature rupture of membranes and neonatal morbidity: a metaanalysis. *Am. Obstet. Gynecol.* 1999; 174:589-597.
42. **Elbourne D, Ayers S, Dellagrammaticas H, et al.(2001):** Randomised controlled trial of prophylactic etamsylate: follow up at 2 years of age. *Arch Dis Child Fetal Neonatal Ed* 2001;84(3):F183–7.
43. **Engelhard HH, Andrews CO, Slavin KV, et al (2003):** Current management of intraventricular hemorrhage . *Surg Neurol* 60; 15-21; 21-22.
44. **Epelman M, Daneman A, Blaser SI, et al.(2006):** Differential diagnosis of intracranial cystic lesions at head US: correlation with CT and MR imaging. *Radiographics* 2006;26(1):173-196.
45. **Fanaroff A A, kiwi R and shah D.M.(2001):** Antenatal and intrapartum care of the high risk infants. In *care of the high risk neonate* by Klaus, M.H. Fanaroff, AA. 5th ed; ch 1, p 1-44.
46. **Filipovic B, Teofilovski-Parapid G.(2004):** Linear parameters of normal and abnormal cava septi pellucidi: a post-mortem study. *Clin Anat.* 2004 Nov; 17(8):626-30.
47. **Fowlie PW, Davis PG.(2003):** Prophylactic indomethacin for preterm infants: a systematic review and meta-analysis. *Arch Dis Child Fetal Neonatal Ed* 2003;88(6): F464–6.

48. **Frankenburg, W. K and Josiah B. Dodds, J. B.(1967):**
The Denver Developmental Screening Test. The Journal of Pediatrics;186:187.
49. **Friese k. (2003):** The Role of Infection in Preterm Labour. Br J Obstet& Gynaecol; 110(20):52-54.
50. **Fukui K, Morioka T, Nishio S, Mihara F, Nakayama H, Tsukimori K, Fukui M(2001):** Fetal germinal matrix and intraventricular haemorrhage diagnosed by MRI. Neuroradiology; 43(1):68-72.
51. **Fulia F, Cordaro S, Meo P, et al.(2003):** Can the administration of antithrombin III decrease the risk of cerebral hemorrhage in premature infants? Biol Neonate2003;83(1):1–5.
52. **Garcia, J.M., Gherpelli, J.L.D. and Leone, C.R.,(2003):**
The role of spontaneous general movement assessment the neurological outcome of cerebral lesions in preterm infants. J. pediatr.(Rio J.); 80(4):296-305.
53. **GerdaVan Wezel-Meijler(2007):** Neonatal Cranial UltrasonographySpringer-Verlag Berlin Heidelberg 2007.
54. **Glenn OA, Goldstein RB, Li KC, Young SJ, Norton ME, Busse RF, Goldberg JD, Barkovich AJ.(2005):** Fetal magnetic resonance imaging in the evaluation of fetuses referred for sonographically suspected abnormalities of the corpus callosum. J Ultrasound Med. 2005 Jun; 24(6):791-804.
55. **Gomella T.L., Cunningham M.D., Eyal F.G. and Zenk K.E. (1999):** Neonatology: Management, Procedures, On-call problems, Disease, and Drugs (4th edition) A Simon & Schuster Company; 21-30
56. **Gomella TL (2004)b:** Neurologic diseases in: Gomella TL, Donn SM et al ,(edt) : fifth edition, Clinical manual neonatology , McGraw Hill Companies ,Appleton and Lange; p644- 678.
57. **Grasby DC, Esterman A, Marshall P. (2003):**
Ultrasound grading of cerebral ventricular dilatation in

- preterm neonates. *J Paediatr Child Health*. 2003 Apr; 39(3):186-90.
58. **Greisen G.(2005):**Autoregulation of cerebral blood flow in newborn babies. *Early Hum Dev*; 81(5):423-8.
 59. **Grether JK, Nelson KB, Emery ES, Cummins SK.(1999):** Prenatal and perinatal factors and cerebral palsy in very low birth weight infants. *J Pediatr* 1999; 128(3):407-14.
 60. **Gustaaf M.Vermeulen , Hein W, Bruinse ,Leo. J.Gerards, Linda S.de Varies(2001):** Perinatal risk factors for cranial ultrasound abnormalities in neonates born after spontaneous labour before 34 weeksEuropean Journal of Obstetrics & Gynecology and Reproductive Biology 94 (2001) 290-295.
 61. **Guyer B, Hoyert DL, Martin JA, et al. Annual summary of vital statistics: 1998.** *Pediatrics*1999;104:1229–46.
 62. **Haataja, L., Mercuri, E., Cowan, F. and Dubowitz, L.,(2000):** Cranial ultrasound abnormalities in full term infants in a postnatal ward: Outcome at 12 and 18 months. *Arch. Dis. Child. Fetal neonatal ed.*;82:128-133.
 63. **Hack, M. and Fanaroff, A., (2000):** Outcomes of age in the 1990s. *Semin. Neonatol.*;5:89-106.
 64. **Hansen TW(2006):** Prophylaxis of intraventricular hemorrhage in premature infants: new potential tools, new potential challenges. *Pediatr Crit Care Med*; 7(1): 90-2.
 65. **Harding DR, Humphries SE, Whitelaw A, et al.(2007):** Cognitive outcome and cyclooxygenase-2 gene (-765 G/C) variation in the preterm infant. *Arch Dis Child Fetal Neonatal Ed* 2007;92(2):F108–12.
 66. **Haque KN, Hayes AM, Ahmed Z, Wilde R, Fong CY(2008):** Caesarean or vaginal delivery for preterm very-low-birth weight (< or =1,250 g) infant: experience from a district general hospital in UK. *Haque KN, Hayes AM, Ahmed Z, Wilde R, Fong CY*;277(3):207-12.
 67. **Hussain N.,Sharma P, McKay K, Rosenkrantz TS,(2004):** Comparisons of mortality and pre-discharge respiratory outcomes in small-for-gestational-age and

appropriate-for-gestational-age premature infants. *BMC Pediatr.* 2004 Jun 8;4(1):9.

68. **Hedderson M M, Ferrara A and Sacks D.(2003):** Gestational diabetes mellitus and lesser degrees of pregnancy hyperglycemia: Association with increased risk of spontaneous preterm birth. *Obst& Gynecol*; 102:850-856.
69. **Hellström-Westas L, Klette H, Thorngren-Jerneck K, Rosén I(2001):** Early prediction of outcome with aEEG in preterm infants with large intraventricular hemorrhages. *Neuropediatrics*; 32(6):319-24.
70. **Herbst A, Källén KI,(2007):** Influence of mode of delivery on neonatal mortality and morbidity in spontaneous preterm breech delivery. *Eur J Obstet Gynecol Reprod Biol*;133(1):25-9.
71. **Hillman K. (1987):** Intrathoracic pressure fluctuations and periventricular hemorrhage in the newborn. *Aust paed J*,23(6): 343-6.
72. **Holling EE, Leviton A.(1999):** Characteristics of cranial ultrasound white-matter echolucencies that predict disability: a review. *Dev Med Child Neurol* 1999;41:136-9.
73. **Inder TE, Anderson NJ, Spencer C, Wells S, Volpe JJ(2003):** White matter injury in the premature infant: a comparison between serial cranial sonographic and MR findings at term. *AJNR Am J Neuroradiol* 2003a; 24: 805–9. *Inflamm Allergy* ; 1:151-165.
74. **Ingemarsson, I., (2003):** Tachykinin effects on gastric secretion and emptying in rats. *Pharmacol. Res.*22:605-610.
75. **Jatinder, S., Goraya, S. and Veena, R.(2002):** Benign familial neonatal convulsions .*Indian pediatrics*,30:292-295.
76. **Jeffrey M.Pearlman ;Nancy Rollins (2000):**Surveillance protocol for the detection of intracranial abnormalities in premature neonates,*Arch Pediatr Adolesc Med* 2000;154:822-26.
77. **John D and Catherin T Mac.(2005):** Low birth weight and preterm neonates: They may be managed at home by

- mother and traveled health worker?.J perinatology ; 25:72-81.
78. **Jun B .Felix WS.Adrian B, (2002):** Parity & pregnancy outcome .Am J Obstetrics & Gynecology; 166(2):274-278.
 79. **Kadri H, Mawla AA, Kazah J et al(2006):**The incidence, timing, and predisposing factors of germinal matrix and intraventricular hemorrhage (GMH/IVH) in preterm neonates.Child Nerv syst.;1086-90.
 80. **Kaineg B, Hudgins PA.(2005):** Images in clinical medicine. Wernicke's encephalopathy N Engl J Med. 2005 May 12;352(19):e18.
 - 81.**Khalifa G. (1986):** Pediatric Ultrasonography Springer-verlag, Berlin H., Delburg, Newyork, Tokyo.
 82. **Khanal, S., W. Zhang, et al. (2009).** "A comparative study of outcome of preterm neonate with and without history of preterm premature rupture of membrane." Nepal Med Coll J 11(2): 99-103.
 - 83.**Khashoggi TY.(2004):** Outcome of pregnancies with preterm premature rupture of membranes. Saudi Med J. 2004 Dec; 25(12):1957-61.
 84. **Kiechl-Kohlendorfer, U., E. Ralser, et al. (2009).** "Adverse neurodevelopmental outcome in preterm infants: risk factor profiles for different gestational ages." Acta Paediatr 98(5): 792-6.
 85. **Kimberly G.Lee & John P cloherty.(2004):** Identifying of the high risk newborn and evaluating gestational age and small for gestational age infants .In Kimberly cloherty.(Eds) Manual of neonatal care, 6th ed philadliphia:p 41-58.
 86. **Kliegman R.(2000):** Prematurity and IUGR .In Nelson W, Behrman R, Kliegman R Nelson textbook of Pediatrics, 18th edition, Philadelphia, London, Sydney, Tokyo, WB. Sauderss company:701-712.
 87. **Kohelet D, Perlman M, Hanna G, BallIn A(1990):**Reduced platelet counts in neonatal respiratory distress syndrome. Biol Neonate.;57(6)334-42.
 88. **Kosuge S, Ohkuchi A, Minakami H, Matsubara S, Uchida A, Eguchi Y, et al.** Influence of chorioaminionitis on survival and morbidity in singletons liveborn at <32

- weeks of gestation. *Acta Obstet Gynecol Scand* 2000;79:861–5.
- 89. Kuban K, Sanocka U , Leviton A (1999):** White matter disorders of prematurity. *J Pediatr*;134:539-46.
 - 90. Laforgia, N., Dofonzo, I., Altomare, M., Mautone, A., (2001):** Cord blood endothelin-1 and perinatal asphyxia. *Acta. Paediatr.*;90(3):351-352.
 - 91. Laptook AR, O'Shea TM, Shankaran S, Bhaskar B, NICHD(2005):** Neonatal Network. Adverse neurodevelopmental outcomes among extremely low birth weight infants with a normal head ultrasound: prevalence and antecedents. *Pediatrics* 2005;115:673-80.
 - 92. Laura R, William OH, Richard A, Alistair GS, and Charles C. (1995):** Antenatal steroids, Delivery mode, and Intraventricular hemorrhage in preterm infants. *Am J of Obs and Gyn*;172: 795-80.
 - 93. Lemmers PM, Toet M, van Schelven LJ, van Bel F(2006):**Cerebral oxygenation and cerebral oxygen extraction in the preterm infant: the impact of respiratory distress syndrome. *Exp Brain Res*; 173(3):458-67.
 - 94. Leviton A, Fenton T, Kuban KC, et al.** Labor and delivery characteristics and the risk of germinal matrix hemorrhage in low birth weight infants. *J Child Neurol*1991;6(1):35–40.
 - 95. Limperopoulos C, Benson CB, Bassan H, et al.(2005):** Cerebellar hemorrhage in the preterm infant: ultrasonographic findings and risk factors. *Pediatrics* 2005;116(3):717-724.
 - 96. Linder Nehama, Orli Haskin, Orli Levit, Gil Klinger, Tal Prince(2003):** Risk factors for intraventricular hemorrhage in very low birth weight premature infants: a retrospective case –control study . *Pediatrics* 2003; 111; 590-595.
 - 97. Lin KL, Wang HS, Lui TN.(2002):** Diagnosis and follow-up of craniopharyngiomas with transcranial Doppler sonography. *J Ultrasound Med* 2002;21(7):801-806.
 - 98. Maitre, N. L., D. D. Marshall, et al. (2009).** "Neurodevelopmental outcome of infants with unilateral or

- bilateral periventricular hemorrhagic infarction." *Pediatrics* 124(6): e1153-60.
99. **McIntire, D.D., Bloom, S.L., Casey, B.M. and Ieveno, K.J., (1999):** Birth weight in relation to morbidity and mortality among newborn infants. *N.E.J.M.*;340(16):1234-1238.
 100. **Ment LR, Oh W, Ehrenkranz RA, et al.(1994):** Low-dose indomethacin and prevention of intraventricular hemorrhage: a multicenter randomized trial. *Pediatrics* 1994; 93(4):543–50.
 101. **Ment LR, Oh W, Ehraenkranz RA, Philip AG, Duncan CC, and Makuch RW. (1995):** Antenatal steroids, delivery mode, and intraventricular hemorrhage in preterm infants. *Am J Obstet Gynecol*; 172:795-800.
 102. **Ment, L.R., Bada,H.S and Barnes,P., (2002):** Practice parameter: Neuroimaging of neonate: Report of the quality standards subcommittee of the American academy of neurology and practice committee of the child neurology society. *Neurology*;58:1726-1738.
 103. **Ment LR, Vohr BR, Makuch RW, Westerveld M, Katz KH, Schneider KC, Duncan CC, Ehrenkranz R, Oh W, Philip AG, Scott DT, Allan WC(2004):** Prevention of intraventricular hemorrhage by indomethacin in male preterm infants.*J Pediatr*; 145(6):832-4. *Mol Cell Biol*.;26:5336-47.
 104. **Mercuri, E., Dubowitz, L., Peterson-Brown, S. and Cown, F.(1998):** Incidence of cranial ultrasound abnormalities in apparently well neonates on a postnatal ward: Correlation with antenatal and perinatal factors and neurological status. *Arch. Dis. Child. Fetal Neonatal Ed.*;79:185-9.
 105. **Mirmiran M, Barnes PD, Keller K, Constantinou JC, Fleisher BE, Hintz SR, Ariagno RL.(2004) :** Neonatal brain magnetic resonance imaging before discharge is better than serial cranial ultrasound in predicting cerebral palsy in very low birth weight preterm infants. *Pediatrics*. 2004 Oct;114(4):992-8.
 106. **Moran GJ, Krishnadasan A, Gorwitz RJ, et al. (2006):**Methicillinresistant *S. aureus* infections among

- patients in the emergency department. *N Engl J Med* 2006;355:666-74.
107. **Morioka T, Hashiguchi K, Nagata S, Miyagi Y, Mihara F, Hikino S, Tsukimori K, Sasaki T (2006):** Fetal germinal matrix and intraventricular hemorrhage. *Pediatr Neurosurg*; 42(6):354-61.
 108. **Moutquin J M.(2003):** Socio-Economic and psychosocial factors in the management and prevention of preterm labour. *Br J Obstet& Gynaecol.* 110(20):56-60.
 109. **Msall ME.** Developmental vulnerability and resilience in extremely preterm infants. *JAMA* 2004;292:2399-401.
 110. **Murphy BP, Inder TE, Rooks V, et al.** Posthae morrhagic ventricular dilatation in the premature infant: natural history and predictors of outcome. *Arch Dis Child Fetal Neonatal Ed* 2002;87(1):F37–41.
 111. **Nehama L, Orli H, Orli L, Gil K, and Nora N. (2003):** Risk Factors for intraventricular Hemorrhage in very low birth weight premature infants: A retrospective case control study. *Pediatrics*, 111 (5): 590-595.
 112. **Neil JJ, Inder TE.(2004):** Imaging perinatal brain injury in premature infants. *Semin Perinatol.* 2004 Dec;28(6):433-43.
 113. **New Ballard score (1991) :** (From Ballard JL, Khoury JC, Wedig K, et al. New Ballard score, expanded to include extremely premature infant. *J pediatr* ;119:417.).
 114. **Nwafor-Anene VN, DeCristofaro JD, Baumgart S (2003):** Serial head ultrasound studies in preterm infants. how many normal studies does one infant need to exclude significant abnormalities? *J Perinatol*; 23(2):104-10.
 115. **Nzeh DA, Erinle SA, Saidu SA, Pam SD. (2004):** Transfontanelle ultrasonography: an invaluable tool in the assessment of the infant brain. *Trop Doct.* 2004 Oct;34(4):226-7.

116. **Ohlsson A, Roberts RS, Schmidt B, et al.** Male/female differences in indomethacin effects in preterm infants. *J Pediatr* 2005;147(6):860–2.
117. **Olaf Dammann, M.D., and Alan Leviton, M.D.(2006):** Neuroimaging and the Prediction of Outcomes in Preterm Infants *n engl j med* 355;7 www.nejm.org august 17, 2006.
118. **Okten A, Ahmetoglu A, Dilber E, Dinc H, Kalyoncu M, Ciftcibais K, Yaris N.(2002):** Cranial Doppler ultrasonography as a predictor of neurologic sequelae in infants with bacterial meningitis. *Invest Radiol.* 2002 Feb;37(2):86-90.
119. **Okumura A. , Fumio Hayakawa, Toru Kato, Kazuya Itomi, Koichi Maruyama, Naoko Ishihara, Tetsuo Kubota, Motomasa Suzuki,(2001):** Hypocarbia in Preterm Infants With Periventricular Leukomalacia: The Relation Between Hypocarbia and Mechanical Ventilation *PEDIATRICS* Vol. 107 No. 3 March 2001 ;469-475.
120. **O’Shea TM, Counsell SJ, Bartels DB, Dammann O.** Magnetic resonance and ultrasound brain imaging in preterm infants. *Early Hum Dev* 2005;81:263-71.
121. **Paneth N. (1999):** Classifying brain damage in preterm infants *J Pediatr*; 134:527-9.
122. **Papile LA, Burstein J, Burstein R, Koffler H.(1978):** Incidence and evolution of subependymal and intraventricular hemorrhage: a study of infants with birth weights less than 1,500 gm. *J Pediatr*\1978;92(4):529-534.
123. **Papile LA, Ment LR, Bada HS, Barnes P, Grant PE, Hirtz D, Pinto-Martin J, Rivkin M, Slovis TL(2006):** Practice parameter. Neuroimaging of the neonate. report of the Quality Standards Subcommittee of the American Academy of Neurology and the Practice Committee of the Child Neurology Society. *Neurology*; 58(12):1726-38.
124. **Paro-panjan, D .,Nenbauer, D., Kodric, J. and Bratanic, B., (2005):** Amiel-Tison neurological assessment at term age: clinical application, correlation with other

- methods, and outcome at 12 to 15 months. *Developmental medicine & child neurology* ; 47:19-29.
- 125. Paul D, Sciscione A, Leef K, and Stefano J. (2002):** Caesarean delivery and outcome in very low birth weight infants. *Aust N Z J Obstet Gynaecol*; 42(1):41-5.
 - 126. Perlman JM, Risser RC, and Gee JB. (1997):** Pregnancy-induced hypertension & reduced intraventricular hemorrhage in preterm infants. *Pediatr neurol*,17(1):29-33.
 - 127. Perlman JM, and volpe JJ. (1986):** Intraventricular haemorrhage in the extremely small premature infants. *AJDC*;140:1122-1124.
 - 128. Perlman JM, Rollins N. (2000):** Surveillance protocol for the detection of intracranial abnormalities in premature neonates. *Arch Pediatr Adolesc Med*. 2000 Aug; 154(8):822-6.
 - 129. Phelps DL, Rosenbaum AL, Isenberg SJ, et al.(1987):** Tocopherol efficacy and safety for preventing retinopathy of prematurity: a randomized, controlled, doublemasked trial. *Pediatrics* 1987;79(4):489–500.
 - 130. Pisani, F., A. L. Barilli, et al. (2008).** "Preterm infants with video-EEG confirmed seizures: outcome at 30 months of age." *Brain Dev* 30(1): 20-30.
 - 131. Pryds O. (1989):** Heterogeneity of cerebral vasoreactivity in preterm infants supported by mechanical ventilation. *J pediatr*;115:638-645.
 - 132. Pursely D.M. and Cloherty J.P. (2004):** Identifying the high risk newborn and evaluating gestational age, prematurity, postmaturity, large for gestational age, small for gestational age infants, In: *Manual of neonatal care*, Chapter III, (5th edition) Lippincott- Rawen Company; 37-49 .
 - 133. Qian JH, Chen HJ, Chen GY, Chen XP, Zhang Q, Ao LM, Wu SM. (2004):** Prospective study on prognosis of infants with neonatal subependymal cysts, *Zhonghua Er Ke Za Zhi*. 2004 Dec;42(12):913-6
 - 134. Rahman S, Teo C, Morris W, et al.(1995):** Ventriculosubgaleal shunt: a treatment option for

- progressive posthemorrhagic hydrocephalus. *Childs Nerv Syst* 1995;11(11): 650–4.
135. **Rasmussen S & Irgens L.(2003):**Fetal growth and body proportion in pre eclampsia .*Obstet& Gynecol J*;101:575-583.
 136. **Ringer S. A.C (1998):** Resuscitation in the delivery room. In: *Manual of Neonatal care*, cloherty J.P. and Stark A.R. (eds.) 4th edition lippincot -raven - Philadelphia; p.p. 53-64.
 137. **Rizzo T, Metzger BE, Burns WJ, Burns K.** Correlations between antepartum maternal metabolism and intelligence of offspring of diabetic mothers.. *N Engl J Med* 1991; 325: 911–916.
 138. **Reinprecht A, Dietrich W, Berger A, Bavinzski G, Weninger M, Czech T. (2001):** Posthemorrhagic hydrocephalus in preterm infants. long-term follow-up and shunt-related complications. *Childs Nerv Syst*; 17(11):663-9.
 139. **Resch B, Gedermann A, Maurer U, et al.** Neurodevelopmental outcome of hydrocephalus following intra/periventricular hemorrhage in preterm infants: short and long-term results.
 140. **Robert L & Goldenberg M.(2002):** High-Risk Pregnancy Series: An Expert's View in the Management of Preterm .*Labor J .Obs & Gynecol*.Vol 100, N 5, Part (1) P 1020 – 1037.
 141. **Roessman V (1999):** Congenital malformations .in Duckett S ed. *Pediatric neuropathology*. Philadelphia :Williams & Wilkin :123-48.
 142. **Roscigno CI (2002):** Periventricular leukomalacia. pathophysiological concerns due to immature development of the brain.*J Neurosci Nurs*; 34(6):296-302.
 143. **Roe S . Lazebnik, M.D., Ph.D. and Laura J. Varich, M.D.(2009):** Neonatal cranial sonography: Modern strategies and Applications,2007 cmpmedica.CME Ultra Sound;1,6.

144. **Rouse DJ, Hirtz DG, Thom E, et al.** A randomized, controlled trial of magnesium sulfate for the prevention of cerebral palsy. *N Engl J Med* 2008;359(9): 895–905.
145. **Roy K.K., Jine B, Kummar S and Malhotra N.(2006):** Maternal antenatal profile and immediate neonatal outcome in VLBW and ELBW infants . *Indian journal of paediatrics*; Vol 73 (8):669-673.
146. **Russel, G., Lydon, Y., and Loyed, D., (1986):** Which deliveries require pediatrician attendance! *B.M.J.*;289(18): 438-453.
147. **Sandler D, Cooper P, Bolton K, and Bental R. (1994):** Periventricular-Intraventricular haemorrhage in low birth-weight infants at Baragwanath Hospital. *S Afr Med J*, 84(1):26-9.
148. **Saurel-Cubizolles, Zeitlin J, Lelong N, Papiernik E, Di Renzo G and Breart G.(2004):** Employment, Working Conditions and Preterm Birth: Results From the J Epidemiol Community Health, Case-Control Survey; 58:395-401Europ.
149. **S.Chaudhari, Deshpande, Kinare , Kumar, Pandit(1994):** Ultrasonography Of The Brain In Prterm Infants and its correlation with neurodevelopmental outcome. *Indian pediatrics volume* 32-July1995.
150. **Schmidt B, Davis P, Moddemann D, et al.(2001):** Long-term effects of indomethacin prophylaxis in extremely-low-birth-weight infants. *N Engl J Med* 2001;344(26): 1966–72.
151. **Schneider A(2004):** Normal values of blood flow velocity in the terminal vein of preterm infants. *Ultraschall Med*; 25(2):137-40.
152. **Selbing A, Hedger and Hung T.(1998):** Gestational age estimation. *Acta Obst&gynecol* ; 64:593-599.
153. **Shah, P.S., Beyene, J., To, T., Ohlsson, A. and Perlman, M., (2006):** Post asphyxial Hypoxic- Ischamic Encephalopathy in neonates outcome prediction rule within 4 hour of birth. *Arch. Pediatr. Adolesc. Med.*;160:729-736

154. **Shirahata A, Nakamura T, Shimono M, et al.(1990):** Blood coagulation findings and the efficacy of factor XIII concentrate in premature infants with intracranial hemorrhages. *Thromb Res* 1990;57(5):755–63.
155. **Sie LT, van der Knapp MS, van Wezel-Meijler G, Taets van Amerongen AH, Lafeber HN, Valk J.(2000):** Early MR features of hypoxic-ischemic brain injury in neonates with periventricular densities on sonograms. *AJNR Am J Neuroradiol.* 2000 May;21(5):852-61
156. **Soraisham, A. S., N. Singhal, et al. (2009).** "A multicenter study on the clinical outcome of chorioamnionitis in preterm infants." *Am J Obstet Gynecol* 200(4): 372 e1-6.
157. **Spinillo, A., L. Montanari, et al. (2009).** "Infant sex, obstetric risk factors, and 2-year neurodevelopmental outcome among preterm infants." *Dev Med Child Neurol* 51(7): 518-25.
158. **Stoll, B.J. and Kliegman, R.M., (2004):** Hypoxia ischemia in: *Nelson textbook of pediatrics.* Behrman, R.M. and jenson (eds). 17thed.W.B.Saunders, Philadelphia.
159. **Strober JB, Bienkowski RS, Maytal J.** The incidence of acute and remote seizures in children with intraventricular hemorrhage. *Clin Pediatr (Phila)* 1997; 36(11):643–7.
160. **Szymonowicz W, Yu VY, Walker A, et al.** Reduction in periventricular haemorrhage in preterm infants. *Arch Dis Child* 1986;61(7):661–5.
161. **Synder, E and Cloherty, J., (2004):** Perinatal asphyxia in : *manual of neonatal care,* Clohertyj.p. and Stank A.R.(eds)5th ed Lippincott-Raven, Philadelphia pp 536-554.
162. **Synnes AR, Macnab YC, Qiu Z, Ohlsson A, Gustafson P, Dean CB, Lee SK; the Canadian Neonatal Network(2006):**Neonatal intensive care unit characteristics affect the incidence of severe intraventricular hemorrhage. *Med Care*; 44(8):754-9.
163. **Taghavi, S., M. S. Melli, et al. (2009).** "Neonatal C-reactive protein value in prediction of outcome of preterm

- premature rupture of membranes: comparison of singleton and twin pregnancies." *J Obstet Gynaecol Res* 35(4): 694-8.
- 164. Terri E, Inder J, and Joseph J. (2000):** Mechanism of perinatal brain injury. *Semin Neonatol*:5:3-16.
- 165. The Northern Neonatal Nursing Initiative [NNNI] Trial Group. (1996):**A randomized trial comparing the effect of prophylactic intravenous fresh frozen plasma, gelatin or glucose on early mortality and morbidity in preterm babies. *Eur J Pediatr* 1996; 155(7):580–8.
- 166. Thilaganathan, B., Athanasiou, S., and Ozmen. S., (1994):**Umbilical cord blood erythroblasts as an index of intrauterine hypoxia. *Arch. Dis. Child.*;70: 192-194.
- 167. Thorne S, Hislop TG, Kuo M, Armstrong E-A.** Hope and probability: patient perspectives of the meaning of numerical information in cancer communication. *Qual Health Res* 2006; 16:318-36.
- 168. Tonni G, Ferrari B, Defelice C, Centini G(2005):** Neonatal porencephaly in very low birth weight infants: ultrasound timing of asphyxial injury and neurodevelopmental outcome at two years of age. *J Matern Fetal Neonatal Med*; 18(6):361-5.
- 169. Van de Bor M, den Ouden L.(2004):** School performance in adolescents with and without periventricular-intraventricular hemorrhage in the neonatal period. *Semin Perinatol.* 2004 Aug;28(4):295-303.
- 170. Vannucci, R.C., (2000):** Hypoxic-Ischemic Encephalopathy.*Am. J. Perinatal.*;17:113-120.
- 171. Vannucci,S.J. and Hagberg, h., (2004):** Hypoxia-ischemia in the immature brain. *Journal of Experimental Biology.*;207:3149-3154.
- 172. Vasileiadis GT, Gelman N, Han VK, Williams LA, Mann R, Bureau Y, Thompson RT (2004):** Uncomplicated intraventricular hemorrhage is followed by reduced cortical volume at near-term age. *Pediatrics*; 114(3):e367-72.
- 173. Vasileiadis GT.(2004):** Grading intraventricular hemorrhage with no grades. *Pediatrics* 2004;113:930-1.

174. **Velisavljev- Filipović G(2007):** Mode of delivery and neurosonographic findings in premature infants] *Med Pregl* ;60(1-2):7-11.
175. **Ventriculomegaly Trial Group.(1990):** Randomised trial of early tapping in neonatal posthaemorrhagic ventricular dilatation. *Arch Dis Child* 1990;65(1):3–10.
176. **Vergani P, Locatelli A, Strobelt N, (1998):** Clinical outcome of mild fetal ventriculomegaly. *AM J Obstet Gynecol*;178:218-22.
177. **Vohr B, Allan WC, Scott DT, Katz KH, Schneider KC, Makuch RW, Ment LR(1999):** Early-onset intraventricular hemorrhage in preterm neonates: incidence of neurodevelopmental handicap. *Semin Perinatol*; 23(3):212-7.
178. **Vohr BR, Allan WC, Westerveld M, et al.(2003):** School-age outcomes of very low birth weight infants in the indomethacin intraventricular hemorrhage prevention trial. *Pediatrics* 2003;111:
179. **Vollmer, B., Roth, S., Riley, K., O'Brien, F., J., De Haan, M., Vargha Khadem, F. and Wyatt, B.G.R.,:(2006):** Long-term neurodevelopmental outcome of preterm children with unilateral cerebral lesions diagnosed by neonatal ultrasound. *J.Early Human Development*;in press.
180. **Volpe JJ (1999):** Germinal matrix hemorrhage in the premature infant in: *Neurology of the newborn*:403-463.
181. **Volpe JJ. (2000):** Neuronal proliferation, migration, organization and myelination. In: Volpe JJ(ed). *Neurology of the newborn*, Philadelphia: WB Saunders, 45-99.
182. **Volpe JJ. Intracranial hemorrhage(2001):** germinal matrix-intraventricular. hemorrhage of the premature infant. In: *Neurology of the newborn*. Philadelphia: WB Saunders,:428–93.
183. **Volpe JJ. (2001):** Neurobiology of periventricular leukomalacia in the premature infant. *Pediatr Res* 2001;50:553–62

184. **Volpe JJ.(2008):** Intracranial hemorrhage: neurology of the newborn. 5th edition. Philadelphia: W.B. Saunders Co.; 2008. p. 481–588.
185. **Weintraub Z, Solovechick M, Reichman B, and Rostchild D. (2001):** Effect of maternal tocolysis on the incidence of severe periventricular/intraventricular haemorrhage in very low birth weight infants. Arch Dis Child Fetal Neonatal Ed; 85:F13-F17.
186. **Welch RA, and Bottoms SF. (1986):** Reconsideration of head compression and intraventricular haemorrhage in the vertex very-low birth-weight fetus. Obstet Gynecol;68:29-34.
187. **Wielenga JM, Smit BJ, Merkus MP, et al.(2007):** Individualized developmental care in a Dutch NICU: short-term clinical outcome. Acta Paediatr 2007;96(10):1409–15.
188. **Wilson-Costello, D., Borawski, E., Friedman, H., Redline., R., Fanaroff, A. and Hack, M., (1998):**Perinatal correlates of cerebral palsy and other neurologic impairment among very low birth weight children. Pediatrics; 102 (2):315-322.
189. **Whitelaw A.(2001):** Intraventricular haemorrhage and posthaemorrhagic hydrocephalus: pathogenesis, prevention and future interventions. Semin Neonatol 2001; 6(2):135–46.
190. **Whitelaw A, Odd D.(2007):** Postnatal phenobarbital for the prevention of intraventricular hemorrhage in preterm infants. Cochrane Database Syst Rev 2007;(4): CD001691.
191. **Wong F, Fraser S, Kelly E, Acton C. (2004):** Clinical significance of isolated paraventricular cysts on cranial ultrasonography. J Paediatr Child Health .2004 Sep-Oct;40(9-10):552-5.
192. **Wu, W.Y., Backstrand, H.K., Zhao, S., Fullerton, J.H. and Jonhnston, S.C., (2004):** Declining diagnosis of birth asphyxia in California 1991-2000. Pediatrics.144: 1584-1590.
193. **Zhang, D., (1996):** Asphyxia of newborn delivered by elective C.S. Chung Hua Fu Chan Ki Tsa Chih.;31(12): 718-719.

- 194. Zotter H, Urlesberger B, Kerbl R, Mueller W, Pichler G, Curzi-Dascalova L(2007):**Cerebral hemodynamics during arousals in preterm infants. *Early Hum Dev*; 83(4):239-46.