

REFERENCES

- Abdelfatah, M.(1999):*** Rapid detection of neonatal sepsis using polymerase chain reaction. M.Sc. thesis (Pediatrics). Faculty of Medicine Cairo University.
- Afroza, S. (2006):*** Neonatal sepsis- a global problem: an overview. Mymensingh Med J; 15(1): 108-114.
- Ahmed, M.; Hamed, Z.; Fouad, H. et al.; (2000):*** Granulocyte colony stimulating factor as a marker for neonatal sepsis. Thesis submitted for partial fulfillment of M.D. Degree in Pediatrics. (Cairo University).
- Alam, T. and Papaconstantinou, J. (1992):*** Biochemistry 31-1928-1936.
- Ali, S.A.; Khan, T.A. and Zaidi, A.K.M. (2004):*** Neonatal sepsis in Peshawar. Arch. Dis. Child.; 89: 568-571.
- Am J Clin Pathol (2002):*** 118(3) 346-353 .PubMed.
- Anderson-Berry, A.; Bellig, L. and Ohning, B. (2006):*** Neonatal sepsis, emedicine. Medscape.com/ article/ 978352-overview.
- Anderson-Berry, A.L.; Bellig, L.L.; RN,et al., (2006):*** neonatal sepsis. Department of Neonatology, Greenville Children's Hospital and Medical University of South Carolina.
- Anwer, S.; Mustafa, S. and Ashraf, S. et al.; (2000):*** Rapid identification of neonatal sepsis. J. Pak. Med. Assoc.; 45 (5): 94-8.
- Athhan, F.; Genel, F.; Bak, M. et al.; (2002):*** PCT: a marker of neonatal sepsis. J. Trop. Pediat.; 48: 10-4.

- Atta, N.A. (2002):** Study of intracellular adhesion molecules-1 (ICAM-1) and E-selection in neonatal infection. MD thesis (Pediatrics). Faculty of Medicine. Zagazig University.
- Bellig, L.L.; Ohning, B.L.; MacGilvray, S. et al.; (2001):** Neonatal sepsis. e-Medicine J. ; 2(11).
- Bellig, L.L.; RN, NNP, Ohning, et al.,(2004):** neonatal sepsis. Department of Neonatology, Greenville Children's Hospital and Medical University of South Carolina. eMedicine.com, Inc.
- Bellig, L.L.R.N.; NNP, Ohning L. (2004):** Neonatal Sepsis. The merck manual. Disturbances In Newborns And Infants. eMedicine.com, Inc.
- Berger, C.; Uchlinger, J.; Ghelfi, D.; et al., (1995):** Comparison of CRP and white blood cell count with differential in neonates at risk for septicemia. Eur J Pediatr 154, 138-144.
- Bonadio, W.A.; Smith, D. et al.,. (1992):** Correlating CBC profile and infectious outcome. A study of febrile infants evaluated for sepsis. Clin. Pediatr.; 31(10): 578-582.
- Bouzada, M.G.F.; Camargose et al ., (2002):** De Araujo APSD. A1-acid glycoprotein for the presumptive diagnosis of neonatal sepsis. Int Pediatr. 17:233-236.
- Ceciliani, F. (2007):** the acute phase protein alpha 1-acid glycoprotein ;a model for altered glycosylation during diseases.cardiovasc Hematol Agents Med Chem.:(6) 272- 274.
- Cell Moll Life Sci.(2009):** 66(1) 9-26. PubMed.

- Chacko, B. and Sohi, I. (2005):** Early onset neonatal sepsis. Indian J. Pediatr. ; 72:23-26.
- Chang, C.J.; Chen, Y.L. and Lee, S.C. (1998):** Mol. Cell Biol. 18, 880-5887.
- Chiesa, C.; Kemp, A. and Avis, A. (2004):** Procalcitonin, CRP, IL-6 and immature to total neutrophil ratio in the diagnosis of early neonatal sepsis. Clin. Intens. Care; 7: 14.
- Chiesa, C.; Panero, A.; Osborn, et al., (2004):** Diagnosis of neonatal sepsis. A clinical and laboratory challenge. Clinical Chemistry; 50(2): 279.
- Chiesa, C.; Panero, A.; Osborn, et al., (2004):** Diagnosis of Neonatal Sepsis: Clinical Chemistry 50: 279-287; 10.1373/clinchem.025171.
- Clin Diagn Lab Immunol(2004): 11(2) 325-329 .PubMed.**
- Colombo, S.; Buclin, T.; Decosterd, et al. (2006):** orosomucoid plasma concentration and genetic variants ,clin. Pharmacol. Ther. 80(4):307-18.
- Crestani, B.; Rolland, C.; Lardeux, B.; et al., (2002):** J. Immunol. 160 4596-4605 .
- Curr Protein Pept Sci.(2007) :8(1) 91-108 .**
- Dewey, M.J.; Rheaume, C.; Beger, et al., (1990):** J. Immunol. 144, 4392-4398 .

- Doellner, H.; Arntzen, K.J.; Haereid, P.E.; et al., (1998):** Interleukin-6 concentrations in neonates evaluated for sepsis. J. Pediatrics; 132: 295-301.
- Drechou, A.; Prez-Gonzalez, N.; Agneray, et al, (1989):** Cell. Biol. 50,111-116.
- Durand, G.; Delranc, C.; Onaly, J.B et al., (1997):** Biochem. Biophys. Res. Commun. 234 (1997) 544-548 .
- Edwards, M.S. and Baker, C.J.L. (2004):** Bacterial infection in the neonate in: Pediatric Infectious Disease (eds, Long SS; Pickering, LKI; Prober CC) Churchill Livingstone, New York, Edinburgh, London, Madrit, Melbourne, San Francisco, Tokyo. Chap 98; P: 605-14.
- El-Kerdani, T.; Abed Wahed, M.A.; Aly, et al., (2001):** The values of neutrophil CD11b expression, IL-6 and soluble receptor of tumor necrosis factor in early diagnosis of neonatal sepsis. J. Egy. Soc. ; 115:125
- El-Maraghi, S.; Mansour, L. and Kaddah, A. (1997):** Renal dysfunction in septicemic neonates. J. Arab. Child; 8(1): 95-105.
- El-Maraghi, S.; Seund, I.; Gad, W. et al., (1994):** Bacteriological study of sepsis in at risk neonates M.Sc. Thesis in Pediatrics Faculty of Medicine Cairo University.
- Engle, W.; Rosenfield, C.; Mouzinho, A. et al.; (1997):** Circulating neutrophil in septic preterm neonates. Pediatrics; 99: 10.
- Erikson, M. (1993):** Neonatal septicemia. Acta Pediatr. Scand; 72: 1-8.

- Fakhr, A.M. (2005):** Comparison between conventional blood culture and PCR in diagnosis of neonatal septicemia. MSc. Thesis in Microbiology Department, faculty of Medicine, Zagazig University, P. 108-112.
- Fournier, T.; Bouach, N.; Delafosse et al.,(1999):** J. Immunol. 163-2883-2890.
- Fournier, T.; Delafosse, C.; Crestani, et al., (2000):** Biochemica et Biophysica Acta 1482 157-171.
- Fournier, T.; Fadok, V. and Henson, P.M. (1997):** J. Biol. Chem. 272 31065- 31072.
- Fournier, T.; Mejdoubi, N.; Monnet, D. et al., (1994):** Hepatology 201584-1588.
- Gahmberg, C.G. and Andersson, L.C. (1978):** J. Exp. Med. 148 507-521.
- Gendrel, D.; Assicot, M.; Raymond, et al., (1996):** Procalcitonin as a marker for the early diagnosis of neonatal infection. J. Pediatr. 128; P: 570-73.
- Gerdes (2006):** clinicopathological approach to cellular and molecular immunology . the diagnosis of neonatal sepsis .clin perinatol, Philadelphia pp:250-277.
- Gerges, M.N. (2001):** Detection of different bacteriological species responsible for early-onset neonatal sepsis. M.Sc. thesis (Pediatrics). Faculty of Medicine. Cairo University.
- Gomella, T.L.; Cunningham, M.D.; Eyal, et al.,(2004):** Infectious diseases and perinatal asphyxia. Neonatology: management,

procedures, on call problems, diseases and drugs(5th edition). McGraw-Hill Companies; 434-469, 512-524.

Guida, J.; Runing, A.; Leaf, K.; et al., (2003): Platelet count and sepsis, is there an organism specific response. *Pediatr.* 11: 310-315.

Haque, K. and Mohan, P. (2003): Pentoxifylline for neonatal sepsis. Epsom & St Helier NHS Trust, Division of Neonatology, Department of Child Health, Queen Mary's Hospital for Children, Wrythe Lane, Carshalton, Surrey UK, SM5 1AA. The Cochrane Library, Issue 4.

Haque, K.H. (2003): Infection and immunity in the newborn. In: Forfar and Aneil's Textbook of Pediatrics. By: Campbell AGM, McIntosh N (eds) 5th ed. Churchill Livingstone. London: p 273-289.

Hodge, G.; Hodge, S.; Haslam, R.; et al., (2004): Rapid simultaneous measurement of multiple cytokines using 100 µl sample volumes- association with neonatal sepsis. *Clinical & Experimental Immunology*. Volume 137 Issue 2 Page 402. doi:10.1111/J.1365-2249.02529.x.

Holladay, J.W.; Dewey, M.J. et al. (1998): *Pham. Res.* 13 1313-1316.

Holladay, J.W.; Dewey, M.J. et al., (2000): *Drug. Metab. Dispos.* 26, 20-24.

Jarrar, D.; Wolf, S.E.; Jeschke et al., (1997): *Arch. Surg.* 132 1171-1175.

Jaye, D.L. and Waites, K.B. (1997): Clinical applications of C-reactive protein in pediatrics. *Pediatr. Infect. Dis. J.*; 16: 735-747.

- Jiang, J.H.; Chiu, N.C.; Huang, et al., (2004):** Neonatal sepsis in the neonatal intensive care unit: characteristics of early versus late onset. *J Microbiol Immunol Infect*;37:301-306.
- Jordan, J.A.; Butchko, A.R. et al., (2004):** Use of Pyrosequencing of 16S rRNA Fragments to Differentiate between Bacteria Responsible for Neonatal Sepsis. American Society for Investigative Pathology & Association for Molecular Pathology. *PEDIATRICS* Vol. 106 No. 1 , p. e4.
- Kalmovarin, N.; Friedrichs, W.E.; 'Brien, et al.,(1991):** *Inflammatin* 15, 369-379.
- Kaseb, A.A.; Abou El-Ela, M.H.; Abou Hussein et al., (2004):** The value of procalcitonin in the diagnosis of neonatal sepsis. M.Sc. thesis (Pediatrics). Faculty of Medicine. Cairo University.
- Kennon, C.; Overturf, G. and Bessman, S. (1996):** Granulocyte colony stimulating factor as a marker for bacterial infection in neonates. *J. Pediatrics*; 128: 765-769.
- Klein, J.O. and Remington, J.S. (1995):** Current concept of infection of the fetus and newborn infants. In: textbook of Infectious Diseases of the Fetus and Newborn Infant. 4th ed., WB Saunders, Philadelphia. P: 1-18.
- Kocabas, E.A Sarikcioglu, N.; Aksaray, et al., (2007):** Role of procalcitonin, C-reactive protein, interleukin-6, interleukin-8 and tumor necrosis factor-alpha in the diagnosis of neonatal sepsis. *Turk J Pediatr*, pp:7-20.
- Kocherlakota, P. and La Gamma, E.F. (1998):** Preliminary Report: rhG-CSF May Reduce the Incidence of Neonatal Sepsis in Prolonged

Preeclampsia-associated Neutropenia. PEDIATRICS Vol. 102 No. 5, pp. 1107-1111.

L. Dente, U.; Ruther, M.; Tripodi, et al., (1988): Cortese Genes Dev. 2, 259-266.

L. Rabehi, F.; Ferriere, L. and Saffar, L. (1995): Gattegno, Glycocoj. J. 12, 7- 16 .

Laforgia, N.; Coppola, B.; Carbone, R.; et al.; (1997): Rapid detection of neonatal sepsis using polymerase chain reaction. Acta Pediatr; 86: 1097-1099.

Lee, Y.M. and Lee, S.C. (1994): DNA Cell Biol. 13 , 1029-1036.

Liao, Y.C.; Taylor, J.M.; Vinnice, J.L et al., (1985): Mol.Cell Biol. 5, 12826-12831.

Libert, C.; Hochepped, T.; Berger, F.G et al., (1998): Transgen. Res. 7, 429- 435.

Lieberman E.; Lang, J.M.; Richardson et al., (1997): Epidural Analgesia, Intrapartum Fever, and Neonatal Sepsis Evaluation. PEDIATRICS Vol. 99 No. 3 March, pp. 415-419.

Lott, W. (2003): Neonatal bacterial sepsis. Crit. Care Nurse Clin. North Am, ; 15 (1): 35-65.

Mathers, N.J. and Pohlandt, F. (2005): Diagnostic audit of C-reactive protein in neonatal infection . Eur J Pediatr, 146:147-151.

McIntosh, K. (1987): The newborn in: Scaffers diseases of the newborn (5th Ed). WB. Saunders. Philadelphia p: 730-734.

- McIntosh, N. (1998):** The newborn. In: Text Book of Pediatrics. Campbell A. G. M. and McIntosh, N. (eds.). 5th ed. Churchill Livingstone. New York. p: 52-192.
- Meisner, M. (2000):** Procalcitonin (PCT): A new, innovative infection parameter. Biochemical and clinical aspects. 3rd ed. New York.
- Mejdoubi, N.; Henriques, C.; Bui, E et al., (1999):** Hepatology 29, 186-194 .
- Morin, D.; Simon, N.; Depres-Brummer, et al.,(1997):** Pharmacology 54, 271-275.
- Mtitimila, E.I. and Deanery, M. (2004):** Antibiotic regimens for suspected early neonatal sepsis. Liverpool Women's Hospital, Crown Street, Liverpool, UK,L8 7SS. The Cochrane Library, Issue 4.
- Nabih, M.; El-Maraghi S. and Sallah, S.; et al., (2001):** Serum intracellular adhesion molecule in neonatal sepsis. Thesis submitted for partial fulfillment of M.D. Degree in pediatrics. Cairo University.
- Ng, P.C.; Cheng, S.H.; Chui, K.M.; et al., (1997):** Diagnosis of late onset neonatal sepsis with cytokines, adhesion molecules, and C-reactive protein in preterm very low birth weight infants. Arch. Dis. Child Fetal Neonatal Ed; 77.P: 221-227.
- Oddie, S. and Embleton, N.D. (2002):** Risk factors for early onset neonatal group B-Streptococcal sepsis: case-control study. BMJ 2: 325(7371): 1037.

- Oddie, S. and Embleton, N.D. (2002):*** Risk factors for early onset neonatal group B streptococcal sepsis: case-control study. Newcastle Neonatal Service, Department of Child Health, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP. *BMJ*; 325:308.
- Osrin, D.; Vergnano, S. and Costello, A. (2004):*** Serious bacterial infections in newborn infants in developing countries. *Curr. Opin. Infect. Dis.*; 17(3): 217-224.
- Ottolini, M.C.; Lundgren, K.; Mirkinson, et al., (2003):*** Utility of complete blood count and blood culture screening to diagnose neonatal sepsis in the a symptomatic at risk newborn. *Pediatr. Infect. Dis. J.*; 22: 430-4.
- Papsidero, L.D.; Nemoto, T. and Chu, T.M. (1978):*** *Cancer Biochem. Biophys.* 3, 15-18.
- Pediatr Res. (2003):*** 54 (2) 276-281. PubMed.
- Pepyes, M.B. and Hirschfield, G.M. (2003):*** C-reactive protein :a critical update. *J Clin Invest* 111(12) :1805-12 .
- Pepyes, M.B.; Hirschfield, G.M.; Tennent, G.A.; et al., (2006):***Targeting C-reactive protein for the treatment of cardiovascular disease. *Nature* 440 (7088) :1217-1221.
- Perez, E.M. and Weisman, L.E. (1997):*** Novel approaches to the prevention and therapy of neonatal bacterial sepsis. Department of Pediatrics, Baylor College of Medicine, Houston, Texas, USA. PMID: 9099511 [PubMed - indexed for MEDLINE]. *Clin Perinatol*; 24(1):213-29.

- Plamer, A.; Carlin, J.B.; Freihorst, J. et al., (2004):** The use of CRP for diagnosing infections in young infants <3 months of age in developing countries. *Ann. Trop. Pediatric.* 24 (3): 205-12.
- Polin, R.A. (2003):** The ins and outs of neonatal sepsis. *J Pediatr*; 143:34.
- Polin, R.A. (2003): The ins and outs of neonatal sepsis.** *J. Pediatr.* 143-214.
- Posen, R. and Delemos, R.A. (1998):** C-reactive protein levels in the extremely premature infant: Case studies and literature review. *J. Perinatol.*; 18(2): 138-41.
- Powell, K.R. and Marcy, S.M. (1995):** Laboratory aids for diagnosis of neonatal sepsis. In: *Infectious Diseases of the Fetus and Newborn Infants*. By: Klein JO and Remington JS (eds.); W. B.: Saunders 4th ed. Philadelphia; p: 1223-1240.
- Pukhalsky, A.L.; Shyian, S.D.; Kalashnikova et al., (1998):** Mediators Inflamm. 7, 1-47.
- Puopolo, K.M. (2004):** Bacterial and fungal infections by Cloherty JP and Stark AR. (eds) in: *Manual of neonatal Care*, 5th ed. Lippincott Ravan. Philadelphia. New York 287-313.
- Rodriguez, A. (1998):** Neonatal sepsis in Santiago de compostela. *Crit. Care Med.*; 22: 673-679.
- Sann, L.; Krawiecki, N.; Bourgeois, et al., (1976):** Study of orosomucoid in the newborn. Value in the diagnosis of bacterial infections; *Arch Fr Pediatr.* 33:961-971.

- Santana, Reyes, C.; Garcia-Munoz, F.; Reyes, et al., (2003):** Role of cytokines (interleukin- 1 β , 6,8 tumour necrosis factor- α and soluble receptor of interleukin-2) and C-reactive protein in the diagnosis of neonatal sepsis. *Acta Paediatr.* 92:221-227.
- Savoldi, G.; Enaroli, A.F.; Ferrari, F. et al., (1997):** *DNA CELL Biol.* 16, 1467-1476.
- Schmid, K. and Am, J. (1953):** *Chem .Soc.* 75, 60-68 .
- Schuchat, A.; Zywicki, S.S. and Sinsmoor, M.J; et al., (2000):** Risk factors and opportunities for prevention of early-onset neonatal sepsis: A multicenter case-control study. *Pediatrics*; 105: 21-26.
- Schuchat, A.; Zywicki, S.S.; Dinsmoor, M.J. et al.,(2000):** Risk Factors and Opportunities for Prevention of Early-onset Neonatal Sepsis: A multicenter Case-Control Study. *Pediatrics*;105:21-26.
- Shwick, H.G. and Haupt, H. (1984):** in: F.W Putnam (Ed.), *The Plasma Proteins* ,Vol. IV ,2nd edn.,Academic Press ,New York, ,pp. 167-200.
- Sörensson, J.; Matejka, G.L.; Ohlson, M. et al., (1999):** *Am. J.Physiol.* 276 H 530-H 534.
- Stoll, B.J. (2002):** Infections of the neonatal infant. In: *Nelson Textbook of Pediatrics.* Behrman, RE, Kleigman, RM and Jenson HB (eds.). 17th ed. W.B. Saunders, Philadelphia P: 623-640.
- Stoll, B.J. (2007):** Infections of the neonatal infant. In: *Kliegman R, Behrman R, Jenson H and Stanton B. Nelson Textbook of Pediatrics* (18th edition). Saunders Elsevier; 794-812.

- Stoll, B.J. and Chapman, I.A. (2007):*** Nervous system disorders. In: Kliegman R, Behrman R, Jenson H and Stanton B. Nelson Textbook of Pediatrics (18th edition). Saunders Elsevier; 713-723.
- Stoll, B.J. and Kliegman, R.M. (2002):*** Prematurity and IUGR. In: Nelson Textbook of Pediatrics. Beharman RE, Kliegman RM and Jenson HB (eds.). 17th ed. W.B. Saunders, . Philadelphia, P: 550-558.
- Stoll, B.J.; Hansen, N.; Fanaroff, A.A.; et al., (2002):*** Late- onset sepsis in very low birth weight neonates: the experience of the NICHD Neonatal Research Network. Pediatrics; 110: 285- 291.
- Stone, R.T. and Maurer, R.A. (1987):*** Dev. Genet. 8295-304 .
- Toh, H.; Hayashida, H.; Kikuno, R. et al.,(1985):*** Nature 314, 199-203 .
- Twining, A.S. (1977):*** Brecher, Clin. Chim. Acta 75, 143-148.
- Urine, S.; Bree, F.; Testa, et al., (1991):*** PH-dependency of basic ligand binding to alpha 1-acid glycoprotein .Biochem. J.280 (pt 1):277-80 .
- Van, W. and Dijk, (1995):*** Glycoconj. J. 12 (1995) 227 -233 .
- Vanliferighen, P.; Peigue-Lafeuille, H.; Gaulme, et al., (1986):*** C-reactive protein and orosomucoid determinations in a neonatal pathology unit. Pediatrie. 41:121-125.
- Verma, S. (2005):*** Adipokines: molecular links between obesity and atherosclerosis. Am J Physiol Heart Circ Physiol 288(5):H2031-41.

Weimer, H.E.; Mehl, J.W. and Winzler, R.J. (1950): J. Biol. Chem.185
561-568.

Williams, J.P.; Weiser, M.R.; Pechet, T.T. et al., (1997): Am. J. Physiol.
273, G 1031 – 1035 .

Woodrum, D.; Hodson, W.A.; Mayock, D. et al.,(2002): Rule out
Guidelines for Possible Neonatal Sepsis. University of
Washington Academic Medical Center Children's Hospital and
Regional Medical Center. NICU-WEO.