

References

- Abd El-Fattah S, El-Kabbany Z, Awad H, El-Kerdany T and Ibrahim H (1993):*** Pediatric and clinical pathology departments. Ain Shams University: Role of exchange transfusion in the treatment of neonatal septicemia. New Egy J Med; 8(6): 1705-1707.
- Abraham E, Reinhart K, Opals, et al (2003):*** Efficacy and safety of tifacogin (recombinant tissue factor pathway inhibitor) in severe sepsis a randomized controlled trial. JAMA 290(2): 256-8.
- Adams-Chapman I and Stoll BJ (2001):*** Systemic inflammatory response syndrome. Semin Pediatr Infect Dis; 12: 5-16.
- Adams-Chapman I and Stoll BJ (2006):*** Early diagnosis of neonatal infections and its long term neurodevelopmental outcome in preterm infants. Curr Opin Infect Dis; 19(3): 290-297.
- Alving BM and Comp C (1992):*** Recent advances in understanding clotting and evaluating patients with recurrent thrombosis. Am J Obstet Gynecol; 167: 1184-91.
- Anane D, Bellissant E and Cavaillon JM (2005):*** Septic shock. Lancet; 365(9453): 63-78.
- Anderson-Berry AL, Bellig LL and Ohning BL (2006):*** Neonatal sepsis. www.emedicine.com. Editor(s): Scott S Macgilvray, Mary L Windle, David A Clark, Carol L Wagner, Neil N Finer.
- Andrew M, Paes B and Johnston M (1990):*** Development of the hemostatic system in the neonate and young infant. Am J Pediatr Hematol Oncol; 12: 95-104.

- Angus DC and Wax RS (2001):** Epidemiology of sepsis: An update. Crit Care Med; 29: 109-116.
- Anwer SK, Mustafa S, Pariyani S, Ashraf S and Taufiq KM (2000):** Neonatal sepsis: an etiology study. J Pak Med Assoc; 50: 91-94.
- Aronis S, Hatzis A, Platokouki H, et al (1999):** Severe protein C deficiency, predictor of lethal outcome in meningococcal septicemia [abstract 132]. Pediatr Res; 45: 765.
- Baker CJ and Edwards MS (1995):** Group B streptococcal infection. In Klein JO and Remington JJS (eds): Infectious Diseases of Fetus and Newborn Infants. WB Saunders, Philadelphia, P: 98-120.
- Balows A and Sonnenwirth AC (1983):** Bacteremia: Laboratory and clinical aspects. Charles C Thomas, Springfield, III.
- Barton P, Kalil AC, Nadel S, et al (2004):** Safety, pharmacokinetics, and pharmacodynamics of drotrecogin alfa (activated) in children with severe sepsis. Pediatrics; 113: 7-17.
- Berger C, Vehlenger J, Ghelfi D, et al (2003):** Comparison of C-reactive protein and white blood cell count with differential in neonates at risk of septicemia. Eur J Pediatr; 154: 138-44.
- Bertina RM, Koeleman BP, Koster T, Roserdaal FR, Dirven RJ, deRonde H, Vander Velden PA and Ratsma PH (1994):** Mutation in blood coagulation factor V associated with resistance to activated protein C. Nature; 369: 6775, 64-7.
- Bithell TC (1993):** Thrombosis and anti thrombin therapy. In: Lee G, et al., eds. Wintrobe's Clinical Hematology; 2: 1515.

- Bizzarro MJ and Gallagher PG (2007):** Antibiotic-Resistant Organisms in the Neonatal Intensive Care Unit. *Seminars in Perinatology*; 31(1): 26-32.
- Bizzarro MJ, Raskind C, Baltimore RS and Gallagher PG (2005):** Seventy-five years of neonatal sepsis at Yale: 1928-2003. *Pediatrics*; 116(3): 595-602.
- Boldt J, Papsdorf M, Rothe A, Kumle B and Piper S (2000):** Changes of the hemostatic network in critically ill patients-is there a difference between sepsis, trauma, and neurosurgery patients? *Crit Care Med*; 28: 445-450.
- Bonduel M, Hepner M, Scuiccati G, et al (1998):** Infectious purpura fulminans in children: a report on 20 patients from Argentina. *Int J Pediatr Hematol Oncol*; 5: 21-27.
- Busch C, Cancilla P and DeBault L (1992):** Use of endothelium cultured on microcarriers as a model for the microcirculation. *Lab Invest*; 47: 498.
- Campbell MJ and Machin D (1998):** Medical statistics. A common sense. Apparel 2nd ed. John Wildy & Sons New York, Chisherter: Brisbon, Toronto & Singapore.
- Cemeca F, Parco S and Simeone R (1995):** A description of two cases of factor V deficiency Haemophilia I; 200-201.
- Chiesa C, Pellegrini G, Panero A, Osborn JF, Signore F, Assumma M, et al (2003):** C-reactive protein, interleukin-6, and procalcitonin in the immediate postnatal period: influence of illness severity, risk status, antenatal and perinatal complications, and infection. *Clin Chem*; 49: 60-68.

- Dahlback B (1995):** New molecular insights into the genetics of thrombophilia resistance to APC caused by Arg 506 to Gln mutation in FV as a pathogenic risk factor for venous thrombosis. *Thromb Haemost*, Jup; 74(1): 139-48.
- Davis RL, Hasselquist MB and Cardenas V (2001):** Introduction of the new centers for disease control and prevention group B streptococcal prevention guideline at a large West Coast health maintenance organization. *Am J Obstet Gynecol*; 184: 603-610.
- DeSilva GD, Kantzanou M, Justice A, Massey RC, Wilkinson AR, Day NP and Peacock SJ (2002):** The ica operon and biofilm production in coagulase-negative staphylococci associated with carriage and disease in a neonatal intensive care unit. *J Clin Microbiol*; 40: 382-388.
- Drossou-Agakidou V, Roilides E, Papakyriakidou-Koliouska P, et al (2004):** Use of ciprofloxacin in neonatal sepsis: lack of adverse effects up to one year. *Pediatr Infect Dis J*; 23(4): 346-9.
- Ehls Gering B, Bartmann P, et al (2001):** C-reactive protein is a useful marker for guiding duration of antibiotic therapy in suspected neonatal bacterial infections. *Pediatrics*; 99: 216-21.
- El-Alfy MS and Awad HA (1993):** The value of intravenous immunoglobulin therapy in the management of septic preterm neonates. *The Egypt J Pediatr*; 10: 227-236.
- El-Beshlawy A, Alaraby I, Abou Hussein H, Abo-Elew HH and Mohamed Abdelkader MS (2010):** Study of protein C, protein S and antithrombin III in newborns with sepsis. *Pediatr Crit Care Med*; 11(1): 52-9.

- Escobar GJ (2003):** Effect of systemic inflammatory response on biochemical markers of neonatal bacterial infections: A fresh look at old cofounders. Clin Chem; 49: 21-22.
- Esmon CT (1993):** The roles of protein C and thrombomodulin in the regulation of blood coagulation. J Biol Chem; 264: 4743.
- Fay PJ (1993):** Factor VIII structure and function: thromb Haemost; 70: 63-67.
- Fay WP, Shapiro AD and Shih JL (1992):** Brief report: complete deficiency of plasminogen-activator inhibitor type 1 due to a frameshift mutation. N Engl J Med; 327: 1729.
- Fidelma RB and Nolan TE (1995):** Inherited disorders of coagulation in pregnancy. Clinical Obstet Gynecol; 38(3): 497-513.
- Fijnvandraat K, Derkx B, Peters M, et al (1995):** Coagulation activation and tissue necrosis in meningococcal septic shock: severely reduced protein C levels predict a high mortality. Thromb Haemost; 73: 15-20.
- Fleisher GR and Ludwig S (2001):** Textbook of pediatric emergency medicine. Philadelphia: Lippincott, Williams & Wilkins.
- Fourrier F, Chopin C, Goudemand J, et al (1992):** Septic shock, multiple organ failure, and disseminated intravascular coagulation: compared patterns of antithrombin III, protein C and protein S deficiencies. Chest; 101: 816-823.
- Garcia-Prats J, Cooper R, Schneider V, Stager C and Hansen T (2000):** Rapid detection of microorganisms in blood cultures of newborn infants utilizing an automated blood culture system. Pediatrics; 105: 523-527.

- Gardiner JE, McGann MA and Berridge CW (1996):*** Protein S as a cofactor for activated protein C in plasma and in the inactivation of purified factor VIII:C. *Circulation*; 2011.
- Goldmann D and Pier GI (1993):*** Pathogenesis of infection related to intravascular catheter. *Clin Microbial Rev*; P: 176.
- Gottof GL (2000):*** Risk diagnosis of neonatal sepsis. *Pediatrics Inf Dis*; 12: 205-5.
- Gregory I, Yin T Hui, Ruth Milner and Alvin Zipursky (2003):*** Haematological changes in neonatal sepsis in pediatrics. Vol 54, No. 38-42.
- Griffin JH, Mosher DF, Zimmerman TS and Kleiss AJ (1992):*** Protein C and antithrombotic protein C, an antithrombotic protein, is reduced in hospitalized patients with intravascular coagulation. *Blood*; 60: 261.
- Guerina NG (1998):*** Bacterial and fungal infection; by Cloherty JP and Stark AR (eds). In manual of neonatal care, 4th ed., Lippincott Raven Publishers, Philadelphia, New York, 271-300.
- Hajiehe Borna and Sedighen Borna (2005):*** Value of laboratory tests and C-reactive proteins in detection of Neonatal sepsis. *The Internet Journal of Pediatric and Neonatology*, Vol. 5, No. 2.
- Haley M, Cui X, Minneci PC, et al (2004):*** Recombinant human activated protein C in sepsis: assessing its clinical use. *Am J Med Sci*; 328(4): 215-9.
- Hamada S, Vearncombe M, McGeer A and Shah PS (2008):*** Neonatal group B streptococcal disease: incidence, presentation, and mortality. *J Matern Fetal Neonatal Med*; 21(1): 53-7.

- Hancock WW, Grey ST, Hau L, et al (1995):** Binding of activated protein C to a specific receptor on human mononuclear phagocytes inhibits intracellular calcium signaling and monocyte-dependent proliferative responses. *Transplantation*; 60: 1525-1532.
- Haque KH (2006):** Management of bacterial infection in the newborn Reader in Neonatal Medicine, University of London, Consultant Neonatologist, St Helier University Hospital, Carshalton, Surrey, UK. *Journal of the Arab Neonatology Forum*, 3, 2.
- Hart S, Ross J, Haslett C and Dransfield I (2003):** Molecular characterisation of the surface of apoptotic neutrophils: implications for functional downregulation and recognition by phagocytes. *Cell Death Differ*; 7: 493-503.
- Hartman DL, Helterbrand JD and Bernard GR (1997):** Protein C (PC) levels in sepsis: association with mortality [abstract]. *Am J Respir Crit Care Med*; 155: A708.
- Hazelzet JA, Risseuw-Appel IM, Kornelisse RF, et al (1996):** Age-related differences in outcome and severity of DIC in children with septic shock and purpura. *Thromb Haemost*; 76: 932-938.
- Hellgren M, Svensson PJ and Dahlback B (1995):** Resistance to activated protein C as a basis for venous thromboembolism associated with pregnancy and oral contraceptives. *Am J Obstet Gynecol*; 273: 210-3.
- Hesseltvik JF, Malm J, Dahlback B and Blomback M (1991):** Protein C, protein S and C4b binding protein in severe infection and septic shock. *Thromb Haemost*; 65: 126-129.

- Jaffe RI, Lane JD, Albury SV and Niemeyer DM (2000):** Rapid extraction from and direct identification in clinical samples of methicillin-resistant staphylococci using the PCR. *J Clin Microbiol*; 38: 3407-3412.
- Jenson H and Pollock B (1998):** The role of intravenous immunoglobulin for the prevention and treatment of neonatal sepsis. *Semin Perinatol*; 22(1): 50-63.
- Jerone DOK and Micheal M (1995):** Bacterial sepsis and meningitis. In: textbook of infectious diseases of the fetus and newborn infants. 4th ed (eds). Lacks Remington, Jeromo, Klein, WB Saunders Company, Philadelphia, London, Toronto, Sydney, Tokyo; PP: 836-879.
- Johnson AE, Esmon NL, Laue TM and Esmon CT (1993):** Structural changes required for activation of protein C are induced by Ca²⁺ binding to a higher affinity site that does not contain gamma-carboxyglutamic acid. *J Biol Chem*; 258: 5554.
- Kisiel W and Davie EW (1991):** Protein C, *Methods Enzymol*; 80: 320.
- Kite P and Millar M (2001):** Diagnosis of neonatal bacteremia using combination of tests. *Midwifery*; 4: 123-9.
- Klein JO (2001):** Bacterial sepsis and meningitis. In: Remington JS, Klein JO, eds. *Infectious Diseases of the Fetus and Newborn Infant*. Philadelphia, PA: WB Saunders, P: 944.
- Kliegman RM, Behrman RE, Jenson HB and Stanton BF (2007):** Nelson Textbook of pediatrics, 18th Edition, Part XVI: infectious Diseases.

- Klinger G, Chin CN, Beyene J and Perlman M (2000):** Predicting the outcome of neonatal bacterial meningitis. *Pediatrics*; 106: 477-482.
- Kosalaraksa P, Janthep P, Jirapradittha J, et al (2004):** Once versus twice daily dose of gentamicin therapy in Thai neonates. *J Med Assoc Thai*; 87(4): 372-6.
- Leng X, Wang C, Song L, et al (2003):** Generation of recombinant human tissue factor pathway inhibitor in E-coli, *Sheng Wu Yi Xue Gong Cheng Xue Za Zhi*; 20(2): 314-7.
- Levi M, Tencate LH, Vander Poll T and Van Denventer SH (2000):** Pathogenesis of thrombocytopenia and disseminated intravascular coagulation in sepsis. *Journal of American Medical Association*; 270: 975-979.
- Long GL, Marshall A, Gardner JC and Naylor SL (1990):** Genes for vitamin K dependent plasma protein C and protein S are located on chromosome 2 and 3 respectively. *Somatcell Molec Genet*; 14: 93.
- Malik A, Hui CPS, Pennie RA and Kirpalani H (2003):** Beyond the complete blood cell count and C-reactive protein. A systematic review of modern diagnostic tests for neonatal sepsis. *Arch Pediatr Adolesc Med*; 157: 511-516.
- Mann RG, Jenny RJ and Krishnaswamy S (1990):** Cofactor proteins in the assembly and expression of blood clotting enzyme complexes. *Anny Rev B Biochem*; 57: 951-956.
- Mcintosh N (1998):** The Newborn. In: Forfar and Arneil's textbook of pediatrics. 5th edition AGM Campbell and N Macintosh (eds). P: 273-309.

- Mehr S and Doyle LW (2000):** Cytokines as markers of bacterial sepsis in newborn infants: a review. *Pediatr Infect Dis J*; 19: 879-887.
- Miletich JP and Broze GJ (1990):** Beta protein C is not glycosylated at asparagine 329: The rate of translation may influence the frequency of usage at asparagine X-Cysteine sites. *J Biol Chem*; 265: 11397.
- Miletich JP, Leykam JF and Broze GJ (1993):** Detection of single chain protein C in human plasma. *Blood*; 62: 306.
- Mohan P and Brocklehurst P (2003):** Granulocyte transfusion for neonates with confirmed or suspected sepsis and neutropenia. *Cochrane Database Syst Rev*; 4: CD 003956.
- Mtitimila E and Cooke R (2004):** Antibiotic regimens for suspected early neonatal sepsis. *Cochrane Database Syst Rev*; 18(4): CD 004495.
- Ng PC (2004):** Diagnostic markers of infection in neonates. *Archives of Disease in Childhood Fetal and Neonatal Edition*; 89(3): 229-35.
- Ng PC and Like Wong RPO (2002):** Neutrophil CD4 expression as sensitive diagnostic marker in culture –ve symptomatic neonatal sepsis in LBW infants. *Pediatr Res*; 51: 296-303.
- Nick NA, Coldern CD, Geraci MW, et al (2004):** Recombinant human activated protein C reduces human endotoxin-induced pulmonary inflammation via inhibition of neutrophil chemotaxis. *Blood*; 104(13): 378-85.
- Nicoles GA, Tans G and Christella M, et al (1995):** Peptide bond cleavages and loss of functional activity during inactivation of

factor V and factor Va 506Q by activated protein C. J Biochem; 270: 21108-21166, Biochem.

Oddo M, Schaller MD, Calandra T and Liaudet L (2004): New therapeutic strategies in severe sepsis and septic shock. Rev Med Suisse Romande; 124(6): 329-32.

Ohlin AK, Linse S and Stenflo J (1993): Calcium binding to the epidermal growth factor homology region of bovine protein C. J Biol Chem; 263: 7411.

Pepys MB (1981): C-reactive protein fifty years on. Lancet; 1: 653-657.

Philip AGS and Herwitt JR (2003): Early diagnosis of neonatal sepsis. Pediatrics; 65: 1036-41.

Phillipe J, Offner F, Declerck PJ, Leroux-Roels G and Vogelaers D (1991): Fibrinolysis and coagulation in patients with infectious disease and sepsis. Thromb Haemost; 65: 291-295.

Plutzky J, Hoskins JH, Long GL and Carbtree GR (1996): Evaluation and organization of the human protein C gene. Proc Natl Acad Sci USA; 83: 546.

Raggio MJ and Morris PE (2004): Drotrectogin alfa. Drugs today (Barc); 40(6): 517-38.

Rice TW and Bernard GR (2004): Drotecogin alfa (activated) for the treatment of severe sepsis and septic shock. Am J Med Sci; 328(4): 205-14.

Roberton NRC (1992): Infection in the newborn. In textbook of neonatology. 2nd edition. Churchill-Livingstone. New York and Tokyo; P: 191-226.

Roberton NRC (1993): Neonatal infection. In: Manual of neonatal intensive care, 1st ed. Churchill-Livingstone, New York and Tokyo; P: 191-226.

Roberton NRC and Rennie JM (1999): Infection in the newborn: by Dear P (ed): In textbook of neonatology, 3rd edition. Churchill-Livingstone; 1109-1139.

Rubin LG, Sanchez PJ, Siegel J, Levine G, Saiman L, Jarvis WR and Pediatric Prevention Network (2002): Evaluation and treatment of neonates with suspected late onset sepsis: A survey of neonatologist practice. Pediatrics; 110: 1-7.

Sakata Y, Loskutoff DJ and Gladson CL (1993): Mechanism of protein C-dependent clot lysis Role of plasminogen activator inhibitor. Blood; 68: 1218.

Salama AA, Metwally AY and Farid FA (1996): M.Sc. Thesis Pediatrics, Ain Shams University.

Schrag S, Gorwitz R, Fultz-Butts K and Schuchat A (2002): Prevention of perinatal group B streptococcal disease. Morbidity and Mortality Weekly Report; 51: 1-22.

Schrag SJ, Zell ER, Lynfield R, Roome A, Arnold KE, Craig AS, Harrison LH, Reingold A, Stefonek K, Smith G, Gamble M, Schuchat A; Active Bacterial Core Surveillance Team (2002): A population-based comparison of strategies to prevent early-onset group B streptococcal disease in neonates. N Engl J Med; 347: 233-239.

St Geme JW 3rd and Polin RA (1996): Prevention and treatment of neonatal sepsis. In Spitzer AR (ed). Intensive care of the fetus and neonates (ch. 76). St. Louis Mosby, Yearbook, Inc.

- Stenflo J and Fenlund P (1992):** Amino acid sequence of the heavy chain of bovine protein C. *J Biol Chem*; 257: 12180.
- Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, Lemons JA, Donovan EF, Stark AR, Tyson JE, Oh W, Bauer CR, Korones SB, Shankaran S, Laptook AR, Stevenson DK, Papile LA and Poole WK (2002):** Changes in pathogens causing sepsis in very low birth weight Infants. *N Engl J Med*; 347: 240-247.
- Stoll BJ, Hansen N, Fanaroff AA, Wright LL, Carlo WA, Ehrenkranz RA, Lemons JA, Donovan EF, Stark AR, Tyson JE, Oh W, Bauer CR, Korones SB, Shankaran S, Laptook AR, Stevenson DK, Papile LA and Poole WK (2002):** Late onset sepsis in very low birth weight neonates: The experience of the NICHD Neonatal Research Network. *Pediatrics*; 110: 285-91.
- Straka M, Dela Cruz W, Blackmon C, Johnson O, Stassen S, Streitman D, Golden S and Stamilio D (2004):** Rapid detection of group B streptococcus and Escherichia coli in amniotic fluid using real-time fluorescent PCR. *Infect Dis Obstet Gynecol*; 12: 109-114.
- Venkataseshan S, Dutta S, Ahluwalia J and Narang A (2007):** Low plasma protein C values predict mortality in low birth weight neonates with septicemia. *Pediatr Infect Dis J*; 26(8): 684-8.
- Webber S, Lindell D and Wilkinson AR (1990):** Neonatal pneumonia. *Arch Dis Child*; 65: 207-211.
- Weber MW, Carlin JB, Gatchalian S, Lehmann D, Muhe L, Mulholland EK, WHO Young Infants Study Group (2003):** Predictors of neonatal sepsis in developing countries. *Pediatr Infect Dis J*; 22: 711-716.

Yurdakok M (1998): Antibiotic use in neonatal sepsis. Turkish Journal of
Pediatr; 40(1): 17-33.

Zipursky A, Palko J, Milner R and Akenzua GI (1976): The hematology
of bacterial infections in premature infants. Pediatrics; 57: 839-
853.