

References

A

- Aida S. (1993):** Alkaline phosphatase isoenzyme activities in rheumatoid arthritis: hepatobiliary enzyme dissociation and relation to disease activity. *Ann. Rheum. Dis.*, 55 : 520 – 524.
- Aitchenson C.; Pebles C. and Joslin H. (1992):** Characteristics of antinuclear antibodies in rheumatoid arthritis. *Arch. Rheumatol.*, 26 : 562.
- Alamanos Y. and Drosos A. (2005):** Epidemiology of adult rheumatoid arthritis. *Autoimmun Rev.*,4:130-6.
- Alamanos Y.; Voulgari P. and Drosos A. (2006):** Incidence and prevalence of rheumatoid arthritis, based on the 1987 American College of Rheumatology criteria: a systematic review. *Arthritis Rheum.*,36:182-8.
- Alasaarela I.; Tervonen O.; Drosos A. et al. (1998):** Evaluation of humeral head erosions in rheumatoid arthritis: a comparison of ultrasonography, magnetic resonance imaging, computed tomography and plain radiography. *Br J Rheumatol.*,37:1152-1156.
- Aletaha D., Neogi T., Silman A.J., et al. (2010):** 2010 rheumatoid arthritis classification criteria: an American College of Rheumatology/European League Against Rheumatism collaborative initiative. *Ann. Rheum. Dis.*, 69 (9): 1580–8.
- Alparslan L.; Yu J.S. and Weissmann B.N. (2005):** Imaging. In Harris E.D., Budd R.C., Firestein G.S., Genovese M.C.,

- Sergent J.S., Ruddy S. and Sledge C.B. (eds.): Kelley's Textbook of Rheumatology, Philadelphia, Pennsylvania. Elsevier, Saunders, 7th ed., PP: 739 – 797.
- Amos C.; Chen W.; Michael F. et al. (2009):** Data for Genetic Analysis Workshop 16 Problem 1, association analysis of rheumatoid arthritis data. BMC Proceedings.,3:1-4.
- Anderson J.; Wells G.; Verhoeven C. et al.(2000):** Factors predicting response to treatment in rheumatoid arthritis: the importance of disease duration. Arthritis and Rheumatism., 43: 22–29.
- Armstrong J. K.; Wenby R. B.; Meiselman H. J. et al.(2004):** Fisher The Hydrodynamic Radii of Macromolecules and Their Effect on Red Blood Cell. Aggregation Biophys J. , 87(6): 4259–4270s.
- Arnett F.; Edworthy S.; Bloch D. et al. (1988):** The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. Arthritis Rheum.,3:315–24.
- Astbury C.; Platt R.; Dixon J. et al. (1993):** Optimizing the assessment of disease activity during treatment with anti-rheumatoid drugs. British Journal of Rheumatology., 32: 467–473.
- Avnon L.S.; Abu-Shakra M.; Flusser D. et al. (2007):** Pleural effusion associated with rheumatoid arthritis: what cell predominance to anticipate? Rheumatol Int., 27:919-25.

B

Baan H.; Haagsma C. and van de Laar M.(2006): Corticosteroid injections reduce size of rheumatoid nodules. Clin Rheumatol., 25:21-23.

Backhaus M.; Kamradt T. and Sandrock D. (1999): Arthritis of the finger joints: a comprehensive approach comparing conventional radiography, scintigraphy, ultrasound, and contrast-enhanced magnetic resonance imaging. Arthritis Rheum.,42:1232-1245.

Backhaus M.; Ohrndorf S.; Kellner H. et al.(2009): Evaluation of a Novel 7-Joint Ultrasound Score in Daily Rheumatologic Practice: A Pilot Project . Arthritis & Rheumatism (Arthritis Care & Research).,9: 1194–1201.

Backhaus M.; Schmidt W.; Mellerowicz H. et al. (2002): Technique and diagnostic value of musculoskeletal ultrasonography in rheumatology. Part 6: ultrasonography of the wrist/hand. Z Rheumatol.,61:674– 87.

Baer A.; Dessypris E. and Goldwasser E. (1993): Blunted erythropoietin response to anemia in rheumatoid arthritis. Medicine.,66 : 599 -602.

Bahremund M. and Schumacher H. (1991): Effect of medication on synovial fluid leukocyte differentials in patients with rheumatoid arthritis. Arthritis and Rheumatism., 34 : 1173 - 1176.

- Balint P.; Kane D. and Sturrock R. (2001):** Modern patient management in rheumatology : interventional musculoskeletal ultrasonography. *Osteoarthritis cartilage* ., 9 : 509 – 111.
- Bellamy N.; Campbell J. and Syrotuik J. (1999):** Comparative study of self-rating pain scales in rheumatoid arthritis patients. *Current Medical Research and Opinion.*, 15: 121–127.
- Bensen W.; Tugwell P.; Roberts R. et al. (2004):** Combination therapy of cyclosporine, methotrexate and gold in rheumatoid arthritis (2 pilot studies). *J. Rheumatol.*, 21 : 2034 – 2038.
- Bernard C. (2009):** Progression in early rheumatoid arthritis. *Best Practice & Research Clinical Rheumatology.*, 23 59–69.
- Blom. and van Riel. (2007):** Management of established rheumatoid arthritis with an emphasis on pharmacotherapy. *Best Practice & Research Clinical Rheumatology* ., 21: 43-57.
- Boumans J. and Tak P. (2009):** Rituximab treatment in rheumatoid arthritis: how does it work?. *Arthritis Research & Therapy* ., 11:134.
- Boutry N.; Morel M.; Flipo R. et al.(2007):** Early rheumatoid arthritis: a review of MRI and sonographic findings. *AJR.*,189:1502–9.
- Braun J.; Ballow M. and Eggens U. (1994):** Use of dynamic magnetic resonance imaging with fast imaging in the detection of early and advanced sacroilitis in spondylarthropathy patients. *Arthritis Rheumatol.*, 37 : 1039.
- Breedveld F.; Weisman M.; Kavanaugh A. et al. (2006):**The PREMIER study: a multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone

- in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis Rheum.*,54:26–37.
- Brower A. (1994):** Rheumatoid Arthritis Imaging. In Klippel, J.H., Dieppe, P.A. (eds.): *Rheumatology*. Mosby. Baltimore, Boston, Chicago, London, Philadelphia, Sydney, Toronte, 3rd ed.,175- 205..
- Brower A.C. (1990):** Use of the radiograph to measure the course of rheumatoid arthritis disease. *Arthritis Rheum.*, 33 : 316 – 24.
- Brown A.; Quinn M.; Karim Z. et al.(2006):** Presence of significant synovitis in rheumatoid arthritis patients with disease-modifying antirheumatic drug-induced clinical remission. Evidence from an imaging study may explain structural progression. *Arthritis Rheum.*,54:3761–3773.
- Bruce B. and Fries J. (2003):** The Stanford health assessment questionnaire (HAQ): a review of its history, issues, progress, and documentation. *J Rheumatol.*,30: 167-78.
- Buchbinder R.; Bombardier C.; Yeung M. et al.(1995):** Which outcome measures should be used in rheumatoid arthritis clinical trials? Clinical and quality-of-life measures' responsiveness to treatment in a randomized controlled trial. *Arthritis and Rheumatism.*, 38: 1568–1580.
- Burtner P.; Anderson J.; Marcum M. et al.(2003):** A comparison of static and dynamic wrist splints using electromyography in individuals with rheumatoid arthritis. *J Hand Ther.*,16: 320–325.

C

- Carole R.; Etchepare F.; Bruno. Et al. (2009):** Diagnosis of synovitis by ultrasonography in RA: A one-year experience is enough for reliability on static images. *Joint Bone Spine.*, 76., 35-38.
- Carotti M.; Salaffi F.; Manganelli P. et al.(2002):** Power Doppler sonography in the assessment of synovial tissue of the knee joint in rheumatoid arthritis: a preliminary experience. *Annals of the Rheumatic Diseases.*,61: 877–882.
- Chan, E. and Cronstein, B. (2002):** Molecular action of methotrexate in inflammatory diseases. *Arthritis Res.*,4:266–73.
- Chen J.; Branstetter B. and Myers E.(2005):** Cricoarytenoid rheumatoid arthritis: an important consideration in aggressive lesions of the larynx. *AJNR Am J Neuroradiol.*, 26:970-2.
- Clair E.; van der Heijde; Smolen J. et al.(2004):** Combination of infliximab and methotrexate therapy for early rheumatoid arthritis: a randomized, controlled trial. *Arthritis Rheum.*,50(11):3432–43.
- Coakley G. (2000):** Major histocompatibility complex haplotypic associations in Felty's syndrome and large granular lymphocyte syndrome are secondary to allelic association with HLA-DRB1*0401, *Rheumatology (Oxford)* 39 (4) 393–398.
- Cohen S.; Emery P.; Greenwald M. et al. (2006):** Rituximab for rheumatoid arthritis refractory to antitumor necrosis factor therapy: Results of a multicenter, randomized, double-blind, placebo-controlled, phase III trial evaluating primary efficacy

- and safety at twenty-four weeks. *Arthritis & Rheumatism*,54(9):2793–806.
- Cohen S.; Hurd E.; Cush J. et al.(2002):** Treatment of rheumatoid arthritis with anakinra, recombinant human interleukin-1 receptor antagonist, in combination with methotrexate. *Arthritis Rheum*.,46(3):614-624.
- Cohen S.; Keystone E.; Genovese M. et al.(2010):** Continued inhibition of structural damage over 2 years in patients with rheumatoid arthritis treated with rituximab in combination with methotrexate. *Ann Rheum Dis*.,69:1158.
- Combe B.; Landewe R.; Lukas C.; et al.(2007):** EULAR recommendations for the management of early arthritis: report of a task force of the European Standing Committee for International Clinical Studies Including Therapeutics (ESCISIT). *Ann Rheum Dis*.,66(1):34–45.
- Condemi J.; Branett E. and Atwater E. (1995):** The significance of antinuclear factors in rheumatoid arthritis. *Arth. Rheum.*, 8 :1075: 1080.
- Cooper C. and Kirwan J. (1990):** The risk of local and systemic corticosteroid administration. *Baillieres Clinical Rheumatol.*, 4 : 305 – 32.
- Cooper C.; Coupland C. and Mitchell M. (1995):** Rheumatoid arthritis, corticosteroid therapy & the hip fracture. *Ann. Rheum. Dis.*, 54 : 49 – 52.
- Costenbader K. and Laden F. (2008):** Geographic variation in rheumatoid arthritis incidence among women in the United States. *Arch Intern Med*.,168:1664-70.
- Costenbader K.; Chang S.; Plenge R.et al. (2008):** Genetic polymorphisms in PTPN22, PADI-4, and CTLA-4 and risk for

rheumatoid arthritis in two longitudinal cohort studies: evidence of gene environment interactions with heavy cigarette smoking. *Arthritis Res Ther.*,10: 52-60..

Cunnane G.; Grehan S.; Geoghegan S. et al. (2000): Serum amyloid A in the assessment of early inflammatory arthritis. *The Journal of Rheumatology.*,27: 58–63.

Cutolo M.; Villaggio B.; Craviotto C. et al. (2002): Sex hormones and rheumatoid arthritis. *Autoimmun Rev.*,1:284-9.

D

D’Agostino MA. ; Said-Nahal R.; Hacquard-Bouder C. et al.(2003) : Assessment of peripheral enthesitis in the spondylarthropathies by ultrasonography combined with power Doppler: a cross-sectional study. *Arthritis Rheum.*,48: 523–33.

Davis D., et al., (1997): Anaemia of chronic disease in rheumatoid arthritis: in vivo effects of tumour necrosis factor alpha blockade, *Br. J. Rheumatol.* 36 (9) 950–956.

De Palma L.; Chillemi C.;Albanelli S. et al. (2000) : Muscle involvement in rheumatoid arthritis: an ultrastructural study. *Ultrastruct Pathol.*;24:151-6.

De Villeroché J.V.; Avouac J.; Ponceau A. et al. (2010): Enhanced late-outgrowth circulating endothelial progenitor cell levels in rheumatoid arthritis and correlation with disease activity. *Arthritis Res Ther.*, 12: 27-33.

Deborah J.; Rubens; Bhatt S.et al.,(2006) : Doppler Artifacts and Pitfalls. *Ultrasound clinics*.,1: 79-109.

Donnelly E.F.; Geng L.; Wojcicki W.E. et al.(2001): Quantified power Doppler US of tumor blood flow correlates with microscopic quantification of tumor blood vessels. *Radiology*;219:166–70.

E

Edmonds J.P. (1997): Rheumatoid arthritis. In Howe H.S. and Feng, P.H. (eds): *Textbook of clinical rheumatology*, Singapore,1st ed., PP: 54 -111.

Edwards J.C.; Szczepanski L.; Szechinski J. et al (2004): Efficacy of B-cell-targeted therapy with rituximab in patients with rheumatoid arthritis. *New England Journal of Medicine*.,350:2572–81.

Elliott J.R. and O'Dell (2002): Rheumatoid Arthritis. In sterling G. (ed.) : *Rheumatology Secrets*, Philadelphia, Hanley and Belfus INC., 2nd ed., PP : 117 – 128.

Emery P.(2006): Treatment of rheumatoid arthritis. *BMJ*.,21:152-5.

Evrin P.E.; Nilsson S.E.; Oberg T. and Malmberg B. (2005): Serum C-reactive protein in elderly men and women and various biochemical values. *Scand J Clin Lab Invest.*, 65:23-31.

Ewa M. and Paleolog. (2009): The vasculature in rheumatoid arthritis: cause or consequence?. *Int. J. Exp. Path.*, 90:249–261.

F

Fasford, J.R. (1990): Electrical Therapy. In Kottke Lehmann (eds.) : Krusen's Handbook of Physical Medicine and Rheabilitation, Sounders Company. Philadilphia, London, Toronto, Sydney, Tokyo W.B., 4th ed .,p. 278.

Felson D.T.; Anderson J.J. and Meenan R.F. (1990): The comparative efficacy and toxicity of second – line drugs in RA. Results of two meta-analysis. Arthritis. Rheum., 33 : 1449 – 1461.

Felson D.T.; Anderson J.J.; Boers M. et al. (1993): The American College of Rheumatology preliminary core set of disease activity measures for rheumatoid arthritis clinical trials. Arthritis and Rheumatism., 36: 729–740.

Felson D.T.; Anderson J.J.; Boers M. et al.(1995): American College of Rheumatology preliminary definition of improvement in rheumatoid arthritis. Arthritis and Rheumatism., 38: 727–735.

Ferraz B. M.; Quaresma M.R.; Aquino L.R.L. et al. (1990): Reliability of pain scales in the assessment of literate and illiterate patients in rheumatoid arthritis. The Journal of Rheumatology., 17: 1022–1024.

Fessell D.P.; Jacobson J.A.; Craig J. et al. (2000): Using sonography to reveal and aspriate joint effusions. Am. Roentgenol., 174 : 1353 – 62.

Filippucci E.; Iagnocco A.; Salaffi F.et al.(2006) :Power Doppler sonography monitoring of synovial perfusion at the wrist

joints in patients with rheumatoid arthritis treated with adalimumab. *Ann Rheum Dis.*,65: 1433–7.

Firestein G.S. (2001): Etiology and pathogenesis of rheumatoid arthritis. In Kelley W.N., Ruddy S., Harris E.D., and Sledge C.B.(eds): *Kelley's Textbook of Rheumatology*, W.B. Saunders Company. Philadelphia, 6th ed., PP. 924-966.

Forslind K.; Ahlmen M.; Eberhardt K. et al.(2004): Prediction of radiological outcome in early rheumatoid arthritis in clinical practice: role of antibodies to citrullinated peptides (anti-CCP). *Ann Rheum Dis.*,63:1090– 5.

Fournié B.; Margarit-Coll N.; Champetier de Ribes T.L.; et al.(2006): Extrasynovial ultrasound abnormalities in the psoriatic finger. Prospective comparative power-Doppler study versus rheumatoid arthritis. *Joint Bone Spine.*,73:527–31.

Fowler N.K. and Nicol, A.C.(2001): Functional and biomechanical assessment of the normal and rheumatoid hand. Elsevier Ireland Ltd.,16:660-666.

Fox R.I. (1998): Safety and effectiveness of leflunomide in the treatment of patients with active rheumatoid arthritis. *Arthritis Rheum.*, 8 (11): 1595.

Freeston E.; Brown A. K.; Hensor A. et al.(2008): Extremity magnetic resonance imaging assessment of synovitis (without contrast) in rheumatoid arthritis may be less accurate than power Doppler ultrasound*Ann Rheum Dis.*,67:1351-1358.

Fukae J.; Shimizu M.; Kon Y. et al. (2009): Screening for rheumatoid arthritis with finger joint power Doppler ultrasonography: quantification of conventional power

Doppler ultrasonographic scoring Mod Rheumatol .,19:502–506.

Fuleihan R.; Ramesh N.; Horner A. et al. (1994): Cyclosporine A inhibits CD40 ligand expression in T lymphocytes. J. Clin. Invest., 93 : 1315.

G

Gabriel J.; Youinou T. and Alain Saraux (2010): The environment, geo-epidemiology, and autoimmune disease: Rheumatoid arthritis. Journal of Autoimmunity.,10:1-5.

Gabriel S.E.; Crowson C.S.; Kremers H.M. et al (2003): Survival in rheumatoid arthritis: a population-based analysis of trends over 40 years. Arthritis and Rheumatism.,48(1): 54–58.

Gaëlle C.; Bessis N. and Boissier M. (2003): Recent data on the role for angiogenesis in rheumatoid arthritis. Joint Bone Spine., 70: 321–326.

Gaffney K. and Scott D.G. (1998): Azathioprine and cyclophosphamide in the treatment of rheumatoid arthritis. Br. J. Rheumatol., 37 : 824.

Ganguly K.S. (1998): A study of the effects of pulsed electromagnetic field therapy with respect to serological grouping in rheumatoid arthritis. J. Indian Med. Assoc., 96 (9): 272 – 275.

García V. (2007): Rheumatoid Nodule. Elsevier Inc.,2:7-15.

Gaschen.; Lorrie D.V.M. and Jan H. (2001): Contribution of Power Doppler Sonography to the Detection of Renal Allograft

- Rejection in the Cynomolgus Monkey Investigative Radiology.,6:335-340.
- Genovese M.C. and Harris E.D. (2005):** Rheumatoid Arthritis (treatment of rheumatoid arthritis). In Harris E.D., Budd R.C., Genovese M.C., Firestein G.S and Sargent J.S (eds.): Kelley's Textbook of Rheumatology, Elsevier Saunders. Philadelphia, Pennsylvania, 7th ed., PP : 1079 – 1100.
- Georg S (2006):** Rheumatoid arthritis: inflammation and bone loss .Wien Med Wochenschr .2: 34–41.
- Gerhard H. and Mostech. (2006):** Doplex and color Doppler imaging of the venous system .Medical radiology .,2:19-7.
- Gerlag D.M.; Tak P.P. (2008):** Novel approaches for the treatment of rheumatoid arthritis: lessons from the evaluation of synovial biomarkers in clinical trials. Best Pract Res Clin Rheumatol ., 22:311-323.
- Ghanem N.; Uhl M.; Pache G. et al.(2007):** MRI in psoriatic arthritis with hand and foot involvement. Rheumatol Int.,27:387–93.
- Goldbach R.;;Lee J.; McCoy A. et al. (2002):** Rheumatoid arthritis associated auto antibodies in patients with synovitis of recent onset. Arthritis Res.,3:236–43.
- Gonzalez A.; Maradit H.; Crowson C.S. et al. (2007):**The widening mortality gap between rheumatoid arthritis patients and the general population. Arthritis Rheum.,56:3583-7.
- Goodson N. (2002):** Coronary artery disease and rheumatoid arthritis. Curr. Opin. Rheumatol., 14 (2) 115–120.

- Gordan and Hasting (1998):** Rheumatoid arthritis: Clinical features of early and progressive disease. *Ann. Rheum. Dis.*, 57: 534-542.
- Gordon D.A. and Klinkhoff A.V. (2005):** Second line agents. In Harris E.D., Budd R.C., Firestein G.S., Genovese M.C., Sargent J.S., Ruddy S. and Sledge C.B. (eds.) : *Kelley's Textbook of Rheumatology*, Elsevier. Philadelphia, Pennsylvania, 7th ed., PP: 877 – 899.
- Goupille P.; Fouquet B.; Goga D. et al. (1990):** The tempromandibular joint in rheumatoid arthritis: Correlations between clinical and computed tomography features. *J Rheumatol.*,17: 1285.
- Grassi W.; Farina A.; Filippucci E. et al. (2002):** Intralesional therapy in carpal tunnel syndrome: a sonographic- guided approach. *Clin Exp Reumatol.*, 20:73-76.
- Grassi W.; Filippucci E.; Farina et al. (2000):** Sonography imaging of the distal phalanx. *Semin Arthritis Rheum.*, 29:379-384.
- Grassi W.; Titarelli E.; Pirani O; et al. (2001):** Ultrasound examination of the metatarsophalangeal joints in rheumatoid arthritis. *Scand J Rheumatol.*,20:163-172.
- Grechenig W.; Clement H.; Bratschitsch G. et al (2002) :** Ultrasound diagnosis of the Achilles tendon. *Orthopade.*, 31:319-325.
- Groll D.L.; To T.; Bombardier C. et al. (2005):** The development of a comorbidity index with physical function as the outcome. *Journal of Clinical Epidemiology.*,58(6): 595–602.

H

Halligan C.S.; Bauch C.D.; Brey R.H. et al. (2006): Hearing loss in rheumatoid arthritis . *Laryngoscope*.,116:2044-9.

Hameed B.; Pilcher J. Heron C. et al.(2008): The relation between composite ultrasound measures and the DAS28 score, its components and acute phase markers in adult RA. *Rheumatology* .,47(4):476–80.

Harris E. D. (2005): Clinical features of rheumatoid arthritis .:In Ruddy S., Harris E.D., Sledge C.B., Kelley W.N. (eds). *Kwilly's Textbook of rheumatology*. WB Saunders. Philadelphia, 7th ed.,PP. 1043-78.

Harris E.D. (2001): Clinical features of rheumatoid arthritis. In Kelley W.N., Ruddy S., Harris E.D. and Sledge C.B (eds). *Kelley's Textbook of Rheumatology*. W.B. Saunders Company. Philadelpha, 6th ed.,PP. 967-1000.

Hart J.E.; Laden F.; Puett R.C. et al. (2009): Exposure to traffic pollution and increased risk of rheumatoid arthritis. *Environ Health Perspect* ., *Rheumatology* .,47:466–60.

Hassell A.B.; Davis M.J. and Fowler P.D. (1995): The relationship between serial measures of disease activity and outcome in rheumatoid arthritis. *Q. J. Med.*.,86: 601-607.

Hau M.; Kneitz C.; Tony H.P. et al.(2002): High resolution ultrasound detects a decrease in pannus vascularisation of small finger joints in patients with rheumatoid arthritis receiving treatment with

soluble tumour necrosis factor alpha receptor (etanercept). *Ann Rheum Dis.*,61:55-8.

Hau M.; Schultz H.; Tony H.P. et al.(1999): Evaluation of pannus and vascularization of the metacarpophalangeal and proximal interphalangeal joints in rheumatoid arthritis by high-resolution ultrasound (multidimensional linear array). *Arthritis Rheum.*,42:2303–8.

Healy P.J.; Groves C. Chandramohan M. et al.(2008): changes in psoriatic dactylitis—extent of pathology, relationship to tenderness and correlation with clinical indices. *Rheumatology* .,47:92–5.

Hench P.S.; Kendall E.C.; Slocumb C.H. et al., (1994): The effect of a hormone of the adrenal cortex (17-hydroxy-11-dehydrocorticosterone; compound E) and of the pituitary adrenocorticotrophic hormone on rheumatoid arthritis. *Proc Staff Meet Mayo Clin.*,24:181.

Henneric M. and Heusler D. (1998): Veins. In Hennerici M and Heusler DN (eds). *Vascular diagnosis with ultrasound*, Stuttgart. Philadelphia, Thieme, 2nd ed., PP: 189-230.

.

Higashida J.; Wun T.; Schmidt S. et al.(2005): Safety and efficacy of rituximab in patients with rheumatoid arthritis refractory to disease modifying antirheumatic drugs and anti-tumor necrosis factor alpha treatment. *The Journal of Rheumatology.*,32:2109– 15.

Hitzel A., Liard A.; Véra P.et al.(2002): Color and Power Doppler Sonography Versus DMSA Scintigraphy in Acute

Pyelonephritis and in Prediction of Renal Scarring. Journal of Nuclear Medicine .,1: 27-32.

Hobbs K. (2002): Laboratory evaluation. In sterling G.W (ed.): Rheumatology Secrets, Hanley and Belfus.. Philadelphia, 2nd ed., PP. 52 – 63.

Houk J. (1998): Joint fluid. In Maddison P.J., Isenberg D.A., Woo P and Glass. D.N. (eds.): Oxford Textbook of Rheumatology, New York Oxford University Press Inc., 2nd ed., PP: 677 – 684.

J

Jacelyn K. and Ian Morris (2002): Cricoarytenoid arthritis: a cause of acute upper airway obstruction in rheumatoid arthritis .Can J anesth., 49: 729–732.

Jacobson J.A. (1999): Musculoskeletal Sonography and MR Imaging. A Role for Both imaging. Radiologic clinics of North America.,37 (4): 713 – 735.

Jessop J.D.; O'sullivan M.M.; Lewis P.A. et al., (1998): A long term five-year randomized trial of hydroxychloroquine, sodium aurothiomalate, auranofin and penicillamine in the treatment of patients with rheumatoid arthritis. Br. J. Rheumatol., 37: 992.

Joel S.; Ronald S.; Ronald A. et al.(1994): Detection of Soft-Tissue Hyperemia: Value of Power Doppler Sonography. AJR.,163:385-389.

Joosten L.A.; Helsen M.M.; Van de Loo, F.A.; et al., (1996): Anticytokine treatment of established type II collagen –

induced arthritis in DBA1 mice : A comparative study using anti-TNF, anti-IL-1B, and IL-1Ra. *Arthritis Rheum.*, 39 : 797 – 809.

K

Kamaya A.; Tuthill T. and Jonathan M. (2003): Twinkling Artifact on Color Doppler Sonography: Dependence on Machine Parameters and Underlying Cause. *AJR.*, 180:215-222.

Kamishima T.; Kazuhide T.; Henmi M.et al. (2009): Power Doppler ultrasound of rheumatoid synovitis: quantification of vascular signal and analysis of interobserver variability. *Skeletal Radiol.*, 38:467–472.

Kane D.; Balint P.V.; Sturrock R. et al. (2004): Musculoskeletal ultrasound – a state of the art review in rheumatology. Part 1 : Current controversies and issues in the development of musculoskeletal ultrasound in rheumatology. *Rheumatology.*, 43 : 823 – 828.

Kane D.; Greaney T.; Shanahan M. et al., (2001): The role of ultrasonography in the diagnosis and management of idiopathic plantar fascitis. *Rheumatology.*, 40 : 1002 – 8.

Kaplan I. and Bonanova P. (2006): Examination of the rheumatoid hand and wrist. *International Congress Series.*, 12: 9 – 26.

Kasukawa R.; Kanno Y.; Homma F.et al. (2005): Comparison in values of color flow signals and vascular resistance of synovial vascularity demonstrated by Doppler sonography

- between knee and metacarpophalangeal joints of patients with rheumatoid arthritis. *Mod Rheumatol* .,15:336–339.
- Kaushik P.; Cooper E.S.; Banda V.R. et al. (2005):** Bronchiolitis obliterans with organizing pneumonia in rheumatoid arthritis a fatal case and short review of literature. *Rheumatol Int.*,25:391-3.
- Kelly W.N.(2005):** Management of rheumatoid arthritis. In: Kelly, W.N.; Harris, E.D.; Ruddy ,S.; and Sledge, C.B. (eds): *Textbook of rheumatology*. WB Saunders. Philadelphia, 3rd ed., 336–339.
- Khurana R. and Berney S. M. (2005) :**Clinical aspects of rheumatoid arthritis. Elsevier Ireland Ltd.,12:153-165.
- Kim D. and Hilibar A. (2005):** Rheumatoid arthritis in the cervical spine. *J Am Acad Orthop surg.*,13:463-74.
- Kirwan J.R. (1995):** The effect of glucocorticoids on joint destruction in rheumatoid arthritis. *N. England J. Med.*, 333 : 142.
- Klauser S.; Moriggl B.; Duftner C.et al. (2009):** Sonography of synovial and erosive inflammatory changes. *Der Radiologe* .,64:365-375.
- Klippel J.H.; Weyand C.M. and Wortmann R.L. (1997):** Rheumatoid arthritis. In Georgia(ed). *Primer on the Rheumatic Disease*, Arthritis Foundation. Atlanta, 11th ed. P. 155-175.
- Kobayashi S.,Okamoto H.; Iwamoto T. et al. (2008):** A role for the aryl hydrocarbon receptor and the dioxin TCDD in rheumatoid arthritis. *Rheumatology*.,47:1317-22.
- Korn T.; Bettelli E.; Oukka M. et al. (2009):** IL-17 and Th17 Cells. *Annual review of immunology*.,27:485–518.

Koski J.M.; Saarakkala S.; Helle M. et al. (2006): Assessing the intra- and inter-reader reliability of dynamic ultrasound images in power Doppler ultrasonography. *Ann Rheum Dis.*, 65: 1658–1660.

Kowalski M. L.; Wolska A.; Grzegorzczak J. et al.(2008): Increased Responsiveness to Toll-Like Receptor 4 Stimulation in Peripheral Blood Mononuclear Cells from Patients with Recent Onset Rheumatoid Arthritis Mediators Inflamm.,13;27-32.

Krix M.;Weber M.A.; Roosen K .H .et al, (2005) :Assessment of skeletal muscle perfusion using contrast- enhanced ultrasonography. *J Ultrasound Med .*,24:431-441.

L

Lacki J.K.; Samborski W.; Mackiewicz S. et al. (1995): Do changes in clinical improvement in rheumatoid arthritis patients treated with immunosuppressive drugs reflect the changes in acute phase proteins? *Upsala Journal of Medical Sciences.*,100: 233–242.

Larché J.; Matthew S. ;Adrian L. et al.,(2010): Quantitative Power Doppler Ultrasonography Is a Sensitive Measure of Metacarpophalangeal Joint Synovial Vascularity in Rheumatoid Arthritis and Declines Significantly Following a 2-week Course of Oral Low-dose Corticosteroids .*The Journal of Rheumatology.*,37:12-20.

- Larsen A. (1995):** How to apply Larsen score in evaluating radiographs of rheumatoid arthritis in long-term studies. *J Rheumatol* .,22:1974–1975.
- Larsen A.; Dali K. and Eek M. (1977):** Radiological evaluation of rheumatoid arthritis and related conditions by standard reference films. *Acta. Radiologica. Diagnosis.*, 18 : 481 – 491.
- Lee J.L.; Kim S.H.; Cho J.Y.et al. (2001):** Color and power Doppler twinkling artifacts from urinary stones: clinical observations and phantom studies. *AJR* .,176:1441 -1445.
- Leirisalo M.; Kautiainen H.; Laasonen L. et al. (2008):** A randomized, double-blind, placebo-controlled study on addition of infliximab to the FIN-RACo DMARD combination therapy for initial six months in patients with early active rheumatoid arthritis. *J Ultrasound Med* .,45:481-491.
- Lim G.Y.; Im S.A.; Jung W.S. et al. (2005):** Evaluation of joint effusion in rabbits by color Doppler, power Doppler, and contrast-enhanced power Doppler ultrasonography. *J Clin Ultrasound*.,33:333 -338.
- Lindqvist E.; Eberhardt K.; Bendtzen K. et al. (2005):** Prognostic laboratory markers of joint damage in rheumatoid arthritis. *Ann Rheum Dis.*, 64:196–201.
- Liuzzo G. (2000):** Monoclonal T-cell proliferation and plaque instability in acute coronary syndromes, *Circulation* 101., (25) 2883–2888.

Lohkampa M.G.; Burrowb T.; McCarronb W. et al. (2006): The prevalence of disabling foot pain in patients with early rheumatoid arthritis. *AJR.*, 16: 201-207.

Luqmani R.A.; Palmer R.G. and Bacon P.A. (1990): Azathioprine, cyclophosphamide and chlorambucil. *Baillieres Clin. Rheumatol.*, 4 : 595 – 619.

M

Mackley C.L.; Ostrov B.E. and Ioffreda M. (2004): Accelerated cutaneous nodulosis during infliximab therapy in a patient with rheumatoid arthritis. *J Clin Rheumatol.*, 10:336-338.

Maetzel A.; Wong A.; Strand P. et al. (2000): Meta-analysis of treatment termination rates among rheumatoid arthritis patients receiving disease-modifying anti-rheumatic drugs. *Rheumatology.*,39:975–81.

Matsuno H.; Yudoh K.; Hanyu T. et al. (2003): Quantitative assessment of hand radiographs of rheumatoid arthritis: interobserver variation in a multicenter radiographic study. *J Orthop Sci.*, 8:467–473.

Matteson E. L.(2003): Cervical spine disease in rheumatoid arthritis: how common a finding? How uncommen a problem ? *Arthritis Rheum.*,48 :1775-8.

Matteson E.L. (2006): Treatment of rheumatoid arthritis. In Klippel J.H. (ed): *Primer on the Rheumatic Disease, Arthritis Foundation.* Georgia, Atlanta, 12th Ed., PP. 225-232.

McDicken W.N. and Hoskins P. (2000): Physics: Principles, Practice and artifacts. In Allan P.L., Dubbins P.A. and Pozniak M.A.

- (eds): Clinical Doppler Ultrasound. Arthritis Foundation. Churchill Livingstone, New York, Edinburgh, London, Philadelphia, ST Louis, Sydney, Toronto, 10th ed., PP: 1-23.
- McInnes I.B. and Schett G.(2007):** Cytokines in the pathogenesis of rheumatoid arthritis. *Nat Rev Immunol.* ,7:429–442.
- Meron M.K.; Amital H.; Shepshelovich D. et al. (2009):** Infectious aspects and the etiopathogenesis of rheumatoid arthritis. *The Journal of Rheumatology.*, 15:12; 18.
- Mevorach D. and Paget S.A. (2000):** Rheumatoid arthritis. In paget S.A., Gibofsky A. and Beary III J.F. (eds): *Manual rheumatology and outpatient orthopedic disorders*, Lippincott Williams and Wilkins. Philadelphia, Baltimore, New York 4th ed., PP: 192 – 229.
- Michaud K. and Wolfe F. (2007):** Comorbidities in rheumatoid arthritis *Best Practice & Research Clinical Rheumatology.*,5: 885–906,.
- Middleton W.D.; Dasyam A. and Teefey, S .(2005):** Diagnosis and Treatment of iatrogenic Femoral Artery Pseudoaneurysms. *Ultrasound Quarterly.*,21(1): 3-17.
- Molenaar E.T.; Voskuyl A.E.; Dinant H.J. et al. (2004):** Progression of radiologic damage in patients with rheumatoid arthritis in clinical remission. *Arthritis Rheum.* ,50:36–42.
- Möller B.; Aeberli D.; Egli S. et al. (2009):** Class-switched B cells display response to therapeutic B-cell depletion in rheumatoid arthritis. *Arthritis Res Ther.*, 11:62-68.
- Moncur C. and Williams H.J. (1995):** RA : Status of drug therapies. *Phys. Ther.*, 75 (6) : 511 – 25.

Murat B.; Kabayel D.; Uzunca K. et al. (2008): The relation of hand functions with radiological damage and disease activity in rheumatoid arthritis .Rheumatol Int .,28:407–412 .

N

Nagashima M, Asano G, Yoshino S (2000): Imbalance in production between vascular endothelial growth factor and endostatin in patients with rheumatoid arthritis. . J Rheumatol. Oct.,27:2339-42.

Nakou M.; Katsikas G.; Sidiropoulos P. et al. (2009): Rituximab therapy reduces activated B cells in both the peripheral blood and bone marrow of patients with rheumatoid arthritis: depletion of memory B cells correlates with clinical response. Arthritis Res Ther., 11:R131.

Naredo E.; G Bonilla F.; Gamero J. et al. (2005): Assessment of inflammatory activity in rheumatoid arthritis: a comparative study of clinical evaluation with grey scale and power Doppler ultrasonography. Ann Rheum Dis., 64:375–381.

Naredo E.; Møller I.; Cruz A. Et al. (2008): Power Doppler Ultrasonographic Monitoring of Response to Anti-Tumor Necrosis Factor Therapy in Patients With Rheumatoid Arthritis .Arthritis& Rheumatism., 8: 2248–2256.

Naredo E.; Moller I.; Moragues C. et al., (2006): Interobserver reliability in musculoskeletal ultrasonography: results from a “Teach the Teachers” rheumatologist course. Ann Rheum Dis.,65: 14–19.

Nell V.P.; Machold K.P. Stamm T.A. et al. (2005): Autoantibody profiling as early diagnostic and prognostic tool for rheumatoid arthritis. *Ann Rheum Dis.*,64(12):1731–6.

Newman J.S.; Bude R.O. et al.(1995): Detection of soft tissue hyperemia: value of Power Doppler sonography. *AJR.*, 163:385-389.

Newman J.S.; Laing T.J.; McCarthy C.J. et al.(1996): Power Doppler sonography of synovitis: assessment of therapeutic response— preliminary observations. *Radiology.*,198:582–4.

Nicholas J.J. (1994): Physical modalities in rheumatological rehabilitation. *Arch. Phys. Rehabil.*, 75 : 994 – 1001.

O

O'Dell J.R. (2005): Methotrexate, leflunomide and combination therapies. In Harris E.D., Budd R.C., Firestein G.S., Genovese M.C., Sargent J.S., Ruddy S. and Sledge C.B. (eds): *Kelley's Textbook of Rheumatology*, Elsevier Saunders.. Philadelphia, Pennsylvania, 7th ed., PP: 900 – 919.

Oae P.J. and Wakefield R.J. (2003): Musculoskeletal US in rheumatology. *Oxford University Press.*,32:25-33.

O'Connor D.; Kortman B.; Smith A. et al. (1999); Correlation between objective and subjective measures of hand function in patients with rheumatoid arthritis. *J Hand Ther.*,12:323-9.

Oliver J.; Sommer M.D.; Andrea M. et al.(2005):Rheumatoid arthritis: A Practical Guide to State-of-the-Art Imaging, Image Interpretation, and clinical importance. *Radiographics.*,25:381-398.

- Ollivier Y.; Saraux A. and Le Goff P. (2004):** Prevalences of rheumatoid arthritis in Roman Catholic nuns and the general female population in Brittany, France: a pilot study. *Clin Exp Rheumatol.*,22:759-62.
- O'Neill G. (1999):** An investigation into the assessment and concept of hand function . *Arthritis Res* 7:11-15.
- Ory P.A. (2003):** Radiograohy in the assessment of musculoskeletal condition. *Best Pract Res Clin Rheumatol.*,1:495-512.
- Ostergaard M.; Ejbjerg B. and Szkudlarek M. (2005):** Imaging in early rheumatoid arthritis: roles of magnetic resonance imaging, Ultrasonography, conventional radiography and computed tomography. *Best Pract Res Clin Rheumatol.*,19:91-116.
- Oyama T.; Kohno N.; Yokoyama A.et al. (1997):** Detection of interstitial pneumonitis in patients with rheumatoid arthritis by measuring circulating levels of KL-6, ahuman MUCI mucin. *LUNG.*,175:379-85.
- Ozgocmen S.; Kiris A.; Kocakoc E. et al. (2004):** Evaluation of metacarpophalangeal joint synovitis in rheumatoid arthritis by power Doppler technique: relationship between synovial vascularization and periarticular bone mineral density *Joint Bone Spine.*, 71:384–388.
- Ozgocmen S.; Ozdemir H.; Kiris A.et al., (2008):** Clinical Evaluation and Power Doppler Sonography in Rheumatoid Arthritis: Evidence for Ongoing Synovial Inflammation in Clinical Remission. *The Southern Medical Association.*,101(3):240-245.

Ozturk A.; Yalcin S.; Kaygusuz I. et al. (2004): High frequency hearing loss and middle ear involvement in rheumatoid arthritis. *Am J Otolaryngol.*, 25: 411-7.

P

Paleolog M. (2002): Angiogenesis in rheumatoid arthritis. *Arthritis Res.*, 4: 81-90.

Palmgren N.; Jensen C.G.; Kaae K. et al., (1998): Low-powered laser therapy in rheumatoid arthritis. *Lasers Med. Sci.*, 4 : 193 – 6.

Patel M. and Moreland W. (2010): Interleukin-6 inhibition for treatment of rheumatoid arthritis: A review of tocilizumab therapy *Drug Design. Development and Therapy.*,4 :263–278.

Pedersen L.M. (1995): Microalbuminuria in patients with rheumatoid arthritis, *Ann. Rheum. Dis.*, 54 189–192.

Peeters H. (1996): Course and characteristics of anemia in patients with rheumatoid arthritis. *Brit. J. Rheum.*, 35 : 879 – 883.

Perry D.; Stewart N.; Benton N. et al. (2005): Detection of erosions in the rheumatoid hand; a comparative study of multidetector computerized tomography versus magnetic resonance scanning. *J Rheumatol* .,32:256–267.

Piet L.C.M. and van Riel (2007): Established rheumatoid arthritis: clinical assessments. *Best Practice & Research Clinical Rheumatology* .,21: 807–825.

- Pinals R.S.; Baum J.; Bland J. et al. (1982):** Preliminary criteria for clinical remission in rheumatoid arthritis. *Bull Rheum Dis.*,32(1):7-10.
- Pincus T. (1998):** Laboratory tests in rheumatic disease. *Pro. Nat. Acad. Sci. U.S.A.*, 95: 125-138.
- Pincus T. and Callahan I.F. (1992):** Early mortality in RA predicted by poor clinical status. *Bull Rheum Dis.*, 41:4.
- Pincus T.; Sokka T. and Stein C.M. (2002):** Are long-term very low doses of prednisone for patients with rheumatoid arthritis as helpful as high doses are harmful? *Ann Intern Med.*,136(1):76–8.
- Pincus T.; Sokka T. and Kautiainen H. (2005):** Patients seen for standard rheumatoid arthritis care have significantly better articular, radiographic, laboratory, and functional status in 2000 than in 1985. *Arthritis Rheum.*,52:1009–19.
- Plant M.J.; Williams A.L.; O’Sullivan M.M. et al. (2000):** Relationship between time-integrated C-reactive protein levels and radiologic progression in patients with rheumatoid arthritis. *Arthritis and Rheumatism.*,43: 1473–1477.
- Poole C.D.; Conway P. and Currie C.J. (2007):** An evaluation of the association between systemic inflammation –as measured by C-reactive protein- and hospital resource use. *Curr Med Res Opin.*,23:2785-2792.
- Posthumus; Limburg P.C.; Westra J. et al. (2000):** Serum matrix metalloproteinase 3 in early rheumatoid arthritis is correlated with disease activity and radiological progression. *The Journal of Rheumatology.*,27: 2761–2768.

Pratt G.; Isaacs D. and Matthey L. (2009): Current concepts in the pathogenesis of early rheumatoid arthritis Best Practice & Research Clinical Rheumatology., 23: 37–48.

Prevoo M.L.; Hof M.A.; Kuper H.H. et al. (1995): Modified disease activity scores that include twenty-eight-joint counts. Development and validation in a prospective longitudinal study of patients with rheumatoid arthritis. Arthritis Rheum., 38:44–48.

Prevoo M.L.; Kuper I.H.; van 't Hof M.A. et al. (1996): Validity and reproducibility of self-administered joint counts. A prospective longitudinal followup study in patients with rheumatoid arthritis. The Journal of Rheumatology., 23: 841–845.

Prevoo M.L.L.; van Riel P.L. Hof M.A. et al. (1993): Validity and reliability of joint indices. A longitudinal study in patients with recent onset rheumatoid arthritis. British Journal of Rheumatology., 32: 589–594.

Q

Quinn M.A. and Emery P. (2005): Are early arthritis clinics necessary ? Best Pract Res Clin Rheumatol.,19:1-17 .

Qvistgaard E.; Kristoffersen H.; Terslev L. et al. (2001): Guidance by ultrasound of intra-articular injections in the knee and hip joints. Osteoarthritis Cartilage ., 9:512-517.

R

- Rall L.C.; Meydani S.N.; Kehayias J.J. et al., (1996):** The effect of progressive resistance training in rheumatoid arthritis. *Arthritis Rheum.*, 39 : 415.
- Ranganathan P.; Perry A, and Gelbart M. (2007):** Localized striated muscle vasculitis in rheumatoid arthritis. *Clin Rheumatol.*,13:35-7.
- Rees J.D.; Pilcher J.; Heron, C.et al. (2007):** A comparison of clinical vs ultrasound determined synovitis in rheumatoid arthritis utilizing gray-scale, power Doppler and the intravenous microbubble contrast agent ‘Sono-Vue’. *Rheumatology.*,46:454–9.
- Ribbens C.; Andre B.; Jaspar J.M. et al. (2000):** Matrix metalloproteinase-3 serum levels are correlated with disease activity and predict clinical response in rheumatoid arthritis. *The Journal of Rheumatology.*, 27: 888–893.
- Richards P. J.; Win T. and Jones P.W. (2005):** The distribution of microvascular response in Achilles tendonopathy . *The Journal of Rheumatology* .,34:336-342.
- Richardson C.; Griffiths B. and Emery B. (1998):** Hematological abnormality in Rheumatoid Arthritis. In Isenberg D.A., Woo P and Glass D.N. (eds.): *Oxford Text book of Rheumatology*, Oxford University press, Inc. New York, 2nd ed., PP: 633 – 646.
- Riedlinger W.F.; Lairmore T.C.; Balfe D.M. et al (2005):** Tumefactive necrobiotic granulomas (nodulosis) of the pancreas in an adult with long-standing rheumatoid arthritis. *Int J Surg Pathol.*,13:207-210.

Riemsma R.P.; Taal E. and Rasker J.J.(2002): patient education programmes for adults with rheumatoid arthritis. *BMJ.*, 14:558-9.

Riente L.; Sedie D. A.; Iagnocco A. et al.(2006): Ultrasound imaging for the rheumatologist V Ultrasonography of the ankle and foot. *Clin Exp Rheumatol.*,24:493–8.

Ritchi, D.M.; Boyle J.A.; Mcinnes J.M. et al., (1968): Clinical studies with an articular index for the assessment of joint tenderness in patients with rheumatoid arthritis. *Quarterly J. Med.*, 37 : 393 – 406.

Rogind H.; Qvistgaard E.; Pedersen S. et al. (2001):Quantitative ultrasonography in rheumatoid arthritis: evaluation of inflammation by Doppler technique. *Ann Rheum Dis.*,60:690–693.

S

Saag K.G.; Teng G.G.; Patkar N.M. et al. (2008): American College of Rheumatology 2008 recommendations for the use of nonbiologic and biologic disease-modifying antirheumatic drugs in rheumatoid arthritis. *Arthritis Rheum.*,59:762–784.

Salisbury M.D. and Nottage W.M. (1985): A new evaluation of gross pathologic changes and concepts of rheumatoid articular cartilage degeneration. *Clin. Orthop. Rel. Res.*, 199: 243-249.

- Salliot C.; Bombardier C.; Saraux A. et al. (2009):** Hormonal replacement therapy may reduce the risk for RA in women who carry HLA-DRB1 01 and/or 04 alleles by protecting against the production of anti-CCP: Results from the ESPOIR cohort. *Ann Rheum Dis.*, 9 ;10:1-5.
- Sarah L. and Gaffen. (2009):** Role of IL-17 in the Pathogenesis of Rheumatoid Arthritis *Curr Rheumatol Rep.* October.,11(5): 365–370.
- Scheel A.K.; Schmidt W.A.; Hermann K.G. et al. (2005):** Interobserver reliability of rheumatologists performing musculoskeletal ultrasonography : results from a EULAR “Train the trainers” course. *Ann Rheum Dis.*,64: 1043-9.
- Schmidt W.A.; Volker L.; Zacher J. et al. (2001):** Colour Doppler ultrasonography to detect pannus in knee joint synovitis, *Clin Exp Rheumatol.*,18:439-444.
- Schumacher H.R. (1993):** Synovial fluid analysis and synovial biopsy. In Kelley WN, Harris ED, Ruddy S, Sledge CB.(eds): textbook of Rheumatology, WB Saunders Company. Philadelphia,PP: 562-578.
- Schwarz E.M.; Friedberg, R.; Stosslein F. et al. (2005):** Rheumatoid arthritis at the cervical spine an underestimated problem. *Dtsch Med Wochenschr.*, 130:1866-70.
- Scott D.L. and Houssien D.A. (1996):** Joint assessment in rheumatoid arthritis. *British Journal of Rheumatology.*, 35: 214–218.
- Sedie D.A.; Riente L. and Bombardieri S. (2008):** Limits and perspectives of ultrasound in the diagnosis and management of rheumatic diseases. *Mod Rheumatol.*,18:125–31.

- Segal N. (2001):** Two configurations of static magnetic fields for treating rheumatoid arthritis of the knee : A double – blind clinical trial. *Arch. Phys. Med. Rehabil.*, 82 : 1453 – 60.
- Shahrara S.; Pickens S.R.; Dorfleutner A. et al. (2009):** IL-17 induces monocyte migration in rheumatoid arthritis. *J Immunol.*,182:3884–3891.
- Sharp J.T.; Strand V.; Lenug H. et al.(2000):** Treatment with leflunomide shows radiographic progression of rheumatoid arthritis. *Arth. Rheum.*, 43: 495-505.
- Shaw k.; Zochling J. and Winzenberg,T.(2007):** Nonpharmacological interventions for rheumatoid arthritis. *Aust fam Physician.*,36:840-1.
- Shmerling R.H. and Liang M.H. (1997):** Rheumatoid arthritis: Evaluation of the patient, laboratory assessment. In Klipfel J.H. (ed): primer on the Rheumatic disease, arthritis foundation. Atlanta, Georgia. 11th ed., PP. 89-97.
- Silva C.; Anamaria P.T.; Paula O.T. et al. (2008):** Effectivness of a night-time hand positioning splint in Rheumatoid Arthritis: a randomized cotrolled trial. *J Rehabil Med.*,40: 749–754.
- Silvestri E.; Biggi E.;Molfetta L. et al (2003):** Power Doppler Analysis of tendon vascularization. *Int J Tissue React* .,25:149-158.
- Simmen B.R.; C. Kolling D.B. and Herren. (2007):** The managment of the rheumatoid wrist. *Current Orthopaedics.*,21: 344–357.
- Singh ;Vishwanath A.; Pramod K. et al. (2009):** Is rheumatoid factor still a superior test for the diagnosis of rheumatoid arthritis. *Rheumatol Int.*,15:1-5.

- Sivakumar B.; Harry L.E. and Paleolog E.M. (2004):** Modulating angiogenesis: more versus less. *JAMA.*, 292:972-7.
- Skoldstam L.; Hagfors L. and Johansson G. (2003):** An experimental study of a Mediterranean diet intervention for patients with rheumatoid arthritis. *Ann Rheum Dis.*,62:208–14.
- Smolen J.S.; Beaulieu A.; Rubbert A. et al. (2008):** Effect of interleukin-6 receptor inhibition with tocilizumab in patients with rheumatoid arthritis. a double-blind, placebo-controlled, randomised trial. *Lancet.*,371:987–997.
- Smolen J.S.; Breedveld F.C.; Eberl G. et al. (1995):** Validity and reliability of the twenty-eight-joint count for the assessment of rheumatoid arthritis activity. *Arthritis and Rheumatism.*, 38: 38–43.
- Soubrier M.; Dougados M. (2005):** Selecting criteria for monitoring patients with rheumatoid arthritis. *Joint Bone Spine* .,72:129–134.
- Spindler A.; Bellomio V.; Berman A. et al. (2002):** Prevalence of rheumatoid arthritis in Tucuman, Argentina. *J Rheumatol.*,29:1166-70.
- Stand V. and Sharp J.T. (2003):** Radiographic data from recent randomized controlled trials in rheumatoid arthritis :What have we learned. *Arthritis Rheum.*,48:21-34.
- Stein C.M. (2005):** Immunoregulatory drugs. In Harris E.D, Budd R.C., Genovese M.C, Firestein G.S. and Sargent J.S. (eds): *Kelley’s Textbook of Rheumatology*, Elsevier Saunders. Philadelphia, Pennsylvania, 7th ed., PP: 920 – 939.
- Steuer A.; Watkins J.; Smith F. et al.(2008):** RF latex and anti-CCP antibodies: a combined strategy for diagnosing RA in primary care? *Rheumatology* .,47:375–6.

- Stone M.; Bergin D.; Whelan B. et al. (2001):** Power Doppler ultrasound assessment of rheumatoid hand synovitis. *J Rheumatol.*,28:1979–82.
- Strunk J.; Heinemann E.; Neeck G. et al.(2004):** A new approach to studying angiogenesis in rheumatoid arthritis by means of power Doppler ultrasonography and measurement of serum vascular endothelial growth factor. *Rheumatology*.,43:1480-3.
- Stucki G. and Langenegger T. (1997):** Management of RA. *Curr. Opin. Rheumatol.*, 9 (3) : 229 – 35.
- Suzuki T.; Tohda E. and Ishihara K. (2009):** Power Doppler ultrasonography of symptomatic rheumatoid arthritis ankles revealed a positive association between tenosynovitis and rheumatoid factor. *Mod Rheumatol.*, 19:235–244.
- Szkudlarek M.; Payen M.; Strandberg C. et al. (2001):** Power Doppler Ultrasonography for Assessment of Synovitis in the Metacarpophalangeal Joints of Patients With Rheumatoid Arthritis. *Arthritis & Rheumatism.*, 9: 2018–2023.

T

- Tak P. and Ferdinand C. (1999):** Current perspectives on synovitis .*Arthritis Res.*, 1:11-15.
- Tak P.P. and Bresnihan B. (2000):** The pathogenesis and prevention of joint af damage in rheumatoid arthritis. *Arthritis Rheum.*,43:2619-2633.
- Taylor P.C.; Steuer A.; Gruber J.et al. (2006):** Ultrasonographic and radiographic results from a two-year controlled trial of

- immediate or one-year delayed addition of infliximab to ongoing methotrexate therapy in patients with erosive early rheumatoid arthritis. *Arthritis Rheum.*,54(1):47-53.
- Tedesco D.; D'Agostino I.; Soriente P.et al. (2009):** ; A new strategy for the early diagnosis of rheumatoid arthritis: A combined approach. *Autoimmunity Reviews* 8: 233–237.
- Teh J.; Stevens K.; Williamson L.et al. (2003):** Power Doppler ultrasound of rheumatoid synovitis: quantification of therapeutic response. *Br J Radiol.*, 76: 875–879.
- Tehranzadeh J.; Ashikyan O.; Anavim A. et al.(2006):** Enhanced MR imaging of tenosynovitis of hand and wrist in inflammatory arthritis. *Skeletal Radiol.*,35:814–22.
- Tehranzadeh J.;Ashikyan O.; Anavim A. et al.(2008):** Detailed analysis of contrast enhanced MRI of hands and wrists in patients with psoriatic arthritis. *Skeletal Radiol.*,37:433–42.
- Terslev L. (2004):** Doppler ultrasound in the detection and quantification of inflammation in rheumatoid arthritis. *Dan Med Bull.*,51:214.
- Terslev L. ;Torp S. ;Qvistgaard E. et al. (2003):**Spectral Doppler and resistive index. A promising tool in ultrasonographic evaluation of inflammation in rheumatoid arthritis. *Acta Radiol.*,44:645– 652.
- Terslev L.; Torp S.; Qvistgaard E. et al. (2004):** Doppler ultrasound findings in healthy wrists and finger joints. *Ann Rheum Dis.*,63:644–8.
- Terslev L.; Torp S.; Bang N. et al. (2005):** Doppler ultrasound findings in healthy wrists and finger joints before and after use of two different contrast agents. *Ann Rheum.*,64:824-827 .

- Thomas E.; Brewster D.H.; Black R.J. et al. (2000):** Risk of malignancy among patients with rheumatic conditions. *International Journal of Cancer*; 88: 497–502.
- Thomas R. and Carrol G.J. (1993):** Reduction of leukocyte and interleukin concentrations in the synovial fluid of rheumatoid arthritis patients treated with methotrexate. *Arthritis Rheumat.*, 36 : 1244 – 1252.
- Thompson P.W. and Kirwan JR. (1995):** Joint counts: a review of old and new articular indices of joint inflammation. *British Journal of Rheumatology.*, 34: 1003–1008.
- Tilley B.; Alarcon G.; Heyse S. et al. (1995):** Minocycline in rheumatoid arthritis : A 48 – week, double blind, placebo – controlled trial. *Ann. Intern. Med.*, 122 : 81.
- Toh M.L.; Miossec P. (2007):** The role of T cells in rheumatoid arthritis: new subsets and new targets. *Curr Opin Rheumatol.*,19:284–288.
- Tugwell P.; Boers M. and Omeract C. (1993):** Developing consensus on preliminary core efficacy endpoints for rheumatoid arthritis clinical trials. *The Journal of Rheumatology.*, 20: 555–561.
- Turesson C. (2002):** Occurrence of extraarticular disease manifestations is associated with excess mortality in a community based cohort of patients with rheumatoid arthritis, *J. Rheumatol.*, 29 (1) 62-67.
- Tutuncu Z. and Kavanaugh A. (2005):** Anticytokine therapies. In Harris E.D., Budd R.C., Firestein G.S., Genovese M.C., Sargent J.S., Ruddy S. and Sledge C.B. (eds) : *Kelley’s Textbook of Rheumatology*, Eklsevier Saunders. Philadelphia, Pennsylvania, 7th ed., PP : 940.

Tuulikki S. and Mañkinen (2009): Drug management of early rheumatoid arthritis. *Best Practice & Research Clinical Rheumatology.*, 23: 93–102.

V

Vallbracht T.and Klaus Helmke (2005): Additional diagnostic and clinical value of anti-cyclic citrullinated peptide antibodies compared with rheumatoid factor isotypes in rheumatoid arthritis. *Autoimmunity Reviews.*, 4: 389– 394.

Vallon R.; Freuler F.; Tsedu N. et al. (2001): Serum amyloid A (apoSAA) expression is up-regulated in rheumatoid arthritis and induces transcription of matrix metalloproteinases. *Journal of Immunology.*, 166: 2801–2807.

Van Riel and Fransen J. (2005): To be in remission or not: is that the question?. *Annals of the Rheumatic Diseases* 64: 1389–1390.

Van Venrooij W.J.; Van Beers J. and Pruijn G.J.M. (2008): Anti-CCP antibody, a marker for the early detection of rheumatoid arthritis. *Ann N Y Acad Sci.*, 1143:268–285.

Van Venrooij W.J.; Zendman A.J. and Pruijn G.J. (2006): Autoantibodies to citrullinated antigens in (early) rheumatoid arthritis. *Autoimmun Rev.*, 6:37–41.

Varol A.; Basa u.k.; Topsakal A. et al. (2008): Assessment of synovial vascularization by power Doppler ultrasonography in TMJ internal derangements treated arthroscopically *British Journal of Oral and Maxillofacial Surgery.*, 46 :625–630.

- Verhoeven A.C.; Boers M. and van der Linden S. (2000):**
Responsiveness of the core set, response criteria, and utilities
in early rheumatoid arthritis. *Annals of the Rheumatic
Diseases.*, 59: 966–974.
- Vicente G (2007):** Rheumatoid Nodule. Elsevier Inc., 2:7-15.
- Visser H.; le Cessie S.; Vos K. et al. (2002):** How to diagnose
rheumatoid arthritis early. A prediction model for persistent
(erosive) Arthritis. *Arthritis Rheum.*,46:357– 65.
- Vlieland T.P. (2003):** Rehabilitation of people with rheumatoid
arthritis. *Best Pract Res Clin Rheumatol.*,17:847-61.
- Vlychou M.; Koutroumpas A.; Malizos K. et al. (2009):**
Ultrasonographic evidence of inflammation is frequent in
hands of patients with erosive osteoarthritis. *Osteoarthritis
Cartilage.*,17:1283-7.
- Voskuyl A.E. (1996):** Factors associated with the development of
vasculitis in rheumatoid arthritis: results of a case-control
study. *Ann. Rheum. Dis.* 55 :190–192.

W

- Wakefield R.J.; Balint P.V.; Szkudlarek M. et al. (2003):**
Musculoskeletal ultrasound including definitions for
ultrasonographic pathology. *J Rheumatol.*,32:2485-7 .
- Wakefield R.J.; D’Agostino M.A.; Iagnocco A. et al. (2007):** The
OMERACT Ultrasound Group: status of current activities and
research directions. *J Rheumatol.*,34(4):848–51.
- Wakefield R.J.; Gibbon W.W. ; Conaghan P.G. et al (2000):** The
value of sonography in the detection of bone erosions in

- patients with rheumatoid arthritis: a comparison with conventional radiography. *Arthritis Rheum.*, 43:2762-2770.
- Wakefield R.J.; Brown A.K.; Connor p. j. et al. (2005):** Power Doppler sonography: improving disease activity assessment in inflammatory musculoskeletal disease. *Arthritis Rheum.*, 48:285-288.
- Wallny T.A.; Schild R.L.; Bertelsbeck D. et al. (2001):** Three dimensional ultrasonography in the diagnosis of rotator cuff lesions. *Ultrasound Med. Biol.*, 27 : 745 – 9.
- Walther J.; van Venrooij W.J.; Albert J.W. et al. (2006):** Autoantibodies to citrullinated antigens in [early] rheumatoid arthritis. *Autoimmun Rev.*, 6:37–41.
- Walther M.; Harms H.; Krenn V. et al. (2001):** Correlation of Power Doppler sonography with vascularity of the synovial tissue of the knee joint in patients with osteoarthritis and rheumatoid arthritis. *Arthritis Rheum.*, 44:331-8.
- Weidekamm C.; Ko" ller M.; Weber M et al.(2003):** Diagnostic value of high-resolution B-mode and Doppler sonography for imaging of hand and finger joints in rheumatoid arthritis. *Arthritis Rheum.*, 48:325–33.
- Weidekamm C.; Krestan C.; Schueller G.et al.(2007):** Power Doppler Sonography and Pulse-Inversion Harmonic Imaging in Evaluation of Rheumatoid Arthritis Synovitis. *American Jornal of Rheumatology.*, 188:504-508.
- Weinblatt M.E.; Kremer J.M.; Coblyn J.S. et al., (1999):** Pharmacokinetics, safety and efficacy of the combination of

- methotrexate and leflunomide in patients with active rheumatoid arthritis. *Arthritis Rheum.*, 48 (7): 1322 – 1327.
- Wener M.H. (2004):** Absence of antibodies to cyclic citrullinated peptide in sera of patients with hepatitis C virus infection and cryoglobulinemia, *Arthritis Rheum.*, 50: 2305–2308.
- Weyand C.M.; Hicok K.C.; Conn D.L. et al. (1992):** The influence of HLADR B1 genes on disease severity in RA. *Ann. Inter. Med.*, 117 : 801-809.
- Wiell C.; Szkudlarek M.; Hasselquist M. et al. (2007):** Ultrasonography, magnetic resonance imaging, radiography, and clinical assessment of inflammatory and destructive changes in fingers and toes of patients with psoriatic arthritis. *Arthritis Res Ther.*,9:119-130.
- Wislowska M.S.; Sypula, I. and Kowalik (1999):** Echocardiographic findings and 24-h electrocardiographic Holter monitoring in patients with nodular and non-nodular rheumatoid arthritis, *Rheumatol. Int.*, 18 : 163–169.
- Wolf M.M.; Lichtenstein D.R. and Singh G. (1999):** Gastrointestinal toxicity of nonsteroidal anti-inflammatory drugs. *N. Engl. J. Med.*, 340: 1888-1892.
- Wolfe F., Hawely D.J. and Cathey M.A. (1993):** Measurement of gold treatment effect in the clinical practice : evidence for effectiveness of intramuscular gold therapy. *J. Rheumatol.*, 20 : 797 – 802.
- Wolfgang A. and Schmidt. (2004):** Doppler sonography in rheumatology. *Best Practice & Research Clinical Rheumatology.*, 18:827–846.

Wright S.; Filippucci E.; Grassi W. et al.(2006): Hand arthritis in systemic lupus erythematosus. an ultrasound pictorial essay. *Lupus*,15:501–6.

Y

Yamada H.; Nakashima Y.; Okazaki K. et al. (2008): Th1 but not Th17 cells predominate in the joints of patients with rheumatoid arthritis. *Annals of the rheumatic diseases*.,67:1299–1304.

Yoshida M. (1999): Prevalence of mutilans-like hand deformities in patients with seropositive rheumatoid arthritis. A prospective 20-year study, *Scand. J. Rheumatol.*, 28 (1) 38–40.

Yu K.H.; Ho H.H.; Chen J.-Y. et al. (2004): Gout complicated with necrotizing fasciitis—report of 15 cases *Rheumatology*., 43: 518-521.

Z

Zielonka T. M.; Barela D.; Zukowska M.et al. (2005): Caplan's syndrome.*Pol Arch Med Wewn.*, 114:779-84.

Zrioual S.; Ecochard R.; Tournadre A. et al. (2009): Genome-wide comparison between IL-17A- and IL-17F-induced effects in human rheumatoid arthritis synoviocytes. *J Immunol.*,182:3112–3120.

Zwiebel, W.J. (2005): Color duplex sonography : Capabilities and limitations. In Lisette B(ed): Introduction to vascular

References

ultrasonography, Saunders Company. Philadelphia, 4th ed.,
PP:157-166.