

REFERENCE

Altman R., Asch E., Bloch D., et al. (1986): Development of criteria for the classification and reporting of osteoarthritis of the knee. *Arthritis Rheum* 29: 1039-1049.

Aglietti P, Inshall J N, Walker P S Trent P (1975): Patellofemoral contact zones. *Clin Orthop*, 107: 175.

Aglietti P, Windsor RE, Buzzi R, et al. (1989): Arthroplasty for the stiff or ankylosed knee. *J Arthro*, 4 (1): 1–5.

Aglietti P. and Baldini A. (2006): Minimally Invasive Total knee Arthroplasty: Is it for every body? *Hospital for Special Surgery Journal*, p. 22-26.

Agur M.R. and Lee M.J. (1996): Grant's atlas of anatomy, the ninth edition,, Williams and Wilkins, p. 300-310.

Alfred J. Tria(2006), Exploring the Depths of Minimally Invasive Quadriceps-Sparing Total Knee Arthroplasty *J Bone Joint Surg Am.*; **88:226-229**

Alfred J Tria, Jr.(2002)total knee arthroplasty in *Surgical Techniques in Total Knee Arthroplasty* edited by R. Scuderi, M.D. Alfred J. Tria, Jr., M.D. Insall Scott Kelly Institute chapter24 p:182- 197 Springer-Verlag New York, Inc.

Berger R.A., Jacobs J.J., Meneghini R.M., Della Valle C., Pa-prosky W., Rosenberg A.G. (2004): Rapid rehabilitation and recovery with minimally invasive total hip arthroplasty. Clin Orthop Relat Res, 429: 239-247

Berger R., Sanders S., and Gerlinger T. (2005): Outpatient total knee arthroplasty with a minimally invasive technique. J Arthroplasty, 20: 33-38.

Bertin CB.(1991) Intramedullary instrumentation for total knee arthroplasty. In: Goldberg VM, ed. *Controversies in Total Knee Arthroplasty*. New York: Raven Press, Ltd., Chap. 18

Bennell K.L., Hinman R.S., Metcalf B.R., et al. (2003): Relationship of knee joint proprioception to pain and disability in individuals with knee osteoarthritis. J Orthop Res 21: 792-797.

Boerger TO, Aglietti P, Mondanelli N, et al(2005). Mini subvastus versus medial parapatellar approaches in total knee arthroplasty. Clin Orthop;440:82.

Bong MR, DiCesare PE, Paul E(2004). Stiffness after total knee arthroplasty. J Am Acad Orthop Surg; 12:164

Bonutti P, Mont M, McMahon M, et al.(2004) Minimally invasive total knee arthroplasty. J Bone Joint Surg Am;86(Suppl 2):26

Bozic K.J., Kinder J., Meneghini R.M. (2005): Implant survivorship and complication rates after total knee arthroplasty with a third-generation cemented system: 5 to 8 years follow-up. Clin Orthop Relat Res, (430): 117-24.

Calligaro K.D. and Dougherty M.J. (2003): Acute arterial complications associated with total hip and knee arthroplasty. J Vasc Surg, 38: 1170-11770.

Carl A., Berger R., and Richard A (2006): A technique for minimally invasive, quadriceps-sparing total knee arthroplasty The Journal Of Knee Surgery January, Vol. 19, No. 1.

Craig S.M. (1994): Soft tissue considerations in the failed total knee arthroplasty. In: Scott WN, ed. The Knee. St. Louis, Mosby, 1279-1295

Curl W.W. (1996): Popliteal cysts. Historical background and current knowledge. J Am Acad Orthop Surg, 4:129-133

Dalury D.F. and Dennis D.A. (2005): Mini-incision total knee arthroplasty can increase risk of component malalignment. Clin Orthop Relat Res, 440: 77-81.

Dark R.L. and Vogal W. (2007): The knee. In: Grays anatomy edited by dark rl et al.,p532-540 ELSEVIER London.

David F. Dalury, Dennis(2005), Mini-incision Total Knee Arthroplasty Can Increase Risk of Component Malalignment Clin Orthop Relat **Res**;440:77

Ecker M.M., Lotke P. (1989): Postoperative care of the total knee patient. Orthop Clin North Am, 20: 55–62.

Eckhoff D.G. (2005): Functional anatomy of the knee in total knee arthroplasty guide to get better performance edited by bellemans. j victor.j Springer Medizin Verlag Heidelberg 2005 chapter 3, p18-25.

Eckhoff D. et al. (2001): Three-dimensional morphology and kinematics of the distal part of the femur viewed in virtual reality, part I. J Bone Joint Surg 83-A [Suppl 2]:43–50.

Eckhoff D. et al. (2003): Three-dimensional morphology and kinematics of the distal part of the femur viewed in virtual reality, part II. J Bone Joint Surg 85-A [Suppl 4]:97–104.

Eckhoff D. et al. (1996): Sulcus morphology of the distal femur. Clin Orthop Rel Res, 331: 23–28

Eckhoff D. et al. (1994): Femoral anteversion and arthritis of the knee. J Pediatr Orthop, 14:608–610

Eckhoff D. et al. (1996): Location of the femoral sulcus in the osteoarthritic knee. J Arthroplasty, 11:163–165

Eckhoff D. et al. (1999): Femorotibial offset. A morphologic feature of the natural and arthritic knee. Clin Orthop Rel Res 368:162–165.

Eckhoff D. et al. (1997): Knee version associated with anterior knee pain. Clin Orthop Rel Res, 339: 152–155.

Eckhoff D. et al. (1994): Version of the osteoarthritic knee. J Arthroplasty, 9:73–79.

Eckhoff D. et al. (1995): Malrotation associated with implant alignment technique in total knee arthroplasty. Clin Orthop Rel Res 321:28–31.

Engh G.A., Holt B.T., Parks N.L. (1997): A midvastus muscle-splitting approach for total knee arthroplasty. J Arthroplasty, 12: 322-331.

England SP, Stern SH, Insall JN, et al. (1990): Total knee arthroplasty in diabetes mellitus. Clin Orthop, 260: 130–134.

Felson D.T., Chaisson C.E., Hill C.L., et al. (2001): The association of bone marrow lesions with pain in knee osteoarthritis. Ann Intern Med 134:541-549.

Figgie H. et al. (1989): The effect of alignment of the implant on fractures of the patella after condylar total knee arthroplasty. J Bone Joint Surg 71-A: 1031–1039.

Geerts W., Heit J., and Clagett G.: Prevention of venous thromboembolism. Chest, 119: 132S

Gerard A. (2002): Engh mid vastus approach. In; Surgical techniques in total knee arthroplasty. By: Giles R. and Alfred J., New York: Springer-Verlag, Chapter 16, pp. 127-131, 162.

Giles R. Scuderi and Alfred J. Tria, Jr(2004). Minimal Incision Total Knee Arthroplasty in mis of and knee editeby Giles R. Scuderi and Alfred J. Tria, Jr chapter 10 p 175 186 Springer-Verlag New York, Inc

Goble EM, Justin DF(2004). Minimally invasive total knee replacement: principles and technique. Orthop Clin North Am;35: 235–45)

Grood E.S., Suntag W.J., Noyes F.R., Butler D.L. (1984): Biomechanics of the knee extension exercise. J Bone Joint Surg, 66A:725

Guyton JL, Crockarell JR and John R. (2003): Arthroplasty of the ankle and knee. In: Campbell's operative orthopedics, edited by **CANALE, st DAUGHERTY k and JONES L** tenth edition Mosby, Inc, chapter 6 p. 250.

Haas S.B. and Lehman A.P. (2005): Minimally invasive total knee arthroplasty. In: Total knee arthroplasty a guide to get better performance. By: Bellemans J, Victor J, (eds.), Springer Medizin Verlag Heidelberg, Chapter 44, p. 228- 293.

Haas S.B., Cook S. and Beksac B. (2004): Minimally invasive total kneereplacement through a mini midvastus approach: a comparativestudy. Clin Orthop, 428: 68-73.

Hofmann A.A., Plaster R.L., Murdock L.E. (1991): Subvastus (southern) approach for primary total knee arthroplasty. Clinical Orthopaedics, 269: 70-77.

Hooper M.M. and Moskowitz R.W. (2007): Osteoarthritis: Clinical Presentations IN Osteoarthritis: Diagnosis and Medical / Surgical Management, edited by Moskowitz, Roland W.; Altman, Roy D et al Lippincott Williams & Wilkins, 4th ed., p. 135-145.

Hoppenfeld S. and de Boer P. (1994): The knee. In: Surgical exposures in orthopaedics - the anatomic approach, 2nd ed, JB Lippincott, Philadelphia, p 429-482.

Insall J.N. (1993): Surgical approaches. In: Surgery of the Kn. By: Insall J.N., Windsor R.E., Scott W.N., Kelly M.A., Aglietti P. (eds.), New York, Churchill Livingstone, p. 135–148.

Insall J., ranawat C., Scott W.N., Walker P. (1976): total condylar knee replacement: preliminary report. Clin Orthop, 120: 149-154.

Insall J.N, Ranawat C.S., Aglietti P., Shine J(1990): A comparison of four models of total knee replacement prosthesis. J Bone Joint Surg, 58-A: 754.

Jason K. and Daniel L. (2007): Postoperative Recuperative Period Total Knee Arthroplasty. Assessment of the Learning Curve and the Minimally Invasive Total Knee Arthroplasty Compared with Traditional. J Bone Joint Surg Am, 89: 1497-1503. doi:10.2106/JBJS.F.00867

Jennifer L.C and Fred D. (2006): Minimal-Incision Total Knee Arthroplasty The Journal Of Knee Surgery January, Vol. 19, No.1

Johnson D.P., Houghton T.A., Radford P. (1986): Anterior midline or medial parapatellar incision for arthroplasty of the knee. A comparative study. J Bone Joint Surg, 68B: 812.

Kapandji I. (1987): The physiology of the joints. Churchill-Livingston, New York

Kelly G. (2002): Vince subvastus approaches. In: Surgical techniques in total knee arthroplasty. By: Giles R. and Alfred J., New York, Springer-Verlag, Chapter 15, p 120-126.

Kenneth M.D. (2006): Minimally Invasive Total Knee Arthroplasty With a Subvastus Approach Oper Tech Orthop 16:153-158, Elsevier Inc

Kettlekamp D.B. (1978): Gait, characteristics of the knee: normal, abnormal and postreconstruction. In American Academy of Orthopaedic Surgeons: Symposium on reconstructive surgery of the knee. St Louis, Mosby.

Kevin R.M and Giles R.S. (2000): Preoperative radiographic assessment in surgical techniques. In: Total knee arthroplasty. By: Giles R. and Tria, Jr., Springer-Verlag New York, Inc., p 9-19. 27: 1513

Krackow K.A. (1990): The technique of total knee arthroplasty St Louis, Mosby.

Krakow K.A. (1990): The Technique of Total Knee Arthroplasty. St. Louis, MO: C.V. Mosby, p. 168–197.

Laskin R.S. (2005): Minimally invasive total knee arthroplasty: the results justify its use. Clin Orthop, 440: 54-59.

Laskin .Richard S. (2006): Reduced-incision total knee replacement through a mini-midvastus technique. The Journal of Knee Surgery, January, Vol. 19, No. 1

Lee D.C. and Kim D.H. (1998): Intraoperative flexion against gravity as an indication of ultimate range of motion in individual cases after total knee arthroplasty. J Arthroplasty, 13(5): 500-503.

Leopold S.S. and McStay C. (2001): Primary repair of intraoperative disruption of the medial collateral ligament during total knee arthroplasty. J Bone Joint Surg Am, 83: 86-91.

Lonner J. H. Booth, Jr.(2005) Assessment and Balancing of Patellar Tracking in total knee arthroplasty A Guide to Get Better Performance By: Bellemans J, Victor J, (eds.), Springer Medizin Verlag Heidelberg, ,chapter36 p. 229-233

Luyten F.P. Westhovens (2005): Arthritis of the knee diagnosis and management in total knee arthroplasty edited by Johan Bellemans Jan M.K. Victor et al Springer Medizin Verlag Heidelberg, p:3-14.

Malkani A.L., Rand J.A., Bryan R.S., Wallrich S.L. (1995): total knee arthroplasty with the kinematic condylar prosthesis. A ten-year follow-up study. J Bone Joint Surg Am, 77: 423-431.

Maric Z, Ott DM, Karpman RR(1991). The standard versus the sub-vastus (southern) approach for total knee arthroplasty: a randomized prospective study. Orthop Trans;15:43 .

Michael S., Rodney K. and Tria, Jr (2006): Minimally Invasive Quadriceps-Sparing Posterior Stabilized Total Knee Arthroplasty Oper Tech Orthop, 16:170-176, Elsevier Inc.

Moreland J.R. (1988): Mechanisms of failure in total knee arthroplasty. Clin Orthop, 226:49.

Pai Y.C., Rymer W.Z., Change R.W., et al. (1997): Effect of age and osteoarthritis on knee proprioception. J Rheumatol 40: 2260-2265.

Patterson B.M., Insall J.N. (1992): Surgical management of patients with poliomyelitis. J Arthro (suppl 2): 419–426.

Rand J.A., Ilstrup D.M. (1991): Survivorship analysis of total kneearthroplasty. Cumulative rates of survival of 9200 total knee arthroplasties. J Bone Joint Surg, 73-A:397-409.

Richard A.D. (2006): Minimally Invasive Quadriceps-Sparing Cruciate-Retaining Total Knee Arthroplasty Oper Tech Orthop 16:177-185, Elsevier Inc.

Richard A.B. and Sanders R.N. (2006): Minimally Invasive Quadriceps-Sparing TKA: Results of a Comprehensive Pathway for Outpatient TKA. The Journal of Knee Surgery, Vol. 19, No.

Ritter M.A., Keating E.M. (1990): Comparison of two anterior medial approaches to total knee arthroplasty. Am J Knee Surgery, 3: 168-171.

Rosenberg A.G., Andriacchi T.P., Barden R., Galante J.O. (1988): Patellar component failure in cementless total knee arthroplasty. Clin Orthop, 238: 106.

Scott M., Sporer M. (2006): The Minimally Invasive Subvastus Approach for Primary Total Knee Arthroplasty. The Journal of Knee Surgery, Vol. 19, No

Scuderi G.R. and Tenholder M. (2004): Surgical approaches in mini-incision total knee arthroplasty. Clin Orthop.

Scuderi and Tria, Jr (2004): Minimal Incision Total Knee Arthroplasty mis of hip and knee Springer, p. 175-186.

Scuderi G.R., Tria A.J. (2005): Minimally invasive total knee arthroplasty. In: Insall J.N., Scott W.N. (eds.): Surgery of the knee. 4th ed., Philadelphia PA: Elsevie, 1631-1639

Scuderi GR, Insall JN (1991) Cement technique in primary total knee arthroplasty. Tech Orthop 6:39-43)

Scuderi G.R and Clarke, H.(2005) Optimizing Cementing Technique Total knee arthroplasty a guide to get better performance. By: Bellemans J, Victor J, (eds.), Springer Medizin Verlag Heidelberg Chapter 35p 223-227)

Sheila Sanders, R.N. Carl Deirmengian, M.D., et al. (2006): Perioperative Protocols for Minimally Invasive Total Knee Arthroplasty The Journal Of Knee Surgery, Vol. 19, No 2.

Siegel JL, Shall LM.,(1991) Femoral instrumentation using the anterosuperior iliac spine as a landmark in total knee arthroplasty.
An anatomic study. *J Arthroplasty*. 6(4):317–320.)

Soudry M., Binazzi R., Johanson N.A., et al. (1986): Total kne arthroplasty in Charcot and Charcot-like joints. Clin Orthop, 208: 199-204.

Sowers M.F., Jamadar D., Capul D., et al. (2003): Magnetic resonance-detected subchondral bone marrow and cartilage defect characteristics associated with pain and X-ray defined knee osteoarthritis. Osteoarthritis Cart, 11: 387-393.

Stephen B. and Gayle W. (2006): Sub-vastus approach versus the medial parapatellar approach in primary total knee. Trials Journal

Stern SH (1994): Surgical exposures in total knee arthroplasty. In: Knee Surgery. By: Fu F., Harner C.D., Vince K.G. (eds.). Baltimore: Williams & Wilkins, p. 1289–1302.

Steven F. Harwin (2002): Indications and Patient Selection in Surgical Techniques. In: Total Knee Arthroplasty. By: Giles R and Tria, Jr., Springer-Verlag New York, Inc., p3-9.

Steven H, Stern (2002): Medial Parapatellar Arthrotomy. In: Surgical techniques in total knee arthroplasty. By: Giles R. and Alfred J (eds.), New York, Springer-Verlag, Chapter14, p. 115-119.

Tanavalee A and Thiengwittayaporn S. (2005): Results of the 136 consecutive minimally invasive total knee arthroplasties. J Med Assoc Thai, 88 (Suppl 4): S74-8.

Tenholder M., Scuderi G.R., Capeci C.(2004): Surgical approaches in mini-incision total knee arthroplasty. Clin Orthop.:428 61_67

Tenholder M, Clarke H, Scuderi G(2005). Minimal-incision total knee arthroplasty: the early clinical experience. Clin Orthop Relat Res;440:67

Tria A (2004): Minimally invasive total knee arthroplasty: the importance of instrumentation. Clin. Orthop. North Am, 35: 227-234

Tria A (2003): MIS TKR: is it feasible? AAOS 70th annual meeting proceedings. New Orleans, 2003, 6: 33-37.

Tria A.J. Jr and Coon T.M. (2003): Minimal incision total knee arthroplasty: early experience. Clin Orthop Relat Res, 416: 185-90.

Walton N.P., Dobson P.J. and Campbell D.G. (2005): Arthrofibrosis following total knee replacement; does therapeutic warfarin make a difference? Knee, 12: 103-6.

Wasielowski R. et al. (1994): Wear patterns on retrieved polyethylene tibial inserts and their relationship to technical considerations during total knee arthroplasty. Clin Orthop Rel Res 299:31–43.

Watanabe, T. MUNETA et al. (2009): MIS vs. conventional approach in TKA the journal of orthopaedics science, 22

Weir DJ, Moran CG, Pinder IM. Kinematic condylar total knee arthroplasty: fourteen year survivorship analysis of 208consecutive cases. J Bone Joint Surg, 78B (6): 907–911.

William C. and Paul J. (2007): Evaluation of Complications Associated with Six Hundred Mini-Subvastus Total Knee Arthroplasties