ACKNOWLEDGEMENT

At first I Thank the Merciful Allah and I'm greatly indebted to **Prof. Dr. Hatem Mostafa Ashour** for suggesting the point of research, generous support during the progress of this work, revising the subject and for his continuous encouragement and valuable advices. I attribute the level of my Masters to his fatherhood manner in teaching me and without him this research would not have been completed.

I hope to express my gratitude to **Prof. Dr. Al Husseiny Mostafa** for his guidance during this work and his grateful advices, simulating discussions and continuous revising and comments during progression of this work.

I would like to acknowledge **Dr. Mohamed Akef** for his continuous supervision, valuable advice, suggestion, and comments that enriched me to work on this research.

My deepest gratitude goes to my beloved family for their unflagging love and support throughout my life, this dissertation is simply impossible without them.

AHMAD HASSAN AL SAYYAD

AIM OF WORK

The purpose of this work is to provide an overview of hip impingement syndrome depending on good understanding of the condition as it was difficult to be diagnosed in the past, showing types, ideal methods of diagnosis and proper management of such cases.

LIST OF FIGURES

- *Fig.* (1) Schematic drawing of hip joint impingement.
- *Fig.*(2) A. Anterosuperior impingement test. B. Posteroinferior impingement test.
- *Fig.*(3) Radiograph showing pistol grip deformity of femoral head.
- *Fig.*(4) Axial view of normal hip offset (OS) and normal alpha angle < 50.
- **Fig.(5)** Radiograph showing decreased femoral head–neck offset (OS) with consecutive increased alpha angle.
- *Fig.*(6) Anteroposterior radiograph of the pelvis illustrating acetabular overcoverage.
- *Fig.*(7) Anteroposterior radiograph showing ossification of labral basis and prominent acetabular bone fragment that separated from adjacent bone margin (os acetabuli).
- Fig.(8) Lateral decubitus position of the patient and firmly secured with specific holding devices.
- Fig.(9) A straight lateral Gibson-style approach.
- Fig.(10) Preparatory steps for the trochanteric osteotomy.
- Fig.(11) Direction of the osteotomy of the internally rotated hip.
- *Fig.*(12) Femoral shaft, the trochanteric fragment advancement anteriorly with a Hohmann retractor.
- *Fig.*(13) A stepwise capsulotomy is performed. (B)Performing the distal leg of the z-shaped capsulotomy
- *Fig.*(14) Femoral head is dislocated. This is performed by flexion and external rotation with the leg secured in a sterile bag.
- Fig.(15) Three retractors are inserted, to view the acetabulum.
- *Fig.*(16) Cranial retroversion of the acetabulum is caused by an anterolateral overhang of the rim.

- *Fig.*(17) Femoral correction usually involves trimming of the aspherical portion of the head, normally located anterosuperiorly.
- *Fig.*(18) Patient in a lateral decubitus position with the right hip marked. Note the trochanter and the anterosuperior iliac spine with the portals marked out.
- *Fig.*(19) Right hip with the arthroscope in the anterolateral portal; the needle is in the posterolateral portal and the anterior portal.