

References

- Abbate M, Zoja C, and Remuzzi G.(2006):** How does proteinuria cause progressive renal damage? J Am Soc Nephrol,; 17:2974-2978.
- Abdel-Hady HE and Zaki ME.(2003):** Evaluation of soluble E-selectin as a marker for neonatal sepsis. The Egyptian journal of neonatology; 4 (2): 69-78.
- Abdel-Rahman W.H.A.(2002):** Premature rupture of membranes: The value of interleukin-1 receptor antagonist in predicting neonatal sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; .Supervised by Awad HAS, El Raggal NM, Youssef MF.
- Acunas, B.A.; Peakman, M. and Lossis, G. (2004).** Effect of fresh frozen plasma and gamma globulin on humoral immunity in neonatal sepsis. Archives of Disease in childhood, 70: 182-187.
- Adel M.,Awad H A., Abdel-Naim A B., Al-Azizi M M.,(2010).** Effects of pentoxifylline on coagulation profile and disseminated intravascular coagulation incidence in Egyptian septic neonates. Journal of Clinical Pharmacy and Therapeutics Volume 35 Issue 3, Pages 257 – 265.
- Ahmed HA(2003):** Early prediction of neonatal sepsis by a hematological scoring system. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain- Shams University; .Supervised by Khazbak MAM, El Kerdany TA, Abou Seif IS.

Ali RA(2006): Serum adiponectin in neonatal sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain-Shams University; Supervised by Soliman NA, Emem EK, Abdel Wahab SEA.

Anderson-Berry AL, Bellig LL, MacGilvray S S, Windle M L, Clark DA, et al.(2006): Neonatal sepsis. Department of Neonatology, Greenville Children's Hospital and Medical University of South Carolina .

Arnon S, Litmanovitz I(2008): Diagnostic tests in neonatal sepsis. Curr Opin Infect Dis;21(3):223-227.

Avery, G.B. (2006). Sepsis neonatorum. Neonatology. Pathophysiology and Management of the Newborn, 4th ed, P. 1088-1096.

Avory, A.; Fanarroff, A. and Richard, J. (2002): Neonatal prenatal Medicine. Infections in the fetus and neonate (31): 763-797.

Austgulen, R.; Arntzeuk, J. and Haereid, P.E. (2005).Infection in neanates delivered at term are associated with increased serum levels of ICAM-1 and E-selectin. Act paediatr, 86 (3): 274-80.

Axelsson L, Bergenfeldt M, Ohlsson K.(1995). Studies of the release and turnover of a human neutrophil lipocalin. Scand J Clin Lab Invest; 55: 577–88.

Badrawi NH, Bashir MM, Iskander IF, Saied DA(2005).: Neonatal infections in NICU: magnitude of the problem. Kasr El-Aini Medical Journal,; 11 (5): 181-195.

Baker, C.; Edwards, M.S.; In Remington, J.S. and Klein, J.O. (2005): Infectious diseases of the Fetus and Newborn infant.

Baley, J. and Fanaroff A.A. (2001). Effective care of the newborn infant. Neonatal infections. J. Infection related to nursery care practices.

Baltimore, R.S. (2003). neonatal sepsis: Epidemiology and management. Pediatr. Drugs (11): 723- 740.

Baltimore, A. (2002): Perinatal bacterial and fungal infections. Pediatric infectious diseases principals and practice. 96: 1119-33.

Barclay, L. (2004). Ciprofloxacin in neonatal sepsis lacks long term adverse effects. The pediatric infectious disease Journal 23 : 346 – 349.

Baudry A, Yang ZZ, Hemmings BA.(2006). PKB is required for adipose differentiation of mouse embryonic fibroblasts. J Cell Sci; 119:889–897.

Bauer M, Eickhoff JC, Gould MN, Mundhenke C, Maass N, Friedl A.(2008). Neutrophil gelatinase-associated lipocalin (NGAL) is a predictor of poor prognosis in human primary breast cancer. Breast Cancer Res Treat, 108(3):389-397.

Bellig, L.L. (2004). Neonatal sepsis. (review). Eur. J. Pediatr., 155: 20-126.

- Benitz, W.E.; Han, M. and Madan, A. (2001).**Serial C-reactive protein levels in the diagnosis of neonatal infection. *Pediatrics*, 102 (4): 5-10.
- Benjamin DK, DeLong ER, Steinbach WJ, Cotton CM, Walsh TJ, and Clark RH(2003):** Empirical therapy for neonatal candidemia in very low birth weight infants. *Pediatrics*; 112 (3): 543-547.
- Berger, C.; Uehlinger, D.J.D. Blau, N. and Fanciness (1995).** Comparison of protein and white blood cell count with differential in at risk for septicemia. *Eur. J. of pediatrics* 154: 138-144.
- Betsuyaky T, Nishimura M, Takeyabu K, Tanino M, Venge P, et al.(2001).:** Neutrophil granule proteins in bronchoalveolar lavage fluid from subjects with subclinical emphysema. *Am J Respir Crit Care Med*; 159:1985-1991.
- Bjorkqvist M, Kallman J, Fjaertoft G, Xu S, Venge P, Schollin J.(2004)** : Human neutrophil lipocalin: normal levels and use as a marker for invasive infection in the newborn. *Acta Paediatr*; 93:534–539.
- Bolignano D, Coppolino G and Campo S(2008):** Urinary neutrophil gelatinase-associated lipocalin (NGAL) is associated with severity of renal disease in proteinuric patients. *Nephrol Dial Transplant*; 23:414-416.
- Brodie SB, Sands KE, Gray JE, Parker Ra, Goldmann DA, Davis RB(2000):** Occurrence of nosocomial bloodstream infections in six neonatal intensive care units. *Pediatr Infect Dis J*; 19 (1): 56-62.

Borregaard N, Cowland JB.(2006). Neutrophil gelatinase- associated lipocalin, a siderophorebinding eukaryotic protein. *Biometals*; 19: 211–215.

Cairo, M.S.; Worcester, C.C. and Rucker, R.W. (2002). Randomized trial of granulocyte transfusion versus IVIG therapy for neonatal neutopenia and sepsis. *J. pediatr.*, 120-281-85.

Carlson M, Raab Y, Seveus L, Xu S, Hallgren R and Venge P(2002): Human neutrophil lipocalin is a unique marker of neutrophil inflammation in ulcerative colitis and proctitis. *Gut*; 50:501-506.

Carr R.(2000). Neutrophil production and function in newborn infants.

Br J Haematol; 110: 18–28.

Carrigan SD, Scott G, Tabrizian M.(2004):. Toward resolving the challenges of sepsis diagnosis. *Clin Chem*; 50:1301–1314.

Chacko B, Sohi I.(2005):. Early onset neonatal sepsis. *Indian J Pediatr.*; 72 (1): 23-26.

Chiesa, C.; Panero, A. Obsborn, J.F. and Pacifice, L. (2004).Diagnosis of neonatal sepsis. A clinical and laboratory challenge. *Clinical chemistry* 50 : 279 – 287.

Christensen EI, Mishra J, Thiessen-Philbrook H and Gburek J(2004): Protein reabsorption in renal proximal tubule function and dysfunction in kidney pathophysiology. *Pediatr Nephrol*; 19:714-721.

Coppa, G.; Gabrielli, O.F. and Gicergi, P. (2001). Preliminary study of breast feeding and bacterial adhesion to epithelial cells. *Lancet*, 1: 569 – 580.

Cotton CM, McDonald S, Stoll B, Goldberg RN , Poole K and Benjamin DK.(2006). The association of third-generation cephalosporin use and invasive candidiasis in extremely low birth weight infants. *Pediatrics*; 118 (2): 717-722.

Cowland J B, Seveus L, Xu S and Borregaard N(2001): Molecular characterization and pattern of tissue expression of the gene for neutrophil gelatinase-associated lipocalin from humans. *Genomics*; 45(1):17-23.

Cowland JB, Borregaard N.(1997). Molecular characterization and pattern of tissue expression of the gene for neutrophil gelatinase associated lipocalin from humans. *Genomics*; 45:17–23.

Da Silva, O.; Ohlsson, A. And Kenyon, A. (2005). Accuracy of leukocyte indices and C-reactive protein for diagnosis of neonatal sepsis: *pediatr. Infect. Dis. J.*, 4363-366.

Dear, P. (1999). Infection in Newborn. In : *Text book of Neonatology* 3rd ed. (eds). P. 1109 – 202.

DeCamp L. R, Dempsey A. F, and Tarini B.A.(2009). Neonatal Sepsis: Looking Beyond the Blood Culture: Evaluation of a Study of

Universal Primer Polymerase Chain Reaction for Identification of Neonatal Sepsis Arch Pediatr Adolesc Med. ;163(1):12-14.

-Devireddy L R, Gazin C, Zhu X and Green M R.(2005).: A cell-surface receptor for lipocalin 24p3 selectively mediates apoptosis and iron uptake. Cell;123(7):1293-1305.

-Ding H, He Y and Li K.(2007).: Urinary neutrophil gelatinase-associated lipocalin (NGAL) is an early biomarker for renal tubulo-interstitial injury in IgA nephropathy. Clin Immunol; 123:227-234.

Dollener, H.; Arntzen, K.J.; Haereid, P.E.; Aag, S. and Austglen, R. (1998). IL-6 Concentrations in neonates evaluated for sepsis. J. Pediatr. , 132 (2) : 295- 9.

Dzwonek AB, Neth OW, Thiebaut R, Gulcynska E, Chilton M, et al.(2008).: The role of mannose-binding lectin in susceptibility to infection in preterm neonates. Ped Res; 63: 680-685.

Ehl, S.; Gehring, B. and Pohlandt, F. (2005). A detailed analysis of changes in serum CRP levels in neonates treated for bacterial infection. Eur. J. Pediatr., 158: 238-42.

El-Khodary AR, Afify MF, Saleem TH and Maher SE.(2001).: Procalcitonin (PCT) and interleukin-6 (IL-6) as markers for neonatal sepsis. The Gaz Egypt Ped; 49 (3): 345-356.

El-Raggal NM, El-Barbary MN, Youssef MF , El-Mansy HA.(2004).: Neutrophil surface antigens CD11b and CD64 expression: a

potential predictor of early-onset neonatal sepsis. Egypt J Pediatr Allergy Immunol; 2 (2): 90-100.

Fanaroff AA, Korones SB, Wright LL, Verter J, Poland RL, et al.(1998).: Incidence, presenting features, risk factors and significance of late-onset septicemia in very low birth weight infants. Pediatr Infect Dis J; 17 (7): 593-598.

Fergany AL.(2006).: Serum cortisol and thyroid hormone levels in neonates with sepsis. Thesis submitted for partial fulfillment of Master degree in Pediatrics, Faculty of Medicine, Ain- Shams University. Supervised by El Barbari MN, Imam SS, Shaheen KY.

Fernandez C A, Yang L, Louis G, Yang J and Kutok J L.(2005).: The matrix metalloproteinase-9/neutrophil gelatinase-associated lipocalin complex plays a role in breast tumor growth and is present in the urine of breast cancer patients. Clin Cancer Res; 11(15):5390-5395.

Figueroa, D.R.; Arredondo,G.J.L.; Mancilla, A. and Ramirez, J. (1999).Aminiotic fluid IL-6 and the risk of early – onset sepsis among preterm infants. Arch. Med. Res., 30 (3): 198- 202.

Fisher AG, Smith K, Seddon B, Purbhoo MA, Zamoyska R, Merkschlager M,(2001). Sensory adaptation in naive peripheral CD4 T cells. J Exp Med.;194(9):1253-61.

Fjaertoft G, Foucard T, Xu S, Venge P.(2005).: Human neutrophil lipocalin (HNL) as a diagnostic tool in children with acute infections: a study of the kinetics. *Acta Paediatr*; 94:661-666.

Flo, T. H., Smith, K. D., Sato, S., Rodriguez, D. J., Holmes, M. A., Strong, R. K., Akira, S.,and Aderem, A. (2004). Lipocalin 2 mediates an innate immune response to bacterial infection by sequestering iron. *Nature* 432, 917–921.

Flower DR.(2000). : The lipocalin protein family: structural and sequence overview.*Biochim Biophys Acta*; 1482:9-24.

Flower DR.(1996). The lipocalin protein family: structure and function. *Biochem J* 318:1–1.

Fournier T, Medjoubi N and Porquet D.(2000).: Alpha-1-acid glycoprotein. *Biochemica et Biophysica. Acta*; 1482(1):157-171.

Friedl A, Stoesz S P, Buckley P and Gould MN.(2000).: Neutrophil gelatinase-associated lipocalin in normal and neoplastic human tissues. Cell type-specific pattern of expression. *Histochem J*; 31(7):433-441.

Gabay, C. and Kushner, I. (2002). Acute phase proteins and other systemic responses to inflammation. *N. Engl. J. Med.* 340: 448-454.

Gathwala G., Chaudhary U.,et al.,(2010). Ten Days vs. 14 days Antibiotic Therapy in Culture-Proven Neonatal Sepsis. *Journal of Tropical Pediatrics*, doi:10.1093/tropej/fmq012.

Goetz DH, Holmes MA, Borregaard N, Bluhm ME, Raymond KN, Strong RK. (2002). The neutrophil lipocalin NGAL is a bacteriostatic agent that interferes with siderophore-mediated iron acquisition. Mol Cell 10:1033–1043.

Gomella, T.; Douglas, M.; Eyal, G. and Zenk, E. (2004). Infectious Diseases, In: Clinical Manual of Neonatology, 5th ed. P.434-468.

Gonzalez, E.; Cecilia, K.; Mercadon, A. and Linda, J. (2004). Early markers of late onset sepsis in premature neonates: clinical, hematological and cytokine profile. J. perinat. Med., 31: 60.

Gonzalez BE, Mercado CK, Johnson L, Brodsky NL, Bhandari V.(2003).: Early markers of late-onset sepsis in premature neonates: clinical, hematological and cytokine profile. J Perinat Med; 31: 60-68.

Griffin MP and Moorman JR.(2001).: Towards the early diagnosis of neonatal sepsis and sepsis-like illness using novel heart rate analysis. Pediatrics;107(1):97–104.

Guerina, N.G. (2002). Bacterial and fungal infection. Manual of neonatal care, 4th ed., 271-300.

Guillois, B.; Berthou, C.; Awad, H. and Youinou, P. (1989). The importance of C3d estimation in the diagnosis of generalized bacterial infections in new born infants. Acta pediatri. 78: 368-372.

Haque K and Mohan P.(2003).: Pentoxifylline for neonatal sepsis. Cochrane Database of Systematic Reviews, Issue4.[DOI: 10.1002/14651858.CD004205].

Hague, R.A.; Rassams and Morgan, G. (1994). Early diagnosis of severe combined immunodeficiency syndrome Archives of disease in childhood 70: 260- 263.

Hanson, L.O. and Lindquist, L. (2005). C-reactive protein: Its role in the diagnosis of and follow up of infectious diseases. Cur. Opin. Infect Dis., 10:196-201.

Hashim MS, Aboulghar HM, El-Gayar DF and Hamam AO.(2004).: Evaluation of serum cortisol and ACTH levels in neonatal sepsis. The Egyptian journal of neonatology; 5 (3): 135-142.

Hebra A., Larson SD (2006). Meconium pseudocyst in a newborn. J Am Coll Surg;203(3):399.

Herzum, I., and Renz, H. (2008). Inflammatory markers in SIRS, sepsis and septic shock. Review. *Curr Clin Chem* 15, 581–87.

Hickey, S.M. and McCracken, G. (2001). postnatal bacterial infections neonatal perinatal Medicine: Diseases of the fetus and Infants. 6th ed.

Hodge, S.; Hodge, G. and Flower, R. (2001). Surface activation markers of T-lymphocyte : Role in the detection of infection in neonates. Clin. Exp. Immunol., 113: 33-38.

Hoffman, D.J. and Harris, M.C. (2002). Diagnosis of neonatal sepsis: in intensive care of the fetus and Neonate. Spitzer, A.R. (Ed): Year Book, Boston, London. P. 940-950.

Isaacs, D. and Moxen, R.E. (1999). Handbook of neonatal infection: A practical guide. Saunders, London.

Iskander I and Morcos H.(2006):. Soluble P-selectin response in FT and PT newborns with neonatal sepsis: impact on aetiology and correlation with other inflammatory parameters. The Egyptian journal of neonatology; 7 (3):92-96.

Jayaraman A, Roberts KA, Yoon J, Yarmush DM, Duan X, Lee K, et al. (2005) .Identification of neutrophil gelatinase-associated lipocalin (NGAL) as a discriminatory marker of the hepatocyte-secreted protein response to IL-1 β : a proteomic analysis. Biotechnol Bioeng;91:502–15.

Jensen , H.B. and Pollock, B.H. (2004).Meta-analysis of the effectiveness of intravenous immunoglobulin for prevention and treatment of neonatal sepsis. *Pediatrics*, 99: 246.

Jiang JH, Chiu NC, Huang F-Y, Kao HA, Hsu CH, Hung HY, Chang JH and Peng CC (2004): Neonatal sepsis in the neonatal intensive care unit: characteristics of early versus late onset. *J Microbiol Immunol Infect*;37:301-306.

Jordan J A, Durso M P,Butchko A R et al. (2006). Evaluating the Near-Term Infant for Early Onset Sepsis Progress and Challenges to Consider with 16S rDNA Polymerase Chain Reaction Testing *J Mol Diagn*. July; 8(3): 357–363.

Kanaoka Y and Urda Y.(2003):. Hematopoietic prostaglandin D synthase. Prostaglandins. Leukotrienes. Essent. Fatty acids; 69:163-167.

Kemp. A.S. and Campbell, D. D. (2002). the neonatal immune system. *Seminar in Neonatology*, 1: 67-75.

Kenon, C.; Overturf, G. and Bessman, S. (1996). Granyocyte colony stimulating factor a marker for bacterial infection in neonates. *Journal of pediatrics* 128: 765-769.

King, J.C.; Berman, E.D. and Wright, P.E. (2000). Evaluaiton of sepsis in infants less than 5 weeks old. *South Med. J.*, 80-998-952.268(14):10425-10432.

Kjeldsen L. (1996). Induction of NGAL synthesis in epithelial cells of human colorectal neoplasia and inflammatory bowel diseases. *Gut* 38 414-20.

Kjeldsen, L., Cowland, J. B., and Borregaard, N. (2000). Human neutrophil gelatinase-associated lipocalin and homologous proteins in rat and mouse. *Biochim. Biophys. Acta* 1482, 272–283.

Klein, J.O. and Marcy, S.M. (1995). Bacterial sepsis and meningitis. *Infectious Diseases of the Fetus and New born Infant*. 4th ed.; 835-878.

Klinger G, Levy I, Sirota L, Boyko V, Reichman B, Lerner-Geva L. Epidemiology and risk factors for early onset sepsis among very-low-birth weight infants. *Am J Obstet Gynecol*. Jul 2009; 201(1):38.e1-6

Kratchmarova I, Kalume DE, Blagoev B, Scherer PE, Podtelejnikov AV, Molina H, et al.(2002). A proteomic approach for identification of secreted proteins during the differentiation of 3T3–L1 preadipocytes to adipocytes. *Mol Cell Proteomics* ; 1:213–222.

Kumar Y, Qunibi M, Neal TJ, Yoxall CW.(2001).: Time to positivity of neonatal blood cultures. *Arch Dis Child Fetal Neonatal Ed*; 85(3):F182-186.

AV, Molina H, et al.(2002). A proteomic approach for identification of 40
Wang et al.: Lipocalin-2 and Obesity-Related Diseases secreted
proteins during the differentiation of 3T3–L1 preadipocytes to
adipocytes. Mol Cell Proteomics;1:213–22.

Lam, H.S. and Ng, P.C. (2008). Biochemical markers of neonatal sepsis.
Pathology 40; 141 – 148.

Lawn JE, Cousens S, Zupan J (2005). 4 million neonatal deaths: when?
Where? Why? Lancet 365: 891–900.

Lin Y, Rajala MW, Berger JP, Moller DE, Barzilai N, Scherer PE.(2001).
Hyperglycemia-induced production of acute phase reactants in
adipose tissue. J Biol Chem; 276:42077–42083.

Linder N, Ohel G, Gazit G, Keidar D, Tamir I, Reichman B.(1995).
Neonatal sepsis after prolonged premature rupture of membranes. J
Perinatol.;15(1):36-8.

Lieberman E, Lang JM, Richardson D K, Ringer SA, and Cohen A
(1997): Epidural Analgesia, Intrapartum Fever, and Neonatal
Sepsis Evaluation. PEDIATRICS Vol. 99 No. 3 March, pp. 415-
419.

Lee S, Lee J, Kim S, Park JY, Lee WH, Mori K, Kim SH, Kim IK, Suk K
(2007).A dual role of lipocalin 2 in the apoptosis and

deramification of activated microglia. J Immunol 179:3231–3241.

Leheste JR, Rolinski B and Vorum H.(1999).: Megalin knockout mice as an animal model of low-molecular-weight proteinuria. Am J Pathol.; 155:1361-1370.

Levy O, Martin S, Eichenwald E, et al.(1999). Impaired innate immunity in the newborn: newborn neutrophils are deficient in bactericidal/permeability-increasing protein. Pediatrics 1999; 104: 1327–33.

LockJ, Craft A, Finer N.(2001). Nosocomial coagulase negative staphylococcal (CoNS) catheter-related sepsis in preterm infants: definition, diagnosis, prophylaxis, and prevention. J Perinatol.;21(3):186-92.

Lott MJ ,Kreuzer-Martin HW, Ehleringer JR, Hegg EL.(2006). Metabolic processes account for the majority of the intracellular water in log-phase Escherichia coli cells as revealed by hydrogen isotopes. Biochemistry. 14;45(45):13622-30.

-Mahieu LM, De Dooy JJ, De Muynck AO, Laroche SM and Van Acker KJ.(2000).: Prediction of nosocomial sepsis in neonates by

means of a computer-weighted bedside scoring system (NOSEP score). Crit Care Med; 28 (6): 2026-2033.

Manroe, B.L.; Weinberg, A.G. and Rosenfeld, C.R. (1979). Neonatal blood count in health and disease. Reference value for neutrophilic cells. J. Pediatr., 95 (1): 89 – 95.

Marten, J.; Samson, W.P.F. and Fetter, A. (1999). TNF- α , IL-6 and IL-1B plasma level in neonatal sepsis. Pediatr. Re. 33 (4): 380- 383.

-Mathai E, Christopher U, Mathai M, Kumar Jana A, Rose D, et al.(2004). Is C-Reactive protein level useful in differentiating infected from uninfected neonates among those at risk of infection. Indian Pediatr; 41:895–900.

McConathy WJ and Alaupovic P.(1973). Isolation and partial characterization of apolipoprotein D: a new protein moiety of the human plasma lipoprotein system. FFBS Lett; 37:178-182.

Mckenna, D.S. and Iams, J.D. (2004): Group B-streptococcal infections. Seminars in perinatology, 22: 26-76.

McPherson RJ and Juul S.(2005). Patterns of thrombocytosis and thrombocytopenia in hospitalized neonates. Journal of perinatology; 2: 166-172.

Meheus LA, Fransen LM, Raymackers JG, Blockx HA, Beeumen JJ, Bun SM, et al.(1993). Identification by microsequencing of lipopolysaccharide-induced proteins excreted by mouse macrophages. *J Immunol*; 151:1535–1547.

Michael, R. and Jeffery, L. (2002).Advances in the therapy for sepsis in children: *Pediatric Clinics of North America*, 44 (1): 179-202.

Miharada K I, Hiroyama t, Sudo k, Nagasawa t, Nakamura y.(2005). Lipocalin 2 functions as a negative regulator of red bloodcell production in an autocrine fashion. *The FASEB Journal* express article 10.1096/fj.05-3809fje.

Mishra J, Ma Q, Kelly C, Mitsnefes M and Mori K.2006):. Kidney NGAL is a novel early marker of acute injury following transplantation. *Pediatr Nephrol*; 21(6):856-863.

Mishra J, Dent C, Tarabishi R, Mitsnefes MM and Ma Q.(2005):. Neutrophil gelatinase-associated lipocalin (NGAL) as a biomarker for acute renal injury after cardiac surgery. *Lancet*; 365(9466):1231-1238.

Mishra J, Mori K, Ma Q et al. (2004).Amelioration of ischemic acute renal injury by neutrophil gelatinase-associated lipocalin. *J Am Soc Nephrol*; 15: 3073–3082

Monneret, G.; Doche, C.; Lepape, A. and Bienvenu, J. (2005):

Procalcitonin as a specific marker of bacterial infection in infants. Clin. Chem. Lab. Med., 36: 67-8.

Mori K, Lee HT, Rapoport D, Drexler IR, Foster K, Yang J, Schmidt-Ott KM, Chen X, Li JY, Weiss S, Mishra J, Cheema FH, Markowitz G, Suganami T, Sawai K, Mukoyama M, Kunis C, D'Agati V, Devarajan P, Barasch J (2005). Endocytic delivery of lipocalin-siderophore-iron complex rescues the kidney from ischemia-reperfusion injury. J Clin Invest. 115:610–621.

Mori K, Nakao K. (2007). Neutrophil gelatinase-associated lipocalin as the real-time indicator of active kidney damage. Kidney Int 71:967–970.

Nagata A, Suzuki Y, Igarashi M, Eguchi N, Toh H, Urade Y and Hayaishi, O. (1991):. Human brain prostaglandin D synthase has been evolutionarily differentiated from lipophilic ligand carrier protein. National Academy of Sciences; 88: 4020-4024.

Nelson AM, Zhao W, Gilliland KL, Zaenglein AL, Liu W, Thiboutot DM (2008). Neutrophil gelatinase-associated lipocalin mediates 13-*cis* retinoic acid-induced apoptosis of human sebaceous gland cells. J Clin Invest 118:1468 –1478.

Nilsen-Hamilton M, Liu Q, Ryon J, Bendickson L, Lepont P, Chang Q. (2003). Tissue involution and the acute phase response. *Ann N Y Acad Sci* 995 94-108.

Ng PC, Lam HS.(2006). Diagnostic markers for neonatal sepsis ,Curr Opin Pediatr. Apr;18(2):125-31.

Ng PC.(2004).: Diagnostic markers of infection in neonates Arch Dis Child Fetal Neonatal Ed; 89 (3): F229-F235.

Ng, PC.; Fok, S.H.; Chenge, G. et al., (1997). Diagnosis of late onset neonatal sepsis with cytokines, adhesion molecule, and C-reactive protein in preterm very low birth weight infants. Arch. Dis. Child., 77:F22 – F227.

Nguyen MT, Devarajan P.(2008).Biomarkers for the early detection of acute kidney injury. Pediatr Nephrol ; 23:2151-2157.

Nuntarumit, P.; Pinkaew, O. and Kitiwan, W. (2002). Predictive values of serial C-reactive protein in neonatal sepsis. J. Med. Assoc.

Nupponen I, Venge P, Pohjavuori M, Lassus P, Andersson S.(2000).: Phagocyte activation in preterm infants following premature rupture of the membranes or chorioamnionitis. Acta Paediatr; 89: 1207–1212.

Oddie S, Embleton ND.(2002).: Risk factors for early onset neonatal group B streptococcal sepsis: case-control study. BMJ; 325: 308-312.

Ohlsson, A. and Lacy, J.B. (2005).Intravenous immunoglobulin for suspected or subsequently proven neonatal infection (Cochrane Review). In: The Cochrane Library, issue 2.

Okano M, Fujiwara T, Sugata Y, Gotoh D, Masaoka Y, et al.(2006).: Presence and characterization of pregnancy protein 14- related molecules in glandular epithelium of the endometrium. Journal of Medicine; 20(3):342-348.

Ortlund E, Parker CL, Schreck SF, Sodetz JM and Minor W.(2002).:

Crystal structure of human complement protein C8 gamma at 1.2 A resolution reveals a lipocalin fold and a distinct ligand binding site. *Biochemistry*;41(22):7030-7037.

Ottolini MC, Lungren K, Mirkinson LJ, Cason S, Ottolini MG.(2003).:

Utility of complete blood count and blood culture screening in the diagnosis of neonatal sepsis. *Pediatr infect Dis J*; 17: 222-235.

Parikh CR, Mishra J, Thiessen-Philbrook H, Dursun B and Ma

Q.(2006).: Urinary IL-18 is an early predictive biomarker of acute kidney injury after cardiac surgery. *Kidney Int*; 70(1):199-203.

Parikh CR, Jani A, Mishra J, Ma Q, Kelly C, et al.(2006).: Urine NGAL

and IL-18 are predictive biomarkers for delayed graft function following kidney transplantation. *Am J Transpl*; 6:1639–1645.

Park JM, Kanaoka Y, Eguchi N, Aritake K, Grujic S, et al.(2007).:

VonEbner's glands protein. *Biochem Biophys Acta*,;1282(1-2):99-108.

Peter Dear, A. (2002).Infection in the Newborn. In *Textbook of*

Neonatology. Part 2: 1109-1138.

Phares CR, Lynfield R, Farley MM.(2008). Epidemiology of invasive

group B streptococcal disease in the United States, 1999–2005. *JAMA*. ;299(17):2056–2065.

Philip, A.G. and Hewitt, J.R. (1990). Early diagnosis of neonatal sepsis. Pediatrics, 65: 1036-41.

Pourcroyous, M.; Bada, H.S.; Orones, S.B. and Wong, S.P. (2004).significance of serial C-reactive protein responses in neonatal infection and other disorders. Pediatrics, 92: 431-35.

Powell, K.R. and Marcy, S.M. (2005). laboratory aids for diagnosis of neonatal sepsis. Infectious Diseases of Fetus and Newborn infants, P. 1223-1240.

Procianoy RS and Silveira RC.(2004): The role of sample collection timing on interleukin-6 levels in early-onset neonatal sepsis. J Pediatr; 80 (5): 407-410.

Quadro L, Blaner WS, Hamberger L, Novikoff PM, Vogel S, et al.(2004): The role of extrahepatic retinol-binding protein in the mobilization of retinoid stores. Journal of Lipid Research; 45:1975-1982.

Ragharen, M.; Mandal, G.P.; Bhat, B.V. and Srinivasen, S. (2003).Perinatal risk factors in neonatal infection. Ind. J. pediatr., 59 (3): 335-340.

Rassart E, Bedirian A, Do Carmo S, et al. (2000). "Apolipoprotein D". Biochim. Biophys. Acta **1482** (1-2): 185–98.

Reier-Nilsen T, Farstad T, Nakstad B, Lauvrak V , Steinbakk M.(2009): comparison of broad range 16S rDNA PCR and

conventional blood culture for diagnosis of sepsis in the newborn: A case control study. *BMC Pediatr*; 9:5-9.

Remington, J.S. and Klein, J.O. (2001). Infectious diseases of the fetus and newborn infant, 5th ed.

Rennie, J.M. and Robertson, N.P.C. (2002). Neonatal infection. In A manual of Neonatal Intensive care, 10th ed., P. 223-56.

Rey C, Los Arcos M, Concha A, Medina A, Prieto S, Martinez P, Prieto B. Procalcitonin and C-reactive protein as markers of systemic inflammatory response syndrome severity in critically ill children. *Intensive Care Med.* 2007).

Rodwell, R.L.; Leslie, A.L. and Tudehope, D.I. (1988). Early diagnosis of neonatal sepsis using a hematologic scoring system. *J. Pediatr.*, 112 (2) : 761-7.

Russel, A.R.; Davies, E.G. and McGuigan, S. (2003). Plasma granulocyte colony stimulating factor concentrations (G-CSF) in the early neonatal transfusion practices: Blood component therapy. *Pediatrics*, 91: 530-36.

Said H.(2006).: Spectrum of neonatal respiratory distress with special reference to clinical and bacteriological profile of early-onset neonatal pneumonia in a tertiary neonatal care unit, Suez Canal University hospital. *The Egyptian journal of neonatology*; 7(1):71-73.

Schmidt-Ott KM, Mori K, Li JY et al. (2007). Dual action of neutrophil gelatinase-associated lipocalin. *J Am Soc Nephrol*; 18: 407–413.

Schmidt-Ott KM, Mori K, Kalandadze A, Li JY and Paragas N.(2006).: Neutrophil gelatinase-associated lipocalin-mediated iron traffic in kidney epithelia. *Curr Opin Nephrol Hypertens*; 15(4):442-449.

Schreck SF, Parker C, Plumb ME and Sodetz JM.(2000).: Human complement protein C8 gamma . *Biochem Biophys Acta*; 1482(1-2): 199-208.

Schuchat A, Zywicki SS, Dinsmoor MJ, Dinsmoor M J, Mercer B, J Romaguera J, O'Sullivan M J, Patel D, Peters M T, Stoll B and Levine O S (2000): Risk Factors and Opportunities for Prevention of Early-onset Neonatal Sepsis: A multicenter Case-Control Study. *Pediatrics*;105:21-26.

Seaward PG ,Barrea C, Hornberger LK, Alkazaleh F, McCrindle BW, Roberts A, Berezovska O, Windrim R , Smallhorn JF, Ryan G.(2006).
Impact of selective laser ablation of placental anastomoses on the cardiovascular pathology of the recipient twin in severe twin-twin transfusion syndrome. *Am J Obstet Gynecol*. 2006 Nov;195(5):1388-95.

Seveus L, Amin K, Peterson CG, Roomans GM, Venge P.(1997). Human neutrophil lipocalin (HNL) is a specific granule constituent of the neutrophil granulocyte. Studies in bronchial and lung parenchymal tissue and peripheral blood cells. *Histochem Cell Biol*; 107: 423–32.

Silveri, R.C. and Procianoy, R.S. (1999). Evaluation of IL-6, TNF- α and IL1 B for early diagnosis of neonatal sepsis. *Acta. Pediatr.*, 88 (6): 647-50.

Skerra, A. (2000).Lipocalins as a scaffold. *Biochim. Biophys. Acta* 1482, 337–350.

Soman KV, Das S, Sierra JC, Suarez G, Mohammad AA, Dang TA, Luxon BA, Reyes VE.(2005). Differential protein expression profiles of gastric epithelial cells following Helicobacter pylori infection using ProteinChips. *J Proteome Res*;4(3):920-30.

Stoll, B.J. (2007).Infection of the neonatal infants. *Nelson text book of pediatrics* (18th ed). Ch., 109: 794-811.

Stoll, B.J. and Weisman, L.E. (2004).Infection in perinatology. *Clin. Perinatol.*, 24:1.

Stoll BJ, Gordon T, Korones SB, Shankaran S, Tyson JE, et al.(1996).: Late-onset sepsis in very low birth weight neonates: a report from

the National institute of child health and human development,
Neonatal Research Network J Pediatr; 129: 63-71.

Stuart, R.L.; Turnidge, J. and Grayson, M.L. (2005). Safety of imipenem in neonates. *Pediatr. Infect. Dis. J.*, 14: 804-5.

Thaver D, Zaidi AK (2009). Burden of neonatal infections in developing countries: a review of evidence from community-based studies. *Pediatr Infect Dis J* 28: S3–9.

Timothy K, Schwartz PJ, Spazzolini C, Crotti L, Bathen J, Amlie JP, Timothy K, Shkolnikova M, Berul CI, Bitner-Glindzicz M, Toivonen L, Horie M, Schulze-Bahr E, Denjoy I.(2006). The Jervell and Lange-Nielsen syndrome: natural history, molecular basis, and clinical outcome. *Circulation.*;113(6):783-90.

Tollner, U. (1982). Early diagnosis of septicemia in newborn. Clinical studies and sepsis score *Eur. J pediatric.* 138 (10): 331-7.

Towbin RB ,Arcement CM, Meza MP, Gerber DA, Kaye RD, Mazariegos GV, Carr BI, Reyes J(2000). Intrahepatic chemoembolization in unresectable pediatric liver malignancies. *Pediatr Radiol.* (11):779-85.

Umana, M.A.; Odio, C.A.; Castro, E.; Salas, J.L. and McCracken, G.H. (2006).Evaluation of a zetronam and ampicillin versus amikacin and ampicillin for treatment of neonatal bacterial infections. *Pediatr. Infect. Dis. J.*, 9: 175-80.

Urade Y, Eguchi N, Aritake K, and Hayaishi O.(2004).: Functional analysis of lipocalin type and hematopoietic prostaglandin D synthase. *Department of Molecular Behavioral Biology*; 123 (1): 5-13.

Urrea M, Iriondo M, Thio M, Krauel X, Serra M, et al.(2003).: A prospective incidence study of nosocomial infections in a neonatal care unit. *Am J Infect Control*; 31 (8): 505-507.

Van-Benneken AM, Wei S, Gamble MV, Vogel S, Piantadosi R and Blaner WS.(2001).: Biochemical basis for depressed serum retinol levels in transthyretin-deficient mice. *J Biol Chem*; 276: 1107-1113

Van-Timmeren MM, Bakker SJ and Vaidya VS.(2006).: Tubular kidney injury molecule-1 in protein-overload nephropathy. *Am J Physiol Renal Physiol*;291:F456-F464.

Venge, (2000) .U.S. Pat. No. 6,136,526; PCT application WO95/29404).

Vogel S, Mendelson CL, Mertz J, Piantadosi R, Blaner WS and Gottesman ME.(2001).: Characterization of a new member of

the fatty acid-binding protein family that binds all-trans-retinol. J Biol Chem; 276:1353-1360.

Wagener G , Jan M, Kim M, Mori K, Barasch JM, Sladen RN, et al. (2006). Association between increases in urinary neutrophil gelatinase associated lipocalin and acute renal dysfunction after adult cardiac surgery. Anesthesiology; 105: 485-491.

Wang Y, Lam KS, Kraegen EW, Sweeney G, Zhang J, Tso AW, et al. (2007). Lipocalin-2 an inflammatory Marker Closely Associated with Obesity, Insulin Resistance and Hyperglycemia in Humans. Clin Chem ; 53(1):34-41.

Weinschen, K.N.P.; Fariona A. and Bianchi, D.W. (2005). Neonatal neutrophil activation is a function of labor length in preterm infants. Pediatr. Res., 44: 42-945.

Xu SY and Venge P.(2000).: Lipocalin as biochemical marker of disease. Biochem Biophys Acta; 18; 1482 (1-2): 298-307.

Xu S, Pauksen K, Venge P.(1995). Serum measurements of human neutrophil lipocalin (HNL) discriminate between acute bacterial and viral infections. Scand J Clin Lab Invest; 55: 125–31.

Xu SY, Petersson CG, Carlsson M, Venge P.(1994).: The development of an assay for human neutrophil lipocalin (HNL) to be used as a specific marker of neutrophil activity in vivo and in vitro. J Immunol Methods; 171: 245–252.

Xu S, Venge P:(2000). Lipocalins as biochemical markers of disease. Biochim Biophys Acta; 1482: 298–307.

Yan QW, Yang Q, Mody N, Graham TE, Hsu CH, Xu Z,Houstis NE, Kahn BB, Rosen ED (2007).The adipokine lipocalin 2 is regulated by obesity and promotes insulin resistance. Diabetes 56:2533–2540

Yang J, Goetz D, Li JY, Wang W, Mori K, Setlik D, Du T, Erdjument Bromage H, Tempst P, Strong R, Barasch J (2002). An iron delivery pathway mediated by a lipocalin. Mol Cell 10:1045–1056.

Yang J, Borregaard N, Kjeldsen L and Moses MA.(2001).: The high molecular weight urinary matrix metalloproteinase (MMP) activity is a complex of gelatinase B/MMP-9 and neutrophil gelatinase-associated lipocalin (NGAL). Modulation of MMP-9 activity by NGAL. J Biol Chem; 276(40):37258-37265.

Yousef ARA.(2003).: Interleukin-1 and phospholipase A2 in septic newborns. Thesis submitted for partial fulfillment of master degree in pediatrics, faculty of medicine, Ain-Shams University. Supervised by Ismail IS, Imam SS, Shaheen KY.

Zaidi AK, Thaver D, Ali SA, Khan TA (2009). Pathogens associated with sepsis in newborns and young infants in developing countries. Pediatr Infect Dis J 28: S10–18.