

REFERENCES

1. **Miedzybrodzka, Z. (2003):** Congenital talipes equinovarus (clubfoot): a disorder of the foot but not the hand. *J. Anat.*;202:37-42.
2. **Dietz, F. (2002):** The genetics of idiopathic clubfoot. *Clinical Orthopaedics and Related Research*:401:39-48.
3. **Strach, E.H. (1986):** Clubfoot Through the Centuries. In *Progress in Pediatric Surgery Vol. 20*. Ed. by P.P. Rickham. Berlin Heidelberg. Springer-Verlag. 215-237.
4. **Turco, V.J. (1981):** Clubfoot. New York. Churchill Livingstone.
5. **Kite, J.H. (1955):** The operative treatment of congenital clubfeet. *Instr Course Lect*;12:100-105.
6. **Kite, J.H. (1963):** Some suggestions on the treatment of club foot by casts. *J Bone Joint Surg Am*;45-A:407-413.
7. **Shimizu, N.; Hamada, S. and Mitta, M. (1993):** Etiological considerations of congenital clubfoot deformity. Tachdjian MO, Simons G, editors. *The clubfoot: the present and a view of the future*. New York:Springer:p 31-8.
8. **Lochmiller, C.; Johnston, D.; Scott, A.; Risan, M. and Hecht, J.T. (1998):** Genetic epidemiologic study of idiopathic talipes equinovarus . *Am J Med Genet*.79:90-96.
9. **Cummings, R.J.; Davidson, R.S.; Armstrong, P.F. and Lehman, W.B. (2002):** Congenital clubfoot. *J Bone Joint Surg.*:84 A (2):290-307.
10. **Haf, G.F.; Walker, C.G. and Crawford, H.A. (2007):** Early clubfoot recurrence after use of the Ponseti method in a New Zealand population. *J Bone Joint Surg Am.* ;89:487–493.

- 11.**Huerta de la, F. (1994):** Correction of the neglected clubfoot by the Ilizarov method. Clin. Orthop.; **301**:89–93.
- 12.**Anand, A. and Sala, D.A.(2008):** Clubfoot: Etiology and treatment. Indian J Orthop;42:22-8.
- 13.**Ponseti, I.V.(1996):** “Congenital clubfoot: Fundamentals of treatment.” Oxford: Oxford University Press.
- 14.**Ponseti, I.V. (1994):** The treatment of congenital clubfoot. J Orthop Sports PhysTher;20:1.
- 15.**Ponseti, I.V. (1992):** Treatment of congenital club foot. J Bone Joint Surg Am; 74:448-454.
- 16.**Stronmeyer, L. (1836):** Division of the tendo achilles in clubfoot. Lancet; 3:645.
- 17.**Kite, J.H. (1972):** Nonoperative treatment of congenital clubfoot. Clin Orthop; 84:29–38.
- 18.**Ponseti, I.V. (1992):** Treatment of Congenital Clubfoot. J Bone Joint Surg; 74A:448-454.
- 19.**Wicart, P.R.; Barthes, X.; Chanem, I. and Sezing, R. (2002):** Clubfoot posteromedial release: advantages of tibialis anterior tendon lengthening. J Pediatr Orthop; 22 : 536-532.
- 20.**Choi, I.H.; Yang, M.S.; Chung, C.Y.; Cho, T.J. and Sohn, J.(2001):** The treatment of recurrent arthrogrypotic clubfoot in children by the Ilizarov method. A preliminary report. J Bone Joint Surg. 83-B : 731-737.
- 21.**Turco, V.J. (1981):** Clubfoot. In: Current Problems in Orthopaedics., Churchill Livingstone, NY, Edinburg and Melbourne.
- 22.**Dimeglio, A.; Bensahel, H.; Souchet, P.; Mazeau, P. and Bonnet, F. (1995):** Classification of clubfoot. J Pediatr Orthop; 4-B : 129-136.

23. **McKay, D.W. (1983):** New concept of an approach to clubfoot treatment evaluation and results. *J Pediatr Orthop*;3:141–8.
24. **Paley, D. (1993):** The correction of complex foot deformities using Ilizarov's distraction osteotomies. *Clin Orthop*; 293:97–111.
25. **Ferreira, R.C.; Costa, M.T.; Frizzo, G.G. et al., (2006):** Correction of neglected clubfoot using the Ilizarov external fixator. *Foot Ankle Int*; 27:266–73.
26. **Grant, A.D.; Atar, D. and Lehman, W.B. (1992):** The Ilizarov technique in correction of complex foot deformities. *Clin Orthop*; 280:94–103.
27. **Ilizarov, G.A.; Shevtsov, V.I. and Kuz'min, N.V. (1983):** Method of treating talipes equinovarus. *Ortop Travmatol Protez*; 5:46–8.
28. **Ferreira, R.C.; Costa, M.T.; Frizzo, G.G.; et al., (2007):** Correction of severe recurrent clubfoot using a simplified setting of the Ilizarov device. *Foot Ankle Int*; 28:557–68.
29. **Bradish, C.F. and Noor, S. (2000):** The Ilizarov method in the management of relapsed club feet. *J Bone Joint Surg Br*;82-B:387–91.
30. **Lehman, W.B. (1996):** The clubfoot. JB Lippincott: New York; ١٧.
31. **Issacs, H.; Handelsman, J.E.; Badenhorst, M. and Pickering, A. (1977):** The muscles in clubfoot: A histological, histochemical and electron microscopic study. *J Bone Joint Surg Br*; 59:465-72.
32. **Irani, R.N. and Sherman, M.S.(1963):** The pathological anatomy of clubfoot. *J Bone Joint Surg Am*;45:45-52.
33. **Loren, G.J.; Karpinski, N.C. and Mubarak, S.J. (1998):** Clinical implications of clubfoot histopathology. *J Pediatr Orthop*;18:765-9.
34. **Shapiro, F. and Glimcher, M.J. (1979):** Gross and histological abnormalities of talus in congenital clubfoot. *J Bone Joint Surg Am*; 61:522-30.

- 35.**Ionasescu, V.; Maynard, J.A.; Ponseti, I.V. and Zellweger, H. (1974):** The role of collagen in the pathogenesis of idiopathic clubfoot: Biochemical and electron microscopic correlations. *Helv Paediatr Acta*; 29:305-14.
- 36.**Ippolito, E. (1995):** Update on pathologic anatomy of clubfoot. *J Pediatr Orthop B*;4:17-24.
- 37.**Hootnick, D.R.; Levisohn, E.M.; Crider, R.J. and Packard, D.S. Jr. (1982):** Congenital arterial malformations associated with clubfoot: A report of two cases. *Clin Orthop Relat Res*; 167:160-3.
- 38.**Turco, V.J. (1981):** Clubfoot. Churchill Livingstone: New York.
- 39.**Porter, R.W. (1987):** Congenital talipes equinovarus: Resolving and resistant deformities. *J Bone Joint Surg Br*;69:822-5.
- 40.**Pryor, G.A.; Villar, R.N.; Ronen, A. and Scott, P.M. (1991):** Seasonal variation in the incidence of congenital talipes equinovarus.*J Bone Joint Surg Br*; 73:632-4.
- 41.**Edwards, M.J.(1971):** The experimental production of clubfoot in guinea pigs by maternal hyperthermia during gestation. *J Pathol*; 103:49-53.
- 42.**Ippolito, E. and Ponseti, I.V.(1980):** Congenital clubfoot in human fetus: A histological study. *J Bone Joint Surg Am*;62:8-22.
- 43.**Ignacio, V. and Ponseti (2002):** Relapsing ClubFoot: Causes, Prevention and Treatment. *The Iowa Orthopedic Journal*.22:55-56.
- 44.**Roye, D.P. and Roye, B.D. (2002):** Idiopathic congenital talipes equinovarus. *J Am Acad Orthop Surg*; 10:239–48.
- 45.**Adams, W. (1866):** Clubfoot: Its causes, pathology and treatment. J and A Churchill: London.
- 46.**Adams, W. and Brockman, E.P. (1930):** Congenital clubfoot. John Wright and Sons: Bristol.

- 47.**Bost, F.C.; Schottstadt, E.R. and Larsen, L.J. (1960):** Plantar dissection-an operation to release soft tissues in recurrent talipes equinovarus. J Bone Joint Surg Am;42:151-64.
- 48.**Pirani, S.; Zeznik, L. and Hodges, D.(2001):** Magnetic resonance imaging study of the congenital clubfoot treated with the Ponseti method. J Pediatr Orthop;21:719 –726.
- 49.**Pirani, S. (2004):** A reliable and valid method of assessing the amount of deformity in the congenital clubfoot. Presented at the Pediatric Orthopaedic Society of North America; May; St. Louis.
- 50.**Penny, A. (2001):** School access: children with motor disabilities in rural Uganda [MA Thesis Dissertation]. University of Victoria.
- 51.**Yadav, S.S. (1981):** Observations on operative management of neglected clubfoot. Int Orthop;5:189 –192.
- 52.**Canale & Beaty (2008):** Campbell's Operative Orthopaedics, 11th ed.2008;26:2-9.
- 53.**Lippincott Williams & Wilkins.(2005):** Techniques In Orthopaedics;20: 153-166.
- 54.**Cummings, R.J.; Davidson, R.S.; Armstrong, P.; et al., (2002):** Congenital clubfoot. AAOS instructional course lecture. J Bone Joint Surg [Am];84:290–308.
- 55.**Ferlic, R.J.; Breed, A.L.; Mann, D.C.; et al., (1997):** Partial wound closure after surgical correction of equinovarus foot deformity. J Pediatr Orthop;17:486–489.
- 56.**Bitariho, D. (2001):** Short term outcome in one stage pantalar soft tissue release with transfixation of talonavicular joint in resistant idiopathic congenital clubfoot [MMed (Orth) Thesis Dissertation]. Makerere University; Uganda; April.
- 57.**Staheli, L.; ed. (2003):** Clubfoot: Ponseti Management. Seattle: Global-HELP Publications.

- 58.**Evans, D. (1961):** Relapsed clubfoot. J Bone Joint Surg [Br]43:722– 733.
- 59.**Scott, W.A.; Hosking, S.W. and Catterall A. (1984):** Club foot. Observations on the surgical anatomy of dorsiflexion. J Bone Joint Surg [Br];66:71–76.
- 60.**Clark, M.W.; D’Ambrosia, R.D. and Ferguson, A.B. (1977):** Congenital vertical talus: treatment by open reduction and navicular excision. J Bone Joint Surg [Am];59:816–824.
- 61.**Koutsogiannis, E. (1971):** Treatment of mobile flat foot by displacement osteotomy of the calcaneus. J Bone Joint Surg [Br];53:96–100.
- 62.**Dwyer, F.C.(1959):** Osteotomy of the calcaneum for pes cavus. J Bone Joint Surg [Br];41:80–86.
- 63.**Ponseti, I.V.(1996):** Congenital Clubfoot: Fundamentals for Treatment. Oxford: Oxford University Press.
- 64.**Davidson, R.S. (2003):** Clubfoot salvage: a review of the past decade’s contributions. J Pediatr Orthop; 23:410–418.
- 65.**Lambrinudi, C. (1927):** New operation on drop-foot. Br J Surg ;15: 193.
- 66.**Heurta, F. (1994):** Correction of the neglected clubfoot by the Ilizarov technique. Clin Orthop;301:89 –93.
- 67.**Yadav, S.S. (1981):** Observations on operative management of neglected clubfoot. Int Orthop;5:189 –192.
- 68.**Angus, P.D. and Cowell, H.R.(1986):** Triple arthrodesis. A critical long term review. J Bone Joint Surg [Br]; 68:260 –265.
- 69.**Heurta, F. (1994):** Correction of the neglected clubfoot by the Ilizarov technique. Clin Orthop; 301:89 –93.

70. **Aronson, J.; Harrison, B.; Cannon, D.J. et al.,** : Mechanical induction of osteogenesis. The importance of pin rigidity. J Pediatr Orthop 1988; 8:396.
71. **Pasyk, K.A.; Argenta, L.C. and Austad, E.D. (1987):** Histopathology of human expanded tissue. Clin Plast Surg; 14:435-445.
72. **Ilizarov, G.A. (1989):** The tension stress effect on the genesis and growth of tissue. Part 2: The influence of the rate and frequency of distraction. Clin Orthop; 238:263-285.
73. **Dyachkova, G.V. and Utenkin, A.A. (1980):** Extensibility of superficial fascia in elongation of the leg in experiment. Ortop Travmatol Protez; 41:44-47.
74. **Appell, H.J.; Forsberg, S. and Hollman, W. (1988):** Satellite cell activation in human skeletal muscle after training. Evidence for muscle fibre neoformation. J Sports Med Phys Fitness; 9:297-299.
75. **Kojimoto, H.; Yasui, N.; Goto, T.; et al., (1988):** Bone lengthening in rabbits by callus distraction. J Bone Joint Surg [Br] 70B:543-549.
76. **Ilizarov, G.A. (1989):** The tension-stress effect on the genesis and growth of tissues. Part 1: The influence of stability of fixation and soft tissue preservation. Clin Orthop; 238:249-281.
77. **Paley, D.; Kovelman, H.F. and Herzenberg, J.E. (1993):** Ilizarov Technology; 1:244-50.
78. **Debastian, I.G.; Aldegheri, R.; Rezni-Brivio, L.; et al., (1987):** Limb lengthening by callus distraction (callotaxis). J Pediatr Orthop; 7:129-134.
79. **Paley, D.; Rumley, T. and Kovelman, H. (1991):** The Ilizarov technique: A method to regenerate bone and soft tissue. Advances in Plastic and Reconstructive Surgery 7:1-40.

80. **Aronson, J.; Good, B.; Stewart, C.; et al., (1990):** Preliminary studies of mineralization during distraction osteogenesis. Clin Orthop 250:43-49.
81. **Delloye, C.; Delafortie, G.; Coutelier, L.; et al., (1990):** Bone regenerate formation in cortical bone during distraction lengthening: An experimental study. Clin Orthop; 250:34-42.
82. **Tajana, G.F.; Morandi, M. and Zembo, M. (1989):** The structure and development of osteogenic repair tissue according to the Ilizarov technique in man. Characterization of the extracellular matrix. Orthopedics; 12:515-523.
83. **Mc.Kibbin, J. (1978):** The biology of fracture healing in long bones. J Bone Joint Surg [Br]; 60B:150-162.
84. **Cross, A.R.; Lewis, D.D.; Murphy, S.T.; et al., (2001):** Effects of ring diameter and wire tension on the axial biomechanics of four-ring circular external fixator constructs. Am J Vet Res; 62:1025–1030.
85. **Gasser, B.; Boman, B.; Wyder, D.; et al., (1990):** Stiffness characteristics of the circular Ilizarov device as opposed to conventional external fixators. J Biomech Eng; 112:15–21.
86. **Podolsky, A. and Chao, E.Y. (1993):** Mechanical performance of Ilizarov circular external fixators in comparison with other external fixators. Clin Orthop; 293:61–70.
87. **Sarpel, Y.; Gulsen, M.; Togrul, E.; et al., (2005):** Comparison of mechanical performance among different frame configurations of the Ilizarov external fixator: experimental study. J Trauma; 58:546–552.
88. **Antoci, V.; Roberts, C.S.; Antoci, V.; Jr, et al., (2005):** The effect of transfixion wire number and spacing between two levels of

- fixation on the stiffness of proximal tibial external fixation. *J Orthop Trauma*; 19:180–186.
89. **Yang, L.; Nayagam, S. and Saleh, M. (2003):** Stiffness characteristics and inter-fragmentary displacements with different hybrid external fixators. *Clin Biomech*; 18:166–172.
 90. **Ilizarov, G.A. (1992):** The Apparatus: components and biomechanical principles of application. In: Green S (Ed) *Transosseus osteosynthesis. Theoretical and clinical aspects of the regeneration and growth of tissue*. Springer-Verlag Berlin, pp 63–136.
 91. **Roberts, C.S.; Antoci, V.; Antoci, V.; Jr, et al., (2005):** The effect of transfixion wire crossing angle on the stiffness of fine wire external fixation: a biomechanical study. *Injury*; 36(9):1107–1112.
 92. **Ilizarov, G.A.; Emilyanova, H.S.; Lebedev, B.E.; et al., (1972):** Some experimental studies. Mechanical characteristics of Kirshner wires. In: *Perosseus compression and distraction osteosynthesis in traumatology and orthopedics*. Kurgan, pp 34–47.
 93. **Piza, G.; Caja, V.L.; Gonzalez-Viejo, M.A.; et al., (2004):** Hydroxyapatite-coated external-fixation pins. The effect on pin loosening and pin-tract infection in lengthening for short stature. *J Bone Jt Surg Br*; 86:892–897.
 94. **Moroni, A.; Faldini, C.; Pegreff, F.; et al., (2002):** Fixation strength of tapered versus bicylindrical hydroxyapatite-coated external fixation pins: an animal study. *J Biomed Mater Res*, 63:61–64.
 95. **Austin, T.; Fragomen and S. Robert Rozbruch (2007):** The Mechanics of External Fixation. *HSS J*. February; 3(1): 13–29.

96. **Sarmiento, A.; Schaeffer, J.F.; Beckerman, L.; et al., (1979):** Fracture healing in rat femora as affected by functional weight-bearing. *J Bone Jt Surg Am*;59:369–375.
97. **Klein, P.; Schell, H.; Streitparth, F.; et al., (2003):** The initial phase of fracture healing is specifically sensitive to mechanical conditions. *J Orthop Res*; 21:662–669.
98. **Kenwright, J.; Richardson, J.B.; Cunningham, J.L.; et al., (1991):** Axial movement and tibial fractures. A controlled randomized trial of treatment. *J Bone Joint Surg Br*; 73:654–659.
99. **Aarnes, G.T.; Steen, H.; Ludvigsen, P.; et al., (2005):** In vivo assessment of regenerate axial stiffness in distraction osteogenesis. *J Orthop Res*; 23:494–498.
100. **Ferreira, R.C. and Costa, M.T. (2009):** Recurrent ClubfootçApproach and Treatment with External Fixation .*Foot Ankle Clin N Am* 14,435–445.
101. **Grill, F. and Franke, J. (1987):** The Ilizarov distractor for the correction of relapsed or neglected clubfoot. *J Bone Joint Surg.* 69B:593-597.
102. **Paley, D. (1993):** The correction of complex foot deformities using Ilizarov's distraction osteotomies. *Clin Orthop*, in press.
103. **Umhanov, H.A.** Method of apparatus correction in orthopedic treatment of children with the cerebral spastic palsies.
104. **Paly, D. (1994):** Principles of foot deformity correction: Ilizarov technique. In, Gould JS (ed) :*Operative foot*. Philadelphia, WB Saunders.;p476-514.
105. **Grant, A.D.; Atar, D. and Lehman, W.B. (1992):** The Ilizarov technique in correction of complex foot deformities. *Clin Orthop*.;94:103.

106. **Ilizarov, G.A.; Shevtson, V.I.; et al., (1983):** Foot form shaping and lengthening methods. *Orthop travmatol protz.*1:49-51.
107. **Paly, D. (1994):** Principles of foot deformity correction :Ilizarov technique. In, Gould JS (ed):*Operative foot.* Philadelphia, WB Saunders. p476-514.
108. **Lehman, W.B.; Paley, D. and Atar, O. (1989):** Operating Room Guide to Cross Sectional Anatomy of the Extremities and Pelvis. New York. Raven Press.
109. **Choi, I.H.; Yang, M.S.; Chung, C.Y.; et al., (2001):** The treatment of recurrent arthrogryptic club foot in children by the Ilizarov method. *J Bone Joint Surg Br*;83-B:731–7.
110. **Ferreira, R.C.; Costa, M.T.; Frizzo, G.G.; et al., (2007):** Correction of severe recurrent clubfoot using a simplified setting of the Ilizarov device. *Foot Ankle Int*;28:557–68.
111. **Ferreira, R.C.; Stefani, K.C.; Fonseca, Filho F.F.; et al., (1999):** Correcao do pe torto congenito inveterado e recidivado pelo metodo de Ilizarov. *Rev Bras Ortop*;34:505–12 [in Portuguese].
112. **Ignacio, V.; Ponseti Jose, A. Morcuend, et al., (2009):** Clubfoot:Ponseti Management.Third Edition.;p24-25.
113. **Sodergard, J. and Ryoppy, S. (1994):** Foot deformities in arthrogryposis multiplex congenita. *J Pediatr Orthop*; 14:768–72.
114. **Choi, I.H.; Yang, M.S.; Chung, C.Y.; Cho, T.J. and Sohn, Y.J. (2001):** The treatment of recurrent arthrogryptic club foot in children by the Ilizarov method: a preliminary report. *J Bone Joint Surg [Br]*; 83-B:731– 7.
115. **Thompson, D.M. and Calhoun, J.H. (2000):** Advanced techniques in foot and ankle reconstruction. *Foot Ankle Clin*; 5 : 417-442.

116. **Neufeld, S.K. and Myerson, M.S. (2001):** Complications of surgical treatments for adults flatfoot deformities. *Foot Ankle Clin*; 6 : 179-191.
117. **Steinwender, G.; Saraph, V.; Zwick, E.B.; Uitz, C. and Linhart, W. (2001):** Complex foot deformities associated with soft-tissue scarring in children. *J Foot Ankle Surg*; 40 : 42-49.
118. **Henrik, W.; Goran, H. and Bjorn, T. (1996):** Correction of persistent clubfoot deformities with the Ilizarov external fixator. *Acta Orthop Scand*; 67 : 283-287.
119. **Choi, I.H.; Yang, M.S.; Chung, C.Y. et al., (2001):** The treatment of recurrent arthrogryptic club foot in children by the Ilizarov method. *J Bone Joint Surg*; 83-B : 731-737.
120. **Hosny, G.A. (2002):** Correction of foot deformities by the Ilizarov method without corrective osteotomies or soft tissue release. *J Pediatr Orthop*; 11-B : 121-128.
121. **Lejman, T.; Sulko, J. and Kowalczyk, B. (1998):** Treatment for recurrent clubfoot with the Ilizarov technique. *Chir Narzadow Ruehu Ortop Pol*; 63: 39-45.
122. **Herzenberg, JE, Paley D. (1992):** Ilizarov applications in foot and ankle surgery. *Adv Orthop Surg*; 16 : 162-174.
123. **Paley, D. (1993):** The correction of complex foot deformities using Ilizarov's distraction osteotomies. *Clin Orthop*; 11 : 238-240.
124. **Kucukkaya, M.; Kabukcuoglu, Y. and Kuzgun, U. (2002):** Management of the neuromuscular foot deformity with the Ilizarov method. *Foot Ankle Int*; 23 : 135-141.
125. **Bradish, C.F. and Noor, S. (2000):** The Ilizarov method in the management of relapsed club feet. *J Bone Joint Surg Br*; 82:387–91.

126. **Kocaoglu, M.; Eralp, L.; Atalar, A. and Bilen, F.E. (2002):** Correction of complex foot deformities using the Ilizarov external fixator. *J Foot Ankle Surg*; 41(1):30–9.
127. **Grant, A. and Lehman, W. (1991):** Clubfoot correction using the Ilizarov technique. *Bull Hosp Jt Dis Orthop Inst*;51(1):84– 7.
128. **Weber, M.; Schwer, H.; Zilkens, K.W. and Siebert, C.H. (2002):** Tibio-calcaneo-naviculo-cuboidale arthrodesis: 6 patients followed for 1 – 8 Years. *Acta Orthop Scand*;73(1):98–103.
129. **de la Huerta F. (1994):** Correction of the neglected clubfoot by the Ilizarov method. *Clin Orthop*; 301:89– 93.
130. **Wallander, H.; Hansson, G. and Tjernstrom, B. (1996):** Correction of persistent clubfoot deformities with the Ilizarov external fixator. *Acta Orthop Scand*; 67(3):283 –7.