

References

- Achenbach T. (1991):*** Manual for the child behavior checklist/4—18 and 1991 profile. Burlington: Department of Psychiatry, University of Vermont; 1991.
- Adam M. (2005):*** Epilepsy Patients More Likely to Exhibit Bipolar Disorder Symptoms. Br J Psychiatry; 109:95–150.
- Adrian J. Harwood and Galila Agam (2003):*** Search for a common mechanism of mood stabilizers. Biochemical Pharmacology; 66: 179–189.
- Alan B. and Charles E. (2007):*** Use of Antiepileptic Drugs for Nonepileptic Conditions: Psychiatric Disorders and Chronic Pain. Neurotherapeutics Volume 4, Issue 1, Pages 75-83.
- Aldenkamp AP, Krom MD, Reijs R. (2003):*** Newer antiepileptic drugs and cognitive issues. Epilepsia; 44(suppl4):21–29.
- Aldenkamp A.P. and Arends J. (2004):*** Effects of epileptiform EEG discharges on cognitive function: Is the concept of “transient cognitive impairment” still valid, Epilepsy Behav **5** (suppl 1), pp. S25–S34.
- Allen Hauser W and Nina Banerjee P. (2008):*** Epidemiology of Epilepsy in Children pediatric epilepsy 3rd edition diagnosis and therapy text book p 156.
- Alvestad S, Lydersen S, Brodtkorb, et al. (2007):*** Rash from antiepileptic drugs: influence by gender, age, and learning disability. Epilepsia; 48: 1360–5.
- Amann B. and Grunze H. (2005):*** Neurochemical underpinnings in bipolar disorder and epilepsy.

Epilepsia; **46** (suppl 4): pp. 26–30.

Anna F, David M, Benedikt A, et al. (2005): Antiepileptic Drugs in Mood Disordered Patients. *Epilepsia*, 46(Suppl.4): 38–44.

Arts WF, Geerts AT, Brouwer OF, et al. (1999): The early prognosis of epilepsy in childhood: the prediction of poor outcome. The Dutch study of epilepsy in childhood. *Epilepsia*; 40:726–734.

Austin JK, Risinger MW, Beckett LA. (1992): Correlates of behavior problems in children with epilepsy. *Epilepsia*; 33:1115–22.

Austin JK, Shelton Smith M, Risinger MW and McNelis AM (1994): Childhood epilepsy and asthma: Comparaison of quality of life. *Epilepsia* 35(3):608–615.

Austin JK, Huster GA, Dunn DW, Risinger MW. (1996): Adolescents with active or inactive epilepsy or asthma: a comparison of quality of life. *Epilepsia*; 37: 1228–38.

Austin JK, J. Harezlak and D.W. Dunn, et al. (2001): Behavior problems in children before first recognized seizures, *Pediatrics* **107**, pp. 115–122.

Austin JK, Mc Nelis A.M. and Shore C.P., et al. (2002): A feasibility study of a family seizure management program: ‘Be seizure smart’, *J Neurosci Nurs* (**34**), pp. 30–37.

Auvin S, Pandit F, Motte J, et al. (2006): Benign Myoclonic Epilepsy in Infants: Electroclinical Features and Long-term Follow-up. *Epilepsia*; 47(2).

- Azam M, Bhatti N, Shahab N (2008):** "Piracetam in severe breath holding spells". *Int J Psychiatry Med* 38 (2): 195–201.
- Bahar E, Arif V, et al.(2008):** Use of lamotrigine in psychiatric disorders. *Anadolu Psikiyatri Dergisi*; 9:169-178
- Ballenger JC, Post RM. (1980):** Carbamazepine in manic – depressive illness: a new treatment. *Am J Psychiatry*; 137(7): 782–90.
- Bauer M. and Pfennig A, (2005):** Epidemiology of bipolar disorders, *Epilepsia* **46** (suppl 4), pp. 8–13.
- Bauer J, Blumenthal S, Reuber M, et al. (2004):** Epilepsy syndrome, focus location, and treatment affect testicular function in men with epilepsy. *Neurology*; 62:243–6.
- Beck B. (2000):** The classification of pediatric epilepsy syndromes. *Clin Nurs Pract Epilepsy*; 1:8-12.
- Becker K, Holtmann M. (2006):** Role of electroencephalography inattention-deficit hyperactivity disorder. *Expert Rev Neurother*; 6 (5): 731–9.
- Begley CE, Famulari M, Annegers JF, et al. (2004):** The cost of epilepsy in the United States: an estimate from population-based clinical and survey data. *Epilepsia*; 41:342-351.
- Beleza P. (2009):** Refractory epilepsy: a clinically oriented review. *Eur Neurol*; 62(2):65- 71.
- Benbadis S (2009):** The differential diagnosis of epilepsy: A critical review *Epilepsy & Behavior*; 15:15 – 21.

- Benbadis SR. (2005):*** A spell in the epilepsy clinic and a history of “chronic pain” or “fibromyalgia” ’independently predict a diagnosis of psychogenic seizures. *Epilepsy Behav*; 6:264–5.
- Benbadis SR, Chichkova R.(2006):*** Psychogenic pseudosyncope: an underestimated and provable diagnosis. *Epilepsy Behav*; 9:106–10.
- Benedikt Amann and Heinz Grunze (2005):*** Neurochemical Underpinnings in Bipolar Disorder and Epilepsy. *Epilepsia*, 46 (Suppl.4): 26 – 30.
- Berg AT, Shinnar S, Levy SR, et al. (2001):*** Defining early seizure outcomes in pediatric epilepsy: the good, the bad and the in-between. *Epilepsy Res* 43:75–84.
- Berg AT (2003):*** Of cabbages and kings: perspectives on classification from the field of systematics. *Epilepsia*; 44(1):8–12.
- Berg AT, Langfitt J, Shinnar S, et al (2003):*** How long does it take for partial epilepsy to become intractable? *Neurology* 60:186–190.
- Berg AT, Smith SN, Frobish D, et al. (2005):*** Special education needs of children with newly diagnosed epilepsy. *Dev Med Child Neurol.*; 47:749–753.
- Berg AT (2006):*** Defining intractable epilepsy. *Adv Neurol* 97:5–10.
- Berg, AT, Vickrey BG, Testa FM, et al. (2006):*** How long does it take for epilepsy to become intractable? A prospective investigation. *Ann Neurol*; 60:73.

- Berg AT, Vickery BG, Testa FM, et al. (2007):** Behavior and social competency in idiopathic and cryptogenic childhood epilepsy. *Dev Med Child Neurol*; 49: 487–92.
- Bergqvist, AG, Schall, JI, Gallagher PR, et al. (2005):** Fasting versus gradual initiation of the ketogenic diet: a prospective, randomized clinical trial of efficacy. *Epilepsia*; 46:1810.
- Berroya AG, McIntyre J, Webster R, et al. (2004):** Speech and language deterioration in benign rolandic epilepsy. *J Child Neurol*. 2004; 19: 53-58.
- Bersudsky Y. (2006):** Phenytoin: an anti – bipolar anticonvulsant? *Int J Neuropsychopharmacol*; 9 (4): 429–84.
- Besag FM.(2002):** When is it inappropriate to prescribe psychotropic medication?. *Epilepsia*; 43 Suppl 2:45-50.
- Besag F.M. (2004):**Behavioral aspects of pediatric epilepsy syndromes, *Epilepsy Behav* 5(suppl 1), pp. S3–S13.
- Besag F.M., Nomayo A. and Pool F. (2005):** The reactions of parents who think that a child is dying in a seizure—In their own words, *Epilepsy Behav* 7, pp. 517–523.
- Beydoun A, Kutluay E. (2003):** Conversion to monotherapy: clinical trials in patients with refractory epilepsy. *Neurology*; 60:S13–25.
- Bhattacharyya M, Kalra, V, Gulati, S. (2006):** Intranasal midazolam vs rectal diazepam in acute childhood seizures. *Pediatr Neurol*; 34:355.

- Blume WT:** Focal seizures: intractability and semiology. *Adv Neurol* 2006; 97:17-25.
- Bialer M, Johannessen SJ, Kupferberg HJ, et al., (2007):** Progress report on new antiepileptic drugs: a summary of the Eighth Eilat Conference (EILAT VIII). *Epilepsy Res* 73:1–52.
- Binnie CD. (1993):** Significance and management of transitory cognitive impairment due to subclinical EEG discharges in children. *Brain Dev*; 15(1):23–30.
- Biraben A, Taussig D, Thomas P, et al. (2001):** Fear as the main feature of epileptic seizures. *J Neurol Neurosurg Psychiatry*; 70:186–91.
- Bishop M. and Boag EM. (2006):** Teachers' knowledge about epilepsy and attitudes toward students with epilepsy: Results of a national survey, *Epilepsy Behav* 8, pp. 397–405.
- Bjornaes H, Stabell K, Henriksen O, Loyning Y. (2001):** The effects of refractory epilepsy on intellectual functioning in children and adults A longitudinal study. *Seizure.* ; 10 : 250–259.
- Blume W, Lüders H, Mizrahi E, et al (2001).**"Glossary of descriptive terminology for ictal semiology: report of the ILAE task force on classification and terminology". *Epilepsia* 42 (9): 1212–8
- Blanco C, Schneier FR, Schmidt A, et al. (2003):** Pharmacological Treatment of social anxiety disorder: ameta-analysis. *Depress Anxiety*; 18: 29–40.
- Blumer D, Montouris G, Davies K .(2004):** The interictal dysphoric disorder: recognition, pathogenesis, and treatment of the

major psychiatric disorder of epilepsy. *Epilepsy Behav*; 5(6):836–40.

Bonilha L, Rorden C, Castellano G, et al., (2005): Voxel-based morphometry of the thalamus in patients with refractory medial temporal lobe epilepsy. *Neuroimage*; 25:1016 – 1021.

Bortolato M, Solbrig MV. (2007): The price of seizure control: dynorphins in interictal and postictal psychosis. *Psychiatry Res*; 151(1–2):139–43.

Bourgeois BF. (1998): Antiepileptic drugs, learning, and behavior in childhood epilepsy. *Epilepsia*; 39: 913–21.

Bough KJ and Rho JM. (2007): Anticonvulsant mechanisms of the ketogenic diet. *Epilepsia*; 48:43.

Boylan LS, FlintL A, Labovitz DL, et al. (2004): Depression but not seizure frequency predicts quality of life in treatment-resistant epilepsy. *Neurology*; 62(2):258-61.

Britton JW, (2004): Syncope and seizures - Differential diagnosis and evaluation. *Clin. Auton Res*; 14:148-159.

Brenda P. (2007): Unraveling Epileptogenesis. *Epilepsy Curr*; 14: 6–11.

Briellmann RS, Hopwood MJ, Jackson GD. (2007): Major depression in temporal lobe epilepsy with hippocampal sclerosis: clinical and imaging correlates. *J Neurol Neurosurg Psychiatry* Jan26

Bromfield EB, Altshuler L, Leideran DB, et al. (1992): Cerebral metabolism and depression in patients with complex partial seizures. *Arch Neurol*; 49(6):617–23.

-
- Caffrey HM, Austin J.K. and Dunn DW., et al. (2003):** A prospective study of teachers' ratings of behavior problems in children with new-onset seizures, *Epilepsy Behav* (4), pp.26–35.
- Camfield and Camfield C. (2005):** The frequency of intractable seizures after stopping AEDs in seizure-free children with epilepsy. *Neurology*; 64:973.
- Caplan R, Arbelle S, Guthrie D, et al. (1997):** Formal thought disorder and psychopathology in pediatric primary generalized and complex partial epilepsy. *Am Acad Child Adolesc Psychiatry*; 36: 1286-1294.
- Caplan R, Austin JK, Dunn D., et al. (2002):** Development of a self-efficacy scale for children and adolescents with epilepsy, *Child Health Care* (31), pp. 295–309.
- Caplan R, Austin JK, Dunn D. (2005):** Depression and anxiety disorders in pediatric epilepsy. *Epilepsia* 46: 720–730.
- Caplan R, Sagun J and Siddarth P., et al. (2005):** Social competence in pediatric epilepsy: Insights into underlying mechanisms, *Epilepsy Behav* 6, pp. 218–228.
- Caplan R, Siddarth P, Stahl L, et al. (2008):** Childhood absence epilepsy: behavioral, cognitive, and linguistic comorbidities. *Epilepsia*. Nov; 49 (11):1838-46.
- Cendes f, (2004):** febrile seizures and mesial temporal sclerosis. *Curr Opin Neurol*; 17: 161-4.
- Cendes F. (2005):** Progressive hippocampal and extrahippocampal atrophy in drug resistant epilepsy. *Curr Opin Neurol.*; 18: 173–177.

Chang BS, et al. (2003): “Epilepsy,” *New England Journal of Medicine*: Vol. 349, No. 13, pp. 1257–66.

Cheetham SC, Crompton MR, Czudek C, et al. (1989): Serotonin concentration and turnover in brains of depressed suicides. *Brain Res*; 502: 332 - 340.

Chen S, Melo M, Benbadis SR. (2008): The differential diagnosis of seizures in the ICU: a video–EEG study. Presented at the 62nd Annual Meeting of the American Epilepsy Society, Seattle, WA, USA.

Commission on Classification and Terminology of the International League Against Epilepsy (1981): Proposal for revised clinical and electroencephalographic classification of epileptic seizures. *Epilepsia*; 22: 489–501.

Commission on Classification and Terminology of the International League Against Epilepsy. (1989): Proposal for revised classification of epilepsies and epileptic syndromes. *Epilepsia*; 30:389–399.

Commission on Neuroimaging of the International League Against Epilepsy. (1997): Recommendations for neuroimaging of patients with epilepsy. *Epilepsia*; 38: 1255-6.

Chun SW, Yuelian S, Mogens V, et al., (2008): Preeclampsia and Risk for Epilepsy in Offspring. *PEDIATRICS* Vol. 122 No. 5, pp. 1072-1078.

- Cormack F, Gadian DG, Vargha-Khadem F, et al., (2005):** Extra-hippocampal grey matter density abnormalities in paediatric mesial temporal sclerosis. *Neuroimage*; 27: 635 – 643.
- Cortez MA, Perez Velazquez JL, Snead OC., (2006):** 3rd Animal models of epilepsy and progressive effects of seizures. *Adv Neurol.*; 97: 293–304.
- Cowan LD. (2002):** The epidemiology of the epilepsies in children. *Ment Retard Dev Disabil Res Rev*; 8: 171–8.
- AI Dai, M Wasay. (2007):** Idiopathic epilepsy of childhood and potassium ion channels. *J Pak Med Assoc* 57: 415-8.
- Dang-Vu TT, Desseilles M, Petit D, et al.(2007):** Neuroimaging in sleep medicine. *Sleep Med*; 8: 349-372.
- Datta SS, Premkumar TS, Chandy S, et al. (2005):** Behaviour problems in children and adolescents with seizure disorder: associations and risk factors. *Seizure*; 14: 190–97.
- Davies S, Heyman I, Goodman R. (2003):** A population survey of mental health problems in children with epilepsy. *Dev Med Child Neurol*; 45:292–5.
- Dean P. (2000):** The treatment of pediatric epilepsies. *Clin Nurs Pract Epilepsy*.1:1-7.
- Dean P. (2003):** Epilepsy surgery in the first three years of life. Paper presented at: Annual Meeting of the American Association of Neuroscience Nurses; April 5-8, Atlanta, Ga.
- Deary IJ, Whiteman MC, Starr JM, et al., (2004):** The impact of childhood intelligence on later life: following up the Scottish

mental surveys of 1932 and 1947. J Pers Soc Psychol.; 86: 130–147.

De Cabo-de la Vega C, Villanueva-Hernández P, Prieto-Martín A. (2006): The neurochemistry of epilepsy, inhibitory neurotransmission and experimental models: new perspectives. Rev Neurol. 2006 Feb 1-15; 42 (3) : 159-68.

Dieter S and Wolfgang L (2005): Drug Resistance in Epilepsy: Putative Neurobiological and Clinical Mechanisms. Epilepsia; 46 (6).

Dieter Schmidt (2009): Drug treatment of epilepsy: Options and limitations. Epilepsy&Behavior15(2009)56–65.

Depiero, A.D. and Teach S.J. (2001): Febrile Seizures. Pediatric Emergency Care; 17(5): p. 384-387.

Dlugos DJ, Samuel MD, et al. (2001): Response to first drug trial predicts outcome in childhood temporal lobe epilepsy. Neurology; 57: 2259 – 2264.

Dodel R, Peter H, Spottke A, et al. (2007): Health-related quality of life in patients with narcolepsy. Sleep Med. Nov;8 (7-8):733-41.

Donner EJ, Snead OC III (2006): New drugs in the treatment of epilepsy in children. Curr Probl Pediatr Adolesc Health Care 35: 398–419.

Dow C, Oyegbile TO, Jones J, et al. (2004): The nature and course of neuropsychological morbidity in chronic temporal lobe epilepsy. Neurology; 62: 1736–1742.

- Drevets W.C. (2003):*** Neuroimaging abnormalities in the amygdala in mood disorders, *Ann N Y Acad Sci* **985**, pp. 420–444.
- Drane DL, Meador KJ. (2002):*** Cognitive and behavioral effects of antiepileptic drugs. *Epilepsy Behav*; 3 (5S): S49–53.
- Duchowny M.(2004):*** Hemispherectomy for epilepsy: when is one half better than two?. *Neurology* 2004; 62:1664.
- Duncan JS, Sander JW, et al. (2006):*** Adult epilepsy. *Lancet* 2006;367: 1087 – 100.
- Dunn DW, Austin JK. (1999):*** Behavioral issues in pediatric epilepsy. *Neurology*; 53(5, Suppl. 2):S96–S100.
- Dunn DW, Austin JK, Caffrey HM, Perkins SM. (2003):*** A prospective study of teachers ratings of behavior problems in children with new-onset seizures. *Epilepsy Behav.* 2003;4(1):26-35.
- Dunn DW, et al. (2004):*** “Differential Diagnosis and Treatment of Psychiatric Disorders in Children and Adolescents with Epilepsy,” *Epilepsy and Behavior: Vol. 5, Suppl. 3*, pp. S10–S17
- Dunn DW, Caffrey HM, Austin JK, et al. (2002):*** Recurrent seizures and behavior problems in children with first recognized seizures: a prospective study. *Epilepsia* ;43 : 1564–73.
- Dunn DW, Austin JK, Harezlak J, et al. (2003):*** ADHD and epilepsy in childhood. *Dev Med Child Neurol* ; 45: 50-54.
- Dunn DW, Austin JK and Huster GA. (1999):*** Symptoms of depression in adolescents with epilepsy, *J Am Acad Child Adolesc Psychiatry* (**38**), pp. 1132–1138.

- Dunn DW, Austin JK, Perkins SM. (2009):** Prevalence of psychopathology in childhood epilepsy: categorical and dimensional measures. *Dev Med Child Neurol.* May; 51(5):364-72.
- Dutta SR, Hazarika I, Chakravarty BP (2007):** "Abdominal epilepsy, an uncommon cause of recurrent abdominal pain: a brief report". *Gut* **56** (3): 439–41.
- Dutta SR, Limdi NA, Knowlton RK, et al., (2007):** Safety of rapid intravenous loading of valproate. *Epilepsia* 48:478–483.
- Dworkin RH, Backonja M, Rowbotham MC, et al.,(2003):** Advances in neuropathic pain: diagnosis, mechanisms, and treatment recommendations. *Arch Neurol.*; 60(11):1524-34.
- Edoardo S and Giulio P (2004):** Antiepileptic drugs: indications other than epilepsy. *Epileptic Disorders.* Volume 6, Numéro 2, 57-75.
- Edwards KR, Sackellares JC, Vuong A, et al. (2001):** Lamotrigine monotherapy improves depressive symptoms in epilepsy: adouble –blind comparison with valproate. *Epilepsy Behav*; 2 (1): 28–36.
- Elger CE, Helmstaedter C, Kurthen M. (2004):** Chronic epilepsy and cognition. *Lancet Neurol.*; 3:663–672.
- Elger G, Hoppe C, Falkai P, et al. (2000):** Vagus nerve stimulation is associated with mood improvements in epilepsy patients. *Epilepsy Res*; 42(2–3):203–10.
- Elger C.E. (2002):** Epilepsy: disease and model to study human brain function. *Brain Pathol*; 12: 193-198.

- Elger CE. (2008):*** Modern management of epilepsy: A practical approach. *Epilepsy and Behavior*; 12: 501.
- Elger CE and Schmidt D (2008):*** Modern management of epilepsy: A practical approach. *Epilepsy& Behavior*; 12: 501–539.
- Engel J. (1996):*** Surgery for seizures. *N Engl J Med.*; 334: 647–652.
- Engel J Jr. (2003):*** Reply to “Of cabbages and kings: some considerations on classifications, diagnostic schemes, semiology and concepts.” *Epilepsia*; 44(1):4–6.
- Engel JrJ, Wiebe S, French J, et al. (2003):*** Practice parameter: temporal lobe and localized neocortical resections for epilepsy. Report of the Quality Standards Subcommittee of the American Academy of Neurology in Association with the American Epilepsy Society and the American Association of Neurological Surgeons. *Neurology*; 60: 538–47.
- Engel J Jr. (2006):*** Report of the ILAE classification core group. *Epilepsia* 2006; 47(9):1558–1568.
- Engel J Jr, Pedley T. (2007):*** editors. *Epilepsy: a comprehensive textbook*, 2nd.Vols.1–3. Philadelphia: Lippincott Williams & Wilkins.
- Ettinger AB. (2006):*** Psychotropic effects of antiepileptic drugs. *Neurology*; 67 (11): 1916–25.
- Ettinger A.B, Reed M.L. and Goldberg J.F, et al. (2005):*** Prevalence of bipolar symptoms in epilepsy vs other chronic health disorders, *Neurology* (65), pp. 535–540.
- Ettinger AB and Argoff CE. (2007):*** Use of antiepileptic drugs for nonepileptic conditions: psychiatric disorders and chronic pain. *Neurotherapeutics*. 2007 Jan; 4(1):75-83.

- Fakhoury TA, Barry JJ, Mitchell Miller J, et al. (2007):** Lamotrigine in patients with epilepsy and co-morbid depressive symptoms. *Epilepsy Behav*; 10(1):155–62.
- Farrell K, Wirrell E, Whiting S (2006):** The definition and prediction of intractable epilepsy in children. *Adv Neurol* 97:435–442.
- Faught E. (2004):** Treatment of refractory primary generalized epilepsy. *Rev Neurol Dis* 2004; 1(suppl 1):S34-S43.
- Fisher R, van Emde Boas W, Blume W, et al (2005):** "Epileptic seizures and epilepsy: definitions proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE)". *Epilepsia* **46** (4): 470–2.
- Fleminger S, Greenwood RJ, Oliver DL. (2006):** Pharmacological management for agitation and aggression in people with acquired brain injury. *Cochrane Database Syst Rev*;4: CD003299.
- Frankel EA, Shalaby TM, Orenstein SR. (2006):** Sandifer syndrome posturing: relation to abdominal wall contractions, gastro-esophageal reflux, and fundoplication. *Dig Dis Sci*; 51: 635–40.
- Frederick Goodwin, Kay Jamison (2007):** "Manic-Depressive Illness: Bipolar Disorders and Recurrent Depression, 2nd Edition;"
- Freeman JM, Kossoff EH, Hartman, AL. (2007):** The ketogenic diet: one decade later. *Pediatrics*; 119:535.
- Friedlander WJ (2000):** The rise and fall of bromide therapy in epilepsy. *Arch Neurol* 57:1782–1785.
- Freilinger M, Reisel B, Reiter E, et al. (2006):** Behavioral and emotional problems in children with epilepsy. *J Child Neurol* 2006; 21: 939–45.

- French JA, Kanner AM, Bautista J, et al. (2004):*** Efficacy and tolerability of the new antiepileptic drugs: I. Treatment of new onset epilepsy .Report of the TTA and QSS Subcommittees of the American Academy of Neurology and the American Epilepsy Society. *Epilepsia*; 45: 401 – 9.
- French JA. (2007):*** Treatment with Antiepileptic Drugs, New and Old. *Continuum Neurol*; 13(4): 71-90.
- Frye M.A, K.D. Denicoff and R.M. Post, et al. (1998):*** A history of the use of anticonvulsants as mood stabilizers in the last two decades of the 20th century, *Neuropsychobiology* **38** , pp. 152–166.
- Gaitatzis A, et al. (2004):*** The Psychiatric Comorbidity of Epilepsy,” *Acta: Neurologic Scandinavica*: Vol. 110, No. 4, pp. 207–20.
- Geerts AT, Niermeijer JM, Peters AC, et al. (2005):*** Four-year outcome after early withdrawal of antiepileptic drugs in childhood epilepsy. *Neurology* 2005; 64:2136.
- Gelisse P. and Genton P. (2008):*** "Cough syncope misinterpreted as epileptic seizure." *Epileptic Disord* 10(3): 223-4.
- Gilliam FG. (2003):*** The impact of epilepsy on subjective health status. *Curr Neurol Neurosci Rep*; 3: 357–62.
- Gonzalez-Heydrich J, Dodds A, Whitney J, et al.(2007):*** Psychiatric disorders and behavioral Characteristics of pediatric patients with both epilepsy and attention-deficit hyperactivity disorder. *Epilepsy Behav*; 10 (3): 384–8.

- Grabowska-Grzyb A, Jedrzejczak J, Naganska E, et al. (2006):** Risk factors for depression in patients with epilepsy. *Epilepsy Behav*; 8(2): 411–7.
- Greenlee BA, Ferrell RB, Kauffman CI, et al. (2003):** Complex Partial seizures and depression. *Curr Psychiatry Rep.*; 5: 410 – 416.
- Gregory L H & Yezekiel BA (2003):** Seizing hold of seizures. *Nature Medicine*; 9, 994 - 996.
- Gucuyener K, Erdemoglu AK, Senol S, et al. (2003):** Use of methylphenidate for attention-deficit-hyperactivity-disorder in patients with epilepsy or electroencephalographic abnormalities. *J Child Neurol* 2003; 18: 109-112.
- Gutierrez-Delicado E, Serratosa JM (2004):** Genetics of the epilepsies. *Curr Opin Neurol*; 17, 147–53.
- Gupta M, Aneja S, Kohli K. (2004):** Add-on melatonin improves quality of life in epileptic children on valproate monotherapy: a randomized, double-blind, placebo-controlled trial. *Epilepsy Behav*; 5(3):316-21.
- Hala sakr. (2003):** Out of the shadows al ahram weekly online 2003 issue no. 629.
- Harden CL, Goldstein MA. (2002):** Mood disorders in patients with epilepsy: epidemiology and management. *CNS Drugs*; 16: 91-3302.
- Hartman AL, Vining EP. (2007):** Clinical aspects of the ketogenic diet. *Epilepsia* 2007; 48:31.
- Harwood AJ, Agam G. (2003):** Search for a common mechanism of mood stabilizers. *Biochem Pharmacol.* 15;66(2):179-89.
- Hartong EG, Moleman P, Hoogduin CA, et al. (2003):** Prophylactic efficacy of lithium versus carbamazepine in treatment bipolar patients. *J Clin Psychiatry*; 64(2):144–51.

- Hee Seo, Mock Lee, Soo Lee, et al. (2007):*** Efficacy and tolerability of the ketogenic diet according to lipid: non lipid ratios-comparison of 3:1 with 4:1 diet. *Epilepsia* 2007; 48:801.
- Helmstaedter C. (2002):*** Effects of chronic epilepsy on declarative memory systems. *Prog Brain Res.*; 135:439–453.
- Hemmer SA, Pasternak JF, Zecker SG, et al. (2001):*** Stimulant therapy and seizure risk in children with ADHD. *Pediatr Neurol* 2001; 24(2):99-102.
- Helen E. (2007):*** The neurobiology of Epilepsy. *Current Neurology and Neuroscience Reports* 2007, 7:348–354.
- Hermann BP, Whitman S, Hughes JR, et al. (1988):*** Multietiological determinants of psychopathology and social competence in children with epilepsy. *Epilepsy Res*; 2:51–60.
- Hermann B, Seidenberg M, Sears L, et al. (2004):*** Cerebellar atrophy in temporal lobe epilepsy affects procedural memory. *Neurology* 2004; 63 : 2129 – 31.
- Hermann BP and Jones JE. (2005):*** Depression in Epilepsy: What is the Extent of the Current Problem? In: *Mood Disorders in Epilepsy: Bridging the Gap Between Psychiatry and Neurology*. American Epilepsy Society and IntraMed. Scientific Solutions.
- Hermann B, Jones J, Sheth R, et al. (2006):*** Children with new-onset epilepsy: neuropsychological status and brain structure. *Brain* (2006), 129, 2609–2619.
- Hermann BP, Seidenberg M, Dow C, et al. (2006):*** Cognitive prognosis in chronic temporal lobe epilepsy. *Ann Neurol*;60:80 – 7.

- Herman ST. (2007):** Classification of Epileptic Seizures. Continuum Neurol. 2007; 13(4): 13-47.
- Hesdorffer DC, Hauser WA, Annegers JF, et al. (2000):** Major depression is a risk factor for seizures in older adults. Ann Neurol ; 47 (2): 246 – 9.
- Hesdorffer DC, Ludvigsson P, Olafsson E, et al. (2004):** ADHD as a risk factor for incident unprovoked seizures and epilepsy in children. Arch Gen Psychiatry; 61 (7):731–6.
- Hesdorffer DC, Hauser WA, Olafsson E, et al. (2006):** Depression and suicide attempt as risk factors for incident unprovoked seizures. Ann Neurol.; 59:35–41.
- Hesdorffer DC, Hauser WA, Olafsson E, et al. (2007):** Epilepsy and suicide risk. Lancet Neurol.; 6:847; author reply 848.
- Hitiris N, Mohanraj R, Norrie J, et al. (2007):** Predictors of pharmaco-resistant epilepsy. Epilepsy Res 2007; 75:192-196.
- Hirtz D, Thurman DJ, Gwinn-Hardy K, et al. (2007):** "How common are the 'common' neurologic disorders?". Neurology **68** (5): 326–37
- Holzer JC and Bear DM. (1997):** Psychiatric considerations in patients with epilepsy. San Diego, CA: Academic Press; p. 131-148.
- Holzhausen S.P., M.M. Guerreiro and C.E. Baccin et al. (2007):** Use of risperidone in children with epilepsy, *Epilepsy Behav* **10**, pp. 412–416.
- Hoare P. (1984):** The development of psychiatric disorder among school children with epilepsy. Dev Med Child Neurol; 26: 3—13.
- Hoare P, Kerley S.(1991):** Psychosocial adjustment of children with chronic epilepsy and their families. Dev Med Child Neurol 1991; 33 (3): 201—15.

- Holmes GL.** (1997): When the first drug does not work: treatment options. J Pediatr. 1997; 131:794-796.
- Homberg V, Kowalik A, Schulze-Bonhage A.** (2001): Ad-hoc Umstellung von Carbamazepin zu Oxcarbazepin. Effizienz und Vertraglichkeit. Eine retrospektive Analyse. Nervenarzt; 72:918–23.
- Hovinga C. (2003):** Developmental pharmacokinetics in epilepsy. Neurology; 57: 2259 – 2264.
- Hubert P, Parain D, Vallée L. (2009):** Management of convulsive status epilepticus in infants and children. Rev Neurol (Paris). 2009; 165(4): 390-7.
- Horrocks IA, Nechay A, Stephenson JB. (2005):** Anoxic-epileptic seizures: observational study of epileptic seizures induced by syncope. Arch Dis Child 2005; 90:1283–7.
- James W, Dave F, Alexis A, et al. (2007):** Treatment of pediatric epilepsy: European expert opinion, 2007. Epileptic Disord 2007; 9 (4) 353- 412.
- Jan MM, Zuberi SA, Alsaihati BA. (2009):** Pregabalin: preliminary experience in intractable childhood epilepsy. Pediatr Neurol. 2009 May; 40(5):347-50.
- Jefferys JG.(2003):** Models and mechanisms of experimental epilepsies. Epilepsia 2003; 44(Suppl.12):44–50.
- Jeremy J.M, Susan RR, Mark S (2009):** Lamotrigine–valproic acid combination therapy for medically refractory epilepsy. Epilepsia, 50 (3):475–479.
- Johnston MV. (2007):** Seizures in Childhood. In: Kliegman RM, Behrman RE, Jenson HB, Stanton BF, eds. Nelson Textbook of Pediatrics. 18th ed. Philadelphia, Pa: Saunders Elsevier; chap 593.

- Jonas R, Nguyen S, Hu B, et al. (2004):** Cerebral hemispherectomy: hospital course, seizure, developmental, language, and motor outcomes. *Neurology* 2004; 62:1712.
- Jokeit H, Ebner A. (2002):** Effects of chronic epilepsy on intellectual functions. *Prog Brain Res.*; 135:455–463.
- Jacoby A., Snape D. and Baker GA. (2005):** Epilepsy and social identity: The stigma of a chronic neurological disorder, *Lancet Neurol* 4, pp. 171–178.
- Jobe PC, Dailey JW, Wernicke JF. (1999):** A noradrenergic and serotonergic hypothesis of the linkage between epilepsy and affective disorders. *Crit Rev Neurobiol*; 13(4): 317–56.
- Jobe PC, Browning RA.(2005):** The serotonergic and noradrenergic effects of antidepressant drugs are anticonvulsant, not proconvulsant. *Epilepsy Behav*; 7: 02- 6619.
- Johnson EK, Jones JE, Seidenberg M, et al. (2004):** The relative impact of anxiety, depression, and clinical seizure features on health-related quality of life in epilepsy. *Epilepsia* ; 45(5): 544–50.
- Jones JA, et al.(2005):** “The Clinical Assessment of Axis I Psychiatric Morbidity in Chronic Epilepsy: A Multicenter Investigation,” *Journal of Neuropsychiatry and Clinical Neuroscience* : Vol. 17, No. 2, pp. 172–79.
- Jones JE, Hermann BP, Woodard JL, et al. (2005):** Screening for major depression in epilepsy with common self-report depression inventories. *Epilepsia* ; 46:731–735.
- Julie EG, Walke and Hannah RC (2006):** Levetiracetam: Antiepileptic Properties and Protective Effects on Mitochondrial Dysfunction in Experimental Status Epilepticus. **Kanazawa**

- O. (2000):** Shuddering attacks : report of four children. *Pediatr Neurol* 2000; 23:421–4.
- Kanazawa O, Tohyama J, Akasaka N, et al., (2005):** Magnetoencephalographic study of patients with Panayiotopoulos syndrome. *Epilepsia*; 46 : 1106-13. *Epilepsia* 2006;47(3).
- Kang HC, Chung da E, Kim DW. (2004):** Early- and late-onset complications of the ketogenic diet for intractable epilepsy. *Epilepsia* 2004; 45:1116.
- Kanner A, Palac S. (2000):** Depression in epilepsy: a common but often unrecognized Comorbid malady. *Epilepsy Behav*; 1:37-51.
- Kanner AM, Kozak AM, Frey M.(2000):** The use of sertraline in patients with epilepsy: is it safe? *Epilepsy Behav*; 1(2):100–5.
- Kanner AM. (2002):** Psychiatric comorbidity in patients with developmental disorders and epilepsy: A practical approach to its diagnosis and treatment. *Epilepsy Behav*; 3 (66S1): 7-13.
- Kanner AM. (2003):** Depression in epilepsy: prevalence, clinical semiology, pathogenic mechanisms, and treatment. *Biol Psychiatry*; 54(3):388–98.
- Kanner A.M. (2003):** Depression in epilepsy is much more than a reactive process, *Epilepsy Curr* (3), pp. 202–203.
- Kanner AM. (2006):** Depression and epilepsy: a new perspective on two closely related disorders. *Epilepsy Curr.*; 6: 141–146.
- Kanner AM, Dunn D. (2004):** Diagnosis and management of depression and psychosis in children and adolescents with epilepsy. *J Child Neurol*; 19 Suppl 1: S65-S72.

-
- Keene DL, Manion I, Whiting S, et al. (2005):** A survey of behavior problems in children with epilepsy. *Epilepsy Behav*; 6: 581–86.
- Kessler RC, Berglund P, Demler O, et al. (2003):** The epidemiology of major depressive disorder: results from the National Comorbidity Study Replication (NCS-R). *JAMA*; 289(23): 3095–105.
- Kessing LV, Nilsson FM, Bolwig TG. (2003):** On the increased risk of developing late-onset epilepsy for patients with major affective disorder. *J Affect Disord*; 76 (1–3):39–48.
- Kliegman RM and Greenbaum LA (2004):** Practical strategies in Pediatric Diagnosis and therapy, Philadelphia Elsevier; 2: 680.
- Kobau R, Gilliam F, Thurman DJ. (2006) :** Prevalence of self-reported epilepsy or seizure disorder and its associations with self-reported depression and anxiety: results from the 2004 Healthstyles Survey. *Epilepsia*; 47(11):1915–21.
- Koh S, Jensen FE. (2001):** Topiramate blocks perinatal hypoxia-induced seizures in rat pups. *Ann Neurol*; 50: 366–372.
- Koponen A., Seppala U., K. Eriksson, et al. (2007):** Social functioning and psychological well-being of 347 young adults with epilepsy only—Population-based, controlled study from Finland, *Epilepsia* (48), pp. 907–912.
- Kotagal P, Costa M, Wyllie E. (2002):** Paroxysmal nonepileptic events in children and adolescents. *Pediatrics* 2002; 110:e46.
- Kossoff EH, Laux LC, Blackford R, et al. (2008):** When do seizures usually improve with the ketogenic diet?. *Epilepsia* 2008; 49:329.

-
- Kossoff EH, Pyzik PL, McGrogan JR, et al. (2002):** Efficacy of the ketogenic diet for infantile spasms. *Pediatrics*; 109 : 780-783.
- Kossoff EH, Pyzik PL, Furth SL, et al. (2000):** Kidney stones, carbonic anhydrase inhibitors, and the ketogenic diet. *Epilepsia* 2002; 43:1168.
- Krishnamoorthy ES, Reghu R. (2007):** The psychoses of epilepsy. In: Ettinger AB, Kanner AM, editors. *Psychiatric issues in epilepsy: apractical guide to diagnosis and treatment*. 2nd edition. Philadelphia: Lippincott Williams & Wilkins; p. 264–71.
- Krishnamoorthy ES, Trimble MR. (1999):** Forced normalization: clinical and therapeutic relevance. *Epilepsia*; 40 (Suppl10):S57–64.
- Krumholz A, Wiebe S, Gronseth G, et al. (2007):** Evaluating an apparent unprovoked first seizure in adults (an evidence-based review): report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Epilepsy Society. *Neurology* 2007; 69: 1996-2007.
- Kutluay E, McCague K, D’Souza J, et al. (2003):** Safety and tolerability of oxcarbazepine in elderly patients with epilepsy. *Epilepsy Behav.*; 4: 175 – 80.
- Kwan P, Sills GJ, Brodie MJ. (2001):** The mechanisms of action of commonly used antiepileptic drugs. *Pharmacol Ther* 2001; 90:21–34.
- Kwan P, Brodie MJ. (2003):** Clinical trials of antiepileptic medications in newly diagnosed patients with epilepsy. *Neurology* 2003 ; 60(suppl4):S2–S12
- Kwan P, Sander JW. (2004):** The natural history of epilepsy: an epidemiological view. *J Neurol Neurosurg Psychiatry* 2004; 75: 1376–81.

- Lachhwani DK, Loddenkemper T, Holland KD, et al. (2008):*** Discontinuation of medications after successful epilepsy surgery in children. *Pediatr Neurol* 2008; 38:340.
- Laurie B, (2009):*** First Simple Febrile Seizure in Some Young Children Not Likely Bacterial Meningitis. *Pediatrics*. 2009; 123: 6-12.
- Lawrence JH, David B, Hilary T, et al., (2006):*** Predictors of Lamotrigine-associated Rash. *Epilepsia* 2006; 47(2).
- Lawson JA, Cook MJ, Vogrin S., et al. (2002):*** Clinical, EEG, and quantitative MRI differences in pediatric frontal and temporal lobe epilepsy. *Neurology*; 58 : 723–729.
- Leach, JP, Chadwick, DW, Miles JB. (1999):*** Improvement in adult-onset Rasmussen's encephalitis with long-term immunomodulatory therapy. *Neurology* 1999; 52:738.
- Lehmkuhl G, Dopfner M, Pluck J, et al (1993):*** Elternfragebogen über das Verhalten von Kindern und Jugendlichen. Deutsche Bearbeitung der CBCL. Köln, Arbeitsgruppe Kinder-, Jugendund Familien diagnostic, 1993.
- Leibenluft E, Charney D.S. and Towbin K.E., et al. (2003):*** Defining clinical phenotypes of juvenile mania, *Am J Psychiatry* 160 (2003), pp. 430–437.
- Leung AK, Robson WL. (2007):*** Febrile seizures. *J Pediatr Health Care*. 2007; 21(4):250-5.
- Levy RH, Mattson RH, Meldrum B, et al., (2002):*** editors. Anti-epileptic drugs. Philadelphia: Lippincott, Williams & Wilkins.

- Levisohn PM (2007):** "The autism-epilepsy connection". *Epilepsia* 48 (Suppl 9): 33–5.
- Lewis JN, Tonge BJ, Mowat DR, et al. (2000):** Epilepsy and associated psychopathology in young people with intellectual disability. *J Pediatr Child Health*; 36:172–5.
- Li Z, Maglione M, Tu W, et al. (2005):** Meta-analysis: pharmacologic treatment of obesity. *Ann Intern Med*; 142(7):532–46.
- Lieven Lagae. (2008):** What’s new in: “Genetics in childhood epilepsy”. *Eur J Pediatr*; 16: 715 – 722.
- Li X, Jope RS. (2002):** Glycogen synthase kinase-3b, mood stabilizers, and neuroprotection. *Bipolar Disord*; 4:137–44.
- Licht EA, Fujikawa DG. (2002):** Nonconvulsive status epilepticus with frontal features: quantitating severity of subclinical epileptiform discharges provides a marker for treatment efficacy, recurrence and outcome. *Epilepsy Res*; 51:13-21.
- Loiseau P, Duche B, Pedespan JM. (1999):** Absence epilepsies. *Epilepsia*. Dec 1999; 36 (12):1182-6.
- Lubrich B, Spleiss O, van Calker D. (2000):** Differential expression, activity and regulation of the sodium/myo-inositol cotransporter in astrocyte cultures from different regions of the rat brain. *Neuropharmacology*; 39: 680–90.
- Mani J, Gupta A, Mascha E, et al. (2006):** Postoperative seizures after extratemporal resections and hemispherectomy in pediatric epilepsy. *Neurology* 2006; 66:1038.

- Marcangelo MJ, Ovsiew F. (2007):*** Psychiatric aspects of epilepsy. Psychiatr Clin North Am.; 30 :781–802.
- Margaret P. Jacobs , Gabrielle G., et al.(2009):*** Curing epilepsy: Progress and future directions. Epilepsy & Behavior 14 (2009) 438–445
- Marianne J. Kjeldsen, Linda A. (2006):*** Analysing the Etiology of Benign Rolandic Epilepsy: A Multi-center Twin Collaboration. Epilepsia;47(3).
- Marson AG, Chadwick DW. (2001):*** New drug treatments for epilepsy. J Neurol Neurosurg Psychiatry 2001; 70: 143–48.
- Marson AG, Al-Kharusi AM, Alwaidh M, et al. (2007):*** The SANAD study of effectiveness of valproate, lamotrigine, ortopiramate for generalized and unclassifiable epilepsy: an unblended randomized controlled trial. Lancet; 24; 369: 1016–1026.
- Marson A, Jacoby A, Johnson A, et al. (2007):*** Immediate versus deferred antiepileptic drug treatment for early epilepsy and single seizures: a randomized controlled trial. Lancet 2005; 365: 2007–13.
- Martin J. Brodie, Elaine A. Wilson, David L. Wesche, et al., (2005):*** Pregabalin Drug Interaction Studies: Lack of Effect on the Pharmacokinetics of Carbamazepine, Phenytoin, Lamotrigine, and Valproate in Patients with Partial Epilepsy. Epilepsia 2005; 46(9).

- Martinez CC, Pyzik PL, Kossoff EH. (2007):** Discontinuing the ketogenic diet in seizure-free children: recurrence and risk factors. *Epilepsia* 2007; 48:187.
- Maton B, Jayakar P, Resnick T, et al. (2008):** Surgery for medically intractable temporal lobe epilepsy during early life. *Epilepsia* 2008; 49:80.
- Maydell BV, Berenson F, Rothner AD, et al. (2001):** *Benign myoclonus of early infancy: an imitator of West's syndrome. J Child Neurol* 2001; 16: 109–12.
- Mazza M, Di Nicola M, Della Marca G, et al. (2007):** Bipolar disorder and epilepsy: a bidirectional relation? Neurobiological underpinnings, current hypotheses and future research directions. *Neuroscientist*; 13: 392 - 404.
- Mazza M, Della Marca G, Di Nicola M, et al. (2007):** Oxcarbazepine improves mood in patients with epilepsy. *Epilepsy Behav* ; 10(3): 397–401.
- Mc Cormick DA, Contreras D. (2001):** On the cellular and network bases of epileptic seizures. *Annu Rev Physiol* 2001; 63: 815–46.
- Mc Dermott S., Mani S., Krishnaswami S. (1995):** A population-based analysis of specific behavior problems associated with childhood seizures. *J Epilepsy*; 8: 110-118.
- Mc Donagh J, Stephen LJ, Dolan FM, et al. (2003):** Peripheral retinal dysfunction in patients taking vigabatrin. *Neurology* 2003; 61: 1690–4.
- Mc Intyre J, Robertson S, Norris E, et al. (2005):** Safety and efficacy of buccal midazolam versus rectal diazepam for emergency treatment of seizures in children: a randomised controlled trial. *Lancet* 2005; 366:205.

- McNelis A., Musick B. and Austin J., et al. (1998):** Psychosocial care needs of children with new-onset seizures. 2, J Neurosci Nurs **30**, pp. 161–165
- Meador KJ. (2006):** Cognitive and memory effects of the new antiepileptic drugs. Epilepsy Res 2006; 68: 63–7.
- Mintzer S. and Lopez F. (2002):** Comorbidity of ictal fear and panic disorder, Epilepsy Behav **3**, pp. 330–337.
- Michael A. (2008):** Common Pathophysiologic Mechanisms in Migraine and Epilepsy. Arch Neurol. 2008; 65(6):709-714.
- Miriam HM and Jennifer AK (2005):** Sodium channel mutations in epilepsy and other neurological disorders. Journal of Clinical Investigation **115** (8): 2010–2017.
- Mitchell WG, Scheier LM, Baker SA. (1994):** Psychosocial, behavioral, and medical outcomes in children with epilepsy: a developmental risk factor model using longitudinal data. Paediatrics; 94:471–7.
- Mishory A, Yaroslavsky Y, Bersudsky Y,et al. (2000):** Phenytoin as an antimanic anticonvulsant: a controlled study. Am J Psychiatry;157(3) : 463–5.
- Mogens V, Anders H, Kreesten M, et al., (2004):** MMR Vaccination and Febrile Seizures. JAMA. 2004 ; 292:351-357.
- Morrell F, Whisler WW, Smith MC, et al. (1995):** Landau-Kleffner syndrome. Treatment with subpial intracortical transection. Brain 1995; 118 (Pt 6):1529.

- Morrell MJ, Pedley TA. (2000):*** "The Scarlet E": epilepsy is still a burden. *Neurology* 2000;54: 1882-1883.
- Morrell MJ, Isojarvi J, Taylor AE, et al. (2003):*** Higher androgens and weight gain with valproate compared with lamotrigine for epilepsy. *Epilepsy Res* 2003; 54:189–99.
- Motamedi G. and. Meador K. (2003):*** Epilepsy and cognition, *Epilepsy Behav* 4 (suppl 2) , pp. S25–S38.
- Motamedi G, Meador K. (2003):*** Epilepsy Behav. 2003 Oct;4 Suppl 2: S25-38.
- Moyer (2004):*** Epilepsy and Mood Disorders: Separate But Overlapping Conditions and Treatments. *Neurology Today*; **Volume 4 - Issue 2 - pp 70, 75.**
- Muzina D.J, Elhaj O, Gajwani P, et al. (2005):*** Lamotrigine and antiepileptic drugs as mood stabilizers in bipolar disorder. *Acta Psychiatr Scand* 2005;111(Suppl.426):21–28.
- Mula M, Trimble MR, Yuen A, et al. (2003):*** Psychiatric adverse events during levetiracetam therapy. *Neurology*; 61(5):704–6.
- Nahas Z, Marangell LB, Husain MM, et al. (2005):*** Two-year outcome of vagus nerve stimulation (VNS) for treatment of major depressive episodes. *J Clin Psychiatry*; 66(9):1097–104.
- Nicolai J., Aldenkamp A.P. and Arends J., et al. (2006):*** Cognitive and behavioral effects of nocturnal epileptiform discharges in children with benign childhood epilepsy with centrotemporal spikes, *Epilepsy Behav* (8), pp. 56–70.

-
- Nishida T, Kudo T. and Nakamura F, et al. (2005):*** Postictal mania associated with frontal lobe epilepsy, *Epilepsy Behav* **6** , pp. 102–110.
- Nilsson L, Ahlbom A, Farahmand B, et al. (2002):*** Risk factors for suicide in epilepsy: a case control study. *Epilepsia*; 43: 644-651.
- Nolan MA, Redoblado MA, Lah S, et al. (2004):*** Memory function in childhood epilepsy syndromes. *J Paediatr Child Health*. 2004; 40(1-2): 20-7.
- Nordli DR Jr, Bazil CW, Scheuer ML, Pedley TA. (1997):*** Recognition and classification of seizures in infants. *Epilepsia* 1997; 38(5):553–560.
- O'Donoghue MF, Duncan JS, Sander JW. (1998):*** The subjective handicap of epilepsy: a new approach to measuring treatment outcome. *Brain* 1998; 121(Pt. 2):317–43.
- Oguni H, Olivier A, Andermann F. (1991):*** Anterior callosotomy in the treatment of medically intractable epilepsies: A study of 43 patients with a mean follow-up of 39 months. *Ann Neurol* 1991; 30:357.
- Oguz A, Kural S, Dirik E. (2002):*** Relationship of epilepsy-related factors to anxiety and depression scores in epileptic children. *J Child Neurol* ; 17: 37-40.
- Opp J, Tuxhorn I, May T, et al. (2005):*** Levetiracetam in children with refractory epilepsy: amulticenter open label study in Germany. *Seizure* ; 14 (7):476–84.

-
- Ott D, Siddarth P, Gurbani S, et al. (2003):** Behavioral disorders in pediatric epilepsy: unmet psychiatric need. *Epilepsia*; 44:591–7.
- Oyegbile T, Hansen R, Magnotta V, et al. (2004):** Quantitative measurement of cortical surface features in localization-related temporal lobe epilepsy. *Neuropsychology*; 18 : 729 – 737.
- Panayiotopoulos CP. (2005):** Idiopathic generalised epilepsies. In: Panayiotopoulos CP, editor. *The Epilepsies: Seizures, Syndromes and Management*. Oxford: Bladon Medical Publishing, 2005:271-348.
- Pande AC, Crockatt JG, Janney CA, et al. (2000):** Gabapentin in bipolar disorder: a placebo controlled trial of adjunctive therapy. Gabapentin Bipolar Disorder Study Group. *Bipolar Disord*; 2(3Pt2):249–55
- Patel H, Dunn DW. (2002):** Epilepsies in children. In: Biller J, ed. *Practical Neurology* (second edition). Philadelphia, Lippincott, Williams & Wilkins, 2002; 469-488.
- Perucca E. (2007):** Treatment of epilepsy in developing countries. *BMJ* 2007; 334:1175.
- Pellock JM. (1999):** Felbamate. *Epilepsia*. 1999; 40(suppl 5):S57-S62.
- Pellock JM, Shinnar S. (2005):** Respiratory adverse events associated with diazepam rectal gel. *Neurology* 2005; 64:1768.

- Pfeifer HH, Thiele EA. (2005):*** Low-glycemic-index treatment: a liberalized ketogenic diet for treatment of intractable epilepsy. *Neurology* 2005; 65:1810.
- Phiel CJ, Zhang F, Huang EY. (2001):*** Histone deacetylase is a direct target of valproic acid, a potent anticonvulsant, mood stabilizer, and teratogen. *J Biol Chem*; 276:36734–41.
- Plioplys S. (2003):*** Depression in children and adolescents with epilepsy, *Epilepsy Behav* 4 (suppl 3), pp. S39–S45.
- Ranganathan LN, Ramaratnam S. (2006):*** Rapid versus slow withdrawal of antiepileptic drugs. *Epilepsia* 2006; 47(7): 1094-1120.
- Ratnapriya R, Satishchandra P, Dilip S, et al., (2009):*** Familial autosomal dominant reflex epilepsy triggered by hot water maps to 4q24-q28. *Hum Genet*. Nov 2009; 126(5):677-83.
- Rayport M, Ferguson SM. (2001):*** Psychosis of epilepsy: An integrated approach. In: Ettinger AB, Kanner AM, editors. *Psychiatric Issues in Epilepsy: A Practical Guide to Diagnosis and Treatment*. Philadelphia (PA): Lippincott Williams & Wilkins; p. 73-94.
- Reijs R, Aldenkamp AP, De Krom M. (2004):*** Mood effects of antiepileptic drugs. *Epilepsy Behav*; 5(Suppl1): S66–76.
- Reith D, Burke C, Appleton DB, et al. (2003):*** Tolerability of topiramate in children and adolescents. *J Paediatr Child Health*; 39(6):416–9.
- Rivinus TM. (1982):*** Psychiatric effects of the anticonvulsant regimens. *J Clin Psychopharmacol*; 2(3):165–92.

- Reynolds JR. (1861):** Epilepsy: its symptoms, treatment and relation to other chronic convulsive diseases. London: Churchill.
- Rianne R , Albert P. Aldenkamp, et al. (2004):** Mood effects of antiepileptic drugs. *Epilepsy & Behavior* Volume 5, Supplement 1, February 2004, Pages 66-76
- Riss J, Cloyd J, Gates J, et al. (2008):** Benzodiazepines in epilepsy: pharmacology and pharmacokinetics. *Acta Neurol Scand.* 2008 Aug; 118 (2): 69-86.
- Riviello JJ, Holmes GL. (2004):** The treatment of status epilepticus. *Semin Pediatr Neurol* 2004; 11: 129–138.
- Robert S. (2009):** Are AEDs Forever? *Epilepsia*. Volum 5 issue 6.
- Robert S. (2009):** Summary of Antiepileptic Drugs. *Spotlight*. Volume 5, Issue 9.
- Rodenburg R., Meijer A.M. and M. Dekovic et al. (2005):** Family factors and psychopathology in children with epilepsy: A literature review, *Epilepsy Behav* 6, pp. 488–503.
- Rodenburg HR, Meijer AM, et al. (2005):** Psychopathology in children with epilepsy: ameta-analysis. *J Pediatr Psychol*; 30:453 – 68.
- Rogawski M, Loscher W. (2004):** The neurobiology of antiepileptic drugs." *Nat Rev* 2004;5: 1–13.
- Rutter M, Tizard J, YuleW, et al. (1979):** 1964—1974 Isle of Wight studies. *Psychol Med* 1979; 6: 313—32.
- Rush AJ, Marangell LB, Sackeim HA, et al. (2005):** Vagus nerve stimulation for treatment-resistant depression: a randomized, controlled acute phase trial. *Biol Psychiatry*;58(5): 347–54.
- Sachdev PS. (2007):** Alternating and postictal psychoses: review and a unifying hypothesis. *Schizophr Bull*; 33(4):1029–37.

- Sackellares JC, Berent S. (1996):** Summary and conclusions. In: Sackellares JC, Berent S, eds. *Psychological Disturbances in Epilepsy*. Boston: Butterworth Heinemann; 1996:259-264.
- Sahin M, Menache C, Holmes GL, Riviello JJ.** Outcome of severe refractory status epilepticus in children. *Epilepsia*; 42: 1461–1467.
- Salgado P.C., and E.A. de Souza. (2006):** A psychosocial view of anxiety and depression in epilepsy, *Epilepsy Behav* (8), pp. 232–238.
- Salpekar J.A., Conry J.A. and W. Doss , et al. (2006):** Clinical experience with anticonvulsant medication in pediatric epilepsy and comorbid bipolar spectrum disorder, *Epilepsy Behav* (9), pp. 327–334.
- Sanchez-Carpintero R, Neville BGR. (2003):** Attention alability in children with epilepsy. *Epilepsia* ; 44(10):1340–9.
- Sarah E, Ingrid E, Leanne M (2007):** in *Idiopathic Epilepsy. Neurotherapeutics* 2007 volum 4, issue 2, pages 295-304.
- Sasha I. Marge Lenane, Judith L. R, et al. (2000):** Gabapentin Prophylaxis of Clozapine-Induced Seizures. *Am J Psychiatry* 157:482-483, March 2000.
- Schipper H, Clinch J and Powell V (1990):** Definitions and conceptual issues. In Spiker B (ed.): “Quality of Life Assessment in Clinical Trials”. New York: Raven Press, pp 11–24.
- Schoenfeld J, Seidenberg M, Woodard A, et al. (1999):** Neuropsychological and behavioral status of children with complex partial seizures. *Dev Med Child Neurol*; 41: 724–31.

- Schmidt D. (2002):** The clinical impact of new antiepileptic drugs after a decade of use in epilepsy. *Epilepsy Res* 2002;50: 21–32.
- Schmitz B (2002):** Antidepressant drugs: indications and guidelines for use in epilepsy. *Epilepsia*. 2002; 43 Suppl 2:14-8.
- Schmitz B. (2005):** Depression and mania in patients with epilepsy. *Epilepsia* 2005; 46 (Suppl 4): 45–9.
- Schubert R. (2005):** Attention deficit disorder and epilepsy. *Pediatr Neurol*; 32(1):1–10.
- Schuele SU, et al. (2008):** Intractable epilepsy: Management and therapeutic alternatives. *Lancet Neurology*. 2008 ;7(6):514-524.
- Scottish Intercollegiate Guidelines Network (SIGN). (2001):** Attention Deficit and Hyperkinetic Disorders in Children and Young People. Edinburgh: SIGN; 2001. (SIGN publication no. 52).
- Scottish Intercollegiate Guidelines Network SIGN, (2005):** Diagnosis and management of epilepsies in children and young people.
- Seethalakshmi R. and E.S. Krishnamoorthy (2007):** Depression in epilepsy: Phenomenology, diagnosis and management, *Epileptic Disord* 9, pp. 1–10.
- Selai C, Bannister D, Trimble M. (2005):** Antiepileptic drugs and the regulation of mood and quality of life (QOL): the evidence from epilepsy. *Epilepsia*; 46(Suppl4):50–7.
- Seidenberg M, Geary E, Hermann B. (2005):** Investigating temporal lobe contribution to confrontation naming using MRI quantitative volumetrics. *J Int Neuropsychol Soc.*; 11: 358 – 366.
- Senol V, Soyuer F, Arman F, et al. (2007):** Influence of fatigue, depression, and demographic, socioeconomic, and clinical

variable on quality of life of patients with epilepsy. *Epilepsy Behav*; 10(1):96–104.

Shinnar S and Berg AT (1996): Does antiepileptic drug therapy prevent the development of "chronic epilepsy"? *Epilepsia* 37:701–708.

Shinnar S, Berg AT, Mosh SL, et al. (1994): Discontinuing antiepileptic drugs in children with epilepsy: A prospective study. *Ann Neurol* 1994; 35:534.

Siddarth P. , Caplan R.and S. Gurbani, et al. (2005): Depression and anxiety disorders in pediatric epilepsy, *Epilepsia* **46** , pp. 720–730.

Siddhartha Nadkarnia and Orrin Devinsky. (2005): Psychotropic Effects of Antiepileptic Drugs. *Epilepsy Curr.* 2005 September; 5(5): 176–181.

Sillanpaa M, Schmidt D. (2006): Prognosis of seizure recurrence after stopping antiepileptic drugs in seizure-free patients: A long-term population-based study of childhood-onset epilepsy. *Epilepsy Behav* 2006; 8:713.

Sillanpaa M. and D. Schmidt (2006): Natural history of treated childhood-onset epilepsy: Prospective, long-term population-based study, *Brain* **129**, pp. 617–624.

Smith D, Defalla BA, Chad wick DW. (1999): The misdiagnosis of epilepsy and the management of refractory epilepsy in a specialist clinic. *QJ Med* 1999; 92: 15–23.

Smith PEM. (2001): Clinical recommendations for oxcarbazepine. *Seizure* 2001; 10:87–91.

-
- Stone JL, Mc Daniel KD, Hughes JR, et al. (1986):** Episodic dyscontrol disorder and paroxysmal EEG abnormalities: successful treatment with carbamazepine. *Biol Psychiatry*; 21(2): 208–12.
- Stores G (2003):** Medication for sleep-wake disorders. *Arch Dis Child*; 88(10):899-903.
- Suncica Lah, Teresa Lee, Sandra Grayson,et al. (2003):** Effects of Temporal Lobe Epilepsy on Retrograde Memory. *Epilepsia* 2006; 47(3).
- Swinkels WA, Kuyk J, van DR, et al. (2005):** Psychiatric comorbidity in epilepsy. *Epilepsy Behav*; 7:37–50.
- Theodore WH, Fisher RS. (2004):** Brain stimulation for epilepsy. *Lancet Neurol* 2004; 3: 111–8.
- Theodore WH, Hasler G. and G. Giovacchini et al. (2007):**Reduced hippocampal 5HT1A PET receptor binding and depression in temporal lobe epilepsy, *Epilepsia* **48** , pp.1526–1530.
- The National Society for Epilepsy (2009):** What is Epilepsy?. Available from <http://www.epilepsynse.org.uk/AboutEpilepsy/Whatisepilepsy> (Accessed on 15 February 2009).
- Theodore H. M and Alan R. B (2004):** Delirium from Valproic Acid With Lamotrigine. *Am J Psychiatry* 161:1128-1129.
- Thomas P. (2000):** Status epilepticus with confusional symptomatology [French]. *Neurophysiol Clin* 2000; 30:147-54.
- Thomas P (2004):** Mechanisms of epilepsy progression: current theories and perspectives from neuroplasticity in adulthood and

development. *Epilepsy Research*. Volume 60, Issue 2, Pages 161-171.

Thomas P, Valton L, Genton P. (2006): Absence and myoclonic status epilepticus precipitated by antiepileptic drugs in idiopathic generalized epilepsy. *Brain* 2006; 129:1281.

Thome-Souza S, Kuczynski E, Assumpcao F, et al. (2004): which factors may play a pivotal role on determining the type of psychiatric disorder in children and adolescents with epilepsy?. *Epilepsy Behave* ;5(6):988-94.

Torta R, Keller R. (1999): Behavioral, psychotic, and anxiety disorders in epilepsy: etiology, clinical features, and therapeutic implications. *Epilepsia*; 40 Suppl 10:S2-20.

Urbach H. (2005): Imaging of the epilepsies. *Eur Radiol*. Mar 2005;15 (3): 494-500.

Vaisleib II, Buchhalter JR, Zupanc ML. (2004): Ketogenic diet: outpatient initiation, without fluid, or caloric restrictions. *Pediatr Neurol* 2004; 31:198.

Vasudev K, Macritchie K, Geddes J, et al. (2006): Topiramate for acute affective episodes in bipolar disorder. *Cochrane Database Syst Rev*; 1: CD003384.

Vining EP, Freeman JM, Pillas et al. (1968): Why would you remove half a brain? The outcome of 58 children after hemispherectomy-The Johns Hopkins experience: 1968 to 1996. *Pediatrics* 1997; 100:163.

Vuilleumier P, Jallon P. (1998): Epilepsy and psychiatric disorders: epidemiological data. *Rev Neurol (Paris)*. 154(4):305-17.

- Warren CR. (2003):** Evaluation and management of febrile seizures in the out-of-hospital and emergency department settings. *Ann Emerg Med.* 2003; 41(2): 215-222.
- Watemberg N, Leitner Y, Fattal-Valevski A, et al., (2009):** Epileptic negative myoclonus as the presenting seizure type in rolandic epilepsy. *Pediatr Neurol.* 2009; 41(1):59-64.
- Waugh J, Goa KL. (2003):** Topiramate: as monotherapy in newly diagnosed epilepsy. *CNS Drugs* 2003; 17:985–92.
- Werz MA, Meador KJ,et al. (2006):** Subjective preference for Lamotrigine or Topiramate in healthy volunteers:Relationship to cognitive and behavioral functioning. *Epilepsy Beav* ; 8(1): 181-191
- Westbrook LE, Bauman LJ and Shinnar S. (1992):** Applying stigma theory to epilepsy: A test of a conceptual model, *J Pediatr Psychol* (17), pp. 633–649.
- WHO: Fact Sheet N° 168, Revised February (2001):** Epilepsy: an historical overview.
- Whiting S, Farrell K, Wirrell E. (2006):** Diseases and syndromes associated with intractable epilepsy. *Adv Neurol* 97:443–462.
- Wiebe S (2006):** Outcome measures in intractable epilepsy. *Adv Neurol* 97:11–15.
- Wiebe s, blum wt , girvin jp et al., (2001):** effectiveness and efficiency of surgery for temporal lobe epilepsy study group. A randomized, controlled trial of surgery for temporal lobe epilepsy. *N Engl J Med* ;345:311-8.

- Wilkins A, Emmett J, Harding G. (2005):** Characterizing the patterned images that precipitate seizures and optimizing guidelines to prevent them. *Epilepsia*. Aug 2005; 46(8):1212-8.
- Wilfong A, et al. (2009):** Overview of the classification, etiology, and clinical features of pediatric seizures and epilepsy. <http://uptodate.com/home/index.html>. Accessed Jan. 6, 2009.
- Williams RS, Cheng L, Mudge AW, (2002):** A common mechanism of action for three mood-stabilizing drugs. *Nature* 2002;417:292–5.
- William O, Peter W, Pierre J (2009):** Antiepileptogenesis. *Epilepsy A to Z* Second Edition.
- William O, Peter W, Pierre J (2009):** Syndromes (EPILEPTIC). *Epilepsy A to Z* Second Edition.
- William O, Peter W, Pierre J (2009):** *Progabide*. *Epilepsy A to Z* Second Edition.
- Winkler AS, Schaffert M, Schmutzhard E. (2007):** Epilepsy in resource poor countries suggestion of an adjusted classification. *Epilepsia* 2007; 48:1029-1030.
- Wiznitzer M. (2005):** Buccal midazolam for seizures. *Lancet* 2005; 366:182.
- Wyllie E, Lachhwani DK, Gupta A, et al. (2007):** Successful surgery for epilepsy due to early brain lesions despite generalized EEG findings. *Neurology* 2007; 69:389.
- Wolf P. (2003):** Of cabbages and kings: some considerations on classifications, diagnostic schemes, semiology, and concepts. *Epilepsia* 2003; 44(1):1–4, 13.
- World Health Organization. (2001):** "Epilepsy: aetiology [sic], epidemiology and prognosis".

<http://www.who.int/mediacentre/factsheets/fs165/en/>.

Retrieved on 2007-06-14.

Williams J., Steel C. and Sharp G.B., et al. (2003): Parental anxiety and quality of life in children with epilepsy, *Epilepsy Behav* (4), pp. 483–486.

Yan QS, Jobe PC, Dailey JW. (1995): Further evidence of anticonvulsant role for 5–hydroxy tryptamine in genetically epilepsy-prone rats. *Br J Pharma-col.*; 115:1314–1318.

Yuan PX, Huang LD, Jiang YM, et al. (2001): The mood stabilizer valproic acid activates mitogen-activated protein kinases and promotes neurite growth. *J Biol Chem*; 276: 31674 – 83.

Yudofsky (2002): Do the epilepsies, pain syndromes, and affective disorders share common kindling-like mechanisms? *Epilepsy Res* 2002; 50(1-2):203-219.

Zaccara G, Franciotta D, Perucca E. (2007): Idiosyncratic adverse reactions to antiepileptic drugs. *Epilepsia*; 48: 1223–44.

Zelnik N, Isler N, Goetz H, et al., (2008): Vigabatrin, lamotrigine, topiramate and serum carnitine levels. *Pediatr Neurol.* 2008; 39 (1):18-21.

Zeina Chemali (2006): Psychotropic effects of antiepileptic drugs. *Neurology*; 67:1916-25.