

References

- Abdullah Kurt, A. Denizmen Aygun, A. Nese Citak Kurt, Ahmet Godekmerdan, Saadet Akarsu, Erdal Yilmaz.(2009);** Use of Phototherapy for Neonatal Hyperbilirubinemia Affects Cytokine Production and Lymphocyte Subsets: Neonatology;95:262–266
- Aderka D., Engelmann H., Shemer A.Y., Hornik V., Galil A., Sarvo B. and Wallach D. (1992):** Variation of serum levels of soluble TNF receptors among healthy individuals. Lymphokine Cytokine Res; 11: 157-159.
- Ahlfors CE, Wennberg RP, Ostrow JD, Tiribelli C. (2009).** Unbound (free) bilirubin: improving the paradigm for evaluating neonatal jaundice. Clin Chem. Jul; 55(7):1288-99. Epub May 7
- AlOtaibi SF, Blaser S, MacGregor DL (2005):** Neurological complication of kernicterus. Can J Neurol Sci Aug; 32(3): 311-5.
- Ameloot P., Fiers W., Bleser P.D., Ware C.F., Vandenabede P. and Broukaert P. (2001) :** Identification of TNF amino acid crucial for binding to murine p75 TNF JR. and construction of receptor selective mutants. J Biol Chem; 276(4): 37426-30.
- American Academy of Pediatrics (2004) (a):** Practice Parameter: Management of hyperbilirubinemia in the healthy term newborn. Pediatrics; 94(4): 558-65.
- American Academy of Pediatrics Subcommittee on Hyperbilirubinemia (2004):** Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. Pediatrics; 114(1):297-316.
- Andreas KÖck, Thomas Schwarz, Reinhard Kirnbauer, Agatha Urbanski, Patricia Perry, S John C. Ansel,§ and Thomas A. Luger (1990):** Human Keratinocytes Are a Source for Tumor Necrosis Factor α Evidence for, Synthesis and Release upon Stimulation with Endotoxin or Ultraviolet Light The journal of Experimental Medicine Volume 172 December 1609-1614

- Andrew Thorburn, (2004).** Cellular Signalling, Volume 16, Issue 2, February, Pages 139-144
- Balagangadhar R. Totapally, MD; Dan Torbati, PhD (2005):** Neonatal Jaundice Practical Biology Int Pediatr. 20(1):47-54.
- Barefield ES, Dwyer MD, Cassady G (2000):** Association of patent ductus arteriosus and phototherapy in infants weighting less than 1000 grams. J Perinatol; 13 (5):376-80.
- Bazzoni F. and Beutler B. (1996):** Tumour necrosis factor ligand and receptor families. N Engl J Med; 334: 1717-1725.
- Bekker L.G., Moreira A.L., Bergtold A., Freeman S., Ryffel B. and Kaplan G. (2000):** Immunopathologic effect of TNF-a in murine mycobacterial infection are dose depenent. Infection and Immunity; 68: 6954-6961.
- Benders MJ, Van Bel F, Van de Bor M (1991):** Hemodynamic consequences of phototherapy in term infants. Eur J Pediatr; 158:323-8.
- Berg CL, Crawford JM, Gollan JL (1999):** Bilirubin metabolism and pathophysiology of jaundice. In: Schiff ER, Sorrell MF, Maddrey WC (Eds.). Schiff s diseases of the liver (8thEd). Lippincott-ravin. Philadelphia, pp 147-60.
- Benjafield A.V., Glenn C.L., Wang X.L., Colagiuri S. and Morris B.J. (2001):** TNFR SFIB in genetic predisposition to clinical neuropathy and effect on HDL, cholesterol and glycosylated hemoglobin in type 2 diabetes. Diabetes care; 24: 753-757.
- Bhutani VK, Johnson L (2009).** Kernicterus in the 21st century: frequently asked questions. J Perinatol. Eb; 29 Suppl 1:S20-4.

- Boonyarittipong P, Kriangburapa W, Booranavanich K.(2008)** Effectiveness of double-surface intensive phototherapy versus single-surface intensive phototherapy for neonatal hyperbilirubinemia- J Med Assoc Thai. Jan; 91(1):50-5.).
- Boutati E, Lambadiari V, Tsegka A, Raptis A, Tountas N, Economopoulos T, Raptis S, Dimitriadis G.(2009):** Insulin resistance in hyperthyroidism: the role of IL-6 and TNF alpha. Eur J Endocrinol. Oct 16. [Epub ahead of print]
- Braun J. and Sieper J. (2003):** Overview of the use of anti TNF agent infliximal in chronic inflammatory diseases. Expert Opin Bio Ther; 3(1): 141-168.
- Brien D.P., Briles D.E., Szalar A.J., Tu A.H., Sanz I. And Nahm M.H. (1999):** TNF & RI is important for survival from streptococcus pneumonia infections. Infection and Immunity; 67: 595-601.
- Camilia R, Martin and John PC (2007):** Neonatal hyperbilirubinemia. In: **Cloherly JP, Eichenwal EC, Stark AR (Eds):** Manual of neonatal care, (6th edition), Lippincott Williams&Wilkins. ppl85-221
- Cairns and A. J. Taggart. (2003).** Tumour necrosis factor inhibitors: maximizing patient safety. Rheumatology; 42: 188-189.
- Cartier A, Côté M, Bergeron J, Alméras N, Tremblay A, Lemieux I, Després JP.(2009):** Plasma soluble tumour necrosis factor-alpha receptor 2 is elevated in obesity: specific contribution of visceral adiposity. Clin Endocrinol (Oxf). Jul 24. [Epub ahead of print]
- Cetingul, N., E. Yener, S. Oztop, and G. Nisli. (1994).** Serum soluble interleukin-2 receptors and tumor necrosis factor-alpha in hematological malignancies of childhood. Acta Paediatr. Jpn. 36:49–52.

- Chang FY, Lee CC, Huang CC, Hsu KS.(2009).** Unconjugated bilirubin exposure impairs hippocampal long-term synaptic plasticity. *PLoS One*. Jun 11; 4(6):e5876.
- Chen G, Goeddel DV (2002)** "TNF-R1 signaling: a beautiful pathway". *Science* 296 (5573): 1634–5.
- Chen HN, Lee ML, Tsao LY. (2008).** Exchange transfusion using peripheral vessels is safe and effective in newborn infants. *Pediatrics*. Oct; 122(4):e905-10. Epub Sep.
- Chen X, Xun K, Chen L, Wang Y.(2009).** TNF-alpha, a potent lipid metabolism regulator. *Cell Biochem Funct*. Sep 15.
- choy E.H. and Panayi G.S. (2001):** Cytokine pathway and joint inflammation in rheumatoid arthritis. *N Eng J Med*; 344: 907-916.
- Cohen SM (2006):** Jaundice in the Full-Term Newborn.*Pediatr Nurs*; 32(3):202-8.
- Costantino Romagnoli, Enrico Zecca, Patrizia Papacci, Giovanni Vento, Pietro Girlando, Caterina Latella (2006)** Which Phototherapy System Is Most Effective in Lowering Serum Bilirubin in Very Preterm Infants?-*Fetal Diagn Ther*;21:204-209
- CREMER RJ, PERRYMAN PW, RICHARDS DH (1958):** Influence of light on the hyperbilirubinaemia of infants. *Lancet*. May 24; 1094-1097.
- De-Carvalho, M.; De-Carvalho, D.; Tizmielin, S.; Lopes, J.M. And Hansen, T.W. (1999):** Intensified phototherapy using day light fluorescent lamps. *Acta. Pediatr*. Jul. 88: (7): 768-71.
- Dennerly PA, Seidman DS, and Stevenson DK (2001):** Neonatal Hyperbilirubinemia; *NEJM*; 344:581-90.
- Deshpande PG (2006);** Breast Milk Jaundice. *Eur J Pediatr*; 158:323-8.
- Dunia M. Jawdat, Geoffrey Rowden, and Jean S. Marshall (2006).** Mast Cells Have a Pivotal Role in TNF-Independent Lymph Node Hypertrophy and the

- Mobilization of Langerhans Cells in Response to Bacterial Peptidoglycan1. *Journal of Immunology*, 177: 1755–1762.
- El-Beshbishi SN, Shattuck KE, Mohammad AA, Petersen JR. (2009).** Hyperbilirubinemia and transcutaneous Clin Chem. Jul; 55(7):1280-7. Epub May 14.
- Elzbieta Waszczykowska, Ewa Robak, Anna Wozniacka, Joanna Narbutt, Jolanta Dorota Torzecka and Anna Sysa-JedrzejowskaCA. (1999).** Estimation of SLE activity based on the serum level of chosen cytokines and superoxide radical generation. *Mediators of Inflammation*, 8, 93–100
- Engel, W.P.; Baumgart, S.; Schwartz J.S. et al. (1981):** Insensible water loss in critically ill neonate. Combined effect of radiant warmer *Am.J.Child.* 135:S16-20.
- Erduran E, Arslan MK, Alperen A, Tekelioglu Y (1997):** In vitro investigation of the effect of ultraviolet-C light on lymphocytes. *Turk Hematol Onkol Derg*; 7: 30–32.
- Fok TF, Wong W, Cheng AF (1995):** Use of eye patches in phototherapy: effects on conjunctival bacterial pathogens and conjunctivitis. *Pediatr Infect Dis J*; 14:1091-4.
- Francis Ka-Ming Chan, Joanna Shisler, Jacqueline G. Bixby, Martin Felices, Lixin Zheng, Michael Appel, Jan Orenstein, Bernard Moss, and Michael J. Lenardo (2003).** A Role for Tumor Necrosis Factor Receptor-2 and Receptor-interacting Protein in Programmed Necrosis and Antiviral Responses. *THE JOURNAL OF BIOLOGICAL CHEMISTRY*; Vol. 278, No. 51, Issue of December 19, pp. 51613–51621.

- Gaur U, Aggarwal BB (2003).** "Regulation of proliferation, survival and apoptosis by members of the TNF superfamily". *Biochem. Pharmacol.* 66 (8): 1403–8.
- Gottstein R, Cooke RWI (2003):** Systematic review of intravenous immunoglobulin in hemolytic disease of the newborn. *Arch Dis Child Fetal Neonatal*; 88:F6-10.
- Gounaris, A.; Alexiou, N.; Costalos, C; Daniilidou, M.; Bakoleas, V.; Comstantellou, E. (1998):** GutHormone levels in neonates under phototherapy. *Early. Hum. Dev.* 17: 51 (1): 57-60.
- Gulcan H, Tiker F, Kilicdag H. (2007)** Effect of feeding type on the efficacy of phototherapy. - *Indian Pediatr.* Jan; 44(1):32-6.
- Haider S, Knöfler M.(2009)** Human tumour necrosis factor: physiological and pathological roles in placenta and endometrium. *Placenta.* Feb; 30(2):111-23. Epub Nov 22.
- Hansen TW (1997):** Acute management of extreme Neonatal jaundice-the potential benefits of intensified phototherapy and interruption of enterohepatic bilirubin circulation. *Acta Paediatr*; 86:843-6.
- Hansen TW (2007):** Jaundice, neonatal. Extracted from <http://www.emedicine.com/ped/neonatology>. Last updated Oct 18.Extracted on January.
- Harashima S.I., Horiuchi T., Hatta N., Morita C, Hignchi M., Sawabe T., Tsukamoto H., Tahera T., Hayashi K., Fujito S. nd NihoY. (2001):** Outside to inside signal through the membrane TNF- α induce E-selectin (CD 62 E) expression on activated CD4⁺ T cells. *J Immunol*; 166: 130-136.

- Harris M, Bernbaum J, Polin J, et al (2001):** developmental follow up of breastfed term and near term infant with marked hyperbilirubinemia. *Pediatrics* 107:1075-1080.
- Henriksen P.A. and Newly D.E. (2003):** Therapeutic inhibition of TNF- α in patients with heart failure. *Heart*; 89: 14-18.
- Hill G.R., Teshima T., Rebel V.L, Krijanovski O.I., Coake K.R., Brinson Y.S. and Ferrara J.L. (2000):** P55 TNF- α receptor plays a critical role in T-cell alloreactivity. *J Immunol*; 164: 656-663.
- Himmerich H, Sheldrick AJ. (2009).** TNF- α and Ghrelin Opposite Effects on Immune System, Metabolism and Mental Health. *Protein Pept Lett*. Nov 10.
- Hong Zhou, Graciela Andonegui, Connie H. Y. Wong and Paul Kubes (2009).** Role of Endothelial TLR4 for Neutrophil Recruitment into Central Nervous System Microvessels in Systemic Inflammation. *The Journal of Immunology*, doi:10.4049/jimmunol.0901309.
- Hu X., Tang M., Fishes A.B., Olashaw N. and Zuckerman K.S. (1999):** TNF- α induced growth suppression of CD 34+ myeloid leukemic cell lines signals through TNF RI is associated with NF-KB. *J Immunol*; 163: 3106-3115.
- Ioannis Xinias, Vasiliki Demertzidou, Antigoni Mavroudi, Konstantinos Kollios, Panagiotis Kardaras, Fotis Papachristou, Georgios Arsos, Ioannis Tsiouris (2009).** Bilirubin levels predict renal cortical changes in jaundiced neonates with urinary tract infection *World J Pediatr*; 5(1):42-45.
- Ip S, Chung M, Kulig J, O'Brien R, Sege R, Glick S, Maisels MJ, Lau J (2004):** An Evidence-Based Review of Important Issues Concerning Neonatal Hyperbilirubinemia. *Pediatr*; 114: e130-e15

- Javeed Travadi, Karen Simmer, James Ramsay, Dorota Doherty & Ronald Hagan. (2006).** Patent ductus arteriosus in extremely preterm infants receiving phototherapy: Does shielding the chest make a difference? A randomized, controlled trial: *Acta Pædiatrica*, 95: 1418_1423
- Jeffrey P. Walterscheid, Dat X. Nghiem, Stephen E. Ullrich. (2002):** Determining the role of cytokines in UV-induced immunomodulation. *Methods* 28: 71–78
- Jeferey S. Flier, M.D., Editor L ISA H. UNDERHILL, FLAVIA BAZZONI, PH .D. AND B RUCE BEUTLER, M.D.Assistant Editor (1996):** The Tumor Necrosis Factor Ligand and receptor families ; the New England Journal of Medicine Vol. 334 No. 26, Massachusetts Medical Society.
- Joanna Narbutt,1 Aleksandra Lesiak,1 Malgorzata Skibinska,2 Anna Wozniacka Anna Sysa-Jedrzejowska,1 Joalanta Lukamowicz,3 and Henk van Loveren4 (2005) :** Repeated Doses of UVR Cause Minor Alteration in Cytokine Serum Levels in Humans j *Mediators of Inflammation* :5 298–303
- Johnson LH, Brown AK, Bhutani VK (2002) :** system-based approach to management of neonatal jaundice and prevention of kernicterus. *J Pediatr*; 140:397-386.
- Johnson LH, Bahutani VK and Brown AK (2002):** System based approach to management of neonatal jaundice and prevention of kernicterus. *J Pediatr* 140:396-403.
- Kadalraja R, Thomas E, Patole S, Muller R (2005):** Is phototherapy a risk factor for ileus in high-risk neonates?. *J Maternal fetal Neonatal Med*; 18(2): 129-31.

- Kamada H., Tsutsumi Y., Yamamoto Y-, Kihira T., Kaneda Y., Mu Y., Kodaira H., Tsunoda S.I., Nakagawa S. and Mayumi T. (2000):** Antitumour activity of TNF- α conjugated with polyvinylpyrrolidone on solid tumours in mice. *Cancer Research*; 60: 6416-6420.
- Kaplan M, Muraca M, Hammerman C, Rubatelli FF, Vilei MT, Vreman HJ, Stevenson DK (2002):** Imbalance between production and conjugation of bilirubin; a fundamental concept in the mechanism of neonatal jaundice. *Pediatr*; 110: e 47.
- Kroll-Palhares K, Silvério JC, Silva AA, Michailowsky V, Marino AP, Silva NM, Carvalho CM, Pinto LM, Gazzinelli RT, Lannes-Vieira J. (2008)** TNF/TNFR1 signaling up-regulates CCR5 expression by CD8⁺ T lymphocytes and promotes heart tissue damage during *Trypanosoma cruzi* infection: beneficial effects of TNF- α blockade. *Mem Inst Oswaldo Cruz*. Jun; 103(4):375-85.
- Kumar P, Murki S, Malik GK, Chawla D, Deorari AK, Karthi N, Subramanian S, Sravanthi J, Gaddam P, Singh SN. (2009).** Light-emitting Diodes versus Compact Fluorescent Tubes for Phototherapy in Neonatal Jaundice: A Multi-center Randomized Controlled Trial. *Indian Pediatr*. May 20.
- Kumar A, Kumar K. (2003):** Meningitis, Bacterial. *Prim Care Nov*; 9(5): 587-601.
- Kumral A, Genc S, Genc K, et al (2005):** Hyperbilirubinemic serum is cytotoxic and induces apoptosis in murine astrocyte. *Biol Neonate*; 87(2): 99-104.
- Kuprash D.V., Alimzhanov M.B., Tumanov A.V., Grivennikov S.I., Shakhov A.N., Drutskaya L.N., Marino M.W., Turetskaya R.L., Ajaderson A.O., Rajewsky K., Pfeffer K. and nedospasov S.A. (2002):** Redundancy in tumour necrosis factor (TNF) and lymphotoxin (LT)

signaling in vivo: mice with in-activation of the entire TNF/LT locus versus single-knockout mice. *Mol Cell Biol*; 22(24): 8626-8634.

Lale Naderi-Hachtroudi; Thorsten Peters, MD; Peter Brenneisen, PhD; Christian Meewes, MD; Christina Hommel; Ziba Razi-Wolf, PhD; Lars A. Schneider, MD; Jutta Schüller, PhD; Meinhard Wlaschek, PhD; Karin Scharffetter-Kochanek, MD (2002); Induction of Manganese Superoxide Dismutase in Human Dermal Fibroblasts A UV-B–Mediated Paracrine Mechanism With the Release of Epidermal Interleukin 1 α , Interleukin 1 β , And Tumor Necrosis Factor α : *Arch Dermatol.*;138:1473-1479.

Lee J.Y. and Sullivan K.E. (2001): Gamma interferon and lipopolysaccharide interact at the level of transcription to induce TNF- α expression: *Infection and Immunity* 69: 2847-2852.

Lipsky P.E., Vander Heijde D.M., Clair E.W., Furst D.E., Breedveld F.C., Kalden J.R., Smolen J.S., Weisman M., Emerg P., Feldmann M., Harriman G.R. and Maini R.N. (2000): Infliximal and methotrexate in the treatment of rheumatoid arthritis. *N Engl J Med*; 343: 1594-1602.

Liu R.Y., Liu C.G., Olashaw N.E. and Zuckerman K.S. (2000): Activation of p38 mitogen-activated protein kinase is required for TNF- α supported proliferation of leukemia and lymphoma cell lines. *J Biol Chem*; 275(28): 21086-31093.

Lodha R, Deorari AK, Janata V (2002): Non-invasive estimation of total serum bilirubin by multi-wavelength spectral reflectance in neonates. *Indian Pediatr*: 37:771-775.

Lopez S., Halbwachs MX., Ravaud P., Bessou G., Dougados M. and Parteu F. (1995): Neutrophil expression of TNF-R and of activation markers (CD11 b, CD43, CD63). *Clin Exp Immunol*; 101: 25-32.

- Madan A, MacMahon JR, and Stevenson DK (2005):** Neonatal hyperbilirubinemia, in: Taeusch HW, Ballard RA and Gleason CA (Eds). Avery's diseases of the newborn, 8th edition, Saunders, 79:1226-55.
- Maisels MJ (1999):** jaundice in Avery GB, Fletcher MA, Macdonald MG (Eds): Neonatology pathophysiology and management in newborn, Lippincott William and Wilkins (5th Edt) 38:765-818.
- Maisels MJ, Kring E (2002)** Rebound in serum bilirubin level following intensive phototherapy. -Arch Pediatr Adolesc Med. Jul; 156(7):669-72.
- Maisels MJ et al., Kring EA, and DeRidder J (2007)** Randomized controlled trial of light - emitting diode phototherapy Sep; 27(9):565-7.
- Maisels MJ (2005):** Jaundice. In: MacDonald MG, Mullett MD, Seshia MMK, (Eds). Avrey's Neonatology. 6th ed. Philadelphia, Pa: Lippineott, Williams & Wilkins: 768-846.
- Maisels MJ and Watchko JF (2003):** Treatment of jaundice in low birth weight infants. Arch Dis Child Fetal Neonatal Edition; 88:F459.
- M. Malec, Margareta SÖDERQVIST, ALLAN SIRSJÖ, BARBARA MACNAMARA, NONGNIT LEWIN, JAN SJÖBERG, MAGNUS BJÖRKHOLM, ANNA PORWIT-MACDONALD.(2004).** Real-time polymerase chain reaction determination of cytokine mRNA expression profiles in Hodgkin's lymphoma. haematologica; 89(6):June.
- Mc Donagh, A.F.(2001):** Phototherapy: from ancient Egypt to new millennium". J. Perinatol. Dec; 21 Suppl 1-.S7-S 12.
- Mehta S, Kumar P, Narang A.(2005)** A randomized controlled trial of fluid supplementation in term neonates with severe hyperbilirubinemia.- : J

Pediatr. Dec; 147(6):781-5.

Meldrum, (1989). Tumor necrosis factor in the heart. Am J Physiol Regul Integr Comp Physiol 274: R577-R595,

Melton K, Akinbi HT (1999): Neonatal jaundice. Strategies to reduce bilirubin-induced complications. Post grad med; 106-178.

Michael Kennedy, Kyu Han Kim, Brad Harten, Jeffrey Brown, Stephen Planck Charles Meshul, Henry Edelhauser, James T. Rosenbaum, Cheryl A. Armstrong, and John C. Anselj1 (1997) : Ultraviolet Irradiation Induces the Production of Multiple Cytokines by Human Corneal Cells Investigative Ophthalmology & Visual Science, November, Vol. 38, No. 12

Mori M, Rothman AL, Kurane I, Montoya JM, Nolte KB, Norman JE, Waite DC, Koster FT, Ennis FA.(1999). High levels of cytokine-producing cells in the lung tissues of patients with fatal hantavirus pulmonary syndrome. J Infect Dis 179: 295-302.

Mukai Y, Shibata H, Nakamura T, Yoshioka Y, Abe Y, Nomura T, Taniai M, Ohta T, Ikemizu S, Nakagawa S, Tsunoda S, Kamada H, Yamagata Y, Tsutsumi (2009) Structure-function relationship of tumor necrosis factor (TNF) and its receptor interaction based on 3D structural analysis of a fully active TNFR1-selective TNF mutantJ. Mol. Biol. v385, p.1221-1229

Narang A,Kumar P, Kumar R (2001). Neonatal jaundice in very low birth weight. Indian Pediatr; 68:307-32.

Joanna Narbutt,1 Aleksandra Lesiak,1 Malgorzata Skibinska,2 Anna Wozniacka,1 Anna Sysa-Jedrzejowska,1 Joalanta Lukamowicz,3 and Henk van Loveren4. (2005). Repeated Doses of UVR

- CauseMinor Alteration in Cytokine Serum Levels in Humans. *Mediators Inflamm.* 2005 October 24; (5): 298–303.
- Newman TB, Liljestrand P, Jeremy RJ (2006):** Outcomes among newborn with total serum bilirubin levels of 25mg per deciliter or more. *N Engl J Med.* May 4; 354(18):1974-9.
- O'Malley, W., Achinstein, B. & Shear, M. (1962)** *J. Natl. Cancer inst.* 29, 1169-1175.
- Olusanya BO, Somefun AO. (2009):** Sensorineural hearing loss in infants with neonatal jaundice in Lagos: a community-based study. *Ann Trop Paediatr.* Jun; 29(2):119-28.
- Oshima T, Lee S, Sato A, Oda S, Hirasawa H, Yamashita T. (2009).** TNF-alpha contributes to axonal sprouting and functional recovery following traumatic brain injury. *Brain Res.* 22; 1290:102-10. Epub Jul 16.
- Pan S, An P, Zhang R, He X, Yin G, Min W. (2002)** Etk/Bmx as a tumor necrosis factor receptor type 2-specific kinase: role in endothelial cell migration and angiogenesis. *Mol Cell Biol.* Nov;22(21):7512-23.
- Pelton BK, HyltonW, Denman AM. (1992)** Activation of IL-6 production by UV irradiation of blood mononuclear cells from patients with systemic lupus erythematosus. *Clin Exp Immunol* ;89:251–4.
- Pezzati M, Biagiotti R, Vangi V, Lmbardi E, Wiechmann L, Rubatelli F(2000):** Changes in mesenteric blood flow response to feeding : conventional versus FO phototherapy. *Pediatr,* 105: 350-53.
- Sabry A, Sheashaa H, El-Husseini A, Mahmoud K, Eldahshan KF, George SK, Abdel-Khalek E, El-Shafey EM, Abo-Zenah H.(2006):** Proinflammatory cytokines (TNF-alpha and IL-6) in Egyptian

- patients with SLE: its correlation with disease activity. *Cytokine*. Aug;35(3-4):148-53. Epub Oct 5
- Sarin M, Dutta S, Narang A (2006):** Randomized controlled trial of compact fluorescent lamp versus standard phototherapy for the treatment of neonatal hyperbilirubinemia. *Ind Pediatr*; 43: 583-90.
- Sarici, S.U., Alpay, F., Unay, B-, Ozean, O. and Gokcay,E. (2000):** Double versus single phototherapy in term newborns with significant hyperbilirubinemia. *J. Trop. Pediatr*.Feb.46 (1): 36-9.
- Sarsat V.W., Rieu P., Descamps - Latscha B., Lesavre P. and Halbwachs - Mecarelli L. (2000):** Neutrophils: Molecules, Functions and Pathophysiological Aspects. *Laboratory Investigation*; 80: 617-653.
- Schulz U, Munker R, Ertl B, Holler E, Kolb HJ. (2001):** Different types of human leukemias express the message for TNF-alpha and interleukin-10. *Eur J Med Res*. Aug 27;6(8):359-63.
- Shapiro SM (2003):** Bilirubin toxicity in the developing nervous system. *Pediatric neurology* 29(5): 410-421.
- Shapiro SM (2005):** Definition of the clinical spectrum of kernicterus and bilirubin induced neurologic dysfunction (BIND). *J Perinatol* Jan; 25(1):54-9.
- Sheu W., Lee W. J., Chang R.L. and Chen Y.T.(2000):** Plasma TNF-a levels and insulin sensitivity in hypertensive subject. *Clin Exp Hypertens*; 22(6): 595-606.
- Shibata H, Yoshioka Y, Ohkawa A, Minowa K, Mukai Y, Abe Y, Tanai M, Nomura T, Kayamuro H, Nabeshi H, Sugita T, Imai S, Nagano K, Yoshikawa T, Fujita T, Nakagawa S, Yamamoto A, Ohta T, Hayakawa T, Mayumi T, Vandenabeele P, Aggarwal BB, Nakamura T, Yamagata Y, Tsunoda S, Kamada H, Tsutsumi Y. (2008)** Creation and X-ray structure analysis of

the tumor necrosis factor receptor-1-selective mutant of a tumor necrosis factor-alpha antagonist. *J Biol Chem.* Jan 11; 283(2):998-1007. Epub Nov 14.

Shibata H, Yoshioka Y, Ohkawa A, Abe Y, Nomura T, Mukai Y, Nakagawa S, Taniai M, Ohta T, Mayumi T, Kamada H, Tsunoda S, Tsutsumi Y (2008) The therapeutic effect of TNFR1-selective antagonistic mutant TNF-alpha in murine hepatitis models. *Cytokine.* Nov; 44(2):229-33. Epub Sep 23.

Shu H.B., Takeuchi M. and Goeddel D.V. (1996):Tumour necrosis factor signaling. *Proc Natl Acad Sci*; 93: 13973-13978.

Sirota L, Straussberg R, Gurary N, Aloni D, Bessler H (1999): Phototherapy for neonatal hyperbilirubinemia affects cytokine production by peripheral blood mononuclear cells. *Eur J Pediatr*; 158: 910–913.

Smith J (2006): Bilirubin, Impaired Conjugation. Extracted from:<http://www.emedicine.com/ped/neonatology>. Last updated Jul 24. Extracted on November.

Springer SC (2006): Kernicterus. Extracted from <http://www.emedicine.com/ped/neonatology> Last updated Jul 31,. Extracted on January.

Stokowski LA, Short M, Witt CL (2006): Fundamentals of phototherapy for neonatal jaundice. *Adv neonat care*; 6 (6): 303-12.

Stoll JB, Kliegman RM: Digestive system disorders; in Behrman RE, Kliegman RM, Jenson HB (eds)(2004): Nelson Textbook of Pediatrics, ed 17. Philadelphia, Saunders, pp 588–599.

Tan, K.L. and Boey, K.W. (1986): Efficacy of phototherapy in nonhaemolytic hyperbilirubinemia. *Br. Med. J.* 293:1361.

References

- Tan, K.L. (1991):** Phototherapy for neonatal jaundice. Clinics in perinatology 18: 423-39.
- Tan, K.L. (1996):** Phototherapy for neonatal jaundice. Acta. Pediatr. Mar. 85(3):277-9.
- Thaithumyanon P, Visutiratmanee C. (2002):** Double phototherapy in jaundiced term infants with hemolysis. -J Med Assoc Thai.Nov; 85(11):1176-81.
- Thor WR Hansen, MD, PhD, MHA, Professor, Section of Neonatology, Department of Pediatrics, Rikshospitalet, University of Oslo Contributor Information and Disclosures updated: Oct 18, 2007
- Tracey K.J. and Cerami A. (1993):** TNF in malnutrition and cancer. Am J Trop Med Hyg; 49: 2-17.
- Trentin L., Pizzolo G., Zambello R., Agostini C, Moroso L., Sancetta R., Adami F., Vinante F., Chilosi M. and Gallati H. (1995) :** Leukemic cell in hairy cell leukemia and B-CLL release soluble TNF receptors. Leukemia; 9(6): 1051-1055.
- Tsigos C, Kyrou I., Chala E., Tsapogas P., Stavridis J.C., Raptis S.A. and Kalsilambros N. (1999):** Circulating TNF-a concentrations are higher in abdominal versus peripheral obesity. Metabolism; 48 (10): 1332-1335.
- Van Deuren M. (1998):** Kinetics of TNF-a, soluble TNF receptors, IL-1B, and its receptor antagonist during serious infection. Eur I. Clin Microbiol; 20: 5-12.
- Vreman HJ, Wong RJ, Stevenson DK (2004):** Phototherapy: current methods and future directions. Sem Perinatol; 28:326-33

- Wagle S (2006):** Hemolytic Disease of Newborn. Extracted from
<http://www.emedicine.com/ped/neonatology> Last updated Jun 22,
Extracted on January.
- Wajant H, Pfizenmaier K, Scheurich P (2003).** "Tumor necrosis factor signaling".
Cell Death Differ. 10 (1): 45–65
- Walther FJ, Wu PY, Siassi B (1985):** Cardiac output changes in newborns with
hyperbilirubinemia treatment with phototherapy. *Pediatr*; 76:918-21.
- Weissman A, Berkowitz E, Smolkin T, Blazer S. (2009):** Effect of phototherapy
on neonatal heart rate variability and complexity. *Neonatology*.
95(1):41-6.
- Wellmer, A., J. Gerber, J. Ragheb, G. Zysk, T. Kunst, A.Smirnov, W. Bruck,
and R. Nau. 2001.** Effect of deficiency of tumor necrosis factor alpha
or both of its receptors on *Streptococcus pneumoniae* central nervous
system infection and peritonitis. *Infect. Immun.* 69:6881–6886.
- Witt H. (2003):** Chronic pancreatitis and cystic fibrosis. *Gal*, 52:31.
- Wong V, Chen WX, Wong KY (2006):** Short- and Long-Term Outcome of Severe
Neonatal Nonhemolytic Hyperbilirubinemia. *J Child Neurol*;
21(4):309-15.
- Yamamoto Y. and Gaynon R.B. (2001):** Therapeutic potential of inhibition of NF-
kB pathway in the treatment of inflammation and cancer. *J Clin Invest*;
107 (2): 135-142.
- Zecca E, Barone G, De Luca D, Marra R, Tiberi E, Romagnoli C. (2009)** Skin
bilirubin measurement during phototherapy in preterm and term newborn
infants. - *Early Hum Dev.* May 28.