

REFERENCECES

- Abu-Haweleh AF.*** Acute renal failure in newborn: etiology and mortality rate in Jordan patients. Saudi J Kidney Dis Transpl. 1998;9:18-21.
- Agras PI, Tarcan A, Baskin E et al. (2004),*** Acute renal failure in the neonatal period. Ren Fail;26:305-9.
- Airede A, Bello M, Weerasinghe HD (1997),*** Acute renal failure in newborn: Incidence and outcome. J Pediatr Child Health;33(3):246-9.
- Anand SK (1995):*** acute renal failure in neonates Ped.Clin.North, Am.J; 9; 361-368.
- Anand SK, Northway JD, Smith JA (1977),*** Neonatal renal papillary and cortical necrosis. Am J Dis Child 131:773-777.
- Andreoli SP (2004),*** Acute renal failure in the newborn. Semin Perinatol;28:112-23.
- Andreoli SP (2002),*** Acute renal failure. Current Opinion in Pediatrics 17:713-717.
- Andreoli SP (2001),*** Acute and chronic renal failure in children, in Gearhart JP, Rink RC, Mouriquand PDE (eds):pediatricUrology, Philadelphia, PA, Saunders, pp 777-789.

Andreoli SP (2000), Renal failure in the newborn unit, in Polin RA, Yoder MC, Berg FD (eds): Workbook in Practical Neonatology. Philadelphia, PA, Saunders, pp 322–337.

Andreoli SP, Dunson JW, Bergstein JM (1987), Calcium carbonate is an effective phosphorus binder in children with chronic renal failure. Am J Kidney Dis 9:206-210.

Annabelle NC and Minnie MS(2005), Acute renal failure management in the neonate. NeoReviews, 6: No.8 e369 - e376.

Arnold W, Holliday M. (1976): Renal failure in the newborn. Read before the American Society of Pediatric Nephrology, St. Louis, April.

Avery GB, Flectcher MA, Mac Donald MG (2005), Neonatology: Pathophysiology and management of newborn. 6th ed. Philadelphia. Lippincott Williams & Wilkins;2:1000-9.

Badr KF and Ichikawa I (1988), Prerenal failure: A deleterious shift from renal compensation to decompensation. N Engl J Med 319:623-628

Baluarte HJ, Gruskin AB, Hiner LB et al (1977), Encephalopathy in children with chronic renal failure. Proc Clin Dial Transplant Forum;7:95–8.

Barratt TM, Avner ED, Harman WE, eds (1999), Pediatric nephrology. 4th ed. Baltimore, Md: Lippincott Williams & Wilkins.

Barrington K and Brion LP (2002), Dopamine versus no treatment to prevent renal dysfunction in indomethacin-treated preterm newborn infants. Cochrane Database of Systematic Reviews, Issue 3. Art. No.:CD003213.

Barza M, Ioannidis JP, Cappelleri JC, et al (1996), Single or multiple doses of aminoglycosides. A Meta Analysis. BMJ 312:338-345.

Basile DP, Donohoe D, Roethe K, et al (2001), Renal ischemic injury results in permanent damage to peritubular capillaries and influences long-term function. Am J Physiol 281:F887-F899.

Brater DC (1999), Effects of nonsteroidal anti-inflammatory drugs on renal function: Focus on cyclooxygenase-2 selective inhibitors. Am J Med 107:65S-71S.

Brenner BM, Lawler EV, Mackenzie HS (1996), The hyperfiltration theory: A paradigm shift in nephrology. Kidney Int 49:1774-1777.

Brion LP, Bernstein J, Spitzer A (1997), : Kidney and urinary tract. In Neonatal- perinatal medicine: Diseases of the Fetus and Infant, Fanaroff AA, Martin RJ eds. St. Louis, Mosby-Year Book Inc, pp 1586.

Bunchman TE, Wood EG, Schenck MH, et al (1991), Pretreatment of formula with sodium polystyrene sulfonate to reduce dietary potassium intake. Pediatr Nephrol 5:29- 32.

Cataldi L, Leone R, Moretti U, et al (2005), Potential risk factors for the development of acute renal failure in preterm newborn infants: a case-control study. Arch Dis Child Fetal Neonatal Ed;90:F514-9.

Chawla D, Agarwal R, Deorari AK et al (2008), Fluid and electrolyte management in term and preterm neonates. AIIMS-NICU protocols 2008, www.newbornwhocc.org

Chevalier RL (1991), What treatment do you advise for bilateral or unilateral renal thrombosis in the newborn, with or without thrombosis of the inferior vena cava? Pediatr Nephrol 5:679.

Chevalier R (1984), Prognostic factors in neonatal acute renal failure. Pediatrics; 74: 165-272.

Christensen AM, Daouk GH, Norling LL, et al (1999), Postnatal transient renal insufficiency in the feto-fetal transfusion syndrome. Pediatr Nephrol 13:117-120.

Cifuentes RF, Olley PR, Balfe JW, et al (1979), Indomethacin and renal function in premature infants with persistent patent ductus arteriosus. J Pediatr 95:583-587.

Coulthard MG and Vernon B (1995), Managing acute renal failure in very low birthweight infants. Arch Dis Child; 73:F187–92.

Cuzzolin L, Fanos V, Pinna B, et al (2006), Postnatal renal function in preterm newborns: a role of diseases, drugs and therapeutic interventions. *Pediatr Nephrol*;21:931-8.

David Moskowitz, (2003), Personal communication. Rajesh, V.T. Manual of neonatal care / Joint program in Neonatology, Harvard Medical School, – 5th editions.

Denton MD, Chertow GM, Brady HR (1996), “Renal-dose” dopamine for the treatment of acute renal failure: Scientific rationale, experimental studies and clinical trials. *Kidney Int* 49:4-14.

Diamond JR and Yoburn DC (1982), Nonoliguric acute renal failure. *Arch Intern Med.* Oct; 142(10):1882-4.

Doherty E G, and Simmons C F Jr. (2008): Fluid and electrolyte management. In Cloherty J P, Eichenwald E C, and Stark A R (eds.): *Manual of neonatal care*, sixth ed. Lippincot Williams & Wilkins; 100-113.

Drukker A and Guignard JP (2002), Renal aspects of the term and preterm infant: a selective update. *Curr Opin Pediatr* 14:175-182.

Ellis D, Kaye RD, Bontempo FA (1997), Aortic and renal artery thrombosis in a neonate: Recovery with thrombolytic therapy. *Pediatr Nephrol* 11:641-644.

Ellis EN and Arnold WC (1982), Use of urinary indices in renal failure in the newborns. Am J Dis Child 136:615.

Feld LG, Springate JE, Feldes RD (1986), Acute renal failure: II. Pathophysiology and diagnosis. J Pediatr 109:567.

Friedlich PS, Evans JR, Tulassay T, et al (2005), Acute and chronic renal failure. In: Taeusch HW, Ballard RA, Gleason CA. 8th ed. Avery's diseases of the newborn. Philadelphia: Elsevier Saunders; p. 1298-305.

Gallego N, Perez-Caballero C, Estepa R, et al (2001), Prognosis of patients with acute renal failure without cardiomyopathy. Arch Dis Childhood 84:258-260.

Gerstman BB, Kirkman R, Platt R (1992), Intestinal necrosis associated with postoperative orally administered sodium polystyrene sulfonate in sorbitol. Am J Kid Dis 20:159-161.

Gharehbaghi MM and Peirovifar A MD (2007), Evaluating causes of ARF in newborn infants, Pak J Med Sci Vol. 23 No. 6

Golej J and Kitzmueller E (2002), Low-volume peritoneal dialysis in 116 neonatal and paediatric critical care patients. Eur. J. Pediatr. Jul; 161(7):385-9. Epub 2002 May 09.

- Golej J, Kitzmueller E, Herman M, et al (2002)**, Low-volume peritoneal dialysis in 116 neonatal and paediatric critical care patients. Eur J Pediatr 161:385-389.
- Gouyon J B and Guignard J P (2000)**, Management of acute renal failure in newborns. Pediatr Nephrol,14:1037-1044.
- Gupta B.D, Pramod Sharma, Jyoti Bagla ,et al (2005)**, Renal Failure in Asphyxiated Neonates Indian Pediatr;42:428-934.
- Hammarlund K and Sedin G (1979)**, Transepidermal water loss in newborn infants. III. Relation to gestational age. Acta Paediatr Scand;68:795–801.
- Haycock GB (2003)** Management of acute and chronic renal failure in the newborn. Semin Neonatol;8:325-34.
- Haycock GB (1998)**, Development of glomerular filtration and tubular sodium reabsorption in the human fetus and newborn. Br J Urol;81:33–8.
- Hencz P, Deverall PB, Crew AD et al (1979)**, Hyperuricemia of infants and children: a complication of open heart surgery. J Pediatr;94:774–6.
- Hentschel R, Lodige B, Bulla M (1996)**, Renal insufficiency in the neonatal period. Clin Nephrol: 46:54–8.

- Heyman E, Morag I, Batash D et al (2003)**, Closure of patent ductus arteriosus with oral ibuprofen suspension in premature newborns: a pilot study. *Pediatrics* 112:354–358
- Humes HD (1988)**, Aminoglycoside nephrotoxicity. *Kidney Int* 33:900-911.
- Itabashi K, Ohno T, Nishida H (2003)**, Indomethacin responsiveness of patent ductus arteriosus and renal abnormalities in preterm infants treated with indomethacin. *J pediatr* 143:203-207.
- Jamro S and Abbasi KA (2000)**, Acute renal failure in neonates: clinical presentation, cause and outcome. *Pak Paed J*;24:57-60.
- Jayashree G, Dutta AK, Sarna MS et al (1991)**, Acute renal failure in asphyxiated newborns. *Ind J Pediatr*; 28: 19-23.
- John P. and Ann R.Starket (1997)**, Manual of neonatal care / Joint program in Neonatology, Harvard Medical school 4th edition.
- Kaneko K, Suzuki Y, Fukuda Y et al (1995)**, Abnormal contralateral kidney in unilateral multicystic dysplastic kidney disease. *Pediatr Radiol*; 25:275–277.
- Karlowicz MG and Adelman RD (1995)**, Nonoliguric and oliguric acute renal failure. *Pediatr Nephrol* 9:718-722.
- Karlowicz MG and Adelman RD (1992)**, Acute renal failure in the newborn. *Clin Perinatol* 19:139.

Karmazyn B and Zerlin JM (1997), Lower urinary tract abnormalities in children with multicystic dysplastic kidney. *Radiology*; 203:223–226.

Kellum JA (1998), Use of diuretics in the acute care setting. *Kidney Int* 53:67-70.

Kraus RA, Gaisie G, Young LW (1990), Increased renal parenchymal echogenicity: causes in pediatric patients. *RadioGraphics*; 10:1009–1018.

Malone TA (1991), Glucose and insulin versus cation-exchange resin for the treatment of hyperkalemia in very low birth weight infants. *J Pediatr* 118:121-123.

Martin-Ancel A, Garcia-Alix A, Gaya F, et al (1995), Multiple organ involvement in perinatal asphyxia. *J Pediatr* 127: 786-793.

Mathews DE, West KW, Rescorla FJ, et al (1990), Peritoneal dialysis in the first 60 days of life. *J Pediatr Surg* 25:110 116.

Mathew OP, Jones AS, James E, et al (1980), Neonatal renal failure: Usefulness of diagnostic indices. *Pediatrics* 65: 57-60.

Mathur NB, Agarwal HS, Maria A (2006), Acute renal failure in neonatal sepsis. *Indian J Pediatr*;73:499-502.

- Mattoo TK and Ahmad GS (1994)**, Peritoneal dialysis in neonates after major abdominal surgery. *Am J Nephrol*;14:6–8.
- McClure RJ, Prasad VK, Brocklebank JT (1994)**, Treatment of hyperkalaemia using intravenous and nebulised salbutamol. *Arch Dis Child*;70:126–8.
- McCourt M (1996)**, Acute renal failure in the newborn. *Crit Care Nurse*; 16:84–94.
- Mocan H, Beattie TJ, Murphy AV (1991)**, Renal venous thrombosis in infancy: Long-term follow-up. *Pediatr Nephrol* 5:45-49.
- Mogal NE ,Brocklebank JT, Meadow SR (1998)**, A review of acute renal failure in children: incidence, etiology and outcome. *Clin Nephrol* 49:91-95.
- Mortazavi Fakhrossadat, Sedigeh Hosseinpour Sakha, Nayyer Nejati (2009)**, Acute Kidney Failure in Neonatal Period; *Iranian Journal of Kidney Diseases*, Volume 3 , Number 3 . *IJKD*;3:136-40.
- Murdoch IA, Dos Anjos R, Haycock GB (1991)**, Treatment of hyperkalaemia with intravenous salbutamol. *Arch Dis Child*; 66:527–8.
- Norman ME, Asadi FK. A, (1979)**, prospective study of acute renal failure in the newborn infant. *Pediatrics*; 63: 475-479.

- Nouri S, Mahdhaoui N, Beizig S, et al (2008).** [Acute renal failure in full term neonates with perinatal asphyxia. Prospective study of 87 cases]. Arch Pediatr;15:229-35. French.
- Pereira S, Pereira BJB, Bhakoo ON et al (1988),** Peritoneal dialysis in neonates with acute renal failure. Ind. J. Pediatr, 58: 973-978.
- Perlman JM and Tack ED 1988:** Renal injury in the asphyxiated newborn infant: Relationship to neurologic outcome. J Pediatr 113:875-879.
- Philippe SF Jacquelyn RE, Tivadar T et al (2005),** Avery's diseases of newborn In: Taeusch HW, Ballard RA, and Gleason CA. (ed) Acute and chronic renal failure, 8 edition, Philadelphia: Saunders; 1298-1306.
- Polito C, Papale MR, LaManna AL (1998),** Long-term prognosis of acute renal failure in the full term newborn. Clin Pediatr 37:381-386.
- Ravindra LM, John AK, Sudhir VS et al (2007),** Acute Kidney Injury Network: report of an initiative to improve outcomes in acute kidney injury. Critical care, 11:R31.
- Riesenfeld T, Hammarlund K, Sedin G (1995),** Respiratory water loss in relation to gestational age in infants on their first day after birth. Acta Paediatr;84:1056-9.
- Risdon RA and Woolf AS (1998),** Development of the Kidney. In: Hepinstall's pathology of the kidney, Olson and Schwartz (eds), 5th ed., Lippincott-Raven, New York.

Roberts DS, Haycock GB, Dalton RN et al (1990), Prediction of acute renal failure after birth asphyxia. Arch Dis Child; 65:1021–8.

Rooks VJ and Lebowitz RL (2001), Extrinsic ureteropelvic junction obstruction from a crossing renal vessel: demography and imaging. Pediatr Radiol; 31:120–124.

Salah Al-Din D, El-Badawy A, Helmy S et al (2010), Acute kidney injury in neonates , M.sc thesis, Faculty of medicine, Cairo University.

Seri I (1995), Cardiovascular, renal, and endocrine actions of dopamine in newborns and children. J Pediatr 126:333.

Seri I (1993), Effects of low dose dopamine infusion on cardiovascular and renal functions cerebral blood flow and plasma catecholamines levels in sick preterm neonates. Pediatric Res, 34:742-749.

Siegel MJ, ed (1995), Pediatric sonography. 2nd ed. New York, NY: Raven.

Soliman A, Abou El-Ela I, Abd El-Rahman A, (2009), Evaluation of occurrence of ARF and factors associated with it in neonatal sepsis, M.sc thesis, Faculty of medicine, Benha University.

Stapleton FB, Jones DP, Green RS (1987), Acute renal failure in neonates: Incidence, etiology and outcome. Pediatr Nephrol 1:314-320.

Steele BT, Paes B, Towell ME et al (1988), Fetal renal failure associated with intrauterine growth retardation Am J Obstet Gynecol. Nov; 159(5):1200-2.

Subramanian S, Agarwal R, Deorari AK, et al (2008), Acute renal failure in neonates. Indian J Pediatr;75:385-91.

Suchy FJ (1996), Cystic diseases of the biliary tract and liver. In: Behrman RE, Kliegman RM, Arvin AM, Nelson WE, eds. Nelson textbook of pediatrics. 15th ed. Philadelphia, Pa: Saunders; 1153.

Suhas M Nafday (1998), In Renal Disease – Avery's Neonatology pathophysiology and management of newborn, 6th e, editors M G MacDonald; Lippincott Williams and Wilkins. 981-1065.

Tack ED and Perlman JM (1998), Renal failure in sick hypertensive premature infants receiving captopril therapy. J Pediatr 112:805-810.

Thomas DFM, ed (1997), Urological disease in the fetus and infant: diagnosis and management. Oxford, England: Butterworth-Heinemann.

Tiker F and Yildirim SF (2007), Acute renal impairment after oral ibuprofen for medical closure of patent ductus arteriosus. Indian Pediatr 44:54–55.

Toth-Heyn P, Drukker A, Guignard J P (2000), The stressed neonatal kidney; from pathophysiology to clinical management of neonatal vasomotor nephropathy. *Pediatr Nephrol*, 14:227-239.

Trichtman H (2004), Sodium & water : pediatric Nephrology 5th ed Williams & Wikins.

Tulassy T and Seri I (1986), Acute oliguria in preterm infants with hyaline membrane disease; interaction of dopamine and frusemide. *Acta Pediatr Scand*, 75:420-424.

Van Bel F, Guit GL, Schipper J, et al (1991), Indomethacin induced changes in renal blood flow velocity waveform in premature infants investigated with color Doppler imaging. *J Pediatr* 118:621-626.

Vanpee M, Blennow M, Linne T, et al (1992), Renal function in very low birth weight infants: Normal maturity reached during childhood. *J Pediatr* 121:784-788.

Vogt BA and Avner ED(2006), The kidney and urinary tract. In: Fanaroff AA, Martin RJ. *Neonatal-Perinatal Medicine: Diseases of Fetus and Newborn Infant*. 8th ed. St, louis Mosby;2:1666-70.

Watanabe T (2001), Rhabdomyolysis and acute renal failure in children. *Pediatr Nephrol* 16:1072-1075.

Wood EG, Bunchman TE, Lynch RE (1991), Captopril-induced reversible acute renal failure in an infant with coarctation of the aorta. Pediatrics 88:816-818.

Young TE and Mangum OB (2002), Neofax. A manual of drugs used in neonatal care, 11th edn. Acorn Publ, Raleigh, N.C.