

Araki C, Hayashi N, Moriyama M, Morinobu S, Mukai M, Koshiba M, Kawano S, Kumagai S (2004): Usefulness of anti-cyclic citrullinated peptide antibodies (anti-CCP) for the diagnosis of rheumatoid arthritis. *Rinsho Byori*. Dec;52(12):966-72.

Arnett FC, Edworthy SM, Bloch DA, (1988) : The American Rheumatism Association of 1987 revised criteria for the classification of RA, *Arthritis Rheum* 31 : 315 - 324.

Auger I, Escola JM, Gorvel JP, Roudier J. (1996): HLA-DR4 and HL-DR10 motifs that carry susceptibility to rheumatoid arthritis bind 70-kD heat shock proteins. *Nature Medicine*, 2:306-10.

Avcin T, Cimaz R, Falcini F, Zulian F, Martini G, Simonini G, Cecchini G, Borghi M, and Meroni P. (2002): Prevalence and clinical significance of anti-cyclic citrullinated peptide antibodies in juvenile idiopathic arthritis. *Ann Rheum Dis*, 61: 608-611.

Ball J. (1952): Sheep cell agglutination test for Rheumatoid arthritis, Clinico - pathological study . *Ann Rheum Dis*. Jun ; 11(2) : 97 - 111.

Berry R, and Huskisson M. J. (1972): Treatment of rheumatoid arthritis. *Clinical Trials J*, 9: 13-15.

Bienenstock H. (1975): Rheumatoid planter synovial cysts. Ann Rheum Dis, 34:98.

Bizzaro N, Mazzanti G, Tonutti E, Villalta D, and Tozzoli R. (2001): Diagnostic accuracy of the anti-citrulline antibody assay for rheumatoid arthritis. Clin Chem. Jun; 47(6): 1089-93.

Bonfiglio T, and Atwater EC. (1969) : Heart disease in patients with seropositive RA. Arch Intern Med, 124:714.

Brewerton, D. A. (1957): Hand deformities in rheumatoid disease. Ann Rheum Dis, 16:183.

Campbel M. J, and Machin D. (1993): Statistical inference. In medical statistics. A common sense approach, Campel, M. J, and Machin D. (Eds). John Wily, Sosh Chischessler, New York, Brisbane, Toronto.

Caplan A. (1953): Certain unusual radiographic appearances in the chest of coal miners suffering from RA. Thorax 8:29.

Chamberlain A, Corbett M. (1970): Carpel tunnel syndrome in early rheumatoid arthritis. Ann Rheum Dis, 29:149-52.

- Chang LW, Gowans JDC, Granger CV. (1972):** Entrapment neuropathy of the posterior interosseous nerve. *Arthritis rheum*, 15:350-2.
- Cohen IR, Holschits J, and Van Eden W. (1985):** T- lymphocyte clones illuminate pathogenesis and effect of therapy of experimental arthritis. *Arth Rheum*, 28: 841-5.
- Combe B, Andary M, Caraux J. (1986):** Characterization of an expanded sub-population of large granular lymphocytes in a patient with RA. *Arthritis Rheum*, 29:675.
- Comstock GW, Burke AE, Hoffman SC. (1997):** Serum concentrations of tocopherol, carotene, and retinal preceding the diagnosis of rheumatoid arthritis and systemic lupus erythematosus. *Ann Rheum Dis*, 56:457–64.
- Cooper C, Coupland C, and Mitchell M. (1995):** Rheumatoid arthritis, corticosteroid therapy and hip fracture. *Ann Rheum Dis*, 54; p49-52.
- Copeman W. S. C. (1963):** Rheumatoid oto-arthritis. *Br Med J*, 2:1536.
- Dacie JV. and Lewis SM. (1975):** *Practical Hematology* , 5th ed Churchill Living stone , Edinburg .

Rycke L, Verhelst X, Kruithof E. (2004): Rheumatoid factor, but not anti-citrullinated protein antibodies, is modulated by infliximab treatment in Rheumatoid Arthritis. *Ann Rheum Dis* doi: 10.1136/ard 023523

Deighton CM, Walker DJ, Griffiths ID, Roberts DF. (1989): The contribution of HLA to rheumatoid arthritis. *Clin Gen*, 36:178–82.

DeSmet A. A, Ting Y. M, and Weiss J. J (1975): Shoulder swelling in rheumatoid arthritis. *Diagn Radiol*,116:601.

Dixon A. St. J. (1971): The rheumatoid foot. In Hill A. G. S. (Ed): *Modern trends in Rheumatology*. Vol. 2. London, Butterworths, PP.158-173.

Dubucquoi S, Solau-Gervais E, Lefranc D, Marguerie L, Sibilia J, Goetz J, Dutoit V, and Prin L. (2004): Evaluation of anti-citrullinated filaggrin antibodies as hallmarks for the diagnosis of rheumatic diseases. *Ann Rheum Dis*, 63: 415-419.

Duke-Elder S, Soley RE. (1976): Summary of systemic ophthalmology. In Duke-Elder S, ed. *System of ophthalmology* VOL XV. St. Louis: CVMosby, 139.

- Ericson S, and Lundberg M. (1967):** Alterations in the tempromandibular joints at various stages of rheumatoid arthritis. *Acta Rheuma Scand*, 13: 257.
- Fassbender HG. (1983):** Histopathologic basis of articular cartilage destruction in rheumatoid arthritis. *Coll Relat Res*, 3:141-5.
- Firestein GS. (2001):** Etiology and pathogenesis of rheumatoid arthritis. In: *Kelly's Textbook of Rheumatology*; Ruddy S, Harris E, Sledge C; 6th ed., W. B. Saunders, Philadelphia, PA,:921-966.
- Firestein GS, and Zvaifler NJ. (1990):** How important are T-cells in chronic rheumatoid synovitis? *Arthritis Rheum*, 33 : 768-73.
- Fisher, RA. (1964):** Statistical methods of research workers 10th ed., Edinburgh, Pliver and Boyed.
- Flatt A. E. (1959):** Surgical rehabilitation of the arthritic hand. *Arthritis Rheum*,11:278.
- Forslind K, Ahlmen M, Eberhardt K, Hafstrom I, and Svensson B. (2004):** Prediction of radiological outcome in early rheumatoid arthritis in clinical practice: role of antibodies to citrullinated peptides (anti-CCP). *Ann Rheum Dis*, 63: 990-95.
- Gan SQ, McBride OW, Idler WW, Nediaalka M, Steinert PM.**

(1990): Organization, structure, and polymorphisms of the human profilaggrin gene. *Biochemistry*, 29:9423-9440.

Gardner D. L, Duthie J. R, Macleod J, and Allan W. S. H. (1957): Pulmonary hypertension in RA: Report of a case study with intimal sclerosis of pulmonary and digital arteries. *Scott. Med, J.* 2: 183.

Genevay S, Hayem G, Verpillant P. (2002): An 8-year prospective study of outcome predilection by antinuclear factor and antikeratin antibodies at onset of RA (in process citation). *Ann Rheum Dis*, p734-6.

Gerber NJ. And Dixon AS TJ. (1974): Synovial cysts and juxta-articular bone cyst. *Semin Arthritis Rheum.* 3:323-48.

Girbal-Neuhauser E, Durieux J-J, Arnaud M, Dalbon P, Sebbag M, Vincent C, Simon M, Senshu T, Masson-Bessière C, Jolivet-Reynaud C, Jolivet M, Serre G. (1999): The epitopes targeted by the rheumatoid arthritis-associated antifilaggrin autoantibodies are posttranslational generated on various sites of (pro) filaggrin by deimination of arginine Residues. *J Immunol*, 162:585-594.

Goupille P, Fouquet B, Goga D, Mateu J, and Valat J-P. (1990):

The tempromandibular joint in rheumatoid arthritis: Correlations between clinical and computed tomography features. *J Rheumatol*, 17: 1285.

Grahame, R, Armstrong R, Simmons N, Wilton J. M. A, Dyson M, Laurent R, Millis R, and Mims C. A. (1983): chronic arthritis associated with the presence of intrasynovial rubella virus. *Ann Rheum Dis*, 42:2.

Green D, Schutte PT, Wallace WH. (1980): Factor VIII antibodies in RA: Effect of cyclophosphamide. *Arch Intern Med*, 140:1232.

Gregersen PK, Silver J, Winchester RJ (1987): The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis. *Arthritis Rheum*, 30:1205–13.

Gupta P. J. (1970): Physical examination of the arthritis patient. *Bull Rheum Dis*, 20:596.

Haase A. T. (1986): pathogenesis of lentiviruses infections. *Nature* 322:130.

Hara KS, Ballard DJ, Llstum DM. (1990): Rheumatoid pericarditis: clinical features and survival. *Medicine*. 69:81.

Harris ED. (2001): Quoted from text book of Rheumatology 6th

Reference

edition volume II chapter 55: P980 edited by Kelly W.N., Harris E, Ruddy S, Sledge C. Toronto, Montreal, Sydney, Tokyo.

Hassfeld W, Steiner G, Sudnicka-Benke A, Skriner K, Graninger W, Smolen JS. (1995): Autoimmune response to the spliceosome. An immunologic link between rheumatoid arthritis, mixed connective tissue disease, and systemic lupus. *Arthritis Rheum*, 38:777-85.

Hastings DE, Parker SM. (1975): Protrusioacetabuli in rheumatoid arthritis. *Clin Orthop*, 108:76-83.

Hazes JM, Dijkmans BAC, Vandenbroucke JP, Vries RRP, Cats A. (1990): Lifestyle and the risk of rheumatoid arthritis: cigarette smoking and alcohol consumption. *Ann Rheum Dis* 49:980–2.

Hearkens AH, Hiemstra PS, Lafeber GJ, Daha MR, and Breedveld FC. (1989): Anti-endothelial cell antibodies in patients with rheumatoid arthritis complicated by vasculitis. *Clinical and Experimental immunology*, 78:7-12

Reference

Henkind P. (1963): Ocular complication of drug treatment of

rheumatic disorders .Ann Physical Med, 7:258.

Hernandez LA, Rowan RM, Kennedy AC. (1975):
Thrombocytosis in RA: A clinical study of 200 patients.
Arthritis Rheum, 6:635.

Hochberg M. C, Chang R. W, Dwosh I, Lindsey S, Pincus T, and Wolfe F. (1992): the American College Of Rheumatology
1991 revised criteria for the classification of global
functional status in rheumatoid arthritis. Arthritis Rheum.
May; 35(5): 498-502.

Hoet RM, Van Venrooij WJ. (1992): The antiperinuclear factor and
antikeratin antibodies in rheumatoid arthritis.
In Rheumatoid Arthritis. Edited by Smolen J, Kalden J &
Maini RN. Berlin: Springer Verlag; 299-318.

Holmdahl R, Malmstrom V, Vuorio E. (1993) : Autoimmune
recognition of cartilage collagens. Ann Med, 25:251-64.

Hurd ER (1979): Extra-articular manifestations of rheumatoid
arthritis. Semin Arthritis Rheum, 8:151-76.

Reference

Huston K. A, Nelson A. M, and Hunder G. G. (1978): Shoulder
swelling in rheumatoid arthritis secondary to subacromial

bursitis. Arthritis Rheum, 21:145.

Iveson, J. M. I, Hill A. G. S, and Wright V. (1975): Wrist cysts and fistulae: An orthographic study of the rheumatoid wrist. Ann Rheum Dis, 34:388.

Jacob, J. R, Sartoris, D., Kursunoglu, S (1986): Distal interphalangeal joint involvement in rheumatoid arthritis. Arthritis Rheum. 29:10.

Jacob J. R, Weisman, M. H. Murk J. H. (1986): Reversible cause of back pain and sciatica in rheumatoid arthritis: An apophysial joint cyst. Arthritis Rheum, 29:431.

Janssens X, Veys E, Verbruggen G, Declerq L. (1998): The diagnostic significance of the antiperinuclear factor for rheumatoid arthritis. J Rheumatol, 15:1346-50.

Jayson MIV, and Jones DEP. (1971): Scleritis and RA .Ann Rheum Dis, 30:343.

Jorizzo JL, and Daniels JC. (1983): Dermatologic conditions reported in patients with rheumatoid arthritis. J Am Acad Dermatol; 8:439-57.

Kals J, and hall N. (1991): Feedback regulation of antibody production: role in RA. Ann Rheum Dis, 50:833-5.

Reference

Kaye BR, Kaye RL, Bobrove A. (1984): Rheumatoid nodules.

Review of the spectrum of associated conditions and proposal of a new classification, with a report of four seronegative cases. Am J Med, 76:279-92.

Kellgren J. H, and Ball J. (1950): Tendon lesion in rheumatoid arthritis: A clinicopathological study. Ann Rheum Dis, 9:48.

Khong TK, and Rooney PJ. (1982): Manubriosternal joint subluxation in rheumatoid arthritis. J Rheumatol, 9:712-15.

Kim JK, and MH Weisman. (2000): When does rheumatoid arthritis begins and why do we need to know? Arthritis Rheum, 43:473-84.

Klippel J. H, and Paul D. P. (1998): Quoted from text book of rheumatology edited by Klippel J.H and Paul D.P. published by Mofby 2nd edition section 5 RA chapter laboratory test in rheumatic disorder.

Kraag G, Thevathasan EM, Gordan DA, Walker IH. (1976): The hemorrhage crescent sign of acute synovial rupture. Ann
 Reference Intern Med, 85:477-8.

Krane SM, Conco W, Stephenson ML, and Golding MB. (1990): Mechanisms of matrix degradation in rheumatoid arthritis. Ann NY Acad. Sci, 580:340-54.

Kroot E. J, Jong B. W, Leeuwen M. A, and Hoogen F. H. (2000):

The prognostic value of anti-cyclic citrullinated peptide protein antibody in patients with recent onset rheumatoid arthritis. *Arthritis Rheum*, 43:1831-5

Larsen A, Dale K, and Eek M. (1977): Radiographic evaluation of rheumatoid arthritis and related conditions by standard reference films. *Acta Radiol Diag*, 18: 481-491

Lebowitz W. B. (1963): The heart in rheumatoid arthritis: A clinical and pathological study of 62 cases. *Ann Intern Med*, 58:102.

Lee DM, and Schur PH. (2003): Clinical utility of the anti-CCP assay in patients with rheumatic diseases. *Ann Rheum Dis*, 2003; 62:8704.

Lienesch D, Morris R, Metzger A, Debuys P, Sherman K. (2005): Absence of cyclic citrullinated peptide antibody in nonarthritic patients with chronic hepatitis C infection. *J Rheumatol*. Mar; 32(3): 489-93.

Linos A, Worthington JW, O'Fallon WM, Kurland LT. (1980):

The epidemiology of rheumatoid arthritis in Rochester, Minnesota: a study of incidence, prevalence, and mortality. *Am J Epidemiol*, 111:87-98.

Lowthian P. J, and Calin A.(1985): geode development and multiple fractures in rheumatoid arthritis. *Ann Rheum Dis*, 44:130.

MacDonald WJ JR, Crawford MH, Klippel JH.kn (1977): Echocardiographic assessment of cardiac structure and function in patients with RA. *Am J Med*, 63:890.

Maddison P. J, and Bacson P. A. (1974): Vitamin D deficiency, spontaneous features and osteopenia in rheumatoid arthritis. *Br. Med. J.* 2:433.

Maini RN, and Zyberk C. (1988): The significance of CD5+ B-cells in rheumatic diseases. *Scand. J Rheumatol*, 76:237.

Mallya, R.K; and Mace, B.E.W. (1981): the assessment of disease activity in RA using the multivariate analysis. *Rheum. Rehabi.* 20: 14-17.

Martin L, and Benson W. G. (1987): An unusual synovial cyst in
~~Reference~~ rheumatoid arthritis. *J Rheumatol*, 14:139.

Masi AT. (1967): Quoted from Dippe PA, Doehrty M., MacFarlane D. and Maddison P. (1985): RA in *Rheumatological Medicine*, ed. By Dippe PA., Doehrly M., MacFarlane D. and Maddison P. :41-64, Churchil Lvingstone, Edinburg,

London and New York.

Masson-Bessière C, Sebbag M, Durieux JJ. (2000) : in the rheumatoid pannus, Anti-filaggrin autoantibodies are produced by local plasma cells and constitute a higher proportion of IgG than in synovial fluid and serum. Clin Exp Immunol 119, 544-552

Masson-Bessière C, Sebbag M, Girbal- Neuhauser E. (2001): The major synovial targets of the rheumatoid arthritis- specific antifilaggrin autoantibodies are deiminated forms of the alpha- and beta- chains of fibrin. J Immunol 166, 4177-4184.

McCarty DJ, O'Duffy JD, Pearson L, and Hunter JB. (1985): Remitting seronegative symmetrical synovitis with pitting edema. (RS3PE) syndrome. JAMA. 254:3763-276.

Reference

McColl GJ, Carm DS, Harrison LC. (1995): Antibodies to synovial antigens in recent onset rheumatoid arthritis. Arthritis Rheum, 38:1418-1428.

McGuigan, L, Burke D, and Fleming A. (1983): Tarsal tunnel syndrome and peripheral neuropathy in rheumatoid disease.

Ann Rheum Dis, 42:128.

Mediawake R, Isenberg D. A Schellekens G. A, and Van Venrooij W. J. (2001): Use of anti-citrullinated peptide and anti-RA33 antibodies in distinguishing erosive arthritis in patients with systemic lupus erythematosus and rheumatoid arthritis. Ann Rheum Dis.; Jan, 60(1): 67-8.

Menard HA, Paquette D. (1980): Skin temperature of the knee: and unrecognized physical sign of the inflammatory disease of the knee. Can Med Assoc J, 122:439-40.

Meyer O, Labarre C, Dougados M.(2003): Anti-citrullinated protein/peptide antibody assays in early rheumatoid arthritis for predicting five year radiographic damage. Ann Rheum Dis, 62:120-6.

Millis M. B, Millender L. H, and Nalebuff E. A. (1976): Stiffness of the proximal interphalangeal joints in rheumatoid arthritis. J. Bone joint Surg. 58A: 801.

Mody G. M, and Meyers O. L. (1988): Resorptive arthropathy in rheumatoid arthritis. J Rheumatol, 15:1075.

Moffat DA, Ransden RT, Rosenberg JN (1977): Otoadmittance measurments in patients with RA. J Laryngol Otol, 91:917.

Mollenhauer J, Von der mark K Burmester G, Gluckert K,

Lutjen-Dercol E, Brune K. (1988): Serum antibodies against chondrocyte cells surface proteins in osteoarthritis and rheumatoid arthritis J Rheum, 15:1811-7.

Mottonen, T.; Paimela, L.; Leirisalo-Repo, M.; Kautiainen, H.; Ilonen, J. and Hannonen, P. (1988): Only high disease activity and positive rheumatoid factor indicate poor prognosis in patients with early rheumatoid arthritis treated with "sawtooth" strategy. Ann Rheum Dis, 57: 533-9.

Mowat, A. G. (1971): Hematological abnormalities in RA. Semin. Arthritis Rheum. 1:195.

Murayama T, Jisaki F, Ayata M. (1992): Cytomegalovirus genomes demonstrated by polymerase chain reaction in synovial fluid from RA patients. Clinic Exp Rheumatol, 10:161.

Nakano KK, Schoene WC, Baker RA, and Dowson DU. (1978): The cervical myelopathy associated with RA. AM Neuro Assoc., 3:144.

Nakashima K, Hagiwara T, Yamada M. (2002): Nuclear localization of peptidylarginine deiminase V and histone

deimination in granulocytes. *J Biol Chem* 277, 49562-49568.

Nalebuff E. A. (1986): Diagnosis, classification, and management of rheumatoid thumb deformities. *Bull Hosp Joint Dis*, 24:119.

Nepom NT, Nepom B. (1998): Genetics of the major histocompatibility complex in rheumatoid arthritis. In: Klippel J, Dieppe P, eds. *Rheumatology*. London: Mosby-Wolfe, 7.1.

Nielen MM, van Schaardenburg D, Reesink HW, van de Stadt RJ, van der Horst-Bruinsma IE, de Koning MH, Habibuw MR, Vandenbroucke JP, Dijkmans BAC (2004): Specific autoantibodies precede the symptoms of rheumatoid arthritis: a study of serial measurements in blood

Reference

donors. *Arthritis Rheum*, 50:380–386.

Nienhuis RLF, Mandena EA. (1964): A new serum factors in patients with rheumatoid arthritis. The antiperinuclear factor. *Ann Rheum Dis*, 23:302-305.

Ollier W, and MacGregor A. (1995): Genetic epidemiology of rheumatoid disease. *Br Med Bull*, 51:267–85.

Ollier W, Thomson W. (1992): Population genetics of rheumatoid arthritis. *Rheum Dis Clin North Am*, 18:741–9.

Pincus T, and Callahan IF. (1992): Early mortality in RA predicted by poor clinical status. Bull Rheum Dis, 41:4.

Polisar I, Burbank B, Levitt L. M, Katz H. M, and Morrione T. G. (1960): Bilateral midline fixation of cricoarytenoid joints as a serious medical emergency. J.A.M.A.172: 901-6.

Poon AC, Geerling G, Dart JK, Fraenkel GE, Daniels JT. (2001): Autologous serum eye drops for dry eyes and epithelial defects: clinical and in vitro toxicity studies.

Pope RM, Mannik M, Gilliland BC. (1975): The hyperviscosity syndrome in RA due to intermediate complexes formed by self association of IgG-rheumatoid factors. Arthritis Rheum, 18:97.

Radoux V, Menard H. A, Begin R, Decary F, and Koopman W. J. (1987): Airway disease in rheumatoid arthritis patients: One element of a general exocrine dysfunction. Arthritis rheum, 30: 249.

Rantapaa-Dahlquist S , de Jong BA, Berglin E, Hallmans G, Wadell G, Stenlund H.(2003): Antibodies against cyclic citrullinated peptide and IgA rheumatoid factor predict the

development of rheumatoid arthritis. *Arthritis Rheum*, 48:2741–9.

Rask, M. R. (1978): Achilles tendon rupture owing to rheumatoid disease. *J. A. M. A.* 239:435.

Reimer K. A, Rodgers R. F, and Oyasu R. (1976): Rheumatoid arthritis with rheumatoid heart disease and granulomatus aortitis. *J. A. M. A.* 235: 2510.

Reparon-Schuijt CC, van Esch WJE, van Kooten C, Schellekens GA, de Jong BAW, van Venrooij WJ, Breedveld FC, Verweij CL. (2001): Secretion of anti-citrulline-containing peptide antibody by B lymphocytes in rheumatoid arthritis. *Arthritis Rheum*, 44:41-47

Reference

Rimon R. A. (1969): Psychosomatic approach to rheumatoid arthritis: a clinical study of 100 patients. *Acta. Rheum. Scand*, 13 (suppl. 1): 1–154.

Ritchie, D. M.; Boyle, J. A.; Mcinnes, J. M.; Jasani, K. M.; Dalakas, J. G. and Buchanan, W. (1968): Clinical studies with an articular index for the assessment of joint tenderness in patients with rheumatoid arthritis. *Quarterly J. Med*, 37:

393-406.

Roberts WC, Kehoe JA, Carpenter DF. (1968): Cardiac valvular lesions in RA. Arch Intern Med, 122:141.

Romain P, Burmester G, Enlow R. (1982): Multiple abnormalities in immunoregulatory function of synovial compartment in RA. Rheumatol Int., 2:121-7.

Ronnelid J, Lysholm J, Engstrom-Laurent A, Klareskog L. B. (1994): Local anti-type II collagen antibody production in rheumatoid arthritis synovial fluid. Evidence for a HL-DR4-restricted IgG response, arthritis Rheum, 37:1023-9.

Ronnelid J, Wick MC, Lampa J, Lindblad S, Nordmark B, Klareskog L, van Vollenhoven RF.(2005) : Longitudinal analysis of citrullinated protein/ peptide antibodies

(anti-CCP) during 5 year follow up in early rheumatoid arthritis: anti-CP status predicts worse disease activity and greater radiological progression. Ann Rheum Dis. Dec; 64(12): 1744-9. Epub 2005 Apr 20.

Samanci N, Ozdem S, Akbas H, Mutlu D, Gultekin M, Arman M, Donmez L.(2005): Diagnostic value and clinical

significance of anti-CCP in patients with advanced rheumatoid arthritis. J Natl Med Assoc. Aug; 97(8):1120-6.

Schellekens GA, de Jong BAW, van den Hoogen FHJ, van de Putte LBA (1998): Citrulline is an essential constituent of antigenic determinants recognized by rheumatoid arthritis-specific autoantibodies. J Clin Invest, 101:273-281.

Schellekens GA, Visser H, de Jong BAW, van den Hoogen FH, Hazes JM, Breedveld FC, van Venrooij WJ (2000): The diagnostic properties of rheumatoid arthritis antibodies recognizing a cyclic citrullinated peptide. Arthritis Rheum, 43:155-163

Schroder AE, Greiner A, Seyfert C, Berek C. (1996) : Differentiation of B-Cells in the non lymphoid tissue of the synovial membrane of patients with rheumatoid arthritis. **Reference** Proc Nat Acad Sci. (USA): 93:221-5.

Schumacher HR, Klippel JH, and Koopman WJ. (1993): Rheumatoid arthritis; clinical features and laboratory parameters of disease activity. In; Schumacher HR, Klippel JH, and Koopman WJ. (Eds) Primer on the rheumatic disease, 10th Ed, arthritis foundation, (publ.), Atlanta, Georgia; 90-94.

Scott D. L, Grindulis K. A, Coulton B. L, Popert A. F. and Bacon

P. A. (1984): Progression of radiological changes in rheumatoid arthritis. *Ann Rheum Dis*, 43: 8-17.

Sebbag M, Simon M, Vincent C, Masson-Bessière C, Girbal E,

Durieux JJ, Serre G. (1995): The antiperinuclear factor and the so-called antikeratin antibodies are the same rheumatoid arthritis-specific autoantibodies. *J Clin Invest*, 95:2672-2679.

Serre G. (2001): Autoantibodies to filaggrin/deiminated fibrin

(AFA) are useful for the diagnosis and prognosis of rheumatoid arthritis, and are probably involved in the pathophysiology of the disease. *Joint Bone Spine*, 68:103-105.

Reference

Sharpino J. S. (1970): A new factor in the aetiology of ulnar drift.

Clin Orthop., 68:32.

Shimizu S, Shiozawa K, Imura S, and Fujita T. (1985):

Quantitative histological studies on the pathogenesis of periarticular osteoporosis in rheumatoid arthritis. *Arthritis Rheum*, 28:25.

Sibbitt WL JR, Williams RC JR(1982): Cutaneous manifestations of rheumatoid arthritis. *Int J Dermatol*, 21:563-72.

Silman AJ, Kay A, Brennan P (1992): Timing of pregnancy in relation to the onset of rheumatoid arthritis *Rheum Arthritis*, 35; 152.

Simpson R. W, McGinty L, Simon L, Smith C. A, Godzeski C. W, and Boyed R. J.(1984): Association of paroviruses with rheumatoid arthritis of humans. *Science*, 233:1425,

Smoleroff JW. (1943): Scleral disease in RA Report of three cases in one of which both eyes were studied postmortem. *Arch ophthalmol*. 29: 98.

Sokoloff L, McCluskey RT, Bunim JJ. (1953): Vascularity of the early subcutaneous nodule of rheumatoid arthritis. *Arch Pathol*; 55:475-95.

Solanki K, Spellerberg M, Chapman P, Moller P, O'Donnell J. (2004): Anti-cyclic citrullinated antibodies: complementary to IgM rheumatoid factor in the early diagnosis of rheumatoid arthritis. *N Z Med J*. Oct 8;117(1203):U1095.

Stack B. H. R., and Grant I. W. B. (1965): Rheumatoid interstitial lung disease. *Br. J. Dis. Chest* 59:202.

Stevens JC, Cartilage NEF, and Saunders M. (1971): Atlantoaxial

subluxation and cervical myelopathy in rheumatoid arthritis.
Q J Med 40:341.

Swezey R. L. (1967): Myocardial infarction due to rheumatoid arthritis. J. A. M. A. 199:191

Symmons DPM, Barrett EM, Bankhead CR, Scott DGI, Silman AJ (1994): The incidence of rheumatoid arthritis in the United Kingdom: results from the Norfolk Arthritis Register. Br J Rheumatol, 33:735–9.

Trentham, D. E, and Masi A. T. (1976): Carpo:metacarpal ratio: A new quantitative measure of radiological progression of wrist involvement in rheumatoid arthritis. Arthritis Rheum, 19:939.

Vallbracht I, Rieber J, Oppermann M, Forger F, Siebert U, Helmke K. (2004): Diagnostic and clinical value of anti-cyclic citrullinated peptide antibodies compared with rheumatoid factor isotopes in rheumatoid arthritis. Ann Rheum Dis. Sep;63(9):1079-8

Reference

Van Gaalen FA, van Aken J, Huizinga TW. (2004): Association between HLA class II genes and autoantibodies to cyclic citrullinated peptides (CCPs) influences the severity of rheumatoid arthritis. Arthritis Rheum, 50:2113-21.

Van Jaarsveld CHM, ter Borg EJ, Jacobs JWG, Schellekens GA, Gmelig-Meyling FHJ, van Booma-Frankfort C, de Jong BAW, van Venrooij WJ, Bijlsma JWJ.(1999): The prognostic value of the antiperi-nuclear factor,

anti-citrullinated peptide antibodies and rheumatoid factor in early rheumatoid arthritis. *Clin Exp Rheumatol*, 17:689-697.

Van Venrooij WJ. (2000): Citrullination: a small change for a protein with great consequences for rheumatoid arthritis. *Arthritis Res*, 2:249-51.

Vencovsky J, Machacek S, Sedova L, Kafkova J, Gatterova J, and Pesakova V. (2003): Autoantibodies can be prognostic markers of an erosive disease in early rheumatoid arthritis. *Ann Rheum Dis*, May, 62(5):427-30.

Vidigal E, Jacoby R, Dixon A. St. J, Rattiff A. H, and Kirkup J. (1975): The foot in chronic rheumatoid arthritis. *Ann Rheum Dis*, 34:292.

Vincent C. (1989): High diagnostic value in rheumatoid arthritis of antibodies to the stratum corneum of rat esophagus epithelium, so-called 'antikeratin antibodies'. *Ann Rheum Dis*, 48:712-22.

Visser H, Le Cessie S, Vos K, Breedveld F. C, and Hazes J. M. (2002): How to diagnose rheumatoid arthritis early: A prediction model for persistent erosive arthritis. *Arthritis Rheum*, 46: 357-65

Voigt LF, Koepsell TD, Nelson JL, Dugowson CE, Daling JR.

(1994): Smoking, obesity, alcohol consumption, and the risk of rheumatoid arthritis. *Epidemiology*, 5:525–32.

Vossenaar ER, Smeets TJ, Kraan MC, Raats JM, van Venrooij WJ, Tak PP (2004): The presence of citrullinated proteins is not specific for rheumatoid synovial tissue. *Arthritis Rheum*; 50:3485–3494.

Vossenaar ER, Zendman AJW, van Venrooij WJ, Pruijn GJ (2003): PAD, a growing family of citrullinating enzymes: Genes, features and involvement in disease. *Bioessays* 25, 1106-1118.

Waalder E (1940) : On the Occurrence of a factor in human serum , activating the specific agglutination of sheep blood corpuscles . *Acta Pathol Microbiol Scand* , 17 : 172 - 88 .

Walker W. C, and Wrigth V. (1968): Pulmonary lesions and rheumatoid arthritis. *Medicine* 47:501.

Walker W. C, and Wrigth V. (1969): Diffuse interstitial pulmonary and rheumatoid arthritis. *Ann. Rheum. Dis.* 28:252.

Weiner S, Bassett L, and Spiegel T. (1982): Superior, posterior and lateral displacement of C1 in RA. *Arthritis Rheum*, 25:1378.

Westergren A. (1926): The Technique of the red cell sedimentation

reaction. American Review of the Tuberculosis, 14:94- 101.

Weyand CM, Hicok KC, Conn DL, Goronzy JJ. (1992): The influence of HLADR B1 genes on disease severity in RA. Ann Intern Med, 117:801.

Winfield J, Young A, Williams P. (1983): Prospective study of the radiological changes in hands, feet and cervical spine in adult rheumatoid disease. Ann Rheum Dis,42:613.

Wolfe F, Cathey MA, Roberts FK. (1991): The latex test revisited, Rheumatoid factor testing in 8,287 rheumatic disease patients. Arthritis Rheum; 34:951-60.

Yellin E, Wanke I.A. (1999): An assessment of the annual and long-term direct costs of rheumatoid arthritis. Arthritis Rheum, 42:1209.

Young BJJ, Mallya RK, Leslie RDG, Clark CJM, Hamblin TJ. (1979): Anti-keratin antibodies in rheumatoid arthritis. Br Med J, 2:97-99.

Zeng X. Ai. M, Tian X, Gan X, Shi Y, Song Q, and Tang F. (2003): Diagnostic value of anti-cyclic citrullinated peptide antibody in patients with rheumatoid arthritis. J.

Rheumatol., Jul.; 30(7): 1451-5.

Ziff M (1990): The rheumatoid nodule. *Arthritis Rheum*; 33:761-7.

Zvaiflar NJ. (1995): Etiology and pathogenesis of RA. In McCarty.
D. J. and Koopman .J. (Eds) *arthritis and allied conditions*.
Ed. 12. Philadelphia. Lea and Febiger, p.659.