
References

- Adelman, R.D. and Solhaug, M.J. (2000):** Pathophysiology of body fluids and fluid therapy. In Behrman R.E.: Kliegman, R.M. and Genson, H.B. (eds.) : Nelson textbook of pediatrics. 16ed., W.B., Saunders, Co.P. 189-227.
- Ahlfors CE, Wennberg RP. (2004):** Bilirubin-albumin binding and neonatal jaundice. *Semin Perinatol*; 28:334–9.
- American Academy of Pediatrics Subcommittee on Hyperbilirubinemia. (2004) :** Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics* ;114(1):297–316
- Amin SB. (2004):** Clinical assessment of bilirubin-induced neurotoxicity in premature infants. *Semin Perinatol* ;28(5):340–7.
- Ashish Jain, Ramesh Agarwal, M. Jeeva Sankar, Ashok K. Deorari and Vinod K. Paul. (2008):** Hypocalcemia in the Newborn.; *Indian J Pediatr*, 75 (2) : 165-169.
- Aziz S, Kotal P, Leroy P, Servaes R, Eggermont E, Fevery J.(2001):** Bilirubin-IXalpha and -IXbeta pigments, coproporphyrins and bile acids in meconium and stools from full-term and preterm neonates during the first month of life. *Acta Paediatr*; 90: 81–7.
- Bancroft JD, Kreamer B, Gourley GR. (1998):** Gilbert syndrome accelerates development of neonatal jaundice. *J Pediatr*; 132(4):656–60.
- Behrman, R.E. Vaughan, V.C. and Nelson ,W.E. (2000) :** jaundice and hyperbilirubinemia in the newborn infant. Nelson textbook of pediatrics , 16th ed. W.B. Saunders Co. Philadelphia. 476-481.
- Behrman, R.E., Kliegman R.M. ,Avrin , A.M. ,Nelson, W.E. and Zrvn, A.M. (eds) .(2000):** Haemolytic disease of the newborn. In Nelson Textbook of pediatrics 16th ed .W.B. Saunders Co., p.500-504.

- Behrman, R.E., Kliegman R.M. and Arvin , A.M. (2000):** Jaundice and hyperbilirubinemia in the newborn. In : Nelson textbook of pediatrics 16th ed.; W.B.Saunders Co.; p.493-499.
- Behrman, R.E., Kliegman R.M. and Arvin , A.M. (2000):** Metabolic disturbance in neonates. Nelson textbook of pediatrics. 16th eds. Philadelphia. W.B. Saunders. Chapter 92.
- Benders,M.J.; Van. Bel. F.;Van de Bor, M. (1999):** Hemodynamic consequences of phototherapy in term infants . Eur. J. Pediatr.158(4) :323-8.
- Benders,M.J.;Van.Bel.F.;Van de Bor,M.(1998):** The effect of phototherapy on cerebral blood flow velocity in preterm infants.Acta. Pediatr. 87 (7): 786-91
- Bhutani VK, Johnson L, Sivieri EM. (1999):** Predictive ability of a predischage hour-specific serum bilirubin for subsequent significant hyperbilirubinemia in healthy term and near-term newborns. Pediatrics .; 103 : 6-14.
- Blackburn ST. (2003):** Bilirubin metabolism. In:Maternal, fetal,neonatal physiology: a clinical perspective. 2nd edition. St.Louis (MO): WB Saunders ; . p.652–76
- Bohm, F.; Drygalla F. and Charlesworth, P. (1995) :** Bilirubin phototoxicity to human cell by green light phototherapy in vitro. Photochem Photobiol. Dec; 62 (6) 980-3. care. 6th edition. St. Louis (MO): Mosby Elsevier; . p. 548–68.
- Cashore WJ , Polin RA, Fox WW, Abman SH,editors. (2004):** Bilirubin metabolism and toxicity in the newborn. In: Fetal and neonatal physiology. 3rd edition.Philadelphia: Saunders; . p. 1199–210.
- Cashore WJ. (2000):** Bilirubin and jaundice in the micropremie. Clin Perinatol J; 27 : 171-179.
- Centers for Disease Control and Prevention. (2001) :** Kernicterus in full-term infants–United States, 1994-1998. Morbid Mortal Wkly Rep. 50:491– 494.

- Chen C, Liu S, Lai C, Hwang C, Hsu H. (2002):** Changing position does not improve the efficacy of conventional phototherapy. *Acta Paediatr Tw.* ;43:255–258.
- Chowdhury JR, Chowdhury NR, Strom SC, et al.(1998):** Human hepatocyte transplantation; gene therapy and more. *Pediatrics*; 102(3 Pt 1):647–8
- Cloherly, J.P. (1998):** Neonatal hyperbilirubinemia. In *Manual of neonatal care*, 3 rd ed. Little Brown and Company Boston \Toronto, p. 298-334.
- Csoma Z, Hencz P, Orvos H, et al. (2007):** Neonatal bluelight phototherapy could increase the risk of dysplastic nevus development. *Pediatrics* , ;119(6):1036–7
- CsomaZ, KemenyL, OlahJ. (2008):** Phototherapy for neonatal jaundice. *N Engl J Med*,; 358(23):2523–4.
- Dagoglu T, Ovali F, Samanci N, et al.(1995):** High dose intravenous immunoglobulin therapy for haemolytic disease. *J Int Med Res*.; 23(4):264–71.
- De Almeida MF, Draque CM.. (2007):** Neonatal jaundice and breastfeeding, *NeoReviews*.;8(7):e282–8.
- Dollberg S, Atherton HB, Hoath SB. (1995):** Effect of different phototherapy lights on incubator characteristics and dynamics under three modes of servocontrol. *Am J Perinatol*.;12:55– 60.
- Doumas BT, Eckfeldt JH.(1996):** Errors in measurement of total bilirubin: a perennial problem. *Clin Chem* ;42(6 Pt 1):845–8.
- Drummond GS, Kappas A. (2004):** Chemoprevention of severe neonatal hyperbilirubinemia. *Semin Perinatol* ; 28:365–8.
- Eaton AP.(2001):** Early postpartum discharge: recommendations from a preliminary report to Congress. *Pediatrics*.;107(2):400–3.
- Ebbesen F.(2000):** Recurrence of kernicterus in term and near-term infants in Denmark. *Acta Paediatr*.; 89(10):1213–7.

- Eghbalian F, Monsef A. (2002):** Phototherapy-Induced Hypocalcemia In Icteric Newborns .Iran J Med Sci.; 27(4):169-171.
- Fevery J.(1997):**Fasting hyperbilirubinemia: Gastroenterology 113(5):1798–800.
- Fok TF, Wong W ,Chenug KL. (1997):** Eye protection for newborn under phototherapy: Comparison between a modified headbox and the conventional eyepatches .Ann Trop Paediatr ; 17:349-354.
- Fok TF, Wong W, Cheng AF.(1995):** Use of eye patches in phototherapy: effects on conjunctival bacterial pathogens and conjunctivitis. Pediatr Infect Dis J.; 14: 1091–1094.
- Fox IJ, Chowdhury JR, Kaufman SS, et al (1998):** Treatment of the Crigler-Najjar syndrome type I with hepatocyte transplantation. N Engl J Med; 338(20):1422–6.
- Garg, A.K; Ward, D.C. (1998):** Double surface phototherapy on a fluid bed .Ann.Pediatr. 14 (1): 81-4.
- Garg AK, Prasad RS, Hifzi IA.(1995):** A controlled trial of high-intensity double-surface phototherapy on a fluid bed versus conventional phototherapy in neonatal jaundice. Pediatrics; 95(6):914–6.
- Gartner LM, Herschel M. (2001):** Jaundice and breast-feeding. Pediatr Clin North Am; 48 : 389-399.
- Gartner LM, Lee KS. (1999):** Jaundice in the breast-fed infant. Clin Perinatol .; 26 : 431-445.
- Gartner LM. (1994):** Neonatal jaundice. Pediatr Rev; 15(11):422–32
- Gartner U, Goeser T, Wolkoff AW. (1997):** Effect of fasting on the uptake of bilirubin and sulfobromophthalein by the isolated perfused rat liver. Gastroenterology .; 113 (5) :1707–13
- Gottstein R, Cooke RW. (2003):** Systematic review of intravenous immunoglobulin in haemolytic disease of the newborn. Arch Dis Child Fetal Neonatal Ed ;88(1):F6–10.

- Gounaris A, Alexiou N, Costalos C, Daniilidou M, Bakoleas V, Constantellou E. (1998):** Gut hormone levels in neonates under phototherapy. *Early Human Development* ; 51:57-60.
- Gourley GR (2002):** Breast-feeding, neonatal jaundice and kernicterus. *Semin Neonatol*; 7(2):135–41.
- Gourley GR.(1997):** Bilirubin metabolism and kernicterus. *Adv Pediatr.*; 44:173–229.
- Govaert P, Lequin M, Swarte R, et al.(2003):** Changes in globus pallidus with (pre)term kernicterus. *Pediatrics*; 112(6 Pt 1):1256–63.
- Grohmann K, Roser M, Rolinski B, et al. (2006):** Bilirubin measurement for neonates: comparison of 9 frequently used methods. *Pediatrics* ;117(4):1174–83.
- Grunhagen DJ, De Boer MGJ, De Beaufort AJ, Walther FJ.(2002):** Transepidermal water loss during halogen spotlight phototherapy in preterm infants. *Pediatr Res.* ;51:402–405.
- Gutcher, G.R. and Odell, G.B (1993):** Hypocalcaemia associated with phototherapy in newborn rats: light source dependence. *Photochem. Potobiol.* 37(2): 177-80.
- Guttyton, A.C. (1998):** Parathyroid hormone, calcitonin, calcium and phosphate metabolism, vitamin D, bone and teeth. In: *Human physiology and mechanism of disease E. Guyton, A.C., 8th eds .W.B. Saunders Co., Philadelphia, London, Toronto, P. 868-884.*
- Hakanson, D.O., and Bergstrom, W.H. (1991):** Phototherapy induced hypocalcemia in newborn rats ; prevention by melatonin .*Science* 13; 214 (4522): 807-9.
- Hammermann C, Kaplan M. (2008):** Recent developments in the management of neonatal hyperbilirubinemia. *NeoReviews* 2000;1(2):e19–24.
- Hanko E. (2006):** Unbound bilirubin and risk assessment in the jaundiced newborn: possibilities and limitations.*Pediatrics*; 117(2):526–7.

- Hansen TW.(1997):** Acute management of extreme neonatal jaundice: the potential benefits of intensified phototherapy and interruption of enterohepatic bilirubin circulation. *Acta Paediatr.*; 86(8):843–6.
- Hart G, Cameron R. (2005):** The importance of irradiance and area in neonatal phototherapy. *Arch Dis Childhood Fetal Neonatal Ed.*; 90:F437–F440.
- Huttner, K.M. (1998):** Hypocalcaemia, hypercalcemia and hypermagnesmia. In Cloherty J.P. and Stark, A.R. (eds): *Manual of neonatal care*, 4th ed. Lippincott, Philadelphia, p.553-562.
- Ip S, Chung M, Kulig J, et al. (2004):** An evidence-based review of important issues concerning neonatal hyperbilirubinemia. *Pediatrics*; 114(1):e130–53.
- Jahnukanien, T.; Lindqvist, A.; Jalonen, J.,Kero,P;Valimaki, I. (1999)** : Responsiveness of cutaneous vasculature to thermal stimulation during phototherapy in neonatal aundice. *Eur . J. Pediatr.* 158 (9).
- JainBK,Singh D, Toor NS(1999):**Hypocalcemic effect of phototherapy. *Indian Pediatr* 36(2):208-9.
- Jaundice. Centers for Disease Control and Prevention. (2008):** Web site.Availableat:<http://www.cdc.gov/breastfeeding/disease/jaundice.htm>. Accessed June 21,(2008)
- Johan Fevery. (2008):** Bilirubin in clinical practice: *Liver International* .The Authors. Journal compilation _c 2008 Blackwell Munksgaard, p: 592-605.
- Johan Fevery. (2008):** Bilirubin in clinical practice. *Liver International* ISSN; 1478-3223.
- Johnson L, Bhutani VK . (1998):** Guidelines for management of the jaundiced term and near-term infant .*Clin Perinatol* ;25(3):555–74, viii
- Johnson L, Brown AK, Bhutani VK. Bind . (1999):** A clinical score for bilirubin induced neurologic dysfunction in newborns. *Pediatrics*; 104:746–7

- Johnson LH, Bhutani VK, Brown AK.(2002):** System-based approach to management of neonatal jaundice and prevention of kernicterus. *J Pediatr*; 140(4):396–403.
- Kaplan M, Hammerman C. (2002):** Glucose-6-phosphate dehydrogenase deficiency: a worldwide potential cause of severe neonatal hyperbilirubinemia *NeoReviews*; 1:e32–9.
- Kaplan M, Hammerman C. (2004):** Glucose-6-phosphate dehydrogenase deficiency: a hidden risk for kernicterus. *Semin Perinatol*; 28(5):356–64
- Kaplan M, Kaplan E, Hammerman C, Algur N, Bromikder R, et al. (2006) :** Post-phototherapy neonatal bilirubin rebound: a potential cause of significant hyperbilirubinemia. *Arch Dis Child.*; 91:31–34.
- Karamifar H. , Pishva N. and Amirhakim GH. (2002) :** Prevalence of phototherapy-induced hypocalcemia . *Iran J Med Sci* 2002; 27 (4): 166-168.
- Keren R, Luan X, Friedman S, et al. (2008):** A comparison of alternative risk-assessment strategies for predicting significant neonatal hyperbilirubinemia in term and near-term infants. *Pediatrics.*;121(1):e170–9.
- Kernicterus in full-term infants. (2001):** United States, 1994–1998. *MMWR Morb Mortal Wkly Rep*(2001);50(23):491–4.
- Laforgia N, Faienza MF, Rinaldi A, et al. (2002):** Neonatal hyperbilirubinemia and Gilbert's syndrome. *J Perinat Med*; 30(2):166–9.
- Laura A, Stokowski, RN, MS. (2006):** Advances In Neonatal Care, Vol 6, No 6 :pp 303-312
- Li R, Zhao Z, Mokdad A, et al. (2003):** Prevalence of breastfeeding in the United States: the 2001 national immunization survey. *Pediatrics* 2003; 111(5 Part 2):1198–201.
- Lo SF, Doumas BT, Ashwood ER. (2004):** Performance of bilirubin determinations in US laboratories: revisited. *Clin Chem*; 50(1):190–4.

- Maayan-Metzger A, Yosipovitch G, Hadad E, Sirota L. (2001):** Transepidermal water loss and skin hydration in preterm infants during phototherapy. *Am J Perinatol.* ;18(7):393-6.
- Maden A, MacMahon JR, Stevenson DK. Taeusch HW, Ballard RA, Gleason CA, editors. (2005):** Neonatal hyperbilirubinemia. In: *Avery's diseases of the newborn*. 8th edition. Philadelphia: Elsevier Saunders; . p. 1226–56
- Maisels MJ, Klaus MH, Fanaroff AA, editors. (2001):** Neonatal hyperbilirubinemia. In: *Care of the high-risk neonate*. 5th edition. Philadelphia: WB Saunders Company; . p. 324–62.
- Maisels MJ, Watchko JF. (2003):** Treatment of jaundice in low birthweight infants. *Arch Dis Child Fetal Neonatal Ed.*;88(6):F459–63.
- Maisels MJ, Baltz RD, Bhutani VK et al., (2004):** Management of hyperbilirubinemia in the newborn infant 35 or more weeks of gestation. *Pediatrics.*; 114 : 297-316
- Maisels MJ, Kring E. (2002):** Rebound in serum bilirubin level following intensive phototherapy. *Arch Pediatr Adolesc Med*; 156:669-72.
- Maisels MJ, MacDonald MG, Mullett MD, Seshia MMK, editors. (2005):** Jaundice In: *Avery's neonatology: pathophysiology & management of the newborn*. 6th edition. Philadelphia: Lippincott Williams & Wilkins;. p. 768–846.
- Maisels MJ, McDonagh AF. (2008):** Phototherapy for neonatal jaundice. *N Engl J Med*; 358(9):920–8.
- Maisels MJ, Newman TB. (1995):** Kernicterus in otherwise healthy, breast-fed term newborns. *Pediatrics.*;96(4 Pt 1):730–3.
- Maisels MJ, Watchko JF, eds. (2000):** Neonatal jaundice. London, UK; Harwood Academic, 169-176.
- Maisels MJ. (2001):** Phototherapy: traditional and nontraditional. *J Perinatol* ;21(Suppl 1):S93–97.
- Maisels MJ. (2005):** A primer on phototherapy for the jaundiced newborn. *Contemp Pediatr.*; 22(6):passim. (or, 38,40,44,47,48,53,54,57).

- Maisels MJ. (2006):** What's in a name? Physiologic and pathologic jaundice: the conundrum of defining normal bilirubin levels in the newborn. *Pediatrics*;118(2):805–7.
- Martin CR, Cloherty JP ,Cloherty JP, Eichenwald EC, Stark AR, eds. (2004):** Neonatal hyperbilirubinemia. In *Manual of Neonatal Care*, 15th ed. Philadelphia: Lippincott Williams & Wilkins. ; 185-221
- Martinez JC, Garcia HO, Otheguy LE, et al. (1999):** Control of severe hyperbilirubinemia in full-term newborns with the inhibitor of bilirubin production Sn-mesoporphyrin. *Pediatrics*; 103(1):1–5.
- Marx SJ. (2000):** Hyperparathyroid and hypoparathyroid disorders. *N Engl J Med* ; 343 : 1863-1875.
- McDonagh AF, Maisels MJ. (2006):** , Bilirubin unbound: déjà vu all over again? *Pediatrics*; 117(2):523–5.
- McDonagh AF. (2001):** Phototherapy: from ancient Egypt to the new millenium. *J Perinatol.*; 21:S7–S12
- Mckiernan P. (2002):** Neonatal cholestasis. *Semin Neonatal.*; 7:(2)153-156,with permission.
- McLin VA, BalistreriWF, Walker WA, Goulet O, Kleinman RE, et al, editors. (2004):** Approach to neonatal cholestasis .In: *Pediatric gastrointestinal disease: pathophysiology, diagnosis, management*. 4th edition. Hamilton (Ontario, Canada): BC Decker, Inc.; p. 1079–93.
- Mehta S, Kumar P, Narang A. (2005):** A randomized controlled trial of fluid supplementation in term neonates with severe hyperbilirubinemia. *J Pediatr.* ; 147:781–785
- Mimouni FB, Root AW, Sperling MA(ed). (1996):** Disorders of calcium metabolism in the newborn. *Pediatric Endocrinology*, Philadelphia,W.B. Saunders Co,; 95-111.
- Moyer VA, Ahn C, Sneed S. (2000):** Accuracy of clinical judgment in neonatal jaundice. *Arch Pediatr Adolesc Med*; 154(4):391–4.

- Murki S, Kumar P, Majumdar S, et al. (2001):** Risk factors for kernicterus in term babies with non-hemolytic jaundice. *Indian Pediatr.*;38(7):757–62.
- Nair PAK, Al Khussiby SM. (2003):** Kernicterus and G6PD deficiency: a case series from Oman. *J Trop Pediatr.*; 49:74–7.
- Nakisa Hooman, Ali Honarpisheh. (2005):** The effect of phototherapy on urinary calcium excretion in newborns. *Pediatr Nephrol.*; 20: 1363-1364.
- Newborn Emergency Transport Service. Jaundice in the first two weeks of life. (2003):** In: Neonatal Handbook. Newborn Emergency
- Newman TB, Liljestrand P, Jeremy RJ, et al. (2006):** .Outcomes among newborns with total serum bilirubin levels of 25 mg per deciliter or more. *N Engl J Med*; 354(18):1889–900.
- Newman, T.B. and Maisles, M.J. (1998):** Evaluation and treatment of jaundice in term newborn. *Pediatrics*.809-18.
- Nishioka T, Hafkamp AM, Havinga R, et al. (2003):** Orlistat treatment increase fecal bilirubin excretion and decrease plasma bilirubin concentrations in hyperbilirubinemic guinea rats. *J Pediatr* ; 143:327-334.
- Oden J, Bourgeois M. (2000):** Neonatal endocrinology. *Indian J Pediatr*; 67: 217-223.
- Okazaki, I.;Igarashi, T. and Kroneberg, H.M. (1998):** 5'- Flanking region of parathyroid hormone gene mediated negative regulation by 1,25- (OH)₂ D3. *J. Biol. Chem.*263-2203.
- Ostrow JD, Mukerjee P, Tiribelli C. (1994):** Structure and binding of unconjugated bilirubin: relevance for physiological and pathophysiological function. *J Lipid Res*; 35:1715–37.
- Penn AA, EnzmannDR, Hahn JS, et al.(1994):** Kernicterus in a full term infant. *Pediatrics.*; 93(6 Pt 1):1003–6
- Pezzati M, Fusi F, Dani C, Piva D, Bertini G, et al.(2002):** Changes in skin temperature of hyperbilirubinemic newborns under

phototherapy: conventional versus fiberoptic device. Am J Perinatal; 19:439–444.

Pritchard MA, Beller EM, Norton B.(2004): Skin exposure during conventional phototherapy: a randomized controlled trial. J Paediatr Child Health.; 40:270 –274.

Richron and Alleyne. (1998): Management of sick newborn. In manual of pediatric therapeutic 4 th ed. Little Brown and Company. Boston \Toronto.p. 148-154.

Rigo J, Curtis MD. (2006): Disorders of Calcium, Phosphorus and Magnesium metabolism. In Richard J, Martin, Avory A, Fanaroff, Michele C Walsh, eds. Neonatal Perinatal Medicine -Diseases of the fetus and infant, 8th ed. Elsevier; Philadelphia,; 1508.

Robin L.Watson, RN,MN, CCRN. (2009): Hyperbilirubinemia. Pediatric\Neonatology, Crit Care Nurs Clin N Am 21, 97–120,

Romagnoli C, Zecca E, Pappaci P, Vento G, Girlando P,et al. (2006): Which phototherapy system is most effective in lowering serum bilirubin in very preterm infants? Fetal Diagn Ther.21:204 –209.

Rosenstein B.S., Ducore J.M (1996): Enhancement by bilirubin of “DNA” damage induced in human cells exposed to phototherapy light. Pediatr. Res.; 18 (1) 3-6.

Roy M.S, Caramelli C, Orquin J, et al. (1999): Effect of early reduced light exposure on central visual development in preterm infants. Acta Paediatr ;88: 459-461.

Salle BL, Delvin EE, Lapillonne A, Bishop NJ, Glorieux FH. (2002): Perinatal metabolism of vitamin D. Am J Clin Nutr .; 71(5 suppl) : 1317S-1324S.

Sarah K. Moerschel, MD; Lauren B. Cianciaruso, DO; and Lloyd R. Tracy, MD. (2008): A Practical Approach to Neonatal Jaundice, American Family Physician, Volume 77, Number 9 ,1255-1262

Satish Mishra, Ramesh Agarwal, Ashok K. Deorari and Vinod K. Paul. (2008): Jaundice in the Newborns. Indian J Pediatr ; 75 (2) : 157-163

- Satish Mishra, Ramesh Agarwal, Ashok K.Deorari and Vinod K.Paul (2008):** Indian Journal Of Pediatric, Volume 75 .
- Schwartz R, Teramo KA. (2000):** Effects of diabetic pregnancy on the fetus and newborn. Semin Perinatol ; 24 : 120-135.
- Sethi, H., Saili A., Dutta, A.K. (1998):** Phototherapy induced hypocalcaemia in Indian pediatric. (30), 1: 403-6.
- Sharma Jyothi, Bajpai Anurag, Kabra Madhulika, Menon PSN. (2002):** Hypocalcaemia – Clinical Biochemical, Radiological Profile and follow-up in a Tertiary hospital in India. Indian Pediatrics; 39: 276-282.
- Shinki T, Ueno Y, Deluca HF,, et al. (1999):** Calcitonin is a major regulator for the expression of renal 25-hydroxyvitamin D3-1 alpha hydroxylase gene in normocalcemic rats. Proc Natl Acad Sci USA ;96: 8253-8258.
- Shinwell ES, Sciaky Y, Karplus M.(2002):** Effect of position changing on bilirubin levels during phototherapy. J Perinatol.; 22:226 –229.
- Shortland DB, Hussey M, Chowdhury AD.(2008):** Understanding neonatal jaundice: UK practice and international profile.J R Soc Promot Health. 2008 Jul;128(4):202-6.
- Slusher TM, Angyo IA, Bode-Thomas F, et al. (2004):** Transcutaneous bilirubin measurements and serum total bilirubin levels in indigenous African infants. Pediatrics; 113(6):1636–41
- Slusher TM, Vreman HJ, McLaren DW, et al.(1995):** Glucose-6-phosphate dehydrogenase deficiency and carboxyhemoglobin concentrations associated with bilirubin-related morbidity and death in Nigerian infants. J Pediatr.; 126(1):102–8.
- Stanley TV.(1997):** A case of kernicterus in New Zealand:a predictable tragedy? J Paediatr Child Health.;33:451–3.
- Steiner LA, Bizzarro MJ, Ehrenkranz RA, et al. (2007):** A decline in the frequency of neonatal exchange transfusions and its effect on exchange-related morbidityand mortality. Pediatrics; 120(1):27–32.

- Stephanie C. Hsu, Michael A. Levine.(2004):** Perinatal calcium metabolism: physiology and pathophysiology.Seminars in Neonatology 9, 23–36.
- Stevenson DK, Fanaroff AA, Maisels MJ, et al. (2001):** Prediction of hyperbilirubinemia in near-term and term infants. Pediatrics; 108(1):31–9.
- Stokowski LA.(2006) :** Fundamentals of phototherapy for neonatal jaundice. Adv Neonatal Care 6(6):303–12
- Tan KL. (1994):** Comparison of the efficacy of fiberoptic and conventional phototherapy for neonatal hyperbilirubinemia. J Pediatr. 125:607– 612
- The Joint Commission. Kernicterus threatens healthy newborns. (2008):** sentinel event alert. Available at: http://www.jointcommission.org/SentinelEvents/SentinelEventAlert/sea_18.htm. Accessed June 12, (2008).
- Tiribelli C, Ostrow JD.(2005):** The molecular basis of bilirubin encephalopathy and toxicity: report of an EASL Single Topic Conference Hepatol ; 43: 156–6.
- Tsang, R.C.and Steichen, J. (1998):** Disorders of calcium and magnesium metabolism. In :Neontal perinatal medicine. 3rd ed. Fanoroff,A.A., Montein, R.J.and Markatz, I.R. the C.V. Mosby Company. P.870-874.
- Valaes T, Petmezaki S, Henschke C, et al. (1994):** Control of jaundice in preterm newborns by an inhibitor of bilirubin production: studies with tin-mesoporphyrin. Pediatrics.;93(1):1–11.
- Van der Veere CN, Jansen PL, Sinaasappel M, et al. (1997):** .Oral calciumphosphate: anewtherapy for Crigler-Najjar disease. Gastroenterology; 112(2):455–62.
- Venigalla S, Gourley GR. (2004):** Neonatal cholestasis.Semin Perinatol ;28(5):348–55.
- Verkade HJ. (2002):** A novel hypothesis on the pathophysiology of neonatal jaundice. J Pediatr; 141:594-595.

- Volpe JJ. (2000):** Bilirubin and brain injury. In: *Neurology of the newborn*. 4th edition. Philadelphia: WB Saunders Company;. p. 521–64.
- Voto LS, Sexer H, Ferreiro G, et al. (1995):** Neonatal administration of high-dose intravenous immunoglobulin in rhesus hemolytic disease. *J Perinat Med.*; 23(6):443–51.
- Vreman HJ, Verter J, Oh W, et al.(1996):** Interlaboratory variability of bilirubin measurements. *Clin Chem*;42(6 Pt 1):869–73.
- Vreman HJ, Wong RJ, Stevenson DK (2004):** Phototherapy;current methods and future directions. *Sem Perinatol.*;28:326 –333
- Watchko JF. (2006):** Neonatal hyperbilirubinemia: what are the risks? *N Engl J Med .*; 354(18):1947–9.
- Watchko JF.(2005):** Vigintiphobia revisited. *Pediatrics*; 115(6):1747–53.
- Wong RJ, De Sandre GH, Sibley E, et al. (2006):** Neonatal jaundice and liver disease. In: *Fanaroff and Martin’s neonatal-perinatal medicine: diseases of the fetus and infant*. 8th edition. Philadelphia: Mosby Elsevier; p. 1419–65.
- Wong RJ, Stevenson DK, Ahlfors CE, et al. (2008):** Neonatal jaundice: bilirubin physiology and clinical chemistry. *Neo-Reviews*, 8:e58–67.
- Yetman, R.J., Parks, D.K., Huseby, V., Mistry, K. and Garicia, J. (1999):** Rebound bilirubin level in infants receiving phototherapy. *J. Pediatr. Nov.* 133 (5): 705-7.