

References

- Ahumada M, Cabadés A, Valencia J, et al.** Reinfarction as a Complication of Acute Myocardial Infarction. *Revista Espanola de Cardiologia* 2005; 58:13-19.
- Al-Delaimy WK, Manson JE, Solomon CG, et al.** Smoking and risk of coronary heart disease among women with type 2 diabetes mellitus. *Arch Intern Med* 2002; 162:273.
- AlHabib KF, Hersi A, AlFaleh H, Kurdi M, Arafah M and Youssef M:** The Saudi Project for Assessment of Coronary Events (SPACE) registry: design and results of a phase I pilot study. *Can J Cardiol.* 2009; 25(7):e255-8.
- Al-Khatib S, Stebbins AL, Califf RM, et al:** Sustained ventricular arrhythmias and mortality among patients with acute myocardial infarction. *Am Heart J* 2003;145: 515-21.
- Almdal T, Scharling H, Jensen JS, et al:** The independent effect of type 2 diabetes mellitus on ischemic heart disease, stroke, and death: a population-based study of 13,000 men and women with 20 years of follow-up. *Arch Intern Med* 2004; 164: 1422–1426.
- Alpert JS, et al:** Definition of myocardial infarction, a global consensus document of The Joint ESC/ACC/AHA/WHF/ WHO Task Force for the Redefinition of Myocardial Infarction. 2007.

Alpert JS, Thygesen K, Antman E, et al: Myocardial infarction redefined--a consensus document of The Joint European Society of Cardiology/American College of Cardiology Committee for the redefinition of myocardial infarction. 2000; J Am Coll Cardiol; 2000 36 (3): 959-69.

American Diabetes Association: Report of the expert Committee on the diagnosis and classification of Diabetes Mellitus. Diabetes Care; 1999; 20:1183-1197.

American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care - Part 8: Stabilization of the Patient With Acute Coronary Syndromes." Circulation 2005; 112: 89 -110.

American Heart Association. 2004 Heart and Stroke Statistical Update. Dallas, Texas: American Heart Association; 2004.

Anderson Elliot M, et al: ACC/AHA 2007 guidelines for the management of patients with Unstable Angina/NSTEMI. J Am Coll Cardiol 2007; 50:677.

Anderson JL, Horne BD, Jones HU, et al. Intermountain Heart Collaborative (IHC) Study which features of the metabolic syndrome predict the prevalence and clinical outcomes of angiographic coronary artery disease? Am J Cardiol; 2004; 101: 185- 93.

Anderson, Elliot M, et al. ACC/AHA guidelines for the management of patients with Unstable Angina/NSTEMI. J Am Coll Cardiol; 2007; 50:677.

- Antithrombotic Trialists Collaboration.*** Collaborative meta-analysis of randomized trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. BMJ 2002; 324:71-86.
- Antman EM.*** General hospital management. In: Julian DG, Braunwald E, eds. Management of acute myocardial infarction. London, England: WB Saunders Co Ltd:1994; 42-44.
- Aroney C, Boyden AN, Jelinek MV, et al.*** Management of unstable angina: guidelines 2000. Med J Aust; 2000; 173:S65-S88.
- Babaev A, Frederick PD, Pasta DJ, et al.*** Trends in management and outcomes of patients with acute myocardial infarction complicated by cardiogenic shock. JAMA 2005; 294:448-54.
- Bahl VK, Prabhakaran D and Karthikeyan, G.*** Coronary artery disease in Indians. Indian Heart J., 2001, 53: 707–713.
- Barrett-Connor EL, Cohn BA, Wingard DL, et al.*** Why is diabetes mellitus a stronger risk factor for fatal ischemic heart disease in women than in men? The Rancho Bernardo Study. JAMA 1991;265:627-31
- Barua RS, Ambrose JA and Srivastava S, et al.*** Reactive oxygen species are involved in smoking- induced dysfunction of nitric oxide synthase: An invitro demonstration in human coronary artery endothelial cells. Circulation 2003; 107:2342.
- Bassand JP, Hamm CW, Ardissino D et al.*** Guidelines for the diagnosis and treatment of non-ST-segment elevation acute coronary syndromes. Eur. Heart J. 2007; 28(13),1598-1660.

- Bazzano LA, He JA, Muntner P, et al.** Relationship between cigarette smoking and novel risk factors for cardiovascular disease in the United States. *Ann Inter Med* 2003; 138:891.
- Beckman JA, Creager MA and Libby P.** Diabetes and atherosclerosis: epidemiology, pathophysiology, and management. *JAMA*. 2002; 287:2570-2581.
- Bedinghaus J, Leshan L and Diehr S.** Coronary Artery Disease Prevention: What's Different for Women? *Am Fam Physician* 2001; 63:1393-1400.
- Bertuzzi M, Negri E, Tavani A, et al.** Family history of ischemic heart disease and risk of acute myocardial infarction. *Prev Med* 2003; 37:183–187.
- Bock JL, Brogan GX, McCuskey CF, et al.** Evaluation of CK-MB isoform analysis for early diagnosis of myocardial infarction. *J Emerg Med* 1999; 17:75-9.
- Boersma E, Harrington RA, Moliterno DJ, et al.** Platelet glycoprotein IIb/IIIa inhibitors in acute coronary syndromes: A meta-analysis of all major randomized clinical trials. *Lancet*; 2002; 359: 189-198.
- Boersma E, Mercado N, Poldermans D, et al.** Acute myocardial infarction. *Lancet* 2003; 361:847.
- Braunwald E, Antman EM, Beasley JW, et al.** ACC/AHA guidelines for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee on the Management of Patients With Unstable Angina) [published correction appears in *J Am Coll Cardiol* 2001;38:294-5]. *J Am Coll Cardiol* 2000; 36:970-1062.

- Brewer HB Jr.** increasing HDL cholesterol level. *N Engl J Med* 2004; 350:1491.
- Brieger D, Eagle KA, Goodman SG, et al.** GRACE Investigators. Acute coronary syndromes without chest pain, an underdiagnosed and undertreated high-risk group. *Chest* 2004;126:461–64.
- Burazeri G, Goda A, Sulo G, et al.** Conventional risk factors and acute coronary syndrome during a period of transition: population-based Case- control study in Tirana, Albania. *Croat Med J.* 2007; 48:225-33.
- Carneiro AV.** Coronary heart disease in diabetes mellitus: risk factors and epidemiology. *Rev Port Cardiol.* 2004; 23(10):1359-66.
- Chesebro JH, Knatterud G, Roberts R, et al.** Thrombolysis in Myocardial Infarction (TIMI) Trial, Phase I: a comparison between intravenous tissue plasminogen activator and intravenous streptokinase: clinical findings through hospital discharge. *Circulation* 1987; 76:142–54.
- Colditz GA and Coakley E.** Weight, weight gain, activity, and major illnesses: the Nurses' Health Study. *Int J Sports Med* 1997; 18(suppl 3): S162-70.
- Collins R, MacMahon S, Flather M, et al.** Clinical effect of anticoagulant therapy in suspected acute myocardial infarction: *BMJ* 1996; 313(7058):652-9.
- Conroy RM, Pyorala K, Fitzgerald AP, et al.** Estimation of ten-year risk of fatal cardiovascular disease in Europe: the SCORE project. *Eur Heart J* 2003; 24: 987.

- Dauerman HL, Lessard D, Yarzebski J, et al.** Ten-year trends in the incidence, treatment, and outcome of Q-wave myocardial infarction. *Am J Cardiol* 2000; 86:730–5.
- Davey-Smith G, Shipley MJ, Marmot MG and, et al.** Plasma cholesterol and mortality: The Whitehall Study. *JAMA* 1992; 267: 70–6.
- David Wood, Guy De Backer, Ole Faergeman, et al.** "ra"la" together with members of the Task Force: Task Force Report Prevention of coronary heart disease in clinical practice. *European Heart Journal* 1998; 19:1434–1503.
- Dawson B and Trapp R G:** Basic & Clinical Biostatistics. Lange Medical Book/ McGraw-Hill, Medical Publishing Division. 3rd ed., Ch. 7-9, PP 161-218, 2001.
- De Luca G, Suryapranata H, Ottervanger JP, et al.** Time delay to treatment and mortality in primary angioplasty for acute myocardial infarction: every minute of delay counts. *Circulation*; 2004; 109:1223-1225
- Deepa R, Arvind K and Mohan V.** Diabetes and risk factors for CHD. *Current Science*, 2002; 83, 1479-1505.
- Eikelboom JW, Anand SS, Malmberg K, et al.** Unfractionated heparin and low-molecular-weight heparin in acute coronary syndrome without ST elevation: A meta-analysis. *Lancet*; 2000; 355:1936-1942.
- Eisenman A.** "Troponin assays for the diagnosis of myocardial infarction and acute coronary syndrome: where do we stand?". *Expert Rev Cardiovasc Ther.* 2006; 4 (4): 509-14.

- Ellis SG, da Silva ER, Heyndrickx G, et al.** Randomized comparison of rescue angioplasty with conservative management of patients with early failure of thrombolysis for acute anterior myocardial infarction. *Circulation*; 1994; 90: 2280-2284.
- Esteghamati A, Abbasi M, Nakhjavani M, et al.** Prevalence of diabetes and other cardiovascular risk factors in Iranian population with acute coronary syndrome. *Cardiovascular Diabetology* 2006; 5;15.
- Ezzati M, Henley SJ, Thun MJ , et al.** Role of smoking in global and regional cardiovascular mortality. *Circulation*. 2005; 112: 489.
- Fang J, Mensah GA, Alderman MH, et al.** Trends in acute myocardial infarction complicated by cardiogenic shock, 1979-2003, United States. *Am Heart J Cardiol* 2000; 85: 1185-8.
- Flegal KM, Carroll MD, Kuczmarski RJ , et al.** Overweight and obesity in the United States: Prevalence and trends, 1960-1994. *Int J Obes Relat Metab Disord* 1998; 22:39.
- Fox CS, Evans JC, Larson MG, et al.** Temporal trends in coronary heart disease mortality and sudden cardiac death from 1950 to 1999: the Framingham Heart Study. *Circulation*. 2004; 110: 522–527.
- Fox CS, Sullivan L, D'Agostino RB Sr , et al.** The significant effect of diabetes duration on coronary heart disease mortality: the Framingham Heart Study. *Diabetes Care* 2004; 27:704-708.

- Fox KA, Mehta SR, Peters R, Zaho F, et al** Benefits and risks of the combination of clopidogril and aspirin in patients undergoing surgical revascularization for non-ST-elevation acute coronary syndrome: the Clopidogril in Unstable angina to prevent Recurrent ischemic events (CURE) Trial 2004; 110(10): 1202-8.
- Freedman DS, Khan LK, Serdula MK, et al.** Relationship of childhood BMI to adult adiposity: the Bogalusa Heart Study. *Pediatrics* 2005;115:22-27.
- Fuller JH, Stevens LK and Wang SL.** Risk factors for cardiovascular mortality and morbidity: the WHO multinational study of vascular disease in diabetes. *Diabetologia* 2001; 44(Suppl. 2): S54–S64.
- Gan SC, Beaver SK, Houck PM, et al.** Treatment of acute myocardial infarction and 30-day mortality among women and men. *N Engl J Med* 2000; 343:8-15.
- Ghaffar A, Reddy KS and Singhi M.** "Burden of non-communicable diseases in South Asia." *BMJ* 2004; 328: 807-810.
- Gibson CM, Cannon CP, Daley WL, et al.** TIMI frame count: a quantitative method of assessing coronary artery flow. *Circulation* 1996; 93:879–88.
- González I, Gaietà P, Jaume M, et al.** MASCARA (Manejo del Síndrome Coronario Agudo. Registro Actualizado) Study. General Findings. *Rev Esp Cardiol* 2008; 61(8):803-816.

- Goodman SG, Barr A, Soltchouk A, et al.** Low molecular weight heparin decreases rebound ischemia in unstable angina or non-Q-wave myocardial infarction: the Canadian ESSENCE ST segment monitoring substudy. *J Am Coll Cardiol*; 2000; 36:1507-1513.
- Grech ED.** ABC of interventional cardiology Pathophysiology and investigation of coronary artery disease *BMJ*. 2003; 326 (7397): 1027–1030.
- Gregg C.** Fonarow In-Hospital Initiation of Statin Therapy in Acute Coronary Syndromes Maximizing the Early and Long-term benefits *Chest* 2005;128:3641-3651.
- Gregg EW, Cauley JA, Stone K, et al.** Relationship of changes in physical activities and mortality among older women. *JAMA* 2003; 289:2379.
- Harvey d. Whit.** Thrombolytic Therapy and Equivalence Trials *JACC* vol. 31, No. 3 1998; 494-6.
- He J, Vupputuri S, Allen K, et al.** Passive smoking and the risk of coronary heart disease a meta-analysis of epidemiologic studies. *N Engl J Med* 1999; 340: 920-6.
- Hedley AA, Ogden CL, Johnson CL, et al.** Prevalence of overweight and obesity among US children, adolescents, and adults, 1999-2002. *JAMA* 2004; 291:2847-2850.
- Henkel DM, Witt BJ, Gersh BJ, et al.** ventricular arrhythmias after acute myocardial infarction: A 20-year community study. *Am Heart J*. 2006; 151:806-12.

- Hochman JS, Tamis JE, Thompson TD, et al.** Sex, clinical presentation, and outcome in patients with acute coronary syndromes. Global Use of Strategies to Open Occluded Coronary Arteries in Acute Coronary Syndromes IIB Investigators. N Engl J Med; 1999; 341:226-232.
- Holli A. DeVon, Catherine J Ryan, Amy L. Ochs, et al.** BSN and Moshe Shapiro Symptoms Across the Continuum of Acute Coronary Syndromes: Differences Between Women and Men. American Journal of Critical Care. 2008;17: 14-24
- Ibrahim MM.** Coronary Artery Disease. Egyptian Hypertension Society Guidelines Cairo - September 1998. Summary p. 4 eds. El-Tobgy S and Rizk H.
- Iribarren C, Sidney S, Bild DE, et al.** Association of hostility with coronary artery calcification in young adults: the CARDIA study. Coronary Artery Risk Development in Young Adults. JAMA 2000; 283: 2546–2551.
- Jackson G.** Primary angioplasty in acute myocardial infarction the preferred option, but time is of the essence. Int J Clin Pract; 2006; 60:379
- Jacobs AK.** Coronary revascularization in women 2003. Sex revisited. Circulation 2003; 107: 375-7.
- Joint National Committee.** The seventh report of the Joint Committee on prevention, Detection. Evaluation and Treatment of High Blood Pressure (JNC-7 Express). JAMA; 2003; 289:2560.

- Kannel W, McGee D and Castelli W.** Latest perspectives on cigarette smoking and cardiovascular disease: the Framingham Study. J Card Rehabil 1984; 4:267–277.
- Kannel WB, Dannenberg AL and Abbott RD.** Un-recognized myocardial infarction in hypertension. The Framingham Study. Am Heart J. 1985;109: 581-585.
- Kannel WB.** Blood pressure as a cardiovascular risk factor: prevention and treatment. JAMA 1996; 275: 1571–6.
- Kannel WB.** Coronary Heart Disease Risk Factors in the Elderly Am J Geriatr Cardiol 2002; 11(2):101-107.
- Karras DJ and Kane DL.** Serum markers in the emergency department diagnosis of acute myocardial infarction. Emerg Med Clin North Am 2001; 19:321-37.
- Kawachi J, Colditz GA, Stampfer MJ, et al.** Smoking cessation in relation to total mortality rates in women. Ann Intern Med 1993; 119: 992–1000.
- Kearney PM, Whelton M, Reynolds K, et al.** Global burden of hypertension: Analysis of worldwide data. Lancet 2005; 365: 217.
- Keeley EC, Boura JA and Grines CL.** Primary angioplasty versus intravenous thrombolytic therapy for acute myocardial infarction: a quantitative review of 23 randomised trials. Lancet 2003; 361:13-20.
- Kirsten Bibbins-Domingo, Pamela Coxson, et al.** Adolescent Overweight and Future Adult Coronary Heart Disease. NEJM 2007; 357: 2371-2379.

- Klag MJ, Ford DE, LA. Mead LA, et al.** Serum cholesterol in young men and subsequent cardiovascular disease. *N Engl J Med* 1993; 328:313–318.
- Klein LW.** Coronary artery disease in young adults *Journal of the American College of Cardiology* Volume 41, Issue 4 19 2003; 529-531.
- Kristian Thygesen, Joseph S. Alpert and Harvey D.** White on behalf of the Joint ESC/ACCF/AHA/WHF Task Force for the Redefinition *European Heart Journal* 2007; 28: 2525–2538.
- Kuper H, Marmot M and Hemingway H.** Systematic review of prospective cohort studies of psychosocial factors in the aetiology and prognosis of coronary heart disease. *Semin Vasc Med* 2002; 2: 267-314.
- Law MR, Morris JK and Wald NJ.** Environmental tobacco smoke exposure and ischaemic heart disease: an evaluation of the evidence. *BMJ* 1997; 315: 973-80.
- Lincoff AM and Topol EJ.** The illusion of reperfusion. Does anyone achieve optimal myocardial reperfusion? *Circulation*; 1993; 87: 1792-1805.
- Lincoff AM, Callif RM, Van de Werf F, et al.** Mortality at 1 year with combination platelet glycoprotein IIb/IIIa inhibition and reduced dose fibrinolytic therapy for acute myocardial infarction GUSTO randomised trials. *JAMA* 2002; 288(17): 2130:5.
- Lloyd-Jones DM, Leip EP, Larson MG, et al.** Prediction of lifetime risk for cardiovascular disease by risk factor burden at 50 years of age. *Circulation*. 2006; 113:791-798.

- Lopez AD and Murray C.** The global burden of disease 1990-2020. *Nat Med.* 1998; 4:1241-1243.
- Lori M, Alex B, Valentina B, et al.** The second Euro Heart Survey on acute coronary syndromes: characteristics, treatment, and outcome of patients with ACS in Europe and the Mediterranean Basin in 2004. *Eur Heart J* 2006; 10: 1093-1096.
- Manson JE, Tosteson H, Ridker PM, et al.** The primary prevention of myocardial infarction. *N Engl J Med* 1992; 326: 1406–16.
- Masoudi FA, Magid DJ, Vinson DR, et al.** "Implications of the Failure to Identify High-Risk Electrocardiogram Findings for the Quality of Care of Patients With Acute Myocardial Infarction." *Circulation* 2006; 114: 1565-1571.
- McGovern PG, Jacobs DR Jr, Shahar E, et al.** Trends in acute coronary heart disease mortality, morbidity, and medical care from 1985 through 1997: The Minnesota heart survey. *Circulation* 2001;104:19–24.
- Mehler PS, Coll JR, Estacio R, et al.** Intensive blood pressure control reduces the risk of cardiovascular events in patients with peripheral arterial disease and type 2 diabetes. *Circulation.* 2003; 107:753.
- Mikhail GW.** Coronary heart disease in women. Is underdiagnosed, undertreated, and under-researched. *BMJ* 2005; 331: 467-468.
- Moise P and Jacobzone S.** Outcomes of interventions for IHD. OECD Study of cross-national differences in the treatment, costs and outcomes of ischaemic heart disease. Paris: OECD. 2003: 70-79.

- Morgan KP, Kapur A and Beatt KJ.** Anatomy of coronary disease in diabetic patients. An explanation for poorer outcomes after percutaneous coronary interventions and potential target for intervention. *Heart*, 2004; 90:732-738.
- Musaiger AO.** Diet and Prevention of Coronary Heart Disease in the Arab Middle East Countries *Med Princ Pract* 2002; Vol. 11 (Suppl. 2):9-16.
- Naghavi M, Libby P, Falk E, et al.** From vulnerable plaque to vulnerable patient: a call for new definitions and risk assessment strategies: Part 1. *Circulation* 2003; 108: 1664-72.
- Nallamothu BK, Wang Y, Magid DJ, et al.** Relation between hospital specialization with primary percutaneous coronary intervention and clinical outcomes in ST-segment elevation myocardial infarction: National Registry of Myocardial Infarction-4 analysis. *Circulation* 2006; 113:222-229.
- Natarajan S, Lio Y, Cao G, et al.** Sex differences in risk for coronary heart disease. *Arch Inter Med* 2003; 163:1735.
- National Cholesterol Education Program (NCEP)** Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults(Adult Treatment Panel III) final report. *Circulation* 2002;106: 3143–3421.

- National Cholesterol Education Program.*** Expert Panel on Detection, Evaluation and Treatment of High Blood Cholesterol In Adults). (Adults Treatment Panel III). JAMA; 2001; 285: 2486: 2497.
- National Heart, Lung, and Blood Institute. Morbidity and Mortality:*** Chartbook on Cardiovascular, Lung, and Blood Diseases. Bethesda, MD: US Department of Health and Human Services; 2000.
- Neaton JD and Wentworth D.*** Serum cholesterol, blood pressure, cigarette smoking and death from coronary heart disease. Overall findings and differences by age for 316,099 white men. Multiple risk factor intervention trial Research group. Arch Inter Med. 1992; 152(1): 56-64.
- Nissen SE, Tuzcu EM, Libby P, et al.*** Effect of antihypertensive agents on cardiovascular events in patients with coronary disease and normal blood pressure: the CAMELOT study: A randomized controlled trial. JAMA 2004; 292:2217.
- Ogden CL, Carroll MD, Curtin LR, et al.*** Prevalence of overweight and obesity in the United States, 1999-2004. JAMA 2006; 295: 1549-1555.
- Okraïneç K, Banerjee DK and Eisenberg MJ.*** Coronary artery disease in the developing world. Am Heart J. 2004; 148:7-15.
- Paradiso-Hardy FL, Gordon WL, Jackevicius CA, et al.*** The importance of in-hospital statin therapy for patients with acute coronary syndromes. Pharmacotherapy 2003; 23: 506–513.

- Parish S, Collins R, Peto R, et al.** Cigarette smoking, tar yields, and non-fatal myocardial infarction: 14 000 cases and 32 000 controls in the United Kingdom. The international studies of infarct survival (ISIS) collaborators. *BMJ* 1995; 311: 471-7.
- Peres E, Caidahl K, Herlitz J, et al.** Spectrum of Acute Coronary Syndrome: History and clinical Presentation in Relation to Sex and Age. *Cardiology* 2004; 102:67-76.
- Petersen JL, Mahaffey KW, Hasselblad V, et al.** Efficacy and bleeding complications among patients randomized to enoxaparin or unfractionated heparin for antithrombin therapy in non-ST-segment elevation acute coronary syndromes: A systematic overview. *JAMA*; 2004; 292:89-96.
- Petersen S, Peto V, Scarborough P and British Heart Foundation Health Promotion Research Group.** Coronary heart disease statistics 2005. Oxford: British Heart Foundation, 2005.
- Peterson ED, Roe MT, Mulgund J, et al.** Association between hospital process performance and outcomes among patients with acute coronary syndromes. *JAMA*; 2006; 295: 1912-1920.
- Petrie A and Sabin C.** Medical Statistics at a Glance. 2nd ed., Blackwell Publishing, 2005.
- Pitsavos C, Panagiotakos DB, Antonoulas A, et al.** Epidemiology of acute coronary syndromes in Mediterranean country; aims, design and baseline characteristics of the Greek study of acute coronary syndromes (GREECS). *BMS Public Health* 2005;5:23.

- Pollack CV Jr and Gibler WB.** ACC/AHA guidelines for the management of patients with unstable angina and non-ST-segment elevation myocardial infarction: a practical summary for emergency physicians published corrections appear in *Ann Emerg Med* 2001; 38:468.
- Poloński L, Mariusz G, Marek G, et al.** Characteristics, treatments and outcomes of patients with acute coronary syndromes in Poland: Polish Registry of Acute Coronary Syndromes (PL-ACS). *Kardiologia Polska*; 2007; 65: 8
- Poulter N.** Global risk of cardiovascular disease. *Heart* 2003;89 Supplement (Suppl 2):ii2–5.
- Prineas RJ, Folsom AR and Kaye SA.** Central adiposity and increased risk of coronary mortality in older women. *Ann Epidemiol* 1993; 3: 35–41.
- Purvis JA, McNeill AJ, Siddiqui RA, et al.** Efficacy of 100 mg of double-bolus alteplase in achieving complete perfusion in the treatment of acute myocardial infarction. *J Am Coll Cardiol* 1994; 23:6 –10.
- Rader J.** Lipid disorders. In Topol J. (eds)Text book of cardiovascular medicine. Lippincott-Raven, Philadelphia, 1998; chapter 3; pp 59-91.
- Rauch U, Osende JI, Fuster V, et al.** Thrombus formation on atherosclerotic plaques: pathogenesis and clinical consequences. *Ann Intern Med.* 2001;134(3):224-38.
- Reaven G and Tsao PS:** Insulin resistance and compensatory hyperinsulinemia: the key player between cigarette smoking and cardiovascular disease? *J Am Coll Cardiol* 2003; 41: 1044.

- Ridker M and Libby P:** Risk factors for atherothrombotic diseases. In Libby P, Zipes P, Bonow O and Braunwald E. (eds): Braunwald's heart disease. A text book of cardiovascular medicine 7th edition. Elsevier saunders, Philadelphia, 2005. Chapter 36 : pp 939-959.
- Ridker PM and Libby P.** Risk factors for atherothrombotic disease. In Braunwald's Heart Disease. A text book of cardiovascular medicine 8th Edition. (eds) Libby P, Bonow RO, Mann DL, Zipes DP and Braunwald's E. Elsevier Saunders. Philadelphia, 2008. chapter39; pp 1003-1022.
- Ridker PM, Manson JE and Gaziano JM.** Primary and secondary prevention of coronary heart disease. In Braunwald's Heart Disease. A text book of cardiovascular medicine 8th Edition. (eds) Libby P, Bonow RO, Mann DL, Zipes DP and Braunwald's E. Elsevier Saunders. Philadelphia, 2008. chapter 45; pp1119-1145.
- Ridker PM, Rifai N, Cook NR, et al.** Non HDL cholesterol, apolipoprotein A-1 and B100, standard lipid measures, lipid ratios and CRP as risk factors for cardiovascular diseases in women. JAMA. 2005; 294:326.
- Ridker PM, Rifai N, Rose LL, et al.** Comparison of C-reactive protein and low-density lipoprotein cholesterol levels in the prediction of first cardiovascular events. *N Engl J Med.* 2002; 347:1557 -1565.
- Rigotti NA and Pasternak RC.** Cigarette smoking and coronary heart disease: risks and management. *Cardiol Clin* 1996;14:51-68.

- Rimm EB, Stampfer MJ, Giovannucci E, et al.** Body size and fat distribution as predictors of coronary heart disease among middle-aged and older US men. *Am J Epidemiol* 1995; 141: 1117–27.
- Rioufol G, Finet G, Ginon I, et al.** Multiple atherosclerotic plaque rupture in *acute* coronary syndrome: a three-vessel intravascular ultrasound study. *Circulation*. 2002; 106(7): 804-8.
- Robert J. Chilton.** Pathophysiology of Coronary Heart Disease: A Brief Review. *The Journal of The American Osteopathic Association JAOA*, September 2004 • Vol 104 • No 9 suppl 5-8.
- Roffi M, Chew DP, Mukherjee D, et al.** Platelet glycoprotein IIb/IIIa inhibitors reduce mortality in diabetic patients with non-ST-segment- elevation acute coronary syndromes. *Circulation*; 2001; 104:2767-2771.
- Roglic G, Unwin N, Bennett PH, et al.** The burden of mortality attributable to diabetes: realistic estimates for the year 2000. *Diabetes Care* 2005; 28:2130–2135.
- Rosengren A, Wallentin L, Simoons M, et al.** Cardiovascular risk factors and clinical presentation in acute coronary syndromes *Heart*. 2005; 91(9): 1141–1147.
- Sabatine MS, Cannon CP, Gibson CM, et al.** Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. *N Engl J Med* 2005;352(12):1179-89.

- Sala M, Barcena JIP, Rota JJ, et al.** Sustained ventricular tachycardia as a marker of inadequate myocardial perfusion during the acute phase of myocardial infarction. *Am Heart J* 2001;145:515-21.
- Santos ES, Minuzzo L, Pereira MP, et al.** Acute coronary syndrome registry at a cardiology emergency center. *Arq Bras Cardiol* 2006; 87(5):597-602.
- Schaefer EJ, Moussa PB, Wilson PWF, et al.** Plasma lipoproteins in healthy octogenarians: lack of reduced high-density lipoprotein cholesterol levels: results from the Framingham Heart Study. *Metabolism*. 1989;38: 293-296.
- Scheuner William C. Whitworth, Henraya Mc Gruder, et al.** Familial risk assessment for early-onset coronary heart disease *Genet Med* 2006;8(8): 525–531
- Scirica BN and Morrow BA.** Troponins in acute coronary syndromes. *Semin Vasc Med* 2003; 3:363-74.
- Serum marker analysis in acute myocardial infarction. American College of Emergency Physicians. *Ann Emerg Med* 2000; 35:534-9.
- Sheridan S, Pignone M and Mulrow C.** Framingham-based tools to calculate the global risk of coronary heart disease: a systematic review of tools for clinicians. *J Gen Intern Med* 2003;18:1039–1052.
- Simes RJ, Topol EJ, Holmes DR Jr, et al.** Link between the angiographic substudy and mortality outcomes in a large randomized trial of myocardial reperfusion: importance of early and complete infarct artery reperfusion. *Circulation* 1995; 91:1923– 8.

- Slater DK, Hlatky MA, Mark DB, et al.** Outcome in suspected acute myocardial infarction with normal or minimally abnormal admission electrocardiographic findings. *Am J Cardiol* 1987; 60: 766-70.
- Smalling RW, Bode C, Kalbfleisch J, et al.** More rapid, complete, and stable coronary thrombolysis with bolus administration of reteplase compared with alteplase infusion in acute myocardial infarction. *Circulation* 1995;91:2725–32.
- Somers MP, Brady WJ, Perron AD, et al.** The prominent T wave: electrocardiographic differential diagnosis." *Am J Emerg Med* 2002; 20(3): 243-51.
- Spence JD, Barnett PA, Bulman DE, et al.** An approach to ascertain probands with non-traditional risk factor for carotid atherosclerosis. *Atherosclerosis*. 1999; 144:429-434.
- Stamler J, Neaton JD, Garside DB, et al.** Current status: six established major risk factors – and low risk. In: Marmot MG, Elliott P, editors. *Coronary heart disease epidemiology: from etiology to public health*. Oxford: Oxford University Press; 2005; 32-70.
- Stampfer MJ, Hu FB, Manson JE, et al.** Primary prevention of coronary heart disease in women through diet and lifestyle. *N Engl J Med* 2000; 343:16-22.
- Stone GW, White HD, Ohman EM, et al.** Bivalirudin in patients with acute coronary syndromes undergoing percutaneous coronary intervention: a subgroup analysis from the Acute Catheterization and Urgent Intervention Triage strategy (ACUITY) trial. *Lancet* 2007; 369(9565),907-919.

- Sullivan AK, Holdright DR, Wright CA, et al.** Chest pain in women: clinical, investigative, and prognostic features. *BMJ* 1994; 308: 883-6.
- Tang EW, Wong C, Restieaux NJ, et al.** Clinical outcome of older patients with acute coronary over the last three decades. *Age and Ageing* 2006; 35:280-285.
- Tang EW, Wong C, and Herbison P.** Global registry of acute coronary events (GRACE) hospital discharge risk score accurately predicts long-term mortality post acute coronary syndrome. *Am Heart J* 2007;153:229-35.
- Terkelsen CJ, Lassen JF, Norgaard BL, et al.** Mortality rates in patients with ST-elevation vs. non-ST-elevation acute myocardial infarction: observations from an unselected cohort. *Eur Heart J* 2005; 26:18-26.
- The Direct Thrombin Inhibitor Trialists' Collaborative Group.** Direct thrombin inhibitors in acute coronary syndromes: principal results of a meta-analysis based on individual patients' data. *Lancet*; 2002; 359:294-302.
- Theroux P and Welsh RC.** Meta-analysis of randomised trials comparing enoxaparin versus unfractionated heparin as adjunctive therapy to fibrinolysis in ST-elevation acute myocardial infarction. *Am Cardiol* 2003;91(7):860-4.
- Turcato E, Bosello O, Di Francesco V, et al.** Waist circumference and abdominal sagittal diameter as surrogates of body fat distribution in the elderly: their relation with cardiovascular risk factors. *Int J Obes Relat Metab Disord* 2000;24:1005-10.

- Vaccarino V, Parsons L, Every NR, et al.** Sex-based differences in early mortality after myocardial infarction. National Registry of Myocardial Infarction 2 Participants. *N Engl J Med* 1999; 341:217-225.
- Vaccaro O, Eberly LE, Neaton JD, et al.** Multiple Risk Factor Intervention Trial Research Group: Impact of diabetes and previous myocardial infarction on long-term survival: 25-year mortality follow-up of primary screenees of the Multiple Risk Factor Intervention Trial. *Arch Intern Med* 2004; 164: 1438–1443.
- Volpi A, Cavalli A, Turato R, et al.** incidence and short-term prognosis of late sustained ventricular tachycardia after myocardial infarction. *Am Heart J* 2001;142:87-92.
- Wallentin L, Becker RC, Budaj A, et al.** PLATO Investigators, Freij A, Thorsén MN *Engl J Med*. 2009; 361(11):1045-57.
- Wannamethee SG, Shaper AG and Walker M.** Changes in physical activity mortality, and incidence of coronary heart disease in older men. *Lancet* 1998; 351: 1603–8.
- Warkentin TE, Levine MN, Hirsh J, et al.** Heparin-induced thrombocytopenia in patients treated with low-molecular-weight heparin or unfractionated heparin. *N Engl J Med*; 2004; 332:1330-1335.
- Waters DD, Hsue PY.** What is the role of intensive cholesterol lowering in the treatment of acute coronary syndromes? *Am J Cardiol* 2001; 88:7J–16J.
- Wenger NK.** Coronary heart disease: The female heart is vulnerable. *Prog Cardiovasc Dis* 2003; 46: 199-229.

- White HS, Simes RJ, Aylward PEG, et al.** Thrombin-specific anticoagulation with bivalirudin versus heparin in patients receiving fibrinolytic therapy for acute myocardial infarction the HERO-2 randomised trial. *Lancet* 2001; 358(9296): 1855-63.
- Yusuf S, Mehta SR, Zaho F, et al.** Early and late effect of clopidogrel in patients with acute coronary syndrome: *circulation* 2003; 107(7):966-72.
- Zimetbaum PJ and Josephson ME.** Use of the electrocardiogram in acute myocardial infarction. *N Engl J Med* 2003; 348:933-40.
- Zubaid M, Rashed WA, Al-Khaja N, et al.** Clinical presentation and outcomes of acute coronary syndromes in the gulf registry of acute coronary events (Gulf RACE). *Saudi Med J*; 2008; 29(2): 251-255.