

REFERENCES

- Adams, J.D.; Flora, K.P.; Goldspiel, B.R.; Wilson, J.W.; Arbuck, S.G. and Finley, R. (1993):** Taxol: a history of pharmaceutical development and current pharmaceutical concerns. *J Natl Cancer Inst Monogr.*, 15: 141–147.
- Agarwal, S.K.; and Brenner, M.B. (2006):** Role of adhesion molecules in synovial inflammation. *Curr Opin Rheumatol.*, 18: 268–276.
- Altman, R.; Asch, E. and Bloch, D. (1986):** Development of criteria for the classification and reporting of osteoarthritis .Classification of osteoarthritis of the knee. *Arthritis Rheum.*, 29:1039-1049.
- Arnett, F.C.; Edworthy, S. and Block, D.A. (1988):** The 1987 Revised ACR criteria for classification of RA. *Arthritis Rheum.*, 31: 315-324.
- Athanasou, N.A. and Quinn, J. (1991):** Immunocytochemical analysis of human synovial lining cells: phenotypic relation to other marrow derived cells. *Ann Rheum Dis.*,50:311.
- Backhaus, M.; Burmester, G.R.; Gerber, T.; Grassi, W.; Machold, K.P. and Swen W.A. (2001):** Guidelines for musculoskeletal ultrasound in rheumatology. *Ann Rheum Dis.*, 60:641 – 649.
- Bajaj, S.; Lopez-Ben, R.; Oster, R. and Alarcon, G.S. (2007):** Ultrasound detects rapid progression of erosive disease in early rheumatoid arthritis: a prospective longitudinal study. *Skeletal Radiol.*, 36(2):123–128.
- Badger, A.M.; Blake, S.; Kapadia, R.; Sarkar, S.; Levin, J.; Swift, B.A.; Hoffman, S.J.; Stroup, G.B.; Miller, W.H.; Gowen, M. and Lark, M.W.(2001):** Disease-modifying activity of SB 273005, an orally active, nonpeptide alphavbeta3 (vitronectin receptor) antagonist, in rat adjuvant-induced arthritis. *Arthritis Rheum.*, 44: 128–137.

- Balint, P.V.; Kane, D.; Hunter, J.; McInnes, I.B.; Field, M. and Sturrock, R.D. (2002):** Ultrasound guided versus conventional joint and soft tissue fluid aspiration in rheumatology practice: a pilot study. *J Rheumatol.*, 29:2209–2213.
- Balint, P.V.; Kane, D. and Sturrock, R.D. (2001):** Modern patient management in rheumatology: interventional musculoskeletal ultrasonography. *Osteoarthritis Cartilage*, 9:509–511.
- Balint, P.V.; Kane, D.; Wilson, H.; McInnes, I.B. and Sturrock, R.D. (2002):** Ultrasonography of enthesal insertions in the lower limb in spondyloarthritis. *Ann Rheum Dis.*, 61:905–910.
- Ballara, S.; Taylor, P.C.; Reusch, P.; Marme, D.; Maini, R.N. and Paleolog, E. (2001):** Serum vascular endothelial growth factor (VEGF) and soluble VEGF receptor in inflammatory arthritis. *Arthritis Rheum.*, 44: 2055–2064.
- Bergholm, R.; Leirisalo-Repo, M.; Vehkavaara, S.; Makimattila, S.; Taskinen, M.R. and Yki-Jarvinen, H. (2002):** Impaired responsiveness to NO in newly diagnosed patients with rheumatoid arthritis. *Arterioscler Thromb Vasc Biol.*, 22:1637–1641.
- Berry, R. and Huskisson, M. J. (1972):** Treatment of rheumatoid arthritis. *Clinical Trials J.*, 9: 13-15.
- Bodolay, E.; Koch, A.E.; Kim, J.; Szegedi, G. and Szekanecz, Z. (2002):** Angiogenesis and chemokines in rheumatoid arthritis and other systemic inflammatory rheumatic diseases. *J Cell Mol Med.*, 6: 357–376.
- Boyle, D.L.; Jones, T.J. and Hammaker, D. (2006):** Modulation of peripheral inflammation by spinal p38 MAP kinase. *PLoS Med.*, 3:e338.
- Bradfield, P.F.; Amft, N. and Vernon-Wilson, E. (2003):** Rheumatoid fibroblast-like synoviocytes overexpress the chemokine stromal cell-derived factor 1 (CXCL12), which supports distinct patterns and rates of CD4⁺ and CD8⁺ T

- cell migration within synovial tissue. *Arthritis Rheum.*, 48:2472-2482.
- Brahn, E.; Tang, C. and Banquerigo, M.L.(1994):** Regression of collagen-induced arthritis with taxol, a microtubule stabilizer. *Arthritis Rheum.*, 37: 839–845.
- Braun, A.; Takemura, S.; Vallejo, A.N.; Goronzy, J.J. and Weyand, C.M.(2004):** Lymphotoxin beta-mediated stimulation of synoviocytes in rheumatoid arthritis. *Arthritis Rheum.*, 50:2140-2150.
- Brennan, F. and Beech, J. (2007):** Update on cytokines in rheumatoid arthritis. *Curr Opin Rheumatol.*, 19: 296–301.
- Brown, A.K. (2009):** Using ultrasonography to facilitate best practice in diagnosis and management of RA *Nat Rev Rheumatol.*, 5(12):698-706.
- Bruyn, G.A. and Schmidt, W.A. (2006):** Introductory Guide to Musculoskeletal Ultrasound for the Rheumatologist. Bohn Stafleu & Van Loghum.
- Burman, A.; Haworth, O. and Hardie, D.L. (2005):** A chemokine dependent stromal induction mechanism for aberrant lymphocyte accumulation and compromised lymphatic return in rheumatoid arthritis. *J Immunol.*, 174:1693-1700.
- Campbell, R.S. and Grainger, A.J. (2001):** Current concepts in imaging of tendinopathy. *Clin Radiol.*, 56:253–267.
- Caplan, A.I. (1991):** Mesenchymal stem cells. *J Orthop Res.*, 9:641-650.
- Carotti, M.; Salaffi, F.; Manganelli, P. and Argalia, G. (2001):** Ultrasonography and colour doppler sonography of salivary glands in primary Sjogren's syndrome. *Clin Rheumatol.*, 20:213–219.
- Carotti, M.; Salaffi, F.; Manganelli, P.; Salera, D.; Simonetti, B. and Grassi, W. (2002):** Power Doppler sonography in the assessment of synovial tissue of the knee joint in rheumatoid arthritis: preliminary experience. *Ann Rheum Dis.*, 61:877–882.

- Cheng, K.S.; Tiwari, A. and Boutin, A. (2003):** Differentiation of primary and secondary Raynaud's disease by carotid arterial stiffness. *Eur J Vasc Endovasc Surg.*, 25:336–341
- Chu, C.Q.; Field, M. and Allard, S., (1992):** Detection of cytokines at the cartilage/pannus junction in patients with rheumatoid arthritis: implications for the role of cytokines in cartilage destruction and repair. *Br J Rheumatol.*, 31:653-661.
- Cid, M.C.; Hernández-Rodríguez, J.; Esteban, M.J.; Cebrián, M.; Gho, Y.S.; Font, C.; Urbano-Márquez, A.; Grau, J.M. and Kleinman, H.K. (2002):** Tissue and serum angiogenic activity is associated with low prevalence of ischemic complications in patients with giant-cell arteritis. *Circulation*, **106**: 1664–1671.
- Clavel, G.; Bessis, N. and Lemeiter, D. (2007):** Angiogenesis markers (VEGF, soluble receptor of VEGF and angiopoietin-1) in very early arthritis and their association with inflammation and joint destruction. *Clin. Immunol.*, 124(2), 158-164.
- Clavel, G. and Boissier M.(2008):** Angiogenesis Markers in Rheumatoid Arthritis. *Future Rheumatol.*, 3(2):153-159.
- Cooperberg, P.L.; Tsang, I.; Truelove, L. and Knickerbocker, W.J. (1978):** Gray scale ultrasound in the evaluation of rheumatoid arthritis of the knee. *Radiology*, 126:759-763.
- Corr, M. and Zvaifler, N.J. (2002):** Mesenchymal precursor cells. *Ann Rheum Dis.*, 61:3-5.
- Coxon, A.; Bolon, B.; Estrada, J.; Kaufman, S.; Scully, S. and Rattan, A. (2002):** Inhibition of interleukin-1 but not tumor necrosis factor suppresses neovascularization in rat models of corneal angiogenesis and adjuvant arthritis. *Arthritis Rheum.*, 46:2604–2612.

- Csernok, E.; Szymkowiak, C.H.; Mistry, N.; Daha, M.R.; Gross, W.L. and Kekow, J. (1996):** Transforming growth factor- β (TGF- β) expression and interaction with proteinase 3 (PR3) in anti-neutrophil cytoplasmic antibody (ANCA).associated vasculitis. *Clin Exp Immunol.*, **105**: 104–109.
- Cyteval, C. (2009):** Doppler Ultrasonography and Dynamic Magnetic Resonance Imaging for Assessment of Synovitis in the Hand and Wrist of Patients with Rheumatoid Arthritis. *Semin Musculoskelet Radiol.*, 13:66–73.
- D’Agostino, M.A.; Said-Nahal, R.; Hacquard-Bouder, C.; Brasseur, J.L.; Dougados, M. and Breban, M. (2003):** Assessment of peripheral enthesitis in the spondylarthropathies by ultrasonography combined with power Doppler: a cross-sectional study. *Arthritis Rheum.*,48:523–533.
- Daniel, T.O.; Liu, H.; Morrow, J.D.; Crews, B.C. and Marnett, L.J. (1999):** Thromboxane A2 is a mediator of cyclooxygenase-2-dependent endothelial migration and angiogenesis. *Cancer Res.*,59:4574–4577.
- Debusk, L.M.; Chen, Y.; Nishishita, T.; Chen, J.; Thomas, J.W. and Lin, P.C. (2003):** Tie2 receptor tyrosine kinase, a major mediator of tumor necrosis factor α induced angiogenesis in rheumatoid arthritis. *Arthritis Rheum.*, **48**: 2461–2471.
- De Flaviis, L.; Scaglione, P.; Nessi, R.; Ventura, R. and Calori, G. (1988):** Ultrasonography of the hand in rheumatoid arthritis. *Acta Radiol.*, 29:457-460.
- Desmouliere, A.; Redard, M.; Darby, I. and Gabbiani ,G.(1995):** Apoptosis mediates the decrease in cellularity during the transition between granuloma tissue and scar. *Am J Pathol.*, 146:56–66.
- Distler, J.H.; Gay, S. and Distler, O. (2006):** Angiogenesis and vasculogenesis in systemic sclerosis. *Rheumatology (Oxford)*, 45 Suppl 3: iii26–iii27.
- Drouart, M.; Saas, P.; Billot, M.; Cedoz, J.P.; Tiberghien, P.; Wendling, D. and Toussirot, E.(2003):** High serum

vascular endothelial growth factor correlates with disease activity in spondylarthropathies. *Clin Exp Immunol.*, **132**: 158–162.

Edwards, J.C. and Willoughby, D.A. (1982): Demonstration of bone marrow derived cells in synovial lining by means of giant intracellular granules as genetic markers. *Ann Rheum Dis.*, **41**:177-182.

Ellegaard, K.; Torp-Pedersen, S. and Lund, H. (2008): Quantification of colour Doppler activity in the wrist in patients with rheumatoid arthritis—the reliability of different methods for image selection and evaluation. *Ultraschall Med.*, **29**(4): 393–398.

Etchepare, F.; Fautrel, B. and Foltz, V. (2003): Elbow synovitis in RA: a sonographic approach. *Ann Rheum Dis.*, **62** (Suppl.1):328-331.

Ferrara, N.; Gerber, H. and J. LeCouter J. (2003): The biology of VEGF and its receptors. *Nature Medicine*, **9**, (6): 669–676.

Fessell, D.P.; Jacobson, J.A.; Craig, J. Habra, G.; Prasad, A.; Radliff, A. and van Holsbeeck, M.T. (2000): Using sonography to reveal and aspirate joint effusions. *Am J Roentgenol.*, **174**:1353–1362.

Fiocco, U.; Cozzi, L.; Rubatelli, L.; Rigon, C.; De Candia, A.; Tregnaghi, A.; Gallo, C.; Favaro, M.; Binachi, F.; Baldovin, M. and Todesco, S. (1996): Long term sonographic follow up of rheumatoid and psoriatic proliferative knee joint synovitis. *Br J Rheumatol.*, **35**:155-163.

Fiocco, U.; Ferro, F.; Vezzu, M.; Cozzi, L.; Checchetto, C., Sfriso, P.; Botsios, C.; Ciprian, L.; Armellin, G.; Nardacchione, R.; Piccoli, A.; Todesco, S. and Rubaltelli, L.(2005): Rheumatoid and psoriatic knee synovitis: clinical, grey scale, and power Doppler ultrasound assessment of the response to etanercept. *Ann Rheum Dis.*, **64**:899–905.

- Firestein, G.S. (2008):** Etiology and pathogenesis of rheumatoid arthritis. In: Kelly's Textbook of Rheumatology; Ruddy S, Harris E, Sledge C; 8th ed., W. B. Saunders, Philadelphia, PA,:921-966.
- Firestein, G.S.(2003):** Evolving concepts of rheumatoid arthritis. *Nature*, 423: 356–36.
- Firestein, G.S. (1996):** Invasive fibroblast-like synoviocytes in rheumatoid arthritis: passive responders or transformed aggressors?. *Arthritis Rheum.*,39 (11) :1781– 1790.
- Firestein, G.S.; Echeverri, F.; Yeo, M.; Zvaifler, N.J. and Green, D.R.(1997):** Somatic mutations in the p53 tumor suppressor gene in rheumatoid arthritis synovium. *Proc Natl Acad Sci U S A.*, 94:10895-10900.
- Firestein, G.S.; Nguyen, K.; Aupperle, K.R.; Yeo, M.; Boyle, D.L. and Zvaifler, N.J.(1996):** Apoptosis in rheumatoid arthritis: p53 overexpression in rheumatoid arthritis synovium. *Am J Pathol.*, 149:2143-2153.
- Firestein, G.S.; Paine, M.M. and Littman, B.H. (1991):** Gene expression (collagenase, tissue inhibitor of metalloproteinases, complement, and HLA-DR) in rheumatoid arthritis and osteoarthritis synovium. Quantitative analysis and effect of intraarticular corticosteroids. *Arthritis Rheum.*, 34:1094-1105.
- Fisher, R.A. (1964):** Statistical methods of research workers 10th ed., Edinburgh, Pliver and Boyed.
- FitzGerald, O. and Bresnihan, B. (1995):** Synovial membrane cellularity and vascularity. *Ann Rheum Dis.*, 54:511–515.
- Fornage, B.D. (2000):** The case for ultrasound of muscles and tendons. *Semin Musculoskelet Radiol.*, 4:375–391.

-
- Foster, W.; Carruthers, D.; Lip, G.Y. and Blann, A.D.(2009):** Relationships between endothelial, inflammatory and angiogenesis markers in rheumatoid arthritis: implications for cardiovascular pathophysiology. *Thromb Res.*, 123(4):659-564.
- Franz, J.K.; Pap, T.; Hummel, K.M.; Nawrath, M.; Aicher, W.K.; Shigeyama, Y.; Müller-Ladner, U.; Gay, R.E. and Gay, S. (2000):** Expression of sentrin, a novel antiapoptotic molecule, at sites of synovial invasion in rheumatoid arthritis. *Arthritis Rheum.*, 43:599-607.
- Gelb, B.D.; Shi, G.P.; Chapman, H.A. and Desnick, R.J. (1996):** Pycnodysostosis, a lysosomal disease caused by cathepsin K deficiency. *Science*, 273:1236-1238.
- Georgopoulos, S.; Plows, D. and Kollias, G. (1996):** Transmembrane TNF is sufficient to induce localized tissue toxicity and chronic inflammatory arthritis in transgenic mice. *J Inflamm.*, 46:86-97.
- Giatromanolaki, A.; Sivridis, E.; Maltezos, E.; Athanassou, N.; Papazoglou, D.; Gatter, K.C.; Harris, A.L. and Koukourakis, M.I. (2003):** Upregulated hypoxia inducible factor-1 α and -2 α pathway in rheumatoid arthritis and osteoarthritis. *Arthritis Res Ther.*, 5: R193–R198.
- Goldbach-Mansky, R.; Lee, J.M.; Hoxworth, J.M.; Smith, D.; Duray, P.; Yarboro, C.H.; Klippel, J.; Kleiner, D. and El-Gabalawy, H.S. (2000):** Active synovial matrix metalloproteinase-2 is associated with radiographic erosions in patients with early synovitis. *Arthritis Res.*, 2:145-153.
- Grassi, W.; Farina, A.; Filippucci, E. and Cervini, C. (2002):** Intralesional therapy in carpal tunnel syndrome: a sonographic-guided approach. *Clin Exp Rheumatol.*, 20:73–76.
- Grassi, W.; Filippucci, E.; Farina, A. and Cervini, C. (2000):** Sonographic imaging of tendons. *Arthritis Rheum.*, 43:969–76.

- Grassi, W.; Lamanna, G.; Farina, A. and Cervini, C.(1999):** Sonographic imaging of normal and osteoarthritic cartilage. *Semin Arthritis Rheum.*, 28:398–403.
- Gravallese, E.M.; Handel, M.L.; Coblyn, J.; Anderson, R.J.; Sperling, R.I.; Karlson, E.W.; Maier, A.; Ruderman, E.M.; Formelli, F. and Weinblatt, M.E. (1996):** N-[4-hydroxyphenyl] retinamide in rheumatoid arthritis: a pilot study. *Arthritis Rheum.*, 39:1021-1026.
- Greher, M. and Kapral, S. (2003):** Is regional anesthesia simply an exercise in applied sonoanatomy?: aiming at higher frequencies of ultrasonographic imaging. *Anesthesiology*, 99:250–251.
- Gudbjörnsson, B.; Christofferson, R. and Larsson, A.(2004):** Synovial concentrations of the angiogenic peptides bFGF and VEGF do not discriminate rheumatoid arthritis from other forms of inflammatory arthritis. *Scand J Clin Lab Invest.*, 64(1):9-15.
- Hamann, J, Wishaupt, JO, van Lier, RA, Smeets TJ, Breedveld FC, Tak PP. (1999):** Expression of the activation antigen CD97 and its ligand CD55 in rheumatoid synovial tissue. *Arthritis Rheum.*, 42:650-658.
- Handel, M.L.; McMorrow, L.B. and Gravallese, E.M. (1995):** Nuclear factor-kappa B in rheumatoid synovium. Localization of p50 and p65. *Arthritis Rheum.*, 38:1762-.1870
- Hannig G, Bernier SG, Hoyt JG, et al.** Suppression of inflammation and structural damage in experimental arthritis through molecular targeted therapy with PPI-2458. *Arthritis Rheum* 2007;56:850–60.
- Harada, M.; Mitsuyama, K.; Yoshida, H.; Sakisaka, S.; Taniguchi, E.; Kawaguchi, T.; Ariyoshi, M.; Saiki, T.; Sakamoto, M.; Nagata, K.; Sata, M.; Matsuo, K. and Tanikawa, K. (1998) :** Vascular endothelial growth factor in

patients with rheumatoid arthritis. *Scand J Rheumatol.*, 27: 377-380.

Haringman, J.J.; Kraan, M.C.; Smeets, T.J.; Zwinderman, K.H. and Tak, P.P.(2003): Chemokine blockade and chronic inflammatory disease: proof of concept in patients with rheumatoid arthritis. *Ann Rheum Dis.*, 62:715-721.

Hauschild ,A.; Kroeger, H.; Mitchison, N.A.; Ugrinovic, S. and Zwingenberger, K. (1997): Thalidomide therapy of established collagen-induced arthritis (CIA) not accompanied by an evident Th2 shift. *Clin Exp Immunol.*, 108:428–431.

Hebbar, M.; Peyrat, J.P.; Hornez, L.; Hatron, P.Y.; Hachulla, E. and Devulder, B.(2000): Increased concentrations of the circulating angiogenesis inhibitor endostatin in patient with systemic sclerosis. *Arthritis Rheum.*, 43: 889–893.

Hetland, M.,; Christensen J.I.; Lottenburger,T.; Johansen, J.S.; Svendsen, M.N.; Kim, H.; Nielsen, L. and Nielsen, H.J.(2008): Circulating VEGF as a biological marker in patients with rheumatoid arthritis? Preanalytical and biological variability in healthy persons and in patients. *Disease Markers*, 24(1):1-10.

Hirohata, S. and Sakakibara, J. (1999): Angioneogenesis as a possible elusive trigger factor in rheumatoid arthritis. *Lancet*, 353:1331-1338.

Hochberg, M. C.; Chang, R. W.; Dwosh, I.; Lindsey, S.; Pincus, T. and Wolfe, F. (1992): the American College Of Rheumatology 1991 revised criteria for the classification of global functional status in rheumatoid arthritis. *Arthritis Rheum.*, 35(5): 498-502.

Hou, W.S.; Li, W.; Keyszer, G.; Levy, R.; Klein, M.J.; Gravallesse, E.M.; Goldring, S.R. and Brömme, D. (2002): Comparison of cathepsins K and S expression within the rheumatoid and osteoarthritic synovium. *Arthritis Rheum.*, 46:663-674.

-
- Huizinga, T.W.; Dijkmans, B.A.; van der Velde, E.A.; van de Pouw; Kraan, T.C.; Verweij, C.L. and Breedveld, F.C.(1996):** An open study of pentoxifylline and thalidomide as adjuvant therapy in the treatment of rheumatoid arthritis. *Ann Rheum Dis.*, 55:833–836.
- Imamura, F.; Aono, H.; Hasunuma, T.; Sumida, T.; Tateishi, H.; Maruo, S. and Nishioka, K. (1998):** Monoclonal expansion of synoviocytes in rheumatoid arthritis. *Arthritis Rheum.*, 41:1979-1986.
- Ishikawa, H, Hirata, S, Andoh, Y, Kubo H, Nakagawa N, Nishibayashi Y, Mizuno K. (1996):** An immunohistochemical and immunoelectron microscopic study of adhesion molecules in synovial pannus formation in rheumatoid arthritis. *Rheumatol Int.*, 16:53-60.
- Jacobson, J.A.; Andresen, R.; Jaovisidha, S.; De Maeseneer, M.; Foldes, K.; Trudell, D.R. and Resnick, D. (1998):** Detection of ankle effusions: comparison study in cadavers using radiography, sonography, and MR imaging. *Am J Roentgenol.*, 170:1231–1238.
- Jackson, J.R.; Minton, J.A.; Ho, M.L.; Wei, N. and Winkler, J.D. (1997):** Expression of vascular endothelial growth factor in synovial fibroblasts is induced by hypoxia and interleukin 1beta. *J Rheumatol.*, **24**: 1253–1259.
- Jimenez, B.; Volpert, O.V.; Crawford, S.E.; Febbraio, M.; Silverstein, R.L. and Bouck, N. (2000):** Signals leading to apoptosis-dependent inhibition of neovascularization by thrombospondin-1. *Nat Med.*, 6:41–48.
- Jones, P.B.; Makki, R.J. and Weiss, J.B. (1994):** Endothelial cell stimulating angiogenesis factor□a new biological marker for disease activity in ankylosing spondylitis? *Br J Rheumatol.*, 33: 332–335.
- Jones, A.; Regan, M.; Ledingham, J.; Pattrick, M.; Manhire, A. and Doherty, M. (1993):** Importance of placement of intra-articular steroid injections. *Br Med J.*, 307:1329–1330.

- Jongbloed, S.L.; Lebre, M.C.; Fraser, A.R.; Gracie, J.A.; Sturrock, R.D.; Tak, P.P. and McInnes, I.B. (2005):** Enumeration and phenotypical analysis of distinct dendritic cell subsets in psoriatic arthritis and rheumatoid arthritis. *Arthritis Res Ther.*, 8:R15.
- Joosten, L.A.; van De Loo, F.A.; Lubberts, E.; Helsen, M.M.; Netea, M.G.; van Der Meer, J.W.; Dinarello, C.A. and van Den Berg, W.B. (2000):** An IFN-gamma-independent proinflammatory role of IL-18 in murine streptococcal cell wall arthritis. *J Immunol.*, 165:6553-6558.
- Joshua, F.; Lassere, M.; Bruyn, G.A.; Szkudlarek, M.; Naredo, E.; Schmidt, W.A.; Balint, P.; Filippucci, E.; Backhaus, M.; Iagnocco, A.; Scheel, A.K.; Kane, D.; Grassi, W.; Conaghan, P.G.; Wakefield, R.J. and D'Agostino, M.A. (2007):** Summary findings of a systematic review of the ultrasound assessment of synovitis. *J Rheumatol.*, 34:839-847.
- Kamatani, N. (2001):** Activated human T cells directly induce osteoclastogenesis from human monocytes: possible role of T cells in bone destruction in rheumatoid arthritis patients. *Arthritis Rheum* 2001; 44:1003.
- Kane, D.; Balint, P.V. and Sturrock, R.D. (2003):** Ultrasonography is superior to clinical examination in the detection and localization of knee joint effusion in rheumatoid arthritis. *J Rheumatol.*, 30:966–971.
- Kane D, Balint PV, Sturrock R, et al¹.(2004):** Musculoskeletal ultrasound—a state of the art review in rheumatology. Part 1: Current controversies and issues in the development of musculoskeletal ultrasound in rheumatology. *Rheumatology (Oxford)*, 43:823– 828.
- Kane, D.; Grassi, W.; Sturrock, R. and Balint², P. (2004):** Musculoskeletal ultrasound—a state of the art review in rheumatology. Part 2: Clinical indications for musculoskeletal ultrasound in rheumatology. *Rheumatology(Oxford)*, 43(7):829–838.
- Kane, D.; Greaney, T.; Bresnihan, B.; Gibney, R. and FitzGerald, O. (1999):** Ultrasonography in the diagnosis and

- management of psoriatic dactylitis. *J Rheumatol.*, 26:1746–1751.
- Kane, D.; Greaney, T.; Shanahan, M.; Duffy, G.; Bresnihan, B.; Gibney, R. and FitzGerald, O. (2001):** The role of ultrasonography in the diagnosis and management of idiopathic plantar fasciitis. *Rheumatology (Oxford)*, 40:1002–1008.
- Karim, Z.; Wakefield, R.J.; Conaghan, P.G.; Lawson, C.A.; Goh, E.; Quinn, M.A.; Astin, P.; O'Connor, P.; Gibbon, W.W. and Emery, P. (2001):** The impact of ultrasonography on diagnosis and management of patients with musculoskeletal conditions. *Arthritis Rheum.*, 44:2932–2933.
- Kasama, T.; Kobayashi, K.; Yajima, N.; Shiozawa, F.; Yoda, Y.; Takeuchi, H.; Mori, Y.; Negishi, M.; Ide, H. and Adachi, M.(2000):** Expression of vascular endothelial growth factor by synovial fluid neutrophils in rheumatoid arthritis. *Clin Exp Immunol.*, 121:533-538.
- Kasama, T.; Shiozawa, F.; Kobayashi, K.; Yajima, N.; Hanyuda, M.; Takeuchi, H.T.; Mori, Y.; Negishi, M.; Ide, H. and Adachi, M. (2001):** Vascular endothelial growth factor expression by activated synovial leukocytes in rheumatoid arthritis: critical involvement of the interaction with synovial fibroblasts. *Arthritis Rheum.*, 44(11):2512-2524.
- Kasukawa, R.; Shio, K.; Kanno, Y.; Sato, A.; Takahashi, A.; Yamadera, Y. and Kanno, T.(2007):** Doppler ultrasound measurements of knee joint synovitis in rheumatoid arthritis patients treated with infliximab. *Mod Rheumatol.*, 17(5):376-379.
- Kim, J.M.; Ho, S.H.; Park, E.J.; Hahn, W.; Cho, H.; Jeong, J.G.; Lee, Y.W. and Kim, S. (2002):** Angiostatin gene transfer as an effective treatment strategy in murine collagen-induced arthritis. *Arthritis Rheum.*, 46:793–801.
- Kim, W.; Kang, S. and Yoo, S. (2006):** Interaction of vascular endothelial growth factor 165 with neuropilin-1 protects rheumatoid synoviocytes from apoptotic death by regulating

Bcl-2 expression and Bax translocation. *The Journal of Immunology*, 177(8): 5727–5735.

- Kim, H.R.; Park, M.K.; Cho, M.L.; Yoon, C.H.; Lee, S.H.; Park, S.H.; Leng, L.; Bucala, R.; Kang, I.; Choe, J. and Kim, H.Y. (2007):** Macrophage migration inhibitory factor upregulates angiogenic factors and correlates with clinical measures in rheumatoid arthritis. *J Rheumatol.*, 34: 927–936.
- Kinne, R.W.; Boehm, S.; Iftner, T.; Aigner, T.; Vornehm, S.; Weseloh, G.; Bravo, R.; Emmrich, F. and Kroczeck, R.A.(1995):** Synovial fibroblast-like cells strongly express jun-B and C-fos proto-oncogenes in rheumatoid- and osteoarthritis. *Scand J Rheumatol.*, Suppl 101:121-125.
- Kiselyov, A.; Balakin, K.V. and Tkachenko, S.E. (2007):** VEGF/VEGFR signaling as a target for inhibiting angiogenesis. *Expert Opin Investig Drugs*, 16: 83–107.
- Klauser, A.; Frauscher, F.; Schirmer, M.; Halpern, E.; Pallwein, L.; Herold, M.; Helweg, G. and ZurNedden, D.(2002):** The value of contrast-enhanced color Doppler ultrasound in the detection of vascularization of finger joints in patients with rheumatoid arthritis. *Arthritis Rheum.*, 46(3):647–653.
- Koch, A.E. (2003):** Angiogenesis as a target in rheumatoid arthritis. *Ann Rheum Dis.*, 62 Suppl 2:ii60-ii67.
- Koch, A.E.; Kunkel, S.L.; Harlow, L.A.; Johnson, B.; Evanoff, H.L.; Haines, G.K.; Burdick, M.D.; Pope, R.M. and Strieter, R.M. (1992):** Enhanced production of monocyte chemoattractant protein-1 in rheumatoid arthritis. *J Clin Invest.*, 90:772-779.
- Koch, A.E.; Szekanecz, Z.; Friedman, J.; Haines, G.K.; Langman, C.B. and Bouck, N.P. (1998):** Effects of thrombospondin-1 on disease course and angiogenesis in rat adjuvant-induced arthritis. *Clin Immunol Immunopathol.*, 86:199–208.
- Koller, M.; Hayer, S.; Redlich, K.; Ricci, David, J.; Steiner, J.; Smolen, J.; Wagner, E. and Georg Schett (2005):** JNK1 is not essential for TNF-mediated joint disease. *Arthritis Res Ther.*, 7:R166-173.

- Konttinen, Y.T.; Li, T.F.; Mandelin, J.; Liljeström, M.; Sorsa, T.; Santavirta, S. and Virtanen I. (2000):** Increased expression of extracellular matrix metalloproteinase inducer in rheumatoid synovium. *Arthritis Rheum.*, 43:275-280.
- Koski, J.M. (2000):** Ultrasound guided injections in rheumatology. *J Rheumatol.*, 27:2131–2138.
- Koski, J. M.; Saarakkala, M.; Helle, U.; Hakulinen, J. O.; Heikkinen, H. and Hermunen, M. (2006):** Power Doppler ultrasonography and synovitis: correlating ultrasound imaging with histopathological findings and evaluating the performance of ultrasound equipments. *Ann Rheum Dis.*, 65:1590–1595.
- Kotake, S.; Udagawa, N.; Hakoda, M.; Mogi, M.; Yano, K.; Tsuda, E.; Takahashi, K.; Furuya, T.; Ishiyama, S.; Kim, K.J.; Saito, S.; Nishikawa, T.; Takahashi, N.; Togari, A.; Tomatsu, T.; Suda, T. and Kamatani, N. (2001):** Activated human T cells directly induce osteoclastogenesis from human monocytes: possible role of T cells in bone destruction in rheumatoid arthritis patients. *Arthritis Rheum* 2001; 44:1003-1012.
- Kowanetz, M. and Ferrara, N. (2006):** Vascular endothelial growth factor signaling pathways: therapeutic perspective. *Clinical Cancer Research*, 12 (17):5018–5022.
- Kraan, M.C.; Reece, R.J.; Smeets, T.J.; Veale, D.J.; Emery, P. and Tak, PP.(2002):** Comparison of synovial tissues from the knee joints and the small joints of rheumatoid arthritis patients: Implications for pathogenesis and evaluation of treatment. *Arthritis Rheum.*, 46:2034-2038.
- Lainer-Carr, D. and Brahn, E. (2007):** Angiogenesis inhibition as a therapeutic approach for inflammatory synovitis. *Nature clinical practice Rheumatology*, 3 (8):434:442.
- Larsen, A.; Dale, K. and Eek, M. (1977):** Radiographic evaluation of rheumatoid arthritis and related conditions by standard reference films. *Acta Radiol Diag.*, 18: 481-491

-
- Lee, L.; Buckley, C.; Blades, M.C.; Panayi, G.; George, A.J.; Pitzalis, C.; (2002):** Identification of synovium-specific homing peptides by in vivo phage display selection. *Arthritis Rheum.*, 46:2109-2120.
- Lee, S.S.; Joo, Y.S.; Kim, W.U.; Min, D.J.; Min, J.K.; Park, S.H.; Cho, C.S. and Kim, H.Y. (2001):** Vascular endothelial growth factor levels in the serum and synovial fluid of patients with rheumatoid arthritis. *Clin Exp Rheumatol.*, 19:321–324.
- Lee, A. and Langer, R. (1983):** Shark cartilage contains inhibitors of tumor angiogenesis. *Science*, 221:1185–1187.
- Leonard, L.; Rangan, A.; Doyle, G. and Taylor, G. (2003):** Carpal tunnel syndrome—is high-frequency ultrasound a useful diagnostic tool? *J Hand Surg (Br)*., 28:77–79.
- Liu, L.X.; Lu, H.; Luo, Y.; Date, T.; Belanger, A.J.; Vincent, K.A.; Akita, G.Y.; Goldberg, M.; Cheng, S.H.; Gregory, R.J. and Jiang, C. (2002):** Stabilization of vascular endothelial growth factor mRNA by hypoxia-inducible factor 1. *Biochem Biophys Res Commun.*, **291**: 908–914.
- Lydyard, P.M. and Edwards, J.C. (1994):** The pathophysiology of rheumatoid arthritis. *Clin Exp Rheumatol.*, 12 Suppl 11:S55-S59.
- Mabjeesh NJ, Escuin D, LaVallee TM, Pribluda VS, Swartz GM, Johnson MS, Willard MT, Zhong H, Simons JW, Giannakakou P. (2003):** 2ME2 inhibits tumor growth and angiogenesis by disrupting microtubules and dysregulating HIF. *Cancer Cell*, 3:363–375.
- Mallya, R.K. and Mace, B.E.W. (1981):** the assessment of disease activity in RA using the multivariate analysis. *Rheum. Rehabi.*, 20: 14-17.
- Margheri, F.; Manetti, M.; Serrati, S.; Nosi, D.; Pucci, M. and Matucci-Cerinic, M.; (2006):** Domain 1 of the urokinase type plasminogen activator receptor is required for its morphological, functional β 2 integrin-mediated connection with actin cytoskeleton in human microvascular endothelial

cells: failure of association in systemic sclerosis endothelial cells. *Arthritis Rheum.*, **54**:3926–3938.

Markham, T.; Mullan, R.; Golden-Mason, L.; Rogers, S.; Bresnihan, B.; Fitzgerald, O.; Fearon, U. and Veale, D.J. (2006): Regulation of endothelial activation and down-regulation of Tie2 receptor in psoriatic skin after infliximab therapy. *J Am Acad Dermatol.*, **54**: 1003–1012.

Martinoli, C.; Bianchi, S.; Dahmane, M.; Pugliese, F.; Bianchi-Zamorani, M.P. and Valle, M. (2002): Ultrasound of tendons and nerves. *Eur Radiol.*, 12:44–55.

Martinoli, C.; Bianchi, S.; Gandolfo, N.; Valle, M.; Simonetti, S. and Derchi, L.E. (2000): US of nerve entrapments in osteofibrous tunnels of the upper and lower limbs. *Radiographics* 2000;20:S199–S213.

Martinoli, C.; Bianchi, S.; Dahmane, M.; Pugliese, F.; Bianchi-Zamorani, M.P. and Valle, M. (2002): Ultrasound of tendons and nerves. *Eur Radiol.*, 12:44–55.

Maruotti, N.; Cantatore, F.P.; Crivellato, E.; Vacca, A. and Ribatti, D. (2006): Angiogenesis in rheumatoid arthritis. *Histology and Histopathology*, 21 (4–6): 557–566.

McDonnell, C.H.; Jeffrey, R.B.; Bjorkengren, A.G. and Li, K.C. (1992): Intraarticular sonography for imaging the knee menisci: evaluation in cadaveric specimens. *Am J Roentgenol.*, 159:573–574.

Moreland, L.W.; Baumgartner, S.W.; Schiff, M.H.; Tindall, E.A.; Fleischmann, R.M.; Weaver, A.L.; Ettlinger, R.E.; Cohen, S.; Koopman, W.J.; Mohler, K.; Widmer, M.B. and Bloch, C.M. (1997): Treatment of rheumatoid arthritis with a recombinant human tumor necrosis factor receptor (p75)-Fc fusion protein. *N Engl J Med.*, 337:141–147.

Morand, E.F.; Leech, M. and Bernhagen, J. (2006): MIF: a new cytokine link between rheumatoid arthritis and atherosclerosis. *Nat Rev Drug Discov.*, 5: 399–410.

- Mori, S.; Yoshikawa, H.; Hashimoto, J.; Ueda, T.; Funai, H. and Kato, M. (1998):** Antiangiogenic agent (TNP-470) inhibition of ectopic bone formation induced by bone morphogenetic protein-2. *Bone*, 22:99–105.
- Moser, T.L.; Stack, M.S.; Asplin, I.; Enghild, J.J.; Hojrup, P. and Everitt, L. (1999):** Angiostatin binds ATP synthase on the surface of human endothelial cells. *Proc Natl Acad Sci USA*, 96:2811–2816.
- Moses, M.A. (1993):** A cartilage-derived inhibitor of neovascularization and metalloproteinases. *Clin Exp Rheumatol.*, 11(suppl 8):S67–S69.
- Moss, S.G.; Schweitzer, M.E.; Jacobson, J.A.; Brossmann, J.; Lombardi, J.V.; Dellose, S.M.; Coralnick, J.R.; Standiford, K.N. and Resnick, D. (1998):** Hip joint fluid: detection and distribution at MR imaging and US with cadaveric correlation. *Radiology*, 208:43–48.
- Moses MA, Wiederschain D, Wu I, Fernandez CA, Ghazizadeh V, Lane WS, et al.** Troponin I is present in human cartilage and inhibits angiogenesis. *Proc Natl Acad Sci USA* 1999;96:2645–50.
- Mould, A.W.; Tonks, I.D. and Cahill, M.M.; Pettit, A.R.; Thomas, R.; Hayward, N.K. and Kay, G.F. (2003):** *Vegfb* gene knockout mice display reduced pathology and synovial angiogenesis in both antigen-induced and collagen-induced models of arthritis. *Arthritis Rheum.*, 48(9): 2660–2669.
- Mulherin, D.; Fitzgerald, O. and Bresnihan, B. (1996):** Synovial tissue macrophage populations and articular damage in rheumatoid arthritis. *Arthritis Rheum.*, 39:115–119.
- Muller-Ladner, U.; Kriegsmann, J.; Tschopp, J.; Gay, R.E. and Gay, S. (1995):** Demonstration of granzyme A and perforin

messenger RNA in the synovium of patients with rheumatoid arthritis. *Arthritis Rheum.*, 38:477-484.

Nagashima, M.; Tanaka, H.; Takahashi, H.; Tachihara, A.; Tanaka, K. and Ishiwata, T.(2002): Study of the mechanism involved in angiogenesis and synovial cell proliferation in human synovial tissues of patients with rheumatoid arthritis using SCID mice. *Lab Invest.*, 82:981–988.

Nagashima, M.; Wauke, K.; Hirano, D.; Ishigami, S.; Aono, H.; Takai, M.; Sasano, M. and Yoshino, S.(2000): Effects of combinations of anti-rheumatic drugs on the production of vascular endothelial growth factor and basic fibroblast growth factor in cultured synoviocytes and patients with rheumatoid arthritis. *Rheumatology*, 39:1255–1262.

Nakamichi, K.I. and Tachibana, S. (2000): Enlarged median nerve in idiopathic carpal tunnel syndrome. *Muscle Nerve*, 23:1713–1718.

Naredo, E, Moller, I, Moragues, C, et al. Interobserver reliability in musculoskeletal ultrasonography: results from a Teach the Teachers rheumatologist course. *Ann Rheum Dis* 2006; 65:14.

Nawrocki, B.; Polette, M.; Clavel, C.; Morrone, A.; Eschard, J.P.; Etienne, J.C. and Birembaut, P. (1994): Expression of stromelysin 3 and tissue inhibitors of matrix metalloproteinases, TIMP-1 and TIMP-2, in rheumatoid arthritis. *Pathol Res Pract.*, 190:690-696.

Newman, J. S.; Laing, T. J.; McCarthy, C. J. and Adler, R. S. (1996): Power Doppler sonography of synovitis: assessment of therapeutic response—preliminary observations. *Radiology*, 198, 582–584.

Olivieri, I.; Barozzi, L.; Favaro, L.; Pierro, A.; de Matteis, M.; Borghi, C.; Padula, A.; Ferri, S. and Pavlica, P. (1996): Dactylitis in patients with seronegative spondylarthropathy. Assessment by ultrasonography and magnetic resonance imaging. *Arthritis Rheum.*, 39:1524–1528.

- O'Reilly, M.S.; Boehm, T.; Shing, Y.; Fukai, N.; Vasios, G.; Lane, W.S.; et al.(1997):** Endostatin: an endogenous inhibitor of angiogenesis and tumor growth. *Cell*, 88:277–285.
- Østergaard, M. and Szkudlarek M. (2005):** Ultrasonography: a valid method for assessing rheumatoid arthritis? *Arthritis Rheum.*, 52(3):681–686.
- Ozgocmen, S.; Ozdemir, H.; Kiris, A.; Bozgeyik, Z. and Ardicoglu, O.(2008):** Clinical Evaluation and Power Doppler Sonography in Rheumatoid Arthritis: Evidence for Ongoing Synovial Inflammation in Clinical Remission. *Southern Medical Journal*, 101(3):240-245.
- Ozgonenel, L.; Cetin, E.; Tutun, S.; Tonbaklar, P.; Aral, H. and Guvenen, G. (2010):** The relation of serum vascular endothelial growth factor level with disease duration and activity in patients with rheumatoid arthritis. *Clin. Rheumatol.*, 29(5), 473–477.
- Pablos, J.L.; Santiago, B.; Galindo, M.; Torres, C.; Brehmer, M.T.; Blanco, F.J. and García-Lázaro, F.J.(2003):** Synoviocyte-derived CXCL12 is displayed on endothelium and induces angiogenesis in rheumatoid arthritis. *J Immunol.*, **170**: 2147–2152.
- Paleolog, E. (2002):** Angiogenesis in rheumatoid arthritis. *Arthritis Res.*, 4 (suppl 3):S81-S90.
- Patonay, L.; Nagy, K. and Engelke, W. (2002):** Real-time endoarticular ultrasound imaging of the TMJ—a new diagnostic possibility? A cadaver study. *Int J Oral Maxillofac Surg.*, 31:553–557.
- Patterson, A.M.; Gardner, L.; Shaw, J.; David, G.; Loreau, E.; Aguilar, L.; Ashton, B.A. and Middleton, J. (2005):** Induction of a CXCL8 binding site on endothelial syndecan-3 in rheumatoid synovium. *Arthritis Rheum.*, 52:2331-2342.
- Paavonen, K.; Mandelin, J.; Partanen, T.; Jussila, L.; Li, T.F.; Ristimäki, A.; Alitalo, K. and Konttinen, Y.T. (2002):** Vascular endothelial growth factors C and D and their

- VEGFR-2 and 3 receptors in blood and lymphatic vessels in healthy and arthritic synovium. *J Rheumatol.*, 29(1):39-45.
- Peacock, D.J.; Banquerigo, M.L. and Brahn, E.(1995):** A novel angiogenesis inhibitor suppresses rat adjuvant arthritis. *Cell Immunol.*, 160:178–184.
- Pettit, A.R.; Ji, H.; von Stechow, D.; Müller, R.; Goldring ,S.R.; Choi, Y.; Benoist, C. and Gravallesse, E.M. (2001):** TRANCE/RANKL knockout mice are protected from bone erosion in a serum transfer model of arthritis. *Am J Pathol.*, 159:1689-1899.
- Pinheiro, G.R.; Andrade, C.A.; Gayer, C.R.; Coelho, M.S.; Freire, S.M. and Scheinberg, M.A. (2001):** Serum vascular endothelial growth factor in late rheumatoid arthritis. *Clin Exp Rheumatol.*, 19:721–723.
- Qu, Z.; Garcia, C.H.; O'Rourke, L.M.; O'Rourke, L.M.; Planck, S.R.; Kohli, M.; Rosenbaum, J.T. (1994):** Local proliferation of fibroblast-like synoviocytes contributes to synovial hyperplasia. Results of proliferating cell nuclear antigen/cyclin, c-myc, and nucleolar organizer region staining. *Arthritis Rheum.*, 37:212-220.
- Qvistgaard, E.; Kristoffersen, H.; Terslev, L.; Danneskiold-Samsøe, B.; Torp-Pedersen, S. and Bliddal, H.¹. (2001):** Guidance by ultrasound of intraarticular injections in the knee and hip joints. *Osteoarthritis Cartilage*, 9:512–517.
- Qvistgaard, E.; Rogind, H.; Torp-Pedersen, S.; Terslev, L.; Danneskiold-Samsøe, B. and Bliddal, H.²(2001):** Quantitative ultrasonography in rheumatoid arthritis: evaluation of inflammation by Doppler technique. *Ann Rheum Dis.*, 60(7):690–693.
- Raza, K.; Lee, C.Y.; Pilling, D.; Heaton, S.; Situnayake, R.D.; Carruthers, D.M.; Buckley, C.D.; Gordon, C. and Salmon, M.(2003):**Ultrasound guidance allows accurate needle placement and aspiration from small joints in patients with early inflammatory arthritis. *Rheumatology*, 42:976–979.

- Reimers, C.D. and Finkenstaedt, M.(1997):** Muscle imaging in inflammatory myopathies. *Curr Opin Rheumatol.*, 9:475–485.
- Ritchie, D. M.; Boyle, J. A.; Mcinnes, J. M.; Jasani, K. M.; Dalakas, J. G. and Buchanan, W. (1968):** Clinical studies with an articular index for the assessment of joint tenderness in patients with rheumatoid arthritis. *Quartery J. Med.*, 37: 393-406.
- Robak, E.; Woźniacka, A.; Sysa-Jedrzejowska, A.; Stepień, H. and Robak, T. (2002):** Circulating angiogenesis inhibitor endostatin and positive endothelial growth regulators in patients with systemic lupus erythematosus. *Lupus*, 11: 348–355.
- Rooney, M.; Whelan, A.; Feighery, C. and Bresnihan, B. (1989):** The immunohistologic features of synovitis, disease activity and in vitro IgM rheumatoid factor synthesis by blood mononuclear cells in rheumatoid arthritis. *J Rheumatol.*, 16:459-467.
- Rubaltelli, L.; Fiocco, U.; Cozzi, L.; Baldovin, M.; Rigon, C.; Bortoletto, P.; Tregnaghi, A.; Melanotte, P.L.; di Maggio, C. and Todesco, S.(1994):** Prospective sonographic and arthroscopic evaluation of proliferative knee joint synovitis. *J Ultrasound Med.*, 13(11):855-862.
- Rubin, J.M.; Adler, R.S. and Fowlkes, J.B. (1995):** Fractional moving blood volume: estimation with power Doppler US. *Radiology* 1995;197: 183–90.
- Ruderman, E.M.; Weinblatt, M.E. and Thurmond, L.M. (1995):** Synovial tissue response to treatment with Campath-1H. *Arthritis Rheum.*, 38:254-259.
- Sarkissian, M. and Lafyatis, R. (1999):** Integrin engagement regulates proliferation and collagenase expression of rheumatoid synovial fibroblasts. *J Immunol.*, 162:1772-1779.

-
- Scheel, A.K.; Hermann, K.G.; Kahler, E.; Pasewaldt, D.; Fritz, J.; Hamm, B.; Brunner, E.; Müller, G.A.; Burmester, G.R. and Backhaus, M.(2005):** A novel ultrasonographic synovitis scoring system suitable for analyzing finger joint inflammation in rheumatoid arthritis. *Arthritis Rheum.*, 52(3):733–743.
- Schmidt, W.A. and Gromnica-Ihle, E. (2002):** Incidence of temporal arteritis in patients with polymyalgia rheumatica: a prospective study using colour Doppler ultrasonography of the temporal arteries. *Rheumatology*, 41:46–52.
- Schmidt, W.A.; Nerenheim, A.; Seipelt, E.; Poehls, C. and Gromnica-Ihle, E. (2002):** Diagnosis of early Takayasu arteritis with sonography. *Rheumatology*, 41:496–502.
- Schmidt, W.A.; Kraft, H.E.; Vorpahl, K.; Volker, L. and Gromnica-Ihle, E.J. (1997):** Color duplex ultrasonography in the diagnosis of temporal arteritis. *N Engl J Med.*, 337:1336–1342.
- Schumacher, H.R.; Bautista, B.B.; Krauser, R.E.; Mathur, A.K. and Gall, E.P.(1994):** Histological appearance of the synovium in early rheumatoid arthritis. *Semin Arthritis Rheum* 1994; 23:3-10.
- Scott, D. L.; Grindulis, K. A.; Coulton, B. L.; Popert, A. F. and Bacon, P. A. (1984):** Progression of radiological changes in rheumatoid arthritis. *Ann Rheum Dis.*, 43: 8-17.
- Scott, B.B.; Zaratini, P.F.; Colombo, A.; Hansbury, M.J.; Winkler, J.D. and Jackson, J.R. (2002):** Constitutive expression of angiopoietin-1 and -2 and modulation of their expression by inflammatory cytokines in rheumatoid arthritis synovial fibroblasts. *J Rheumatol.*, 29:230-239.
- Sone, H.; Sakauchi, M.; Suzuki, H.; Inoue, N.; Iida, K.; Shimano, H.; Toyoshima, H.; Kawakami, Y.; Okuda, Y.; Matsuo, K. and Yamada, N. (2001):** Elevated levels of vascular endothelial growth factor in the sera of patients with rheumatoid arthritis correlation with disease activity. *Life Sci.*, 69:1861–1869.
- Suri, C. Jones, P.F.; Patan, S.; Bartunkova, S.; Maisonpierre, P.C.; Davis, S.; Sato, T.N. and Yancopoulos, G.D. (1996):**

Requisite role of angiopoietin-1, a ligand for the TIE2 receptor, during embryonic angiogenesis. *Cell*, **87**: 1171–1180.

Schmidt, W.A. (2001): Value of sonography in diagnosis of rheumatoid arthritis. *Lancet*, 357:1056–1057.

Schoettler, N. and Brahn, E. (2009): Angiogenesis inhibitors for the treatment of chronic autoimmune inflammatory arthritis. *Current Opinion in Investigational Drugs*, 10(5):425-433.

Sone, H.; Sakauchi, M. and Takahashi, A. (2001): Elevated levels of vascular endothelial growth factor in the sera of patients with rheumatoid arthritis correlation with disease activity. *Life Sci.*, 69:1861–1869.

Stone, M.; Bergin, D.; Whelan, B, et al. Maher, M.; Murray, J. and McCarthy, C. (2001): Power Doppler ultrasound assessment of rheumatoid hand synovitis. *J Rheumatol.*, 28(9):1979–1982.

Strunk, J.; Heinemann, H.; Neeck, G. Schmidt, K. L. and Lange U.(2004): A new approach to studying angiogenesis in rheumatoid arthritis by means of power Doppler ultrasonography and measurement of serum vascular endothelial growth factor. *Rheumatology*, 43:1480–1483.

Szekanecz,Z.; Besenyeia,T.; Szentpe' terya, A. and Koch,A. (2010): Angiogenesis and vasculogenesis in rheumatoid arthritis. *Curr Opin Rheumatol.*, 22:299–306.

Szekanecz, Z.; Gaspar, L. and Koch, A.E. (2005): Angiogenesis in rheumatoid arthritis. *Front Biosci.*, 10:1739–1753.

Szekanecz, Z. and Koch, A.E. (2008): Vascular involvement in rheumatic diseases: ‘vascular rheumatology’. *Arthritis Res Ther.*, 10:224-233.

Szekanecz, Z. and Koch, A.E. (2004): Vascular endothelium and immune responses: Implications for inflammation and angiogenesis. *Rheum Dis Clin N Am.*, 30:97–114.

-
- Szekanecz, Z.; Besenyei, T.; Paragh, G. and Koch, A.E. (2009):** Angiogenesis in rheumatoid arthritis. *Autoimmunity*, 42(7): 563–573.
- Szekanecz, Z. and Koch, A.E. (2001):** Chemokines and angiogenesis. *Curr Opin Rheumatol.*, 13: 202–208.
- Szekanecz, Z. and Koch, A.E. (2007):** Mechanisms of Disease: angiogenesis in inflammatory diseases *Nature clinical practice Rheumatology*, 3 (11):635-643.
- Szkudlarek, M.; Narvestad, E.; Klarlund, M.; Court-Payen, M.; Thomsen, H.S. and Østergaard, M. (2004):** Ultrasonography of the metatarsophalangeal joints in rheumatoid arthritis: comparison with magnetic resonance imaging, conventional radiography, and clinical examination. *Arthritis Rheum.*, 50(7):2103–2112.
- Szkudlarek, M.; Court-Payen, M.; Strandberg, C.; Klarlund, M.; Klausen, T. and Ostergaard, M. (2001):** Power Doppler ultrasonography for assessment of synovitis in the metacarpophalangeal joints of patients with rheumatoid arthritis: a comparison with dynamic magnetic resonance imaging . *Arthritis Rheum.*, 44(9):2018-2023.
- Szkudlarek, M.; Court-Payen, M.; Jacobsen, S.; Klarlund, M.; Thomsen, H.S. and Ostergaard, M. (2003):** Interobserver agreement in ultrasonography of the finger and toe joints in rheumatoid arthritis. *Arthritis Rheum.*, 48:955–962.
- Szkudlarek, M.; Court-Payen, M.; Strandberg, C.; Klarlund, M.; Klausen, T. and Ostergaard, M. (2001):** Power Doppler ultrasonography for assessment of synovitis in the metacarpophalangeal joints of patients with rheumatoid arthritis: a comparison with dynamic magnetic resonance imaging. *Arthritis Rheum.*, 44:2018–2023.
- Tak, P.P.; Thurkow, E.W.; Daha, M.R.; et al. (1995):** Expression of adhesion molecules in early rheumatoid synovial tissue. *Clin Immunol Immunopathol* 1995; 77:236.
- Tak, P.P.; Taylor, P.C.; Breedveld, F.C.; Smeets, T.J.; Daha, M.R.; Kluin, P.M.; Meinders, A.E. and Maini, R.N. (1996):** Decrease in cellularity and expression of adhesion

molecules by anti-tumor necrosis factor alpha monoclonal antibody treatment in patients with rheumatoid arthritis. *Arthritis Rheum.*, 39:1077-1081.

Tak, P.P.; Smeets, T.J.; Daha, M.R.; Kluin, P.M.; Meijers, K.A.; Brand, R.; Meinders, A.E. and Breedveld, F.C. (1997): Analysis of the synovial cell infiltrate in early rheumatoid synovial tissue in relation to local disease activity. *Arthritis Rheum.*, 40:217-225.

Tak, P.P.; Gerlag, D.M.; Aupperle, K.B.; van de Geest, D.A.; Overbeek, M.; Bennett, B.L.; Boyle, D.L.; Manning, A.M. and Firestein, G.S. (2001): Inhibitor of nuclear factor kappaB kinase beta is a key regulator of synovial inflammation. *Arthritis Rheum.*, 44:1897-1907.

Taylor, P.C.; Steuer, A.; Gruber, J.; Cosgrove, D.O.; Blomley, M.J.; Marsters, P.A.; Wagner, C.L.; McClinton, C.; Maini, R.N. (2004): Comparison of ultrasonographic assessment of synovitis and joint vascularity with radiographic evaluation in a randomized, placebo-controlled study of infliximab therapy in early rheumatoid arthritis. *Arthritis Rheum.*, 50:1107-1116.

Taylor, P. C. (2002): VEGF and imaging of vessels in rheumatoid arthritis. *Arthritis Res.* 4 Suppl 3:S99-107.

Taylor, P. C. (2005): Serum vascular markers and vascular imaging in assessment of rheumatoid arthritis disease activity and response to therapy. *Rheumatology*, 44:721–728.

Terai, M.; Yasukawa, K.; Narumoto, S.; Tateno, S.; Oana, S. and Kohno, Y. (1999): Vascular endothelial growth factor in acute Kawasaki disease. *Am J Cardiol.*, 83: 337–339.

Terslev, L.; von der Recke, P.; Torp-Pedersen, S.; Koenig, M.J. and Bliddal, H. (2008): Diagnostic sensitivity and specificity of Doppler ultrasound in rheumatoid arthritis. *J Rheumatol.*, 35(1):49–53.

- Thomas, R. and Lipsky, P.E. (1996):** Presentation of self peptides by dendritic cells: possible implications for the pathogenesis of rheumatoid arthritis. *Arthritis Rheum.*, 39:183-186.
- Thompson, P.W.; Silman, A.J.; Kirwan, J.R. and Currey, H.L. (1987):** Articular indices of joint inflammation in rheumatoid arthritis: correlation with the acute-phase response. *Arthritis Rheum.*, 30:618-623.
- Toh, M.L. and Miossec, P. (2007):** The role of T cells in rheumatoid arthritis: new subsets and new targets. *Curr Opin Rheumatol.*, 19:284-287.
- Van Dinther-Janssen, A.C.; Pals, S.T.; Scheper, R.J.; Scheper, R.J. and Meijer, C.J. (1993):** Role of the CS1 adhesion motif of fibronectin in T cell adhesion to synovial membrane and peripheral lymph node endothelium. *Ann Rheum Dis.*, 52:672-576.
- Vazquez, F.; Hastings, G.; Ortega, M.A.; Lane, T.F.; Oikemus, S. and Lombardo, M. (1999):** METH-1, a human ortholog of ADAMTS-1, and METH-2 are members of a new family of proteins with angio-inhibitory activity. *J Biol Chem.*, 274(33):23349-23357.
- Veale, D.J. and Fearon, U. (2006):** Inhibition of angiogenic pathways in rheumatoid arthritis: potential for therapeutic targeting. *Best Pract Res Clin Rheumatol.*, 20: 941-947.
- Verburg, R.J.; Flierman, R.; Sont, J.K.; Ponchel, F.; van Dreunen, L.; Levarht, E.W.; Welling, M.M.; Toes, R.E.; Isaacs, J.D. and van Laar, J.M. (2005):** Outcome of intensive immunosuppression and autologous stem cell transplantation in patients with severe rheumatoid arthritis is associated with the composition of synovial T cell infiltration. *Ann Rheum Dis.*, 64:1397-1405.
- Volin, M.V.; Harlow, L.A.; Woods, J.M.; Campbell, P.L.; Amin, M.A. and Tokuhira, M. (1999):** Treatment with sulfasalazine or sulfapyridine, but not 5-aminosalicylic acid, inhibits basic fibroblast growth factor-induced endothelial cell chemotaxis. *Arthritis Rheum.*, 42:1927-1935.

- Wakefield, R.J.; Balint, P.V.; Szkudlarek, M.; Filippucci, E.; Backhaus, M.; D'Agostino, M.A.; Sanchez, E.N.; Iagnocco, A.; Schmidt, W.A.; Bruyn, G.A.; Kane, D.; O'Connor, P.J.; Manger, B.; Joshua, F.; Koski, J.; Grassi, W.; Lassere, M.N.; Swen, N.; Kainberger, F.; Klauser, A.; Ostergaard, M.; Brown, A.K.; Machold, K.P. and Conaghan, P.G. (2005):** Musculoskeletal ultrasound including definitions for ultrasonographic pathology. *J Rheumatol.*, 32(12):2485–2487.
- Wakefield, R.J.; Gibbon, W.W.; Conaghan, P.G. ; O'Connor, P.; McGonagle, D.; Pease, C.; Green, M.J.; Veale, D.J.; Isaacs, J.D. and Emery, P. (2000):** The value of sonography in the detection of bone erosions in patients with rheumatoid arthritis: a comparison with conventional radiography. *Arthritis Rheum.*, 43:2762–2770.
- Waller, H.A.; Butler, M.G.; McClean, J.G.; Dowd, G.S. and Scott, D.L. (1992):** Localisation of fibronectin mRNA in the rheumatoid synovium by in situ hybridisation. *Ann Rheum Dis.*, 51:735-740.
- Walsh, D.A. (1999):** Angiogenesis and arthritis. *Rheumatology*, 38: 103–112.
- Walther, M.; Harms, H.; Krenn, V.; Radke, S.; Faehndrich, T.P. and Gohlke, F. (2001):** Correlation of power Doppler sonography with vascularity of the synovial tissue of the knee joint in patients with osteoarthritis and rheumatoid arthritis. *Arthritis Rheum.*, 44(2):331–338.
- Walther M, Harms H, Krenn V, Radke S, Kirschner S, Gohlke F. (2002):** Synovial tissue of the hip at power Doppler US: correlation between vascularity and power Doppler US signal. *Radiology*, 225:225–231.
- Warrington, K.J.; Takemura, S.; Goronzy, J.J. and Weyand, C.M. (2001):** CD4, CD28 T cells in rheumatoid arthritis patients combine features of the innate and adaptive immune systems. *Arthritis Rheum.*, 44:13-20.

- Wauke, K.; Nagashima, M.; Ishiwata, T.; Asano, G. and Yoshino, S.(2002):** Expression and localization of vascular endothelial growth factor-C in rheumatoid arthritis synovial tissue. *J Rheumatol.*, 29(1):34-38.
- Weidekamm, C.; Koller, M.; Weber, M. and Kainberger, F. (2003):** Diagnostic value of high-resolution B-mode and Doppler sonography for imaging of hand and finger joints in rheumatoid arthritis. *Arthritis Rheum.*, 48(2):325–333.
- Westergren, A. (1926):** The Technique of the red cell sedimentation reaction. *American Review of the Tuberculosis*, 14:94- 101.
- Weyand, C.M. and Goronzy, J.J. (1999):** T-cell responses in rheumatoid arthritis: systemic abnormalities-local disease. *Curr Opin Rheumatol.*, 11:210- 217.
- Wilkinson, L.S.; Pitsillides, A.A.; Worrall, J.G. and Edwards, J.C. (1992):** Light microscopic characterization of the fibroblast-like synovial intimal cell (synoviocyte). *Arthritis Rheum.*, 35:1179-1184.
- Wolfe, G.C.; MacNaul, K.L.; Buechel, F.F.; McDonnell, J.; Hoerrner, L.A.; Lark, M.W.; Moore, V.L. and Hutchinson, N.I. (1993):** Differential in vivo expression of collagenase messenger RNA in synovium and cartilage: Quantitative comparison with stromelysin messenger RNA levels in human rheumatoid arthritis and osteoarthritis patients and in two animal models of acute inflammatory arthritis. *Arthritis Rheum.*, 36:1540-1547.
- Wong, S.M.; Griffith, J.F.; Hui, A.C.; Tang, A. and Wong, K.S.(2002):** Discriminatory sonographic criteria for the diagnosis of carpal tunnel syndrome. *Arthritis Rheum.*, 46:1914–21.
- Woods, J.M.; Katschke, K.J.; Volin, M.V.; Ruth, J.H.; Woodruff, D.C.; Amin, M.A., Connors, M.A.; Kurata, H.; Arai, K.; Haines, G.K.; Kumar, P. and Koch, A.E. (2001):** IL-4 adenoviral gene therapy reduces inflammation, proinflammatory cytokines, vascularization, and bony destruction in rat adjuvant-induced arthritis. *J Immunol.*, 166:1214–1222.

-
- Wu, H.; Beuerlein, G.; Nie, Y.; Smith, H.; Lee, B.A.; Hensler, M.; Huse, W.D. and Watkins, J.D. (1998):** Stepwise in vitro affinity maturation of Vitaxin, an α v β 3-specific humanized mAb. *Proc Natl Acad Sci USA.*, **95**: 6037–6042.
- Yamashita, A.; Yonemitsu, Y.; Okano, S.; Nakagawa, K.; Nakashima, Y.; Irisa, T.; Iwamoto, Y.; Nagai, Y.; Hasegawa, M. and Sueishi, K. (2002):** Fibroblast growth factor-2 determines severity of joint disease in adjuvant-induced arthritis in rats. *J Immunol.*, 168(1):450-457.
- Yanni, G.; Nabil, M.; Farahat, M.R.; Poston, R.N. and Panayi, G.S. (1994):** Intramuscular gold decreases cytokine expression and macrophage numbers in the rheumatoid synovial membrane. *Ann Rheum Dis.*, 53: 315-322.
- Yeo, E.J.; Chun, Y.S.; Cho, Y.S.; Kim, J.; Lee, J.C.; Kim, M.S. and Park, J.W. (2003):** YC-1: a potential anticancer drug targeting hypoxia-inducible factor 1. *J Natl Cancer Inst.*, **95**: 516–525.
- Yin, G.; Liu, W.; An, P.; Li, P.; Ding, I.; Planelles, V.; Schwarz, E.M. and Min, W. (2002):** Endostatin gene transfer inhibits joint angiogenesis and pannus formation in inflammatory arthritis. *Mol Ther.*, 5:547–554.
- Yoo, S.; Bae, D. and Ryoo, J. (2005):** Arginine-rich antivascular endothelial growth factor (anti-VEGF) hexapeptide inhibits collagen-induced arthritis and VEGF-stimulated productions of TNF- α and IL-6 by human monocytes. *The Journal of Immunology*, 174(9): 5846–5855.
- Yoo, S.; Kwok, S. and Kim, W. (2008):** Proinflammatory role of vascular endothelial growth factor in the pathogenesis of rheumatoid arthritis: prospects for therapeutic intervention. *Mediators Inflamm.*, Epub129873.
- Youssef, P.P.; Haynes, D.R.; Triantafillou, S.; Parker, A.; Gamble, J.R.; Roberts-Thomson, P.J.; Ahern, M.J. and Smith, M.D. (1997):** Effects of pulse methylprednisolone on inflammatory mediators in peripheral blood, synovial fluid, and synovial membrane in rheumatoid arthritis. *Arthritis Rheum.*, 40:1400-1408.
-

-
- Youssef, P.P.; Triantafillou, S.; Parker, A.; Coleman, M.; Roberts-Thomson, P.J.; Ahern, M.J. and Smith MD.(1996):** Effects of pulse methylprednisolone on cell adhesion molecules in the synovial membrane in rheumatoid arthritis. Reduced E-selectin and intercellular adhesion molecule 1 expression. *Arthritis Rheum.*, 39:1970-1979.