

References

- Airaksien PJ , Tuulonen A and Werner EB, (1996):*** Clinical evaluation of the optic disc and retinal nerve fibre layer. In: Ritch R, Shields MB, Krupin T (eds). The Glaucomas. 2nd edition, St Louis : Mosby:617-657.
- AisaKsinen PJ, Mustonen F and Alanko HI,(1981) :*** Optic disc hemorrhages precede retinal nerve fibre layer deflection in ocular hypertension. Acta ophthalmol;59:627 -641.
- Alanko HI , Jaanio E , Airaksinen PJ and Nieminen H, (1980):*** Demonstration of glaucomatous optic disc changes by electronic subtraction. Acta Ophthalmol ;58 : 14-19.
- Anderson, D.R. and Hoyt, W.F., (1969):*** Ultrastructure of intra-orbital portion of human and monkey optic nerve. Arch Ophthalmol; 82: 506.
- Anthony J. Bron, Rsmesh C. Tripathi and Brenda J, (1997):*** Wolff anatomy of the eye and orbit, edition VIII,.Champan and Hall, London, UK. pp: 489-508.
- Asrani S, Challa P, Herndon L, Lee P, Stinnett S, and Allingham RR, (2003):*** Correlation among retinal thickness, optic disc and visual field in glaucoma patients and suspects; a pilot study. J Glaucoma;54:120-128.
- Bagga H, Greenfield DS, Feuer W, et al., (2003):*** Scanning laser polarimetry with variable corneal compensation and optical coherence tomography in normal and glaucomatous eyes. Am J Ophthalmol. Apr;135(4):521-9.

- Bathija R, Zangwill L, Berry CC, Sample PA, Weinreb RN, (1998):** Detection of early glaucomatous structural damage with confocal laser scanning tomography. J Glaucoma;7:121-127.
- Baumal and Caroline R, (1999):** Clinical application of optic coherence tomography. Current opinion in ophthalmology,V.10:PP182-188.
- Baumal CR, (1999):** Clinical applications of optical coherence tomography.Curr Opin Ophthalmol;10:182-8.
- Bechman M ,Thiel MJ, Neubauer AS, (1999):**Central corneal thickness measurements with OCT and Ultrasonic pachymetry.(ARVO Abstract).Invest ophthalmol vis sci, 40: 902.
- Becker D and Shaffer S, (1999):** Clinical evaluation of the optic nerve head atrophy In: Stamper RL, Liberman MF,Drake MV (eds). Diagnosis and therapy of the glaucomas, 7th edition,St Louis: Mosby:191-211.
- Belyea DA , Dan JA and Liberman MF, (1994):** The Glaucoma scope Reproductivity and accuracy of results . Invest Ophthalmol Vis Sci;35:429.
- Benjam FB and Speath G,(2001) :**Unveiling the mysteries of glaucoma. Highlights of Ophthalmology;29:2.
- Bishop KL , Werner EB and Krupin T, (1988) :**Variability and reproducibility of topographic measurements with the Rodenstock optic nerve head analyzer . AmJ Ophthalmol;106: 696.
- Blumenthal EZ, Williams JM, Weinreb RN, et al., (2000):** Reproducibility of nerve fiber layer thickness measurements by use of optical coherence tomography. Ophthalmology; 107:2278.

- Bowd C, Chan K, Zangwill LM, Goldbaum MH, Lee T, Sejnowski TJ, Weinreb RN, (2002):**Comparing neural networks and linear discriminant functions for glaucoma detection using confocal scanning laser ophthalmoscopy of the optic disc. Invest Ophthalmol Vis Sci;43:3444-3454.
- Bowd C, Weinreb RN, Williams JM, et al., (2000):** The retinal nerve fiber layer thickness in ocular hypertensive, normal, and glaucomatous eyes with optical coherence tomography. Arch Ophthalmol 118:22.
- Bowd C, Zangwill LM, Berry CC, et al., (2001):**Detecting early glaucoma by assessment of retinal nerve fiber layer thickness and visual function. Invest Ophthalmol Vis Sci 42:1993.
- Bowd C, Zangwill LM, Medeiros FA, Hao J, Chan K, Lee TW, Sejnowski TJ, Goldbaum MH, Sample PA, Crowston JG, Weinreb RN, (2004):** Confocal scanning laser ophthalmoscopy classifiers and stereophotograph evaluation for prediction of visual field abnormalities in glaucoma-suspect eyes. Invest Ophthalmol Vis Sci;45:2255-2262.
- Bowd C, Zangwill LM, Weinreb RN, (2003):** Association between scanning laser polarimetry measurements using variable corneal polarization compensation and visual field sensitivity in glaucomatous eyes. Arch Ophthalmol. Jul;121(7):961-6.
- Bowd C, Zangwill LM, Weinreb RN, (2003):** Association between scanning laser polarimetry measurements using variable corneal polarization compensation and visual field sensitivity in glaucomatous eyes. Arch Ophthalmol. Jul;121(7):961-6.

- Britton RJ, Drance SM and Schulzer M, (1987)*** :The area of the neuroretinal rim of the optic nerve in normal eyes .AmJ Ophthalmol;103:497-504.
- Bron AJ, Tripathi RC and Tripathi BJ (eds), (1997)***: The visual pathway. In: Wolff anatomy of the eye and orbit. 8th edition, Chapman and Hall : London Uk;15:489-508 .
- Cairns JE, (1986)***: Examination of the optic disc in glaucoma. AmJ Ophthalmol;14:17-35.
- Caprioli J., (1994)***: Clinical evaluation of the optic nerve in glaucoma. Trans Am Ophthalmol Soc;92:589-641.
- Chauhan BC, Blanchard JW, Hamilton DC and LeBlanc RP, (2000)***: Technique for detecting serial topographic changes in the optic disc and peripapillary retina using scanning laser tomography. Invest Ophthalmol Vis Sci;41:775-782.
- Chauhan BC, LeBlanc RP, McCormick TA, Rogers JB, (1994)***: Test-retest variability of topographic measurements with confocal scanning laser tomography in patients with glaucoma and control subjects. Am J Ophthalmol;118:9-15.
- Chauhan BC, McCormick TA, Nicolela MT, LeBlanc RP, (2001)***: Optic disc and visual field changes in a prospective longitudinal study of patients with glaucoma. Comparison of scanning laser tomography with conventional perimetry and optic disc photography. Arch Ophthalmol;119:1492-1499.

- Chi QM, Tomita G, Inazumi K, Hayakawa T, Ido T, Kitazawa Y., (1995):** Evaluation of the effect of aging on the retinal nerve fiber layer thickness using scanning laser polarimetry. J Glaucoma;4:406-413.
- Chi, T., Ritch, R., Stickler, D. et al., (1989):** Racial differences in optic nerve head parameters. Arch Ophthalmol, 107, 83.
- Curcio CA, Allen KA., (1990):** Topography of ganglion cells in human retina. J Comp Neurol;300:5-25.
- Dandona L , Quigley HA and Jampel HD, (1989) :** Reliability of optic nerve head topographic measurements with computerized image analysis. AmJ Ophthalmol;108:414.
- Davies DEN.,Youngquist RC.,Cairs, (1987):**Optical coherence-domain reflectometry: A new optical evaluation technique. Opt Lett.;12: 158- 160.
- Diehl DLC, Quigley HA, Miller NR, Sommer A and Burney EN.,(1990):** Prevalence and significance of optic disc hemorrhage in a longitudinal study of glaucoma. Arch Ophthalmol;108:545-550.
- Drance S, Anderson DR and Schulzer M., (2001):** Collaborative Normal-Tension Glaucoma Study Group. Am J Ophthalmol;131:699-708.
- Drance SM., (1989):** Disc hemorrhages in the glaucomas. Surv Ophthalmol; 33:331-337.
- Drexler W, Sattman H and Hermann B, (2003):**Enhanced visualization of macular pathology with the use of ultrahigh-resolution optical coherence tomography. Arch Ophthalmol; 121:695-706.

- Echelman DA and Shields MB, (1993):** Principles & practice of ophthalmology. Philadelphia:1310-1328.
- Ehlers N, Bramsen T and Sperling S., (1975):** Applanation tonometry and central corneal thickness. Acta Ophthalmol;53:34-43
- Fontana L, Poinsoosawmy D, Bunce CV, et al., (1998):** Pulsatile ocular blood flow investigation in asymmetric normal tension glaucoma and normal subjects. Br J Ophthalmol;82:731-736.
- Garway-Heath DF, Greaney MJ, Caprioli J., (2002):** Correction for the erroneous compensation of anterior segment birefringence with the scanning laser polarimeter for glaucoma diagnosis. Invest Ophthalmol Vis Sci.;43:1465-1474.
- Gilgen H.; Novak RP; Salath RP; Hodel W and Beaud P, (1989):** Submillimeter optical reflectometry. Light wave technology;13:1867- 1869.
- Giovanni Taibbi et al., (2009):** Reproducibility of the Heidelberg Retina Tomograph III Glaucoma Probability Score. Journal of Glaucoma. 18:247-252.
- Grauwiler G, Stumpfig D and Flammer J, (1993):** Do beta-blockers cause vasospasm. Ophthalmologica.; 45: 206.
- Greaney MJ, Hoffman DC and Garway-Heath DF, (2002):** Comparison of optic nerve imaging methods to distinguish normal eyes from those with glaucoma. Invest Ophthalmol Vis Sci ; 43: 140-145.

- Greenfield DS , Zacharia P and Schuman JS, (1993):** Comparison of Nidek 3DX and Donaldson simultaneous stereoscopic disc photography. AmJ Ophthalmol;116:741-747.
- Gross RL, Ji J, Chang P, et al., (2003):** A mouse model of elevated intraocular pressure: retina and optic nerve findings. Trans Am Ophthalmol Soc.;101:163–169.
- Guan K, Hudson C, and Flanagan JG, (2004):** Comparison of Heidelberg Retina Tomograph II and Retinal Thickness Analyzer in the Assessment of Diabetic Macular Edema Invest Ophthalmol Vis Sci;45:610-616.
- Gunderson KG , Heijl A and Bengtsson B, (1995):** Sensitivity & specificity of structural optic disc parameters in chronic glaucoma. Acta Ophthalmol scand; 73 : 1-6.
- Gurses-Ozden R, Ishikawa H, Hoh ST, et al., (1999):** Increasing sampling density improves optical coherence tomography measurement reproducibility. J Glaucoma 8:238.
- Hakki Birinci et al., (2002):** Ocular Blood Flow in Healthy and Primary Open-Angle Glaucomatous Eyes. Ophthalmologica;216:434–437.
- Harris A and Sheets C, (2003):** Blood flow in Glaucoma in Glaucoma, color atlas and synopsis of clinical Ophthalmology, Wills Eye Hospital. Rhee DJ (ed); McGraw-hill, Inc.ch 20 p418-32.
- Harris A, Cantor L and Kagemann L, (1999):** Imaging of blood flow. In: imaging in glaucoma. Schumann JS(ed) Slack, thorofare. N.J, p. 135-154.

- Harris A, Sergott RC and Spaeth K, (1994):**Color Doppler analysis of ocular vessel blood velocity in normal tension glaucoma. AM. J. ophthalmol.; 118: 642-649.
- Harrison R and Kaufman CS, (1980):**Clonidine effects of a topically administrated solution on intraocular pressure and blood pressure in open angle glaucoma. Arch. Ophthalmol.; 95: 1368.
- Hayreh SS., (1985):** Interindividual variation in blood supply of the optic nerve head: its importance in various ischemic disorders of the nerve head and glaucoma, low-tension glaucoma, and allied disorders. Doc Ophthalmol;59:217.
- Hayreh SS., (1989):** Blood supply to the optic nerve head. Current Concepts on Ocular Blood Flow in Glaucoma. The Hague, The Netherlands: Kugler & Ghedini;1-31.
- Hayreh, S.S., (1964):** The orbital vessels of rhesus monkey. Exp Eye Res, 3, 16.
- Hee MR, Puliafito CA, Wong C, et al., (1995):** Optical coherence tomography of macular holes. Ophthalmology 102:748.
- Hee MR; Huang D; Swanson EA and Fujimoto JG, (1992):**Polarization sensitive low-reflectometer for birefringence charecterization and ranging- OCT. Am;9:903-908.
- Hee MR; Puliafito CA; Wong C; Duker JS Richel E; Schuman JS; Swanson Ea and Fujimoto JG, (1995):** Optical Coherence Tomography of macular holes. Ophthalmology, 102:748-756.

Hendrickx KH, van den Enden A, Rasker MT and Hoyng PFJ., (1994): Cumulative incidence of patients with disc hemorrhages in glaucoma and the effect of therapy. *Ophthalmology*;101(7):1165-1172.

Hitchings RA Brown DB and Anderson SA, (1983): An optic disc grid. Its evaluation in reproducibility studies on cup/disc ratio. *Br J Ophth*;67:352-355.

Hoerauf H; Gordes RS; Scholz C; Wribelauer C; Koch P; Engelhardt R; Winkler J and Liqua H, (2000): First experimental and clinical results with transscleral OCT Ophthalmic surgery and Laser;31(3) : 218-222.

Hoerauf H; Wrinkler J; Scholz C; Wirablauer C; Gordes RS; Koch P; Engelhardt R ; Liqua H and Bringruber R, (2002): Transscleral OCT.- An experimental study in ex- viva human eyes. *Laser surgery*;30(3):209-215.

Hoffman DC , Greaney MJ and Garway –health DF, (2002): Comparison of optic nerve imaging methods to distinguish normal eyes from those with glaucoma .*Invest Ophthalmol Vis Sci*;43:40-145.

Hoh ST, Greenfield DS, Mistlberger A, et al., (2000): Optical coherence tomography and scanning laser polarimetry in normal, ocular hypertensive, and glaucomatous eyes. *Am J Ophthalmol*;129:129.

Hoh ST, Greenfield DS, Mistlberger A, Liebmann JM, Ishikawa H and Ritch R., (2000): Optical coherence tomography and scanning laser polarimetry in normal, ocular hypertensive, and glaucomatous eyes. *Am J Ophthalmology*;129(2):129-135.

Holm OL and Krankan CET, (1970): A photographic method for measuring the volume of papillary excavations. *Am J Ophthalmol*;1: 327-332.

- Hoskins HD, Glenday M , Samules SJ and Verdooner SR, (1994):** Repeatability of the glaucoma-scope measurements of optic nerve head topography . J of Glaucoma ; 3 :17-27.
- Huang D; Swanson EA; Lin CP; Schuman JS; Stinson WG; Chang W; Hee MR; Flott T; Gregory K; Puliafito CA and Fujimoto JG, (1991):**Optical Coherence Tomography. Science , 254:1178- 1181.
- Iester M, Mikelberg FS, Courtright P, Drance SM, (1997):** Correlation between the visual field indices and Heidelberg Retina Tomograph parameters. J Glaucoma;6:78-82.
- Iester M, Mikelberg FS, Swindale NV, et al., (2004):** ROC analysis of Heidelberg Retina Tomograph optic disc shape measures in glaucoma. Can J Ophthalmol 32:382, 1997 Invest Ophthalmol Vis Sci;45:2279-2285.
- Ilgaz Sagdic Yalvac, et al., (2009):** The Correlation Between Visual Field Defects and Focal Nerve Fiber Layer Thickness Measured With Optical Coherence Tomography in the Evaluation of Glaucoma. Journal of Glaucoma. 18(1):53-61, January.
- Itay Ben Zion et al., (2009):** Interobserver Repeatability of Heidelberg Retinal Flowmetry Using Pixel-by-Pixel Analysis. Journal of Glaucoma;18(4):280-283.

- Ivan Goldberg, Stuart I Graham and Paul R Healey., (2002):*** early diagnosis of primary open angle glaucoma;v.177:pp535-536.
- Javitt JC, Spaeth GL, Katz LJ, Poryzees E and Addiego R., (1990):*** Acquired pits of the optic nerve: Increased prevalence in patients with low-tension glaucoma. *Ophthalmology*;97(8):1038-1044.
- Jerald A Bell., (2002):*** Glaucoma, primary open angle. *Eye medicine*;vol.29.
- Johnson CA, Keltner JL, Cello KE, et al., (2002):*** Baseline visual field characteristics in the ocular hypertension treatment study. *Ophthalmology* 109:432-437.
- Jonas JB, Budde WM and Panda-Jonas S., (1999):*** Ophthalmoscopic evaluation of the optic nerve head. *Surv Ophthalmol.*;43:293–320.
- Jonas JB, Fernandez MC, Sturmer J., (1993):*** Pattern of glaucomatous neuroretinal rim loss. *Ophthalmology*; Jan;100(1):63-8.
- Jonas JB, Gusek GC, NaumannGOH., (1992):*** Optic disc, cup and neuroretinal rim size, configuration and correlations in normal eyes. *Invest Ophthalmol Vis Sci*;29(7):1151-1158.
- Jonas JB, Zach FM, Gusek GC, Naumann GO., (1989):*** Pseudoglaucomatous physiologic large cups. *American Journal of Ophthalmology*;107(2):137-144.
- Jonas, J.B. and Fernadez, M.C., (1994):*** Shape of the neuroretinal rim and position of the central retinal vessels in glaucoma. *Br J Ophthalmol*;78:99.

- Jonas, J.B. and Naumann, G.O.H., (1991):*** The anatomical structure of the normal and glaucomatous optic nerve, in Glaucoma Update IV (ed. G.K. Krieglstein), Springer- Verlag, Berlin-Heidelberg-New York, p. 66.
- Jonas, J.B. and Schiro D, (1993):*** Normal retinal nerve fiber layer visibility correlated to rim width and vessel caliber. A von Graefes Arch Klin Exp Ophthalmol;231:207 .
- Jonas, J.B. Gusek, G.C. and Naumann and G.O.H., (1988):*** Optic disc, cup and neuroretinal rim size, configuration and correlations in normal eyes. Invest Ophthalmol Vis Sci;29:1151.
- Jonas, J.B., Fernandez, M.C. and Naumann, G.O.H., (1991):*** Correlation of the optic disc size to glaucoma susceptibility. Ophthalmology;98:675.
- Jonas, J.B., Fernandez, M.C. and Naumann, G.O.H., (1992):*** Glaucomatous parapapillary atrophy. Occurrence and correlations. Arch Ophthalmol;110: 214.
- Kalina PH , Hutchinson BT and Netland PA, (1994):*** Quantitative assessment of optic nerve head topography. Invest Ophthalmol; 34:239 -253.
- Kalina PH , Hutchinson BT and Netland PA, (1994):*** Quantitative assessment of optic nerve head topography. Invest Ophthalmol; 34:239 -253.
- Kamal DS, Garway-Heath DF, Hitchings RA, Fitzke FW, (2000):***Use of sequential Heidelberg retina tomograph images to identify changes at the optic disc in ocular hypertensive patients at risk of developing glaucoma. Br J Ophthalmol;84:993-998.

- Kamal DS, Viswanathan AC, Garway-Heath DF, Hitchings RA, Poinoosawmy D, Bunce C., (1999):**Detection of optic disc change with the Heidelberg Retina Tomograph before confirmed visual field change in ocular hypertensives converting to earlyglaucoma. Br J Ophthalmol;83:290-294.
- Kerrigan-Baumrind LA, Quigley HA, Pease ME, Kerrigan DF, Mitchell RS, (2000):** Number of ganglion cells in glaucoma eyes compared with threshold visual field tests in the same persons. Invest Ophthalmol Vis Sci.;41(3):741-748.
- Kim J, Schuman JS, (2000):**Imaging of the optic nerve head and nerve fiber layer in glaucoma. Ophthalmol;13(3):383-406.
- Lemij et al., (2008):** New developments in scanning laser polarimetry for glaucoma. Current Opinion in Ophthalmology;19(2):136-140,.
- Lichter PR, (1972):** Transillumination photography of the eye. AmJ Ophthalmol;75:927.
- Lieberman, M.F., Maumenee, A.E. and Green, W.R., (1976):** Histologic studies of the vasculature of the anterior optic nerve. Am J Ophthalmol;82:405.
- Logan JF, Rankin SJ and Jackson AJ., (2004):** Retinal blood flow measurements and neuroretinal rim damage in glaucoma. Br J Ophthalmol;88:1049-1054,.
- Martinez A, Gonzaleg F and Capeans T, (1999):**Dorzolamide effect on ocular blood flow. Invest. Ophthalmol. Vis. Sci.; 40: 1270-1275.
- May CA and Lütjen-Drecoll E., (2002):** Morphology of the murine optic nerve. Invest Ophthalmol Vis Sci.;43:2206–2212.

- Medeiros FA, Zangwill LM, Bowd C, et al., (2003):** Fourier analysis of scanning laser polarimetry measurements with variable corneal compensation in glaucoma. *Invest Ophthalmol Vis Sci*;44(6):2606-2612.
- Melamed S, Krupsky S and Treister G, (1999):** Confocal Tomographic Angiography of the optic nerve head in glaucoma;P.157.
- Mikelberg FS, Yidegiligne HM, Shulzer M., (1995):** Optic nerve axon count and axon diameter in patients with ocular hypertension and normal visual fields. *Ophthalmology*;102:342–348.[Medline]
- Minckler DS and Mclean IW, (1976):** Distribution of axonal and glial elements in the rhesus optic nerve head studied by electron microscopy. *AmJ Ophthalmol*;82:179.
- Minckler,D.S., (1980):** The organization of nerve of the primate optic nerve head.*arch ophthalmol*;98:1630.
- Mistlberger A, Liebmann JM, Greenfield DS, et al, (1999):** Heidelberg retina tomography and optical coherence tomography in normal, ocular-hypertensive, and glaucomatous eyes. *Ophthalmology*;106:2027.
- Mohammadi K, Bowd C et al., (2004):** Retinal Nerve Fiber Layer Thickness Measurements with Scanning Laser Polarimetry Predict Visual Field Loss. *American Journal of Ophthalmology*;138:592-601
- Nanba K and Schwartz B, (1983):** Fluorescein angiographic defects of the optic disc in glaucomatous visual field loss. In: Greve El. Heijl A. (eds). *Fifth International Visual field symposium*;pp. 67-73.

- Netland PA, (1999):** The Glaucoma scope principle, techniques and applications.
In: Schuman JS (ed) Imaging in Glaucoma 4th edition Slack: Thorofare
NJ;2:17-32.
- O'Connor DJ, Zeyen T and Caprioli J., (1993):** Comparisons of methods to detect
glaucomatous optic nerve damage. Ophthalmology;100(10):1498-1503.
- Olver, J.M. and McCartney, A.C.E., (1989):** Orbital and ocular micro-vascular
casting in man. Eye;3:588.
- Olver, J.M., (1990):** Functional anatomy of the choroidal circulation, methyl
methacrylate casting of human choroids. Eye;4:262.
- Olver, J.M., Spalton, D.J. and McCartney, A.C.E., (1990):** Microvascular study
of the retrolaminar optic nerve in man: the possible significance in anterior
ischaemic optic neuropathy. Eye;4:7.
- Onsy A ,Ahmed T, and Raefat H, (2001):** Assessment of the retinal nerve fibre
layer thickness in normal and glaucomatous eyes using optic coherence
tomography .Bull Ophthalmol.Soc.Egypt;94:347-349.
- Ozdek SC, Onol M, Gurelik G & Hasanreisoglu B, (2000):**Scanning laser
polarimetry in normal subjects and patients with myopia. Br J
Ophthalmology 84: 264–267.
- Paczka JA, Friedman DS, Quigley HA, Barron Y & Vitale S, (2001):**Diagnostic
capabilities of frequency-doubling technology, scanning laser polarimetry,
and nerve fiber layer photographs to distinguish glaucomatous damage. Am
J Ophthalmol;131: 188–197.

- Park KH, Park SJ, Lee YJ, Kim JY and Caprioli, (2001):** Ability of peripapillary atrophy parameters to differentiate normal tension glaucoma from glaucoma like disc. J of Glaucoma;10:95-101.
- Park KH, Tomita G, Liou SY and Kitazawa Y., (1996):** Correlation between peripapillary atrophy and optic nerve damage in normal-tension glaucoma. Ophthalmology;103(11):1899-1906.
- Pederson JE and Gaasterland DE., (1984):** Laser-induced primate glaucoma. I. Progression of cupping. Arch Ophthalmol;102:1689–1692.
- Perry CM, McGavin JK and Culy CR, (2003):** Latanoprost: an update of its use in glaucoma and ocular hypertension. Drug aging.; 20(8): 597-630
- Pieroth L, Schuman JS, Hertzmark E, Hee MR, Fujimoto JG, and Swanson E, (1999):** Evaluation of focal defects of the nerve fibre layer using optical coherence tomography. Ophthalmology;106:570-579.
- Polito, (2007):** Comparison between retinal thickness analyzer and optical coherence tomography for assessment of foveal thickness in eyes with macular disease. American Journal of Ophthalmology;134(2):240-251.
- Puliafita CA, Hee MR, Schuman JS and Fujimoto JC, (1996):** Optical coherence tomography of ocular diseases. Br J Ophthalmol;13:291-344.
- Quaranta L, Manni B, Donato F, et al., (1994):** The effect of increased intraocular pressure on pulsatile ocular blood flow in low tension glaucoma. Sur Ophthalmol;38.
- Quigley HA, Addicks EM, Green WR., (1982):** Optic nerve damage in human glaucoma. III. Quantitative correlation of nerve fiber loss and visual field

defect in glaucoma, ischemic neuropathy, papilledema, and toxic neuropathy. Arch Ophthalmol;100(1):135-46.

Quigley HA, Miller NR, and George T., (1980): Clinical evaluation of nerve fiber layer atrophy as an indicator of glaucomatous optic nerve damage. Arch Ophthalmol.;98:1564-1571.

Quigley, H.A., Coleman, A.L. and Dorman-Pease, M.E., (1991): Larger optic nerve heads have more nerve fibers in normal monkey eyes. Arch Ophthalmol109:1443.

Ravalico G, Toffoli G and Pastori G., (1996): Age related ocular blood flow changes. Invest. Ophthalmol. Vis. Sci; 37: 2645-2650.

Reus NJ, Lemij HG., (2004): Diagnostic Accuracy of the GDX VCC for Glaucoma. Ophthalmology.; 111(10):1860-5.

RobinA L, (1987): Eliminating the acute intraocular pressure rise following argon laser glaucoma surgery. Invest. Ophthalmol. Vis. Sci; 28: 378

Rohrschneider K, Burk ROW, Kruse FE, Völcker HE, (1994): Reproducibility of optic nerve head topography with a new laser tomographic scanning device. Ophthalmology;101:1044-1049.

Rolins AM; Yazdanfar S; Barton JK and Izatt JA, (2002): Real time in vivo color Doppler OCT. J Biomed Opt7 (1): 123.

Rosenthal AR , Falcover DG and Banett P, (1986): Digital measurements of pallor disc ratio . Arch Ophthalmol; 98 : 2027-2031 .

- Salgarello Tommaso et al., (2005):*** Posterior Pole Retinal Thickness in Ocular Hypertension and Glaucoma: Early Changes Detected by Hemispheric Asymmetries. Journal of Glaucoma: October - Volume 14 - Issue 5 - pp 375-383.
- Schuman JS, (1996):*** Imaging of the optic nerve head and nerve fibre layer in glaucoma .In: Epstein DI, Allingham RR, Schuman JS , (eds) Chandler and Grants glaucoma 4th edition Baltimore;15:402-435.
- Schuman JS, (1999):*** Introduction. In: Schuman JS(ed) Imaging in glaucoma 4th edition Slack: Thorofare N J: 15-16.
- Schuman JS, Hee MR, Puliafito CA, et al., (1995):*** Quantification of nerve fiber layer thickness in normal and glaucomatous eyes using optical coherence tomography. Arch Ophthalmol 113:586.
- Schuman JS, Pedut-Kloizman T and Hertzmark E, (1996):*** Reproducibility of nerve fibre layer thickness measurements using optical coherence tomography. Ophthalmology;34:138-144.
- Schuman JS, Pedut-Kloizman T, Hertzmark E, et al., (1996):*** Reproducibility of nerve fiber layer thickness measurements using optical coherence tomography. Ophthalmology;103:1889.
- Schuman JS; Pedut-Kiloizman T; Hee MR; Puliafito CA, Hartzmark E; Swanson EA and Fujimoto JG, (1996):*** Reproducibility of nerve fiber layer thickness measurements using OCT. Ophthalmology;103:11, 1889-98.

- Sherwood MB, (1989):***Long term morphologic effects of antiglaucomatous drugs on the conjunctiva and tenon's capsule in glaucomatous patients. Ophthalmology; 96: 327.
- Shields MB and Tiedman JS (1994):*** Binocular ophthalmoscopic techniques for evaluation of the optic nerve head. Philadelphia,J.B:Lippincott;11: 137 -146.
- Shields MB, (1998):*** The optic nerve head. In: Williams and Wilkins (eds) Textbook of glaucoma 4th edition Baltimore;6: 71-96.
- Shimmyo M, Ross AJ, Moy A, et al., (1981):*** Intraocular pressure , Goldmann applanation tension, corneal site of injury and susceptibility to damage. Arch Ophthalmol;99:635.
- Shimmyo M, Ross AJ, Moy A, et al., (2003):*** Intraocular pressure , Goldmann applanation tension, corneal thickness, and corneal curvature in Caucasians, Asians, Hispanics, and African Americans. Am J Ophthalmol;136:603- 613.
- Speath G, (2001):*** Conducting the optic disc evaluation .Highlights of Ophthalmology;29:1-10.
- Speath G, (2001):*** Evaluation of the optic disc in managing glaucoma. Unveiling the mysteries of glaucoma. Highlights of Ophthalmology; 29 : 4-11.
- Sugiyama K, Kawase C , Yamamoto T and Kitazawa Y, (1999):*** Optic nerve and peripapillary choroidal microvasculature of the rat eye. Invest Ophthalmol Vis Sci;40:3084-3090 .

Swanson EA; Izatt JA; Hee MR; Huang D; Lin CP, Schuman JS; Puliafito CA and Fujimoto JG, (1992): In vivo retinal imaging by OCT. Optt Lett.; 18:1864- 1866.

Swindale NV, Stjepanovic G, Chin A, Mikelberg FS, (2000):Automated analysis of normal and glaucomatous optic nerve head topography images. Invest Ophthalmol Vis Sci;41:1730-1742.

Tan JCH, Hitchings RA, (2003): Approach for identifying glaucomatous optic nerve progression by scanning laser tomography. Invest Ophthalmol Vis Sci;44:2621-2626.

Tan JCH, Poinosawmy D, Hitchings RA, (2006):Tomographic identification of neuroretinal rim loss in high-pressure, normal-pressure, and suspected glaucoma.

Teesalu P and Airaksinin PJ, (1997): The optic disc in glaucoma. Glaucoma world;10:1-2

Teesalu P, Vihanninjoki K, Airaksinen PJ, Tuulonen A, Läärä E, (1997):Correlation of blue-on-yellow visual fields with scanning confocal laser optic disc measurements. Invest Ophthalmol Vis Sci;38:2452-2459.

Tjon-Fo-Sang MJ, Lemij HG, (1997): The sensitivity and specificity of nerve fiber layer measurements in glaucoma as determined with scanning laser polarimetry. Am J Ophthalmol;123:62-69[Medline].

- Varma R, Tielsch JM, Quigley HA, et al., (1994):** Race, age, gender, and refractive error-related differences in the normal optic disc. Arch Ophthalmol 112:1068-1076,.
- Velota CT , Anna B and Stephen A, (2002):** Agreement in assessing optic discs with a digital stereoscopic optic disc camera Discam .Br J Ophthalmol; 86 : 196- 202.
- Velota CT , Anna B and Stephen A, (2002):** Agreement in assessing optic discs with a digital stereoscopic optic disc camera Discam .Br J Ophthalmol 2002 ; 86 : 196- 202.
- Vihanninjoki K, Teesalu P, Burk RO, et al., (2000):** Search for an optimal combination of structural and functional parameters for the diagnosis of glaucoma. Multivariate analysis of confocal scanning laser tomograph, blue-on-yellow visual field and retinal nerve fiber layer data. Graefes Arch Clin Exp Ophthalmol;238:477.
- Vivian Guides, Joel S; Schuman, Elen Hertzmark MA ; Antoni Correnti, Ronald Mancini and David Lederer, (2001):** OCT, measurement of macular and nerve fiber layer thickness in normal and glaucomatous eyes. American academy of ophthalmology.
- Watkins R, Bijan B and Yates M, (2001):** Intraocular pressure and pulsatile ocular blood flow after retrobulbar and peribulbar anesthesia. Br. J. Ophthalmol.; 85: 796-798.
- Weinreb RN, Barch DU and Freeman RW, (1994):** Angiography of the glaucomatous optic nerve head. J Glaucoma: 3(suppl): S55-S60.

- Weinreb RN, Bowd C, Greenfield DS, et al., (2002):** Measurement of the magnitude and axis of corneal polarization with scanning laser polarimetry. Arch Ophthalmol.;120(7):901-906.
- Weinreb RN, Bowd C, Zangwill LM., (2002):** Glaucoma detection using scanning laser polarimetry with variable corneal polarization compensation. Arch Ophthalmol; 120:218-224.
- Weinreb RN, Shakiba S, Zangwill L., (1995):** Scanning laser polarimetry to measure the nerve fiber layer of normal and glaucomatous eyes. Am J Ophthalmol;119:627-636
- Williams ZY, Kloizman TP and Schuman JS, (2004):** Disc Analysis in Ophthalmology. Yanoff M and Duker JS (eds), 2nd edition, Mosby Inc.; Sec 12 Ch. 216: 1458-1463.
- Williamson TH and Harris A, (1994):**Ocular blood flow measurements. BR. J. Ophthalmol; 78: 939-945.
- Witusek W, (1966):** Types of physiological excavation of the optic nerve head. Ophthalmologica, 152, 57.
- Wollstein G, Garway-Heath DF, Fontana L, et al., (2000):** Identifying early glaucomatous changes. Comparison between expert clinical assessment of optic disc photographs and confocal scanning ophthalmoscopy. Ophthalmology;107:2272.
- Wollstein G, Garway-Heath DF, Hitchings RA, (2007):** Identification of early glaucoma cases with the scanning laser ophthalmoscope. Ophthalmology;105:1557-1563.

Yuan L and Neufeld AH ,(2002): Activated microglia in the human glaucomatous optic nerve head. J Neurosci Res;64:523-32.

Yücel YH, Gupta N, Kalichman M, Mizisin AP, Hare W, de Souza Lima M, Zangwill L, Weinreb RN, (1998): Relationship of optic disc topography to optic nerve fiber number in glaucoma. Arch Ophthalmol;116:493-497.

Zangwill LM , Chang CF , Williams JM and Weinreb RN, (1999):New technology for diagnosing and monitoring glaucomatous optic neuropathy Ophthalmol Vis Sci;76(8):526-536.

Zangwill LM, Bowd C and Berry CC, (2001):Discriminating between normal and glaucomatous eyes using the Heidelberg Retina Tomograph, GDX Nerve Fiber Analyzer,and Optical Coherence Tomograph. Arch Ophthalmol; 119: 985-993.

Zangwill LM, Weinreb RN, Berry CC, Smith AR, Dirkes KA, Liebmann JM, Brandt JD, Trick G, Cioffi GA, Coleman AL, Piltz-Seymour JR, Gordon MO, Kass MA, (2004): The confocal scanning laser ophthalmoscopy ancillary study to the ocular hypertension treatment study: study design and baseline factors. Am J Ophthalmol;137:219-227.

Zangwill LM; Williams J; Weinereb RN and Berry CC, (2000): Reproducibility of nerve fiber layer thickness measurements by use of OCT. Ophthalmology; 107:2268-82.

Zeimer R, Asrani S, Zou S, et al., (1998): Quantitative detection of glaucomatous damage at the posterior pole by retinal thickness mapping. A pilot study. *Ophthalmology* ;105(2):224-31.

Zhou Q, Weinreb RN., (2002): Individualized compensation of anterior segment birefringence during scanning laser polarimetry. *Invest Ophthalmol Vis Sci*; Jul;43(7):2221-8.