

REFERENCES

References

1. **Konstantopoulos A, Hossain P, Anderson DF (2007):** Recent advances in ophthalmic anterior segment imaging: A new era for ophthalmic diagnosis? Br J Ophthalmol ;91: 551-557
2. **Dorairaj S, Liebmann JM, Ritch R (2007):** Quantitative evaluation of anterior segment parameters in the era of imaging. Trans Am Ophthalmol Soc ;105: 99-110
3. **Hrynchak P and Simpson T (2000):** Optical Coherence Tomography: An introduction to the technique and its use. Optometry and Vision Science;77: 347-356
4. **Nolan W (2008) :**Anterior segment imaging ultrasound biomicroscopy and anterior segment optical coherence tomography. Curr Opin Ophthalmol.;19:115-121
5. **Pavlin CJ and Foster FS (1995):** Ultrasound biomicroscopic anatomy of the normal eye and adnexa. In: Ultrasound Biomicroscopy of the Eye. 1st ed. New York: Springer Verlag ;47-60
6. **Nolan WP, See JL, Chew PT, et al.,(2007) :** Detection of primary angle closure using anterior segment optical coherence tomography in Asian eyes. Ophthalmology;114: 33-39
7. **Radhakrishnan S, Goldsmith J, Huang D, et al.,(2005) :** Comparison of optical coherence tomography and ultrasound biomicroscopy for detection of narrow anterior chamber angles. Arch Ophthalmol;123: 1053-1059
8. **Ishikawa H ., (2007):** Anterior segment imaging for glaucoma: OCT or UBM? Br J Ophthalmology;91: 1420-1421

REFERENCES

9. **Hayashi K, Hayashi H, Nakao F, Hayashi F.,(2001):** Changes in anterior chamber angle width and depth after intraocular lens implantation in eyes with glaucoma. *Ophthalmology* ;108 : 428-429
- 10.**Izzat JA, Hee MR, Swanson EA, et al.,(1994):** Micron-scale resolution imaging of the anterior eye in vivo with optical coherence tomography. *Arch Ophthalmol*; 112: 1584-1589
- 11.**Huang D, Swanson EA, Lin CP, et al. ,(1991):**Optical coherence tomography. *Science*; 254: 1178–1181
12. **Drexler W, Sattmann H, Hermann B, et al., (2003):** Enhanced visualization of macular pathology with the use of ultrahigh resolution optical coherence tomography. *Arch Ophthalmol*; 121: 695–706
13. **Van Velthoven ME, Faber DJ, Verbraak FD, et al., (2007):** Recent developments in optical coherence tomography for imaging the retina. *Progress in Retinal and Eye Research*; 26: 57–77
14. **Puliafito CA, Hee MR, Schuman JS, Fujimoto JG (1996):** Principles of Operation and Technology. In: *Optical coherence tomography of ocular diseases*. . Slack Inc, Thorofare, NJ, 2nd ed; 3-14
- 15.**Huang D, Wang J, Lin CP, et al., (1991):** Micron-resolution ranging of cornea anterior chamber by optical reflectometry. *Las Surg Med.*; 11:419-425
16. **Jaffe GJ and Caprioli J (2004):** Optical coherence tomography to detect and manage retinal disease and glaucoma. *Am J Ophthalmol* 137: 156–169

REFERENCES

17. **Wollstein G, Paunescu LA, Ko TH, et al., (2005):** Ultrahigh-resolution Optical Coherence Tomography in glaucoma. *Ophthalmology*; 112 : 229-237
18. **Radhakrishnan S, Rollins AM, Roth JE, et al. ,(2001):** Real-time OCT of the anterior segment at 1310 nm. *Arch Ophthalmol.*;119:1179-1185
19. **Schmidt-Erfurth U, Leitgeb RA, Michels S, et al., (2005):** Three dimensional ultrahigh-resolution optical coherence tomography of macular diseases. *Invest. Ophthalmol. Vis. Sci.* 46: 3393–3402
20. **Goldsmith JA, Li Y, Chalita MR, et al., (2005):** Anterior chamber width measurement by high-speed Optical Coherence Tomography. *Ophthalmology*; 112 : 238-244
21. **Memarzadeh F, Li Y, Francis BA, et al.,(2007):** Optical Coherence Tomography of the Anterior Segment in Secondary Glaucoma with Corneal Opacity after Penetrating Keratoplasty. *Br J Ophthalmol.*; 91: 189-192
22. **Kent C (2005):**Any way you slice it :a better AC view. *Review of ophthalmology* ;12: 02
23. **Deepak B (2002) :**Ultrasound Biomicroscopy, *Journal of the Bombay Ophthalmologists Association*; 12 :1-4
24. **Hayden BC, Kelley L, Singh AD (2008):** Ophthalmic Ultrasonography Theoretic and Practical Considerations. *Ultrasound clinics*;3 :179-183
25. **Fledelius HC (1997):** Ultrasound in ophthalmology. *Ultrasound Med Biol*;23 :365–375

REFERENCES

- 26.Pavlin CJ and Foster FS (1998):** Ultrasound Biomicroscopy: High Frequency Ultrasound Imaging of the Eye at Microscopic Resolution. Radiologic Clinics of North America. 36: 1047-1058
- 27.Thijssen MJ, Moll MJ , Timer MR (1985):** Acoustic parameters of ocular tissues. Ultrasound Med Biol. 11:157-161
- 28.Pavlin CJ, and Foster FS (2002):** Ultrasound biomicroscopy. Of the eye. In: Ultrasound of the eye and orbit. Byrne SF, Green RL, (eds). St Louis (MO): Mosby. 2nd ed ; 223–235
- 29.Liebmann JM and Ritch R (1996):** Glaucoma secondary to lens intumescence and dislocation. In: The glaucoma. Ritch R, Shields MB, Krupin T, (eds). St Louis, Mo: Mosby-year book Inc; 1033-1053
- 30.Sherar MD, Starkoski BG, Taylor WB , Foster FS (1989):** A 100 MHz B-scan ultrasound backscatter microscope. Ultrasonic Imaging; 11:95-105
- 31.Pavlin CJ, Simpson ER, Foster FS (2008):**Ultrasound biomicroscopy. Ultrasound clinics ;3 :185-194
- 32.Hoerauf H, Wirbelauer C, Scholz C, et al .,(2000):** Slit-Lamp-Adapted Optical Coherence Tomography of the Anterior Segment. Graefe's Arch Clin Exp Ophthalmol. ; 238: 8-18
- 33.Kanski JJ (2003):** Glaucoma. In: Clinical Ophthalmology, A systematic Approach. Benson K (ed). 5th ed .Butterworth Heinemann, Edinburgh. 193-203
- 34.Radhakrishnan S, Huang D, Smith S D (2005):** Optical Coherence Tomography imaging of the anterior chamber angle. Ophthalmol Clin N Am; 18 : 375-381

REFERENCES

35. **Hoerauf H, Winkler J, Scholz C, et al .,(2002):** Transscleral Optical Coherence Tomography –An Experimental Study in Ex-Vivo Human eyes. *Lasers Surg Med.*; 30: 209-215
36. **Asrani S, Sarunic M, Santiago C, Izatt J (2008):** Detailed Visualization of the Anterior Segment Using Fourier-Domain Optical Coherence Tomography .*Arch Ophthalmol.*;126 :765-771
37. **Friedman DS, and He M (2008):** Anterior Chamber Angle Assessment Techniques. *Surv Ophthalmol* 53:250—273
38. **Wirbelauer C, Karandish A, Häberle H, Pham DT (2005):** Noncontact Goniometry with Optical Coherence Tomography. *Arch Ophthalmol.*; 123: 179-185
39. **Spaeth GL (1971):** The normal development of the human anterior chamber angle: a new system of descriptive grading. *Trans Ophthalmol Soc U K.* ;91:709-739
40. **Pavlin CJ Harasiewicz K , Foster FS (1992):** Ultrasound biomicroscopy of anterior segment structures in normal and glaucomatous eyes. *Am J Ophthalmol* 113: 381-389
41. **Ishikawa H, Liebmann JM, Ritch R (2000):** Quantitative Assessment of the Anterior Segment Using Ultrasound Biomicroscopy. *Curr Opin Ophthalmol.*; 11: 133-139
42. **Baikoff G, Lutun E, Ferraz C, Wei J (2004):** Static and dynamic analysis of the anterior segment with OCT. *J cataract Refract Surg*;30 :1843-1850
43. **Baikoff G, Lutun E, Wei J, Ferraz C (2004):** Anterior Chamber Optical Coherence Tomography Study of Human Natural

REFERENCES

- Accommodation in a 19- year-old Albino. J Cataract Refract Surg.; 30: 696-701
- 44.Lavanya R, Teo L, Friedman DS, et al .,(2007):** Comparison of Anterior Chamber Depth Measurements Using the IOL Master, Scanning Peripheral Anterior Chamber Depth Analyzer, and Anterior Segment Optical Coherence Tomography. Br J Ophthalmol.; 91: 1023-1026
- 45.Baikoff G , Jodai HJ, Bourgeon G (2005):** Measurement of the Internal Diameter and Depth of the Anterior Chamber: IOL Master versus Anterior Chamber Optical Coherence Tomographer. J Cataract Refract Surg.; 31: 1722-1728
- 46.Wirbelauer C , Karandish A, Häberle H, Pham DT (2003):** Optical Coherence Tomography in Malignant Glaucoma Following Filtration Surgery. Br J Ophthalmol.; 87: 952-955
- 47.Medina L, Callen I, Nunez, et al .,(2006):** Exploration of the Anterior Segment by Optical Coherence Tomography -3. Arch Soc Esp Oftalmol; 81: 647-652
- 48.Leung CK, Chan WM, Ko CY, et al., (2005):** Visualization of anterior chamber angle dynamics using Optical Coherence Tomography. Ophthalmology; 112 : 980-984
- 49.Huang D, Li Y, Radhakrishnan S, et al.,(2004):** Optical Coherence Tomography of the anterior segment of the eye. Ophthalmol.Clin. North Am 17 :1-6
- 50.Matonti F , Magnin OP, Coulibaly F, et al., (2006):** Normal and Pathological Imaging of The Anterior Segment Using OCT3. J Francais Ophthalmol.; 29 : 714 – 715

REFERENCES

51. **Liu S, Li H, Dorairaj S, et al.,(2009):** Assessment of Scleral Spur Visibility With Anterior Segment Optical Coherence Tomography. *J Glaucoma* ;1-4
52. **Aslanides I, Libre P, Silverman Rh, et al., (1995):** High Frequency Ultrasound in Pupillary Block Glaucoma. *Br J Ophthalmol.*; 79: 972-976
53. **Chalita MR, Li Y, Smith S , et al., (2005):** High-Speed Optical Coherence Tomography of Laser Iridotomy. *Am J Ophthalmol.*; 140: 1133-1136
54. **Lowe RF (1981):** Plateau Iris. *Australian J Ophthalmol.*; 9: 71-78
55. N
onaka A , Kondo T, Kikuchi M, et al.,(2006): Angle Widening and Alteration of Ciliary Process Configuration after Cataract Surgery for Primary Angle Closure. *Ophthalmology* ;113 :437–341
56. **Strenk SA, Semmlow JL, Strenk LM, et al.,(1999):** Age-related changes in human ciliary muscle and lens: a Magnetic Resonance Imaging Study. *Invest Ophthalmol Vis Sci* ;40:1162-1169
57. **Memarzadeh F, Li Y, Chopra V, et al.,(2007):** Anterior segment optical coherence tomography for imaging the anterior chamber following laser peripheral iridotomy. *Am J Ophthalmol* ;143:877– 879
58. **Memarzadeh F, Tang M, Li Y,et al.,(2007):** Optical Coherence Tomography Assessment of Angle Anatomy Changes After Cataract Surgery. *Am J Ophthalmol* ;144:464–465
59. **Gunning FP and Greve EL (1998):** Lens extraction for uncontrolled angle closure glaucoma: long-term follow-up. *J Cataract Refract Surg*;24 :1347–1356

REFERENCES

- 60. Teekhasaence C and Ritch R (1999):** Combined phacoemulsification and goniosynechialysis for un controlled chronic angle closure glaucoma after acute angle closure glaucoma. *Ophthalmol* ; 106 : 669-675
- 61. PAVLIN CJ (2004):** Ultrasound Biomicroscopy presented in The Rounds of The Department of Ophthalmology and Vision Sciences, Faculty of Medicine, University of Toronto;2:1-6
- 62. Marchini G, Pagliarusco A, Toscano A, et al.,(1998):** Ultrasoundbiomicroscopic and conventional ultrasonographic study of ocular dimensions in primary angle-closure glaucoma. *Ophthalmology* ;105:2091– 2098
- 63. Wilensky JT, Kaufman PL, Frohlichstein D, et al., (1993):** Follow-up of angle-closure glaucoma suspects. *Am J Ophthalmol* 115:338–346.
- 64. Woo EK, Pavlin CJ, Slomovic A, et al., (1999):** Ultrasound biomicroscopic quantitative analysis of light-dark changes associated with pupillary block. *Am J Ophthalmol* ;127 :43-47
- 65. Pavlin CJ, Harasiewicz K, Foster FS (1995):** An ultrasound biomicroscopic dark-room provocative test. *Ophthalmic Surg* ;26 :253-255
- 66. Caronia RM, Liebmann JM, Stegman Z, et al .,(1996):** Increase in iris lens contact after laser iridotomy for pupillary block angle closure. *Am J Ophthalmol* ; 122:53-57
- 67. Garudadri CS, Chelerkar V, Nutheti R (2002):** An ultrasound biomicroscopic study of the anterior segment in Indian eyes with primary angle-closure glaucoma. *J Glaucoma* ; 11: 502–507

REFERENCES

- 68.Friedman DS, Gazzard G, Foster P (2003):** Ultra sonographic biomicroscopy, Scheimpflug photography, and novel provocative tests in contralateral eyes of Chinese patients initially seen with acute angle closure. Arch Ophthalmol ; 121: 633-642
- 69.He M, Friedman DS , Ge J (2007):** Laser peripheral iridotomy in primary angle closure suspects: biometric and gonioscopic outcomes. The Liwan Eye Study. Ophthalmology ; 114 : 494-500
- 70. Kurimoto Y, Park M, Sakaue H, Kondo T (1997):** Changes in the anterior chamber configuration after small-incision cataract surgery with posterior chamber intraocular lens implantation. Am J Ophthalmol ;124: 775– 780
- 71.Nonaka A, Kondo T, Kikuchi M, et al ., (2005) :** Cataract surgery for residual angle closure after peripheral laser iridotomy. Ophthalmology ;112:974 –979
- 72.Tran HV, Liebmman JM, Ritch R (2003):** Iridociliary apposition in plateau iris syndrome persists after cataract extraction. Am J Ophthalmol ;135:40 –43
- 73.Seah SKL, Foster PJ, Chew PTK, et al ., (1997):** Incidence of acute primary angle-closure glaucoma in Singapore. An island-wide survey. Arch Ophthalmol ;115:1436–1440
- 74.Hayashi K, Hayashi H , Nakao F, Hayashi F.,(2000):** Changes in Anterior Chamber Angle Width and Depth after Intraocular Lens Implantation in Eyes with Glaucoma. Ophthalmology ;107 :698–703
- 75.Dada T, Sihota R, Gadia R, et al., (2007):**Comparison of anterior segment optical coherence tomography and ultrasound biomicroscopy

REFERENCES

for assessment of the anterior segment. J Cataract Refract Surg ; 33:837–840

76.Gilgen HH, Novak RP, Salathe RP, et al .,(1989): Submillimeter Optical Reflectometry. IEEE J Lightwave Technol. 7 : 1225-1233

77.Ishikawa H, Inazumi K, Liebmann JM, Ritch R (2000): Inadvertent corneal indentation can cause artifactual widening of the iridocorneal angle on ultrasound biomicroscopy. Ophthalmic Surg Lasers. ;31 :342-345

78.Muller M, Dahmen G, Porksen E, et al., (2006): Anterior chamber angle measurement with Optical Coherence Tomography Intraobserver and Interobserver Variability. J Cataract Refract Surg;32 :1803-1808