

References

A

Allemann N., Chamon W. and Silverman R.H. (1993): High-frequency ultrasound quantitative analysis of corneal scarring following excimer laser keratectomy. Arch Ophthalmol; 111: 968–73.

Alp M.N., Buyuktortop N. and Hosal B.M. et al., (2009): Ultrasound biomicroscopic evaluation of the efficacy of a transillumination technique for ciliary sulcus localization in transscleral fixation of posterior chamber intraocular lenses. J Cataract Refract Surg; 35: 291–296.

Altmeyer P., El Gammal S. and Hoffman K. (1992): Ultrasound in Dermatology. Springer-Verlag Inc. Berlin.

Anton A. and Weinreb R.N. (1997): Recurrent hyphema secondary to anterior chamber lens implant. Surv Ophthalmol; 41: 414–6.

Aptel E., Dumas S. and Denis P. (2009): Ultrasound biomicroscopy and optical coherence tomography imaging of filtering blebs after deep sclerotomy with new collagen implant. Eur. J. Ophthalmol; 19: 223–230.

Arya S.K., Kochhar S. and Kumar S.(2004): Malignant glaucoma as a complication of Nd:YAG laser posterior capsulotomy. Ophthalmic Surg Lasers Imaging; 35: 248–50.

Augsburger J.J., Affel L.L. and Benarosh D.A.(1996): Ultrasound biomicroscopy of cystic lesions of the iris and ciliary body. Trans Am Ophthalmol Soc; 94: 259–71.

Azuara-Blanco A. and Dua H.S.(1999): Malignant glaucoma after diode laser cyclophotocoagulation. Am J Ophthalmol; 127: 467–9.

Azuara-Blanco A., Spaeth G.L. and Araujo S.V. et al., (1997): Ultrasound biomicroscopy in infantile glaucoma. Ophthalmology; 104: 1116-19.

B

Bajaj M.S., Aalok L. and Gupta V. (2007): Ultrasound biomicroscopic appearances of eyelid lesions at 50 MHz. J Clin Ultrasound ; 35: 424-9.

Bao-Qun Y., Ling-Ling W. and Zhang C. (2009): UBM features associated with angle closure in fellow eyes of acute primary angle closure after Laser iridotomy. Ophthalmology; 116: 444-448.

Barash D., Goldenberg-Cohen N. and Tzadok D. et al., (1998): Ultrasound biomicroscopic detection of anterior ocular segment foreign body after trauma. Am J Ophthalmol ; 126: 197-202.

Barkana Y., Dorairaj SK. and Gerber Y. (2007): Agreement between gonioscopy and ultrasound biomicroscopy in detecting iridotrabecular apposition. Arch Ophthalmol; 125:1331–5.

Baum G. and Greenwood I. (1958): The application of ultrasonics locating techniques to ophthalmology. Am J Ophthalmol; 46: 319–29.

Bellows A.R., Chylack L.T. and Hutchinson B.T. (1981): Choroidal detachment. Clinical manifestation , therapy and mechanism of formation. Ophthalmology; 88: 1107–1115.

Bhende M., Biswas J. and Gopal L. (2005): Ultrasound biomicroscopy in the diagnosis and management of intraocular gnathostomiasis. Am J Ophthalmol ; 140: 140–2.

Bloch-Michel E. and Nussenblatt R.B. (1987): International Uveitis Study Group recommendations for the evaluation of intraocular inflammatory disease. Am J Ophthalmol ; 103: 234–5.

Bonfioli A.A., Damico F.M. and Curi A.L. (2005): Intermediate uveitis. Semin Ophthalmol ; 20: 147–54.

Bora N.S. and Kaplan H.J.(2007): Intraocular diseases—anterior uveitis. Chem Immunol Allergy; 92: 213–20.

Breingan P.J., Esaki K. and Ishikawa H. (1999): Iridolenticular contact decreases following laser iridotomy for pigment dispersion syndrome. Arch Ophthalmol; 117: 325–8.

Brian L., Ronald H. and Christopher E.(2009): Bilateral Salzmann-like nodular corneal degeneration after laser in situ keratomileusis imaged with anterior segment optical coherence tomography and high-frequency ultrasound biomicroscopy. *Journal of Cataract & Refractive Surgery* ;35: 785-787.

Brown L.F. (1992): Ferroelectric polymers, Current and future applications. *IEEE Ultrasonics Symposium Proceedings*; 539-545.

C

Cadera W., Viirre . and Karlik S.(1992): Cine magnetic resonance imaging of ocular motility. *J Pediatr Ophthalmol Strabismus*; 29: 120-2.

Carassa R.G., Bettin P. and Fiori M.(1998): Nd:YAG laser iridotomy in pigment dispersion syndrome: an ultrasound biomicroscopic study. *Br J Ophthalmol*; 82: 150–3.

Carassa RG, Bettin P. and Fiori M.(2003):Viscocanalostomy versus trabeculectomy in white adults affected by openangle glaucoma: a 2-year randomized, controlled trial. *Ophthalmology*; 110: 882–887.

Carpel E.F., Sigelman R.J. and Doughman D.J.(1977): Posterior amorphous corneal dystrophy. *Am J Ophthalmol* ; 83: 629–32.

Chan S.M., Hudson M. and Weis E. (2007): Anterior and intermediate uveitis cases referred to a tertiary centre in Alberta. *Can J Ophthalmol*; 42: 860–4.

Chawla B., Agarwal A., Kashyap S. (2007): Diagnosis and management of corneal keloid. *Clin Experiment Ophthalmol* ; 35: 855–7.

Chiou A.G., Mermoud A. and Underdahl J.P. et al., (1998): An ultrasound biomicroscopic study of eyes after deep sclerectomy with collagen implant. *Ophthalmology*; 105: 746–750.

Cobbold R.S.C. (2007): Foundations of biomedical ultrasound. New York: Oxford University Press.

Condon G.P., Masket S. et al., Kranemann C. et al.,(2007) : Small-incision iris fixation of foldable intraocular lenses in the absence of capsule support. *Ophthalmology*; 114:1311–1318.

Corriveau L.A., Nasr Y. and Fanous S.(1986): Choroidal and retinal detachment following argon laser iridotomy. *Can J Ophthalmol* ; 21: 107–108.

Cucevic V., Brown A.S. and Foster F.S.(2005): Thermal assessment of 40-MHz pulsed Doppler ultrasound in human eye. *Ultrasound Med. Biol.*; 31: 565–573.

D

da Costa D.S., Lowder C. and de Moraes H.V.(2006): The relationship between the length of ciliary processes as measured by ultrasound biomicroscopy and the duration, localization and severity of uveitis. *Arq Bras Oftalmol*; 69: 383–8.

Dada T., Sihota R. and Gadia R.et al., (2007): Comparison of anterior segment optical coherence tomography and ultrasound biomicroscopy for assessment of the anterior segment. *J. Cat. Ref. Surg*; 33: 837–840.

Dai S., Kraft S. and Smith D. et al.,(2006): Ultrasound biomicroscopy in strabismus reoperations. *J AAPOS*; 10: 202-5.

Dansingani K. K., Al-Khaier A. and Russell-Eggitt I. M.et al., (2005): Management of Intracorneal Bleb After Trabeculectomy for Congenital Glaucoma. *Cornea*; 24 : 486-488.

Das T. and Namperumalsamy P. (1987): Ultrasound in ocular trauma. *Indian J Ophthalmol* ; 35: 121-5.

Deramo V., Shah G.K. and Bauman C.R. et al., (1999): Ultrasound biomicroscopy as a tool for detecting and localizing occult foreign bodies after ocular trauma .*Ophthalmology* ; 106: 301-305.

Detry-Morel M.L., van Acker E. and Pourjavan S. et al., (2007): Anterior segment imaging using optical coherence tomography and ultrasound biomicroscopy in secondary pigmentary glaucoma associated

with in-the-bag intraocular lens. J. Cataract Refract. Surg : 32; 1866–1869 .

Dietlein T.S, Jacobi P.C. and Krieglstein G.K.(1998): Erbium:YAG laser trabecular ablation (LTA) in the surgical management of glaucoma. Laser Surg Med; 23: 104–110.

Dietlein T.S., Engels B.F. and Jacobi P.C et al., (2000): Ultrasound biomicroscopic patterns after glaucoma surgery in congenital glaucoma. Ophthalmology; 107: 1200-1205.

Dietlein T.S., Engels B.F. and Jacobi P.C. et al., (2003): UBM-guided chamber angle surgery for glaucoma management: an experimental study .Eye; 17: 340–345.

Donaldson M.J., Pulido J.S. and Herman D.C. et al., (2007): Pars planitis: a 20-year study of incidence, clinical features, and outcomes. Am J Ophthalmol; 144: 812–7.

Duck F.A. (2002): Non linear acoustics in diagnostic ultrasound. Ultra Med Biol; 28: 1–18.

E

Ettl A., Kramer J. and Daxer A. et al.,(1997): High-resolution magnetic resonance imaging of the normal extraocular musculature. Eye; 11: 793-7.

F

Fledelius H.C. (1997): Ultrasound in ophthalmology. Ultrasound Med Biol; 23: 365–75.

Forbes M. (1966): Gonioscopy with corneal indentation. A method for distinguishing between appositional closure and synechial closure. Arch Ophthalmol; 76: 488–92.

Foster F.S., Strban M. and Austin G. et al., (1994) : The ultrasound macroscope. Ultrasonic Imaging; 6: 243-261.

Foster F.S., Pavlin C.J. and Harasiewicz K.A. et al., (2000): Advances in ultrasound biomicroscopy. *Ultrasound Med Biol*; 26: 1–27.

Friedman D.S., Gazzard G. and Foster P. et al., (2003): Ultrasonographic biomicroscopy, Scheimpflug photography, and novel provocative tests in contralateral eyes of Chinese patients initially seen with acute angle closure. *Arch Ophthalmol*; 121: 633–42.

Friedman D.S. and He M. (2008): Anterior chamber angle assessment techniques. *Surv. Ophthalmol*; 53: 250–273.

Frieling E. and Dembinsky B. (1995):: Morphometry of the ciliary body using ultrasound biomicroscopy. *Ophthalmology*; 92: 745-749.

Futa R. and Furuyoshi N. (1989): Phacodonesis in capsular glaucoma: a clinical and electron microscopic study. *Jpn J Ophthalmol.*; 33: 311-317.

G

Garcia-Feijoo J., Martin-Carbajo M., Benitez del Castillo J.M. (1996): Ultrasound biomicroscopy in pars planitis. *Am J Ophthalmol*; 121: 214–5.

Garcia J.P. and Rosen R.B. (2008): Anterior segment imaging: optical coherence tomography versus ultrasound biomicroscopy. *Ophthalm. Surg. Laser Imag. ; 39: 476–484.*

Gazzard G., Foster P.J. and Friedman D.S et al., (2004): Light to dark physiological variation in irido–trabecular angle width. *Br. J. Ophthalmol.*; 88:1357–1482.

Gentile R.C., Liebmann J.M. and Tello C. et al.,(1996): Ciliary body enlargement and cyst formation in uveitis. *Br J Ophthalmol*; 80: 895–9.

Gentile R.C., Berinstein D.M. and Liebmann J. et al.,(1998) : Highresolution ultrasound biomicroscopy of the pars plana and peripheral retina. *Ophthalmology*; 105: 478–84.

Gohdo T., Tsumura T. and Hashemian M.N. et al., (2009): Traumatic corneal flap dislocation four years after LASIK. *Iranian Journal of Ophthalmology*; 21: 58-61.

Iijima H. (2000): Ultrasound biomicroscopic study of ciliary body thickness in eyes with narrow angles. *Am J Ophthalmol*; 129: 342—6.

Greiner K.H., Kilmartin D.J. and Forrester J.V. et al., (2002): Grading of pars planitis by ultrasound biomicroscopy—echographic and clinical study. *Eur J Ultrasound*; 15: 139–44.

Grewal S., Laibson P.R. and Cohen E.J. et al., (1999): Acute hydrops in the corneal ectasias: associated factors and outcomes. *Trans Am Ophthalmol Soc*; 97: 187–198.

Guha S., Bhende M. and Baskaran M. et al., (2006): Role of Ultrasound Biomicroscopy (UBM) in the Detection and Localisation of Anterior Segment Foreign Bodies. *Ann Acad Med Singapore*; 35: 536-40.

H

Haddad A.M., Greenfield D.S and Stegman Z. et al., (1997): Peter's anomaly: diagnosis by ultrasound biomicroscopy. *Ophthalmic Surg Lasers*; 28: 311–2.

Harbour J.W., Rubsamen P. and Palmberg P. et al., (1996): Pars plana vitrectomy in the management of phakic and pseudophakic malignant glaucoma. *Arch Ophthalmol*; 114: 1073–8.

Hashemian M.N., Soleimani M. and Moghimi S. et al., (2009): Traumatic corneal flap dislocation four years after LASIK. *Iranian Journal of Ophthalmology*; 21: 58-61.

Hayden B.S., Brandy C. and Kelley L. et al., (2008): Ophthalmic Ultrasonography: Theoretic and Practical Considerations. In: *Ultrasound Clin 3*. Arun D. Singh, Brandy C. Hayden, Charles J. Pavlin editors. Elsevier Inc. 179–183.

He M., Friedman D.S. and Ge J. (2007): Laser peripheral iridotomy in eyes with narrow drainage angles: ultrasound biomicroscopy outcomes. The Liwan Eye Study. *Ophthalmology*; 114: 1513–9.

Heiligenhaus A., Schilling M. and Lung E. et al., (1998): Ultrasound biomicroscopy in scleritisHistorical imag *Ophthalmology*; 105: 527-534.

Helveston E.M. (1979): Reoperations in strabismus. *Ophthalmology*; 86:1379-88.

Heur M. and Jeng B.H. (2008): Anterior Segment Disorders. In: *Ultrasound Clin 3*. Arun D. Singh, Brandy C. Hayden, Charles J. Pavlin editors. Elsevier Inc. 201–206.

Hoops J.P., Ludwig K. and Boergen K.P. et al., (2001): Preoperative evaluation of limbal dermoids using high-resolution biomicroscopy. *Graefes Arch Clin Exp Ophthalmol*; 239: 459–61.

Hu C.H., Xu X.C. and Cannata J.M. (2006): Development of a real-time, high-frequency ultrasound digital beamformer for high-frequency linear array transducers. *IEEE Trans. Ultrason. Ferroelectr. Freq. Control*; 53: 317–323.

Huang D., Swanson E.A. and Lin C.P. (1991): Optical coherence tomography. *Science*; 254: 1178–1181

Huang C., Liu B. and Brezinski M.E. (2008): Ultrasound-enhanced optical coherence tomography: improved penetration and resolution. *J Opt Soc Am*: 938–46.

Hurwitz J.J., Pavlin C.J. and Hassan A. (1998): Proximal canalicular imaging utilizing ultrasound biomicroscopy B: experimental balloon catheter dilation. *Orbit*; 17: 37-9.

Hurwitz J.J., Pavlin C.J. and Rhemtulla E. K. (2004): Identification of retained intracanalicular plugs with ultrasound biomicroscopy. *Can J Ophthalmol.*; 39: 533-7.

I

Irvine W.D., Flynn H.W. and Murray T.G. et al., (1992): Retained lens fragments after phacoemulsification manifesting as marked intraocular inflammation with hypopyon. *Am J Ophthalmol*; 114: 610–614.

Ishikawa H., Esaki K. and Liebmann J.M. (1999): Ultrasound biomicroscopy dark room provocative testing: a quantitative method for estimating anterior chamber angle width. *Jpn J Ophthalmol*; 43: 526–34.

Ishikawa H., Liebmann J.M. and Ritch R. (2000): Quantitative assessment of the anterior segment using ultrasound biomicroscopy. *Curr. Opin. Ophthalmol*; 11: 133–139 .

Ishikawa H. and Schuman J.S. (2004): Anterior segment imaging: ultrasound biomicroscopy. *Ophthalmol Clin North Am*; 17: 7–20.

Izatt J.A., Hee M.R. and Swanson M.S. (1994): Micrometer scale resolution imaging of the anterior eye *in vivo* with optical coherence tomography. *Arch. Ophthalmol.*; 112: 1584–1589 .

J

Jabs D.A., Nussenblatt R.B. and Rosenbaum J.T. (2005): Standardization of uveitis nomenclature for reporting clinical data. Results of the First International Workshop *Am J Ophthalmol*; 140: 509–16.

Jacobi P.C., Dietlein T.S. and Krieglstein G.K. (1999): Microendoscopic trabecular surgery in glaucoma management. *Ophthalmology*; 106: 538–544.

Jeng B. and Meisler D.M. (2006): A combined technique for surgical repair of Descemet's membrane detachments. *Ophthalmic Surg Lasers Imaging*; 37:291–7.

Johnson D.H. and Johnson M. (2001): How does non-penetrating glaucoma surgery work? Aqueous outflow resistance and glaucoma surgery. *J Glaucoma*; 10: 55–67.

K

Kamal A.M., Hanafy M. and Ehsan A. et al., (2009): Ultrasound biomicroscopy comparison of ab interno and ab externo scleral fixation of posterior chamber intraocular lenses. *J Cataract Refract Surg*; 35: 881–884.

Karlen M.E., Sanchez E. and Schnyder C.C. et al., (1999): Deep sclerectomy with collagen implant: medium term results. *Br J Ophthalmol*; 83: 6–11.

- Kashiwagi K., Kashiwagi F. and Toda Y. et al., (2004):** A newly developed peripheral anterior chamber depth analysis system: principle, accuracy, and reproducibility. *Br. J. Ophthalmol.*; 88: 1030–1035.
- Kawana K., Yasuno Y. and Yatagai T. et al., (2007):** High-speed, swept-source optical coherence tomography: a 3D view of anterior chamber angle recession. *Acta Ophthalmol. Scand.* 85, 684–685.
- Kawana K., Kiuchi T. and Yasuno Y. et al., (2009):** Evaluation of trabeculectomy blebs using 3D cornea and anterior segment optical coherence tomography. *Ophthalmology*; 116:848–855.
- Kazakova D., Roters S. and Schnyder C.C. et al., (2002) :** Ultrasound biomicroscopy images: long-term results after deep sclerectomy with collagen implant. *Graefes Arch Clin Exp Ophthalmol* ; 240: 918–923.
- Khairy H.A., Atta H.R. and Green F.D.(2005) :** Ultrasound biomicroscopy in deep sclerotomy. *Eye* ;19: 555–560 .
- Kielty C.M., Davies S.J. and Phillips J.E. et al.,(1995):** Marfan syndrome: fibrillin expression and microfibrillar abnormalities in a patient with predominant ocular defects. *J Med Genet.*; 32: 1-6.
- Kim T., Cohen E.J. and Schnall B.M.(1998):**Ultrasound biomicroscopy and histopathology of sclerocornea. *Cornea* ; 17: 443–5.
- Kim H.H., Cannata J. and Liu R. (In press):** 20-MHz/40-MHz Dual element transducers for high frequency harmonic imaging. *IEEE Trans Ultrason Ferroelectr Freq Control*.
- Klaeger A.J., Tran V.T and Hiroz C.A.(2004):** Use of ultrasound biomicroscopy, indocyanine green angiography and HLA-B51 testing as adjunct methods in the appraisal of Behcet's uveitis. *Int Ophthalmol* ; 25:57–63.
- Kunimatsu S., Tanaka S. and Araie M. (1995):** Ultrasound biomicroscope in breaks of ciliary body associated with atopic dermatitis. *Jpn J Clin Ophthalmol*; 49: 1420-1424.

L

Laroche D., Ishikawa H. and Greenfield D.(1998): Ultrasound biomicroscopic localization and evaluation of intraocular foreign bodies. *Acta Ophthalmol Scand* ;76: 491–5.

Lee B.J., Lowder .CY. and Biscotti C. (2007): Ciliary body metastasis masquerading as scleritis. *Br J Ophthalmol* ; 91: 1582–1649.

Liebmann J.M .and Ritch R.(1996): Glaucoma secondary to lens intumescence and dislocation. In: Ritch R, Shields MB, Krupin T, eds. *The Glaucomas*. 2nd ed. St Louis, Mo: Mosby–Year Book Inc: 1033–1053.

Liebmann J.M., Ishikawa H. and Ritch R. (1999): Ultrasound biomicroscopic features of the glaucomas. In: Sassni J, editor. *Ophthalmic fundamentals*. Thorofare, NJ: SLACK: 73–82.

Liu L., Wang T. and Li Z.(1998): Studies of mechanism of malignant glaucoma using ultrasound biomicroscope. *Zhonghua Yan Ke Za Zhi* ;82:10.

Lizzi F.L. and Feleppa E.J.(1979): Practical physics and electronics of ultrasound. *Int Ophthalmol Clin* ;19:35–63.

Lizzi F.L. and Coleman D.J.(2004): History of ophthalmic ultrasound. *J Ultrasound Med*; 23: 1255–66.

Lockwood G.R., Rayn L.K. and Gotlieb A.I. et al.,(1992): In vitro high imaging in muscular and elastic arteries. *J Am Coll Cardiol* ;20: 153–160.

Lumme P. and Laatikainen L.(1993): Exfoliation syndrome and cataract extraction. *Am J Ophthalmol.*;116:51–55.

Lynch M.G., Brown R.H. and Michels R.G.(1986): Surgical vitrectomy for pseudophakic malignant glaucoma. *Am J Ophthalmol* ;102: 149–53.

Lyshchik A., Fleischer A.C. and Huamani J.(2007): Molecular imaging of vascular endothelial growth factor receptor 2 expression using targeted contrast-enhanced high-frequency sonography. *J Ultra Med* ; 26: 1575–86.

M

Maberley D.A., Pavlin C.J. and McGowan H.D. (1997): Ultrasound biomicroscopic imaging of the anterior aspect of peripheral choroidal melanomas. *Am J Ophthalmol* ;123: 506–14.

Macken P.L., Pavlin C.J. and Tule R. et al., (1995): Ultrasound biomicroscopic features of spherophakia. *Aust N Z J Ophthalmol.*; 23: 217-220.

Madhavan C. D., Basti S. and Naduvilath T.J. et al., (2000): Use of Ultrasound Biomicroscopic Evaluation in Preoperative Planning of Penetrating Keratoplasty Cornea; 19 : 17-21.

Mamou J., Ketterling J.A. and Silverman R.H.(2008): Chirp coded excitation imaging with a high-frequency ultrasound annular array. *IEEE Trans Ultrason Ferroelectr Freq Control* ;55: 508–13.

Mandell M.A., Pavlin C.J. and Weisbrod D.J. et al., (2003) : Anterior chamber depth in plateau iris syndrome and pupillary block as measured by ultrasound biomicroscopy. *Am J Ophthalmol.*; 136 : 900-903.

Mannino G., Perdicchi A. and Medori E. M. et al.,(2007): Ultrabiomicroscopy (UBM) in flap dislocation following LASIK : A case report. *European Journal of Ophthalmology*; 17: 259-261.

Marchini G., Pagliarusco A. and Toscano A. et al., (1998): Ultrasound biomicroscopic and conventional ultrasonographic study of ocular dimensions in primary angle-closure glaucoma. *Ophthalmology*; 105: 2091–2098.

Marchini G., Marraffa M. and Brunelli C. et al., (2001): Ultrasound biomicroscopy and intraocular-pressurelowering mechanisms of deep sclerectomy with reticulated hyaluronic acid implant. *J Cataract Refract Surg* ; 27: 507–517.

McDonnell J.M. (1989): Ocular embryology and anatomy. In: Ryan SJ, ed. *Retina*. Vol 1. St Louis, Mo: CV Mosby Co:5-11.

McWhae J.A., Crichton A.C. and Rinke M. (2003): Ultrasound biomicroscopy for the assessment of zonules after ocular trauma. *Ophthalmology*; 110: 1340–3.

Memmen J.E. and McLean I.W. (1990): The long-term outcome of patients undergoing iridocyclectomy. *Ophthalmology*; 97: 429–432.

Molgat Y.M., Pavlin C.J. and Hurwitz J.J.(1993): Ultrasound biomicroscopy as a diagnostic tool in space-occupying lesions of the superotemporal conjunctival fornix. *Orbit*;12 : 121 – 126.

Mundt G. and Hughes W. (1956): Ultrasonics in ocular diagnosis. *Am J Ophthalmol* ;41: 488–98.

N

Nakagawa T., Maed N. and Okazaki N. et al., (2006): Ultrasound Biomicroscopic Examination of Acute Hydrops in Patients With Keratoconus . *Am J Ophthalmol*; 141: 1134–1136.

Netland P.A. (2001): Ophthalmic Technology Assessment Committee Glaucoma Panel, American Academy of Ophthalmology. Nonpenetrating glaucoma surgery. *Ophthalmology*; 108: 416–421.

Nguyen T.N., Mansour M. and Deschenes J. (2003): Visualization of posterior lens capsule integrity by 20 MHz ultrasound probe in ocular trauma. *Am J Ophthalmol*; 136 : 754–5.

Nightingale K.R., Palmeri M.L. and Nightingale R.W. et al.,(2001): On the feasibility of remote palpation using acoustic radiation force. *J Acoust Soc Am*; 110: 625–34.

Nischal K.K., Naor J. and Jay V. (2002): Clinicopathological correlation of congenital corneal opacification using ultrasound biomicroscopy. *Br J Ophthalmol*; 86 :62–9.

Nishikawa S. and Okisaka S. (1992): The tension of the zonule and aging changes of ciliary bodies. *Nippon Ganka Gakkai Zasshi*; 96: 721-730.

Noble J., Quazi F.A. and Lakosha H.K. (2005): Extrascleral extension in association with cilio-choroidal melanoma: ultrasound biomicroscopy with histopathological correlation. *Can J Ophthalmol* ; 40: 616–8.

Nolan W. (2008): Anterior segment imaging: ultrasound biomicroscopy and anterior segment optical coherence tomography. *Curr. Opin. Ophthalmol.* ;19: 115–121 .

Nouby-Mahmoud G., Silverman R.H. and Coleman D.J. (1993): Using high-frequency ultrasound to characterize intraocular foreign bodies. *Ophthalmic Surg*; 24: 94-9.

O

Oh J., Shin H.H. and Kim J.H. (2007): Direct measurement of the ciliary sulcus diameter by 35-megahertz ultrasound biomicroscopy. *Ophthalmology*; 114: 1685–8.

Oka C., Ideta H. and Nagasaki H. et al.,(1994): Retinal detachment with atopic dermatitis similar to traumatic retinal detachment. *Ophthalmology*; 101: 1050–1054.

Oksala A. and Lehtinen A. (1957): Diagnostic value of ultrasonics in ophthalmology. *Ophthalmologica*; 134 : 387–95.

Oliveira C., Liebmann J.M. and Dodick J.M. (2006): Identification of retained nucleus fragment in the posterior chamber using ultrasound biomicroscopy. *Am J Ophthalmol* ;141: 964–6.

Oscar A.Q., Ishikawa H. M. and Liebmann J.M. et al., (2001): Ultrasound biomicroscopy before and after goniosynechialysis. *AmJ Ophthalmol*; 132: 570-571.

Ostendorf M. and Tost F. (2003): J Imaging the lacrimal canaliculus with 20-MHz ultrasonography: a normal diagnosis . *Fr Ophtalmol.*; 26: 1031-4.

Ozdal M.P., Mansour M. and Deschenes J. (2003): Ultrasound biomicroscopic evaluation of the traumatized eyes. *Eye*; 17 : 467–72.

Ozgen A., Alp M.N. and Ariyurek M. et al., (1999): Quantitative CT of the orbit in graves' disease. Br J Radiol; 72: 757-62.

P

Park M. and Kondo T. (1998): Ultrasound biomicroscopic findings in a case of cyclodialysis. Ophthalmologica; 212: 194-7.

Pavlin C.J., Sherar M.D. and Foster F.S. (1990): Subsurface ultrasound microscopic imaging of the intact eye. Ophthalmology; 97: 244-50.

Pavlin C.J., Harasiewicz K. and Sherar M.D. (1991): Clinical use of ultrasound biomicroscopy. Ophthalmology; 98 : 287-295.

Pavlin C.J., Ritch R. and Foster F.S. (1992a): Ultrasound biomicroscopy in plateau iris syndrome. Am J Ophthalmol; 113: 390-5.

Pavlin C.J., McWhae J.A. McGowan H.D. et al., (1992b): Ultrasound biomicroscopy of anterior segment tumors. Ophthalmology; 99: 1220 – 1228.

Pavlin C.J., Harasiewicz K. and Foster F.S. (1992c): Ultrasound biomicroscopy of anterior segment structures in normal and glaucomatous eyes. Am J Ophthalmol.; 113: 381-389.

Pavlin C.J., Easterbrook M. and Hurwitz J.J. (1993): Ultrasound biomicroscopy in the assessment of anterior sclera disease. Am J Ophthalmol; 116 : 628-35.

Pavlin C.J., Harasiewicz K. and Foster F.S. (1994a): Posterior iris bowing in pigmentary dispersion syndrome caused by accommodation. Am J Ophthalmol; 118 : 114-6.

Pavlin C., Harasiewicz K. and Foster K. (1994b): Ultrasound biomicroscopic analysis of haptic position in late-onset, recurrent hyphema after posterior chamber lens implantation. J Cataract Refract Surg; 20: 182-185.

Pavlin C.J. and Foster F.S. (1995): Ultrasound microscopy of the eye. Springer-Verlag, New York.

Pavlin C.J., Harasiewicz K. and Foster F.S. (1995): An ultrasound biomicroscopic dark-room provocative test. *Ophthalmic Surg*; 26 : 253–5.

Pavlin C.J., Macken P. and Trope G.E. (1996): Accommodation and iridotomy in the pigment dispersion syndrome. *Ophthalmic Surg Lasers*; 27: 113–20.

Pavlin C.J. and Foster F.S. (1998): Ultrasound biomicroscopy: high-frequency ultrasound imaging of the eye at microscopic resolution. *Radiol Clin North Am*; 26: 1047–1058.

Pavlin C.J., Christopher D.A. and Burns P.N. et al., (1998a): High-frequency Doppler ultrasound examination of blood flow in the anterior segment of the eye. *Am J Ophthalmol*; 126: 597-600.

Pavlin C.J., Buys Y.M. and Pathmanathan T. (1998b): Imaging Zonular Abnormalities Using Ultrasound Biomicroscopy Arch *Ophthalmol.*; 116: 854-857.

Pavlin C.J. and Foster F.S. (2004): High-resolution ultrasound. In: Krachmer JH, Mannis MJ, Holland EJ, editors. *Conrea*. St. Louis (MO): Mosby; 299–308.

Pavlin C.H., Simpson E.R. and Foster F.S. (2008): Ultrasound Biomicroscopy, In: *Ultrasound Clinic*,3. Arun D. Singh, Brandy C. Hayden, Charles J. Pavlin editors. Elsevier Inc.; 185–194.

Pierro L., Conforto E., and Resti A.G. (1998): High-frequency ultrasound biomicroscopy versus ultrasound and optical pachymetry for the measurement of corneal thickness. *Ophthalmologica*; 212: 1–3.

Pircher M., Gotzinger E. and Findl O. (2006): Human macula investigated *in vivo* with polarization-sensitive optical coherence tomography. *Invest. Ophthalmol. Vis. Sci.* 47, 5487–5494 .

Potash S.D., Tello C. and Liebmann J. (1994): Ultrasound biomicroscopy in pigment dispersion syndrome. *Ophthalmology*; 101: 332–9.

Potsaid B., Gorczynska I. and Srinivasan V.J. (2008): Ultrahigh speed spectral/Fourier domain OCT ophthalmic imaging at 70,000 to 312,500 axial scans per second. *Opt. Express*;16: 15149–15169.

R

Rabsilber T.M., Khoramnia R. and Auffarth G.U. (2006): Anterior chamber measurements using Pentacam rotating Scheimpflug camera. *J. Cat. Refr. Surg.* 32, 456–459.

Radhakrishnan S., Rollins A.M. and Roth J.E. (2001): Real-time optical coherence tomography of the anterior segment at 1310 nm. *Arch. Ophthalmol.*;119: 1179–1185 .

Rapuan C.J. (2005): Excimer laser phototherapeutic keratectomy in eyes with anterior corneal dystrophies: short-term clinical outcomes with and without an antihyperopia treatment and poor effectiveness of ultrasound biomicroscopic evaluation. *Cornea* ;24: 20–31.

Rathinam S.R. and Namperumalsamy P. (2007): Global variation and pattern changes in epidemiology of uveitis. *Indian J Ophthalmol* ; 55: 173–83.

Rayleigh J.S. (1984): The Theory of Sound; 2nd edition, Macmillan Co., London.

Reed J.E., Thomas J.V. and Lytle R.A. (1990): Malignant glaucoma induced by an intraocular lens. *Ophthalmic Surg*; 21: 177–80.

Reinstein D.Z., Silverman R.H. and Coleman D.J. (1993): High-frequency ultrasound measurement of the thickness of the corneal epithelium. *Refract Corn Surg*; 9: 385–7.

Reinstein D.Z., Silverman R.H. and Rondeau M.J. et al.,(1994): Epithelial and corneal thickness measurements by high-frequency ultrasound digital signal processing. *Ophthalmology*;101:140–46.

Reinstein D.Z., Silverman R.H. and Coleman D.J. (1999): Very high-frequency ultrasound corneal analysis identifies anatomic correlates of optical complications of lamellar refractive surgery. *Ophthalmology*; 105: 474–82.

Reinstein D.Z., Silverman R.H. and Raevsky T. (2000): Arc scanning very high-frequency digital ultrasound for 3D pachymetric mapping of the corneal epithelium and stroma in laser in situ keratomileusis. *J Refract Surg*; 16 (4): 414–30.

Reinstein D.Z., Archer T. and Gobbe M. (2008): Stability of LASIK in corneas with forme fruste or suspect keratoconus, where keratoconus was excluded by epithelial thickness mapping. *American Academy of Ophthalmology Annual Meeting*; Atlanta, GA.

Rieser J.C. and Schwartz B.(1972).: Miotic-induced malignant glaucoma. *Arch Ophthalmol* ; 87: 706–12.

Roche O., Beby F. and Dufier J. et al., (2008): Descemet Membrane Rupture Accompanied by Stromal Clefting in Congenital Glaucoma *Arch Ophthalmol Aug*; 126 : 1163-1164.

Rockwood E.J., Sharma S. and Hayden B.C. et al., (2008): Glaucoma. In: Arun D. Singh, Brandy C. Hayden, Charles J. Pavlin editors. *Ultrasound Clin 3*. Elsevier Inc.; 207–215.

Roscoe M. and Landis T. (2006): How to diagnose the acute red eye with confidence. *JAAPA*; 19 : 4–30, quiz 45–6.

Roters S., Luke C. and Jonescu-Cuypers C.P. et al., (2002): Ultrasound biomicroscopy and its value in predicting the long-term outcome of Viscocanalostomy. *Br J Ophthalmol* ; 86: 997–1001.

Rothman R.F., Sidoti P.A. and Gentile R.C. (2001): Glaucoma drainage tube kink after pars plana insertion. *Am. J. Ophthalmol.* 132, 413–414 .

Rubsamen P.E. and McLeish W.M. (1991): Keratoconus with acute hydrops and perforation: brief case report. *Cornea*; 10: 83– 84.

Russell-Eggitt I.M. (1995): In defence of goniotomy . *Br J Ophthalmol*; 79:709.

Rutnin S.S., Pavlin C.J. and Slomovic A.R. (1997): Preoperative ultrasound biomicroscopy to assess ease of haptic removal before penetrating keratoplasty combined with lens exchange. *J Cataract Refract Surg* ; 23: 239–43.

S

Sakai H., Morine-Shinjo S. and Shinzato M. (2005): Uveal effusion in primary angle-closure glaucoma. *Ophthalmology*; 112 : 413–9.

Sakai H., Sakima N. and Nakamura Y. et al., (2001): A case of ciliochoroidal effusion after laser iridotomy. *Atarashii Ganka*; 18: 237–240.

Sakata L.M., Sakata K. and Susanna R. (2006): Long ciliary processes with no ciliary sulcus and appositional angle closure assessed by ultrasound biomicroscopy. *J Glaucoma*; 15: 371-9.

Sanghvi N.T., Wiersman M. and Reilly C.I. et al., (1990): PC based high resolution, High frequency US system for gastroenterology. *IEEE Ultrasonic Symposium Proceedings*; 1477-1479.

Schein O.D., Kenyon K.R. and Steinert R.F. (1993): A randomized trial of intraocular lens fixation techniques with penetrating keratoplasty. *Ophthalmology*; 100: 1437–1443.

Schweykart N., Reinhard T., Engelhardt S. and Sundmacher R. (1999): Postoperative ultrasound biomicroscopic evaluation of the tangible position of black diaphragm posterior chamber lenses in congenital and traumatic aniridia in comparison with gonioscopy. *Klin Monatsbl Augenheilkd*; 214 :381–390.

Sewelam A., Ismail A.M. and El Serogy H. (2001): Ultrasound biomicroscopy of haptic position after transscleral fixation of posterior chamber intraocular lenses. *J Cataract Refract Surg*; 27: 1418–1422.

Shaffer R.N. (1960): A new classification of the glaucomas. *Trans. Am. Ophthalmol. Soc.* 58, 219–225.

Sherar M.D., Noss M.B. and Foster F.S. (1987): Ultrasound backscatter microscopy images the internal structure of living tumor spheroids multicellular tumor spheroids. *Nature*; 330: 493-495.

Shingleton B., Tetz M. and Korber N. (2008): Circumferential viscodilation and tensioning of Schlemm canal (canaloplasty) with temporal clear corneal phacoemulsification cataract surgery for open-angle glaucoma and visually significant cataract. One-year results. *J. Cataract Refract. Surg.* 34, 433–440.

Shiu-Chen W. and Ken-Kuo L.(2004): Correspondence of the Dark-prone Test to Ultrasound Biomicroscopy in the Diagnosis of Plateau Iris Syndrome: A Pilot Stud . *Journal of Medical Ultrasound*; 12 : 49-52

Shung K.K., Cannata J.M. and Zhou Q.F. (2008): High frequency ultrasonic transducers and arrays. In: *Piezoelectric and Acoustic Materials for Transducer Applications*. Safari A and Akdogan EK, eds. Springer: New York, 431–51.

Shung K.K.(2009): High Frequency Ultrasonic Imaging. *J Med Ultrasound*; 17: 25–30.

Silverman R.H., Kruse D. and Coleman D.J. et al., (1999): High-resolution ultrasonic imaging of blood-flow in the anterior segment of the eye. *Invest Ophthalmol Vis Sci*; 40: 1373–81.

Silverman R.H., Ketterling J.A. and Coleman D.J. (2007): High-frequency ultrasonic imaging of the anterior segment using an annular array transducer. *Ophthalmology* 114, 816–822.

Singh M., Aung T. and Aquino M.C. et al., (2009): Utility of bleb imaging with anterior segment optical coherence tomography in clinical decision-making after trabeculectomy. *J. Glaucoma* 18, 492–495.

Small K.M. and Maslin K.F. (1995): Malignant glaucoma following laser iridotomy. *Aust N Z J Ophthalmol*; 23: 339–341.

Spencer W.H.(1996) : Conjunctiva. In: *Ophthalmic Pathology: An Atlas and Textbook*. 4th ed. Philadelphia, Pa: WB Saunders Co; 38-125.

Steiner A., Steinhorst U.H. and Steiner M. et al.,(1997): Ultrasound biomicroscopy (UBM) for the localization of artificial lens haptics after transscleral suture fixation. *Ophthalmology*; 94: 41–44.

T

Taban M., Lowder C.Y. and Hajj-Ali R. (2008): Anterior scleritis as the presenting sign of metastatic lung cancer. *Br J Ophthalmol*; 92: 147.

Tamburrelli C., Salgarello T. and Vaiano A.S. et al., (2003): Ultrasound of the horizontal rectus muscle insertion sites: Implications in preoperative assessment of strabismus. *Invest Ophthalmol Vis Sci*; 44: 618-22.

Tello C., Liebmann J.M. and Ritch R. (1994a): An improved coupling medium for ultrasound biomicroscopy. *Ophthalmic Surg*; 25: 410–1.

Tello C., Liebmann J. and Potash S.D. et al., (1994b): Measurement of ultrasound biomicroscopy images: intraobserver and interobserver reliability. *Invest Ophthalmol Vis Sci*; 35: 3549–3552.

Topilow H.W., Ackerman A.L. and Zimmerman R.D. (1984): Limitations of computerised tomography in the localisation of intraocular foreign bodies. *Ophthalmology*; 91: 1086-91.

Tost F., Bruder R. and Ostendorf M. (2003): High-frequency ultrasonography applied to disorders of the lacrimal canaliculi. *J Fr Ophtalmol.*; 26: 1035-8.

Tran V.T., LeHoang P. and Herbort C.P. (2001): Value of highfrequency ultrasound biomicroscopy in uveitis. *Eye*; 15: 23–30.

U

Urbak S.F. (1999): Ultrasound biomicroscopy. Accuracy and agreement of measurements. *Acta Ophthalmol Scand*; 77: 293–7.

Urbak S.F., Pedersen J.K. and Thorsen T.T. (1998): Ultrasound biomicroscopy. Intraobserver and interobserver reproducibility of measurements. *Acta Ophthalmol Scand*; 76: 546–9.

Ursea R. and Silverman R.H. (2010): Anterior-segment imaging for assessment of glaucoma. *Ophthalmol.* 5, 59–74.

Uysal Y. , Shields C.L., Mashayekhi A., Shields A.J. and Eagle C.R. (2006): Ultrasound Biomicroscopy of Cystic and Solid Caruncular Oncocytoma. *Arch Ophthalmol.*; 124: 1650-1652.

V

Vasquez L., Pavlin C.J. and McGowan H. (2008): Ring melanoma of the ciliary body: clinical and ultrasound biomicroscopic characteristics. *Can J Ophthalmol*; 43: 229–33.

W

Wagoner M.D., Cox T.A. and Ariyasu R.G. et al., (2003): Intraocular lens implantation in the absence of capsular support: a report by the American Academy of Ophthalmology. *Ophthalmology*; 110: 840–859.

Watts P., Smith D. and Mackeen L. et al., (2002): Evaluation of the ultrasound biomicroscope in strabismus surgery. *J AAPOS.*; 6: 187-90.

Werner L., Izak A.M. and Pandey S.K. (2004) : Correlation between different measurements within the eye relative to phakic intraocular lens implantation. *J Cataract Refract Surg*; 30: 1982–8.

Wilensky J.T., Ritch R. and Kolker A.E. (1996): Should patients with anatomically narrow angles have prophylactic iridectomy? *Surv Ophthalmol*; 41: 31–36.

X

Xu M. and Wang L.V. (2006): Photoacoustic imaging in biomedicine. *Rev Sci Instrum*; 77: 041101-1-21.

Y

Yamamoto T., Sakuma T. and Kitazawa Y. (1995): An ultrasound biomicroscopic study of filtering blebs after mitomycin C trabeculectomy. *Ophthalmology*; 102: 1770–1776.

Yasuno Y., Yamanari M. and Kawana K. et al., (2009): Investigation of post-glaucoma-surgery structures by 3D and polarization sensitive anterior eye segment optical coherence tomography. Opt. Express ;16: 3980–3996.

Yip L.W., Sothornwit N. and Berkowitz J. et al.,(2009): A comparison of intraocular differences in patients with pigment dispersion syndrome. J. Glaucoma; 18: 1–5 .

Yoo C., Oh J.H. and Kim Y.Y. et al., (2007): Peripheral anterior synechiae and ultrasound biomicroscopic parameters in angle-closure glaucoma suspects. Korean J Ophthalmol.; 21 : 106-10.

Yoshida S., Sasoh M. and Arima M. et al., (1997): Ultrasound biomicroscopic view of detachment of the ciliary epithelium in retinal detachment with atopic dermatitis. Ophthalmology; 104: 283–287.

Yu A.Y., Lin Z.D. and Chen X.Q. (2008): Position of myopic irisclaw phakic intraocular lens by Scheimpflug photography and ultrasound biomicroscopy. Eye; 22: 233–9.

Z

Zhang H.F., Maslov K. and Stoica G. et al., (2006): Functional photoacoustic microscopy for high-resolution and noninvasive in vivo imaging. Nat Biotechnol; 24: 848–51.

Zhang Y., Wu Q. and Zhang M. et al., (2008): Evaluating subconjunctival bleb function after trabeculectomy using slit-lamp optical coherence tomography and ultrasound biomicroscopy. Chin. Med. J. 121, 1274–1279 .

Zhou Q., Xu X. and Gottlieb E.J.(2007): PMN-PT single crystal, high-frequency ultrasonic needle transducers for pulsed-wave Doppler application. IEEE Trans Ultrason Ferroelectr Freq Control; 54: 668-75.