

References

1. **Abe T., Stinger R.J. and Marks M.P. (1998):** Coexistence of occult vascular malformations and developmental anomalies of the central nervous system: MR evaluation. *AJNR*; 19:51-57.
2. **Akiba H., Tamakawa M. and Hyodoh H. (2008):** Assessment of Dural Arteriovenous Fistulae of the Cavernous Sinuses on 3D Dynamic MR Angiography. *AJNR*; 29: 1652–57
3. **Allroggen H. and Abbott R. (2000):** Cerebral venous sinus thrombosis. *Postgraduate medicine journal*; 76: 12-15.
4. **Aoki S., Yoshikawa T. and Hori M. (2000):** Two-dimensional thick-slice MR digital subtraction angiography for assessment of cerebrovascular occlusive diseases. *European radiology*; 10: 1858-1864.
5. **Aoki S., Yoshikawa T. and Hori M. (2000):** MR digital subtraction angiography for the assessment of arteriovenous malformations and fistulas. *AJR*; 175: 451-453.

6. **Atkins P.T., Pilgram T.K. and Cross T.D. (1998):** Natural history of stenosis from intracranial atherosclerosis by serial angiography. *Stroke*; 29: 433-438.
7. **Ayanzen R.H., Bird C.R. and Keller P.J. (2000):** Cerebral MR venography: Normal anatomy and potential diagnostic pitfalls. *AJNR*; 21:74-78.
8. **Babiarz L.S., Romero J.M. and Murphy E.K. (2009):** Contrast-Enhanced MR Angiography Is Not More Accurate Than Un-enhanced 2D Time-of-Flight MR Angiography for Determining >70% Internal Carotid Artery Stenosis. *AJNR*; 30: 761 - 768
9. **Baert A.L., Knauth M. and Sartor K. (2007):** Intracranial Vascular Malformations and Aneurysms From Diagnostic Work-Up to Endovascular Therapy, 2nd Edition, Springer.
10. **Balaji P.S., Varma R.G., and Kulkarni RN (2005):** Microsurgical anatomy of the middle cerebral artery. *Neurology India*; 53:186-190.
11. **Barnett H.J., Taylor D.W. and Eliaswiz M. (1998):** Benefit of carotid endarterectomy in patients with symptomatic moderate or severe stenosis: North America

symptomatic carotid endarterectomy trial collaborators.
North England Journal of medicine; 339: 1415-1425.

12. **Benseler S.M., DeVeber G. and Silverman E.D. (2007):** Angiography of Primary Central Nervous System Angiitis of Childhood: Conventional Angiography versus Magnetic Resonance Angiography at Presentation. AJNR; 28: 9 –15

13. **Berlis A., Klisch J. and Spetzger U. (2002):** Carotid Cavernous Fistula: Embolization via a Bilateral Superior Ophthalmic Vein Approach. AJNR; 23: 1736-1738.

14. **Biousse V. (1999):** Cerebral venous Neuroradiology; 5: 326-349.

15. **Biousse V., Tong F. and Newman N.J. (2003):** Cerebral venous thrombosis: Current treatment options in cardiovascular medicine; 5: 181-192.

16. **Bock M., Schoenberg S.O. and Fionier F. (2000):** separation of arteries and veins in 3D MR angiography using correlation analysis. Magnetic resonance in medicine; 43: 481-487.

17. **Borisch I., Horn M. and Butz B. (2003):** Preoperative evaluation of carotid artery stenosis:

Comparison of contrast-enhanced MR angiography and duplex sonography with digital subtraction angiography. AJNR; 24: 1117-1122.

18. **Boulin A. and Pierot L. (2001):** Follow up of intracranial aneurysms treated with detachable coils: Comparison of gadolinium enhanced 3D Time-of-flight MR angiography and digital subtraction angiography. Radiology; 219: 108-113.

19. **Brown R.D. and Wiebers D.O. (1998):** Subarachnoid hemorrhage and unruptured intracranial aneurysms. In: Cerebrovascular disease: Pathophysiology, diagnosis, and management, edited by Ginsberg M.D. and Bogousslavsky J., 3rd edition, Blackwell science; 108: 1508-1518.

20. **Brugieres P., Blustajn J. and LeGuerinel C. (1998):** Magnetic resonance of giant intracranial aneurysms. Neuroradiology; 40: 96-102.

21. **Brunelle F. (1997):** Arteriovenous malformations of the vein of Galen in children. Pediatric radiology; 27: 501-513.

- 22. Caplan L. (2000):** Posterior circulation ischemia: The now, and tomorrow: The Tomas Willis lecture. *Stroke*; 31: 2011-2023.
- 23. Carr J., McCarthy R.M. and Laub G. (2001):** Subsecond contrast enhanced 3 dimensional MR Angiography: A new technique of dynamic imaging of the vasculature. In: *Proceedings in the international society for magnetic resonance imaging in medicine. 9th annual meeting book of abstracts. Berkley, California*; 302.
- 24. Carroll T.J. (2002):** The emergence of time resolved contrast kinetics M.R. imaging for intracranial Angiography. *AJNR*; 23: 346-348.
- 25. Carroll T.J., Korosec F.R. and Peterman G.M. (2001):** Carotid bifurcation: Evaluation of time resolved three dimensional MR Angiography. *Radiology*; 220: 525-532.
- 26. Chaloupka J.C. and Huddle D.C. (1998):** Classification of vascular malformation of the central nervous system. *Neuroimaging clinic of North America*; 8: 295-321.
- 27. Ciceri E.F., KJucznik R.P. and Grossman R.G. (2001):** Aneurysms of the Posterior Cerebral Artery:

Classification and Endovascular Treatment. AJNR; 22: 27-34.

28. Cloft H.J., Joseph G.J. and Dion J.E. (1999): Risk of cerebral angiography in patients with subarachnoid hemorrhage, cerebral aneurysms, and arteriovenous malformations: A Meta analysis. Stroke; 30: 317-320.

29. Coley S.C., Romanowski C.A. and Hodgson T.J. (2002): Dural arteriovenous fistulae: Non-invasive diagnosis with dynamic MR digital subtraction angiography. AJNR; 23: 404-407.

30. Cosottini M., Pingiorte A. and Puglioli M. (2003): Contrast enhanced 3D magnetic resonance angiography of atherosclerotic internal carotid stenosis is as the non-invasive imaging modality in revascularization decision making. Stroke; 34:660-664

31. Cottier J., Couthon A. and Gallas S. (2003): Intracranial aneurysms treated with Guglielmi detachable coils: Is contrast material necessary in the follow up with 3D TOP MR Angiography?. AJNR; 24: 1797-1803.

32. Daikokuya H., Inoue Y., and Yamada R. (2001): Three-dimensional gadolinium enhanced MRA of the

intracranial venous system. Osaka city medical journal; 47:165-175.

33. Detwiler P.A., Porter R.W. and Zambramski J.M. (1998): Radiation induced cavernous malformation. Journal of Neurosurgery; 89: 167-168.

34. Eastwood J.D. (2003): Stroke. In: C.T. and MR imaging of the whole body, edited by Haaga J.R., Lanzieri F.R., and Gilkeson R.C., 4th edition, Mosby; 7(2): 246.

35. Ellison D. and Love S. (1998): Subarachnoid hemorrhage. In: Neuropathology: A reference text of CNS pathology, edited by Ellison D. and Love S., 3rd edition, Mosby; 10: 3-8.

36. Farb R.I., McGregor C. and Kirn J.K. (2001): Intracranial arteriovenous malformations: Real time auto-triggered elliptic centric-ordered 3D gadolinium-enhanced MR Angiography-Initial assessment. Radiology; 220: 244-251.

37. Frosch M. P., (2007): Cerebrovascular diseases. In: basic pathology, edited by Robbins S, 8TH Edition, Elsevier; 23:863 - 869

38. Gauthrit J., Leclerc X. and Caron S. (2006): Intracranial Aneurysms Treated with Guglielmi

Detachable Coils - Imaging Follow- Up With Contrast-Enhanced MR Angiography. Stroke ; 37:1033-1034.

39. **Gauvrit J.Y., Law M. and Xu J. (2007):** Time-Resolved MR Angiography: Optimal Parallel Imaging Method. AJNR ; 28: 835–38

40. **Gibbs, G.F., Hutson J. and Brenstein M. (2001):** "angiography of the intracranial aneurysms: 3.0 T compared with 1.5 T. Proc. Of international society of magnetic resonance in medicine; 9: 300-301.

41. **Gijn J.V. (1999):** Main groups of cerebral and spinal vascular disease. In: Cerebrovascular disease: Pathophysiology, diagnosis, and management, edited by Ginsberg M.D. and Bogousslavsky J., 3rd edition, Blackwell science; 101:1376-1382.

42. **Grandin C.B., Mathurin P. and Duprez T. (1998):** Diagnosis of intracranial aneurysms. Accuracy of MR angiography at 0.5T.AJNR; 19: 245-252.

43. **Grau A.J., Weimar C and Buggie F. (2001):** Risk factors, outcome, and treatment in subtypes of ischemic stroke. Stroke: 32: 2559-2563.

44. **Griffith P., Hoggard N. and Warren D. (2000):** Brain arteriovenous malformations: Assessment with

dynamic MR digital subtraction angiography. *AJNR*; 21: 1892-1899.

45. **Hadizadeh D.R., Falkenhausen M.V. and Gieseke J. (2008):** Cerebral Arteriovenous Malformation: Spetzler-Martin Classification at Subsecond-Temporal-Resolution Four-dimensional MR Angiography Compared with That at DSA. *Radiology*; 246: 205-213.

46. **Halbach V.V., Doud C.F. and Higashida R.T. (1998):** Endovascular treatment of mural type vein of Galen malformation. *Journal of neurosurgery*; 89:74-80.

47. **Hallam D.K. and Russel E.J. (1998):** Imaging of angiographically occult cerebral vascular malformations. *Neuroimaging clinic of North America*; 8: 323-347.

48. **Hamilton M.G. and Spetzler R.F. (1994):** The prospective application of a grading system for arteriovenous malformations. *Neurosurgery*; 34: 2-7 .

49. **Haroun A. (2005):** Utility of contrast enhanced 3D turbo-flash MR angiography in evaluating the intracranial venous system. *Neuroradiology*; 47: 322-327.

50. **Hartmann A., Mast H. and Mohr J.P. (1998):** Morbidity of intracranial hemorrhage in patients with cerebral arteriovenous malformation. *Stroke*; 29: 931-934.

- 51. Hofmeister C., Stapf C. and Hartmann A. (2000):** Demographic, morphological, and clinical characteristics of 1289 patients with brain arteriovenous malformations. *Stroke*; 31: 1307-1310.
- 52. Horie N., Morikawa M. and Kitigawa N. (2006):** 2D-thick section digital subtraction angiography for the assessment of dura) arteriovenous fistulae. *AJNR*; 27:264-269.
- 53. Imbesi S.G. and Kerber C.H. (1998):** Why do ulcerated atherosclerotic artery plaques embolize ? A flow dynamic study. *AJNR*; 19:761-766.
- 54. Inci S. and Spetzler R.F. (2000):** Intracranial aneurysms and arterial hypertension: A review and hypothesis. *Surgical neurology*; 53: 530-540.
- 55. Ishimaru H., Ochi M. and Morikawa M. (2007):** Accuracy of Pre- and Post-contrast 3D Time-of-Flight MR Angiography in Patients with Acute Ischemic Stroke: Correlation with catheter Angiography. *AJNR*; 28: 923–926
- 56. Isoda H., Takehara Y. and Isogai S. (2000):** Software triggered contrast enhanced three dimension MR

angiography of the intracranial arteries. *AJR*; 174:371-374.

57. Jager H.R., and Grives J.P. (2000): Advances in non-invasive imaging of intracranial vascular disease. *Ann. R. Coll. Surg. Eng.*; 82: 1-5.

58. Kato T., Indo T. and Yoshida E. (2002): Contrast enhanced 2D-cine phase MR angiography for measurement of basilar artery blood flow in posterior circulation ischemia. *AJNR*; 23:1346-1351.

59. Kenet G., Sadetzki S. and Murad H. (2000): Factor I leiden & antiphospholipid antibodies are significant risk factors for stroke in children. *Stroke*; 31: 1283-1288.

60. Kim E.J., Halim A.X. and Duod C.F. (2004): The relationship of coexisting extracranial aneurysms to intracranial hemorrhage in patients harboring arteriovenous malformations. *Neurosurgery*; 54:1349-1358.

61. Kumar M. and Kitchen N.D. (1998): Infective and traumatic aneurysms. *Neurosurgical clinic of North America*; 9: 577-586.

- 62. Kumashiru M., Murasi K. and Oda k. (2008):** Assessment of time resolved dynamic contrast enhanced MRDSA using radial sliding –window reconstruction. Magnetic resonance medical science; volume 7, 1: 1-12
- 63. Larson P.S., Reisner A. and Morassutti D.J. (2000):** Traumatic intracranial aneurysms. Neurosurgical focus; 8: 25-32.
- 64. Larson T.C. (2003):** Cerebral aneurysms and vascular malformations. In: C.T. and MR imaging of the whole body, edited by Haaga J.R., Lanzieri F.R., Gilkeson R.C., 4th edition, Mosby: 8: 285-308.
- 65. Lasjaunias P. (1997):** A revised concept of the congenital nature of cerebral Arteriovenous malformations. Interventional neuroradiology; 3: 275-281.
- 66. Lasjaunias P. (1997):** Types and Disease Groups of Vascular Lesions. In: Vascular diseases in neonates and infants. 1st ed., Springer-Yeriag; 1: 3-51.
- 67. Lazar R.M., Connaire K. and Marshall R.S. (1999):** Developmental deficits in adult patients with arteriovenous malformations. Archive of neurology; 56: 303-106.

68. **Lecierc X., Navez, J. and Guavrit J. (2002):** Aneurysms of the anterior communicating artery treated with Guglieimi detachable coils: Follow up with contrast-enhanced MR angiography. *AJNR*; 23:1 121-1 127.
69. **Liang L., Korogi Y. and Sugahara T. (2001):** Evaluation of the intracranial dural sinuses with a 3D contrast enhanced MP-RAGE sequence: Prospective comparison with 2D TOF MR venography and digital subtraction angiography. *AJNR*; 22:481-492.
70. **Liang, L., Korogi Y. and Sugahara T. (2001):** Evaluation of the intracranial dural sinuses with a 3D contrast enhanced MP-PAGE sequence: Prospective comparison with 2D TOF MR venography and digital subtraction angiography. *AJNR*; 22: 481-492.
71. **Lubicz B., Levivier M. and Sadeghi N. (2007):** Immediate intracranial aneurysm occlusion after embolization with detachable coils: a comparison between MR angiography and intra-arterial digital subtraction angiography. *Neuroradiology*; 34: 190–197
72. **Malek A.M., Halbach V.V. and Duod C.F. (1998):** Diagnosis and treatment of arteriovenous fistulas. *Neuroimaging clinic of North America*; 8: 445-468.

73. **Mas V. and Meder J. (1998):** Cerebral venous thrombosis. In: Cerebrovascular disease; Pathophysiology, diagnosis, and management, edited by Ginsberg M.D. and Bogousslavsky J., 3rd edition, Blackwell Science; 107: 1488-1499.
74. **Mathews V.P., Ulmar J.L. and White M.X. (1999):** Depiction of intracranial vessels with MRA: The utility of magnetization transfer saturation and gadolinium. Journal of computer assisted tomography; 23: 597-602.
75. **McLaughlin M.R., Kondziolka D. and Flickinger J.C. (1998):** The prospective natural history of cerebral venous malformations. Neurosurgery; 43: 195-201.
76. **Meckel S. Maier M. and Ruiz D.S. (2007):** MR Angiography of Dural Arteriovenous Fistulas: Diagnosis and Follow-Up after Treatment Using a Time-Resolved 3D Contrast-Enhanced Technique. AJNR; 28: 877– 84
77. **Metens T., Rio F. and Baleriaux D. (2000):** Intracranial aneurysms: Detection with gadolinium enhanced dynamic three-dimensional MR angiography - Initial results. Radiology; 216: 39-46.
78. **Mistretta C.A. Grist T.M. and Korosec F.R. (1998):** 3D time-resolved contrast enhanced MR Digital

subtraction Angiography: Advantages and tradeoffs. Magnetic resonance in medicine; 40: 571-532.

79. **Moore P.M. (2000):** Vasculitis of the central nervous system. Current rheumatology reports; 2: 378-382.

80. **Mori H., Aoki S. and Masumoto T. (2002):** Two-dimensional magnetic resonance digital subtraction angiography using array spatial sensitivity encoding technique in the assessment of intracranial hemodynamics. Radiation medicine; 20:223-229.

81. **Nael K., Laub G. and Finn J.P. (2005):** Contrast enhanced MR Angiography at 3T: Indications, benefits, and outcomes. Journal of imaging technology management; 12: 22-26.

82. **Nael K., Michaely H. and Villablanca P. (2006):** Time-resolved contrast enhanced MRA of the head and neck at 3.0 Tesla - Initial result. Investigative radiology; 41: 116-123.

83. **Nael k., Villablanca J.P. and Pope W.B. (2007):** Supraaortic Arteries: Contrast enhanced MR Angiography at 3.0T— Highly Accelerated Parallel Acquisition for

Improved Spatial Resolution over an Extended Field of View. *Radiology*; 242: 600–609

84. **Nael K., Villablanca J.P. and Mossaz L. (2008):** 3-T Contrast-Enhanced MR Angiography in Evaluation of Suspected Intracranial Aneurysm: Comparison with MDCT Angiography. *AJNR*; 190: 389–395

85. **Nagaraja S., Capener D. and Coley S.C. (2005):** Brain arteriovenous malformation: Measurement of nidal volume using a combination of static and dynamic magnetic resonance angiography technique. *Neuroradiology*; 47: 387-397.

86. **Noguchi K., Melhem E.R. and Kanazawa T. (2004):** Intracranial dural arteriovenous fistulae: Evaluation with combined 3D time-of-flight MR angiography and MR digital subtraction angiography. *AJR*; 182:183-190.

87. **Nussbaum E.S., Heros R.C. and Madiron M.T. (1998):** The pathologies of arteriovenous malformations: Insights provided by a case of arteriovenous malformations developing in relation to developmental venous anomalies. *Neurosurgery*; 43:347-352.

88. **Okumara A., Wamabe Y. and Dohke M. (1999):** Contrast enhanced 3 dimensional M.R. porlography. Radiographics; 39:973-987.
89. **Osborn A.G. (1999):** Atherosclerosis and carotid stenosis. In: Diagnostic cerebral angiography, 2nd edition, Lippineott Williams and Wilkins; 16:359-378.
90. **Parker D.L., Tsuruda J.S. and Goodrich K.C. (1998):** Contrast enhanced magnetic resonance angiography of cerebral arteries. Investigative radiology; 33: 560-5.
91. **Petkova M., Gauvrit J.Y. and Trystram D. (2009):** Three-Dimensional Dynamic Time-Resolved Contrast-Enhanced MRA Using Parallel Imaging and a Variable Rate *k*-Space Sampling Strategy in Intracranial Arteriovenous Malformations. Journal of magnetic resonance imaging; 29: 7–12
92. **Provenzale J.M., Joseph G.J. and Barhoriak D.P. (1998):** Dural sinus thrombosis: Findings on CT and MR imaging and diagnostic pitfalls. AJR; 170: 777-783.
93. **Ramgren B., Siemund R. and Cronqvist M. (2008):** Follow-up of intracranial aneurysms treated with detachable coils: comparison of 3D inflow MRA at 3T and

1.5T and contrast-enhanced MRA at 3T with DSA. *Neuroradiology*; 11: 947-954

94. **Regli L. and DeTribollet N. (1998):** Cerebral vascular malformation. In: *Cerebrovascular disease: Pathophysiology, diagnosis, and management*, edited by Ginsberg M.D. and Bogousslavsky 1, 3rd edition, Blackwell science; 109: 1532-1547.

95. **Reinacher P.C., Stracke P. and Reinges H.T. (2007):** Contrast-enhanced time-resolved 3-D MRA: applications in neurosurgery and interventional neuroradiology. *Neuroradiology*; 49: 3-14

96. **Remonda L., Serin P. and Earth A. (2002):** Contrast enhanced 3D MR angiography of carotid artery: Comparison with digital subtraction angiography. *AJNR*; 23: 213-219.

97. **Rubin G.D. and Rofsky N.M., (2009):** Cerebral arteries and veins .In : *CT and MR Angiography: Comprehensive Vascular Assessment*, 1st Edition , Lippincott Williams & Wilkins ; 11: 282-348

98. **Saba P.R. (1998):** The caput medusa sign. *Radiology*; 207: 599-600

99. **Saleh R.S, Lohan J.P. and Villablanca J.P. 2008:** Assessment of Craniospinal Arteriovenous Malformations at 3T with Highly Temporally and Highly Spatially Resolved Contrast-Enhanced MR Angiography. *AJNR*; 29: 1024 –31
100. **Schneider G., Prince M.R. and Meaney J.F. (2005):** Intracranial MR Angiography. In: *Magnetic Resonance Angiography, Techniques, Indications and Practical Applications*, Springer; 3: 103- 139
101. **Selman W.R., Tarr R.W. and Ratcheson R.A. (2000):** Intracranial aneurysms and subarachnoid hemorrhage. In: *Neurology in clinical practice*, edited by Bradly W.G. et al, edition, Butterworth-Heinemann; 4: 1188-1190.
102. **Smith W.S., Johnston S.C. and Easton J.D. (2004):** Cerebrovascular disease. In: *Harrison's principles of internal medicine*, edited by Kasper D.L., 16th edition, volume 2, McGraw Hill company; 349: 1997-2002.
103. **Spektor S., Weinberger G. and Constantini S. (1998):** Giant lateral sinus percranii. *Journal of Neurosurgery*; 88: 145-147.

104. Standring S., Crossman A.R. and Turlough M.J. (2005): Vascular supply of the brain. In: Gray's anatomy, the anatomical basis of clinical practice, edited by Standring S. associates, 39th edition, Churchill Livingstone; 17: 275-305.

105. Stapf C., Mofar J.P. and Speliman P.J. (2001): Epidemiology and natural history of arteriovenous malformations. Neurosurgical focus; 5:1-5.

106. Strecker R., Arnold S. and Scheffler K. (2001): Multislab projection angiography with subsecond temporal resolution. In: Proceedings in the international society for magnetic resonance imaging in medicine. 9th annual meeting book of abstracts, Berkley, California; 301.

107. Strecker R., Scheffler K. and Klisch J. (2000): Fast functional MR angiography with correlation analysis. Magnetic resonance in medicine: 43: 303-309.

108. Taschner C.A., Gieseke J. and Thuc V.L. (2008): Intracranial Arteriovenous Malformation: Time-resolved Contrast-enhanced MR Angiography with Combination of Parallel Imaging, Keyhole Acquisition, and k-Space Sampling Techniques at 1.5 T. Radiology; 246: 871-879

- 109. Takahashi S., Tomura N. and Walarai J. (1999):** Dural arteriovenous fistula of the cavernous sinus with venous congestion of the brain: report of two cases. *American Journal of Neuroradiology*; 20: 886-888.
- 110. Tartaro A., and Torriero N. (2004):** Advantages of high relaxitivity contrast media in intracranial contrast enhanced angiography. *Highlights in MRI*; 2: 2-5.
- 111. Tsuchiya K., Aoki C. and Fujikawa A. (2004):** Three dimensional MR digital subtraction angiography using par imaging and keyhole data sampling in cerebrovascular disc Initial experience. *European radiology*,
- 112. Tsuchiya K., Katase S. And Yoshino A. (2000):** Digital subtraction angiography of cerebral malformations. *AJNR*; 21:707-711.
- 113. Unlu E., Cakir B. and Gocer B. (2005):** The role contrast enhanced MR angiography in assessment of ruptured intracranial aneurysms: A comparative study *Neuroradiology*; 47: 780-791.
- 114. Vega C, Kwoon J.V. and Lavine S.D. (2002):** Intracranial aneurysms: Current evidence and clinical practice. *Amer. family physician*; 66: 601-608.

- 115. Vinuela F., Duckweier G. and Mawad M. (1997):** Detachable coil embolization of acute intracranial aneurysms: Peri-operative anatomical and clinical outcome in 403 patients. *Journal of neurosurgery*; 86: 475-482.
- 116. VonIngersleben G., Schmeidl U.P. and Hatsukami T.S. (1997):** characterization of atherosclerotic plaques at the carotid bifurcation: correlation of high resolution MR imaging with histological analysis - preliminary study. *Radiographics*; 17: 315-396.
- 117. Wallace R.C., Karis J.P. and Partovi S. (2007):** Noninvasive Imaging of Treated Cerebral Aneurysms, Part I: MR Angiographic Follow-Up of Coiled Aneurysms. *AJNR*; 28:1001– 08
- 118. Wardlaw J.M. and White P.M. (2000):** The detection and management of unruptured intracranial aneurysms. *Brain*, 123. 205-221.
- 119. Warren B.J., Hoggard N. and Walton L. (2001):** Cerebral arteriovenous malformations: Comparison of novel magnetic resonance angiography technique and conventional catheter angiography. *Neurosurgery*; 48: 973-982.

- 120. Weiger M., Pruessman K.P. and Kassner A. (2000):** Contrast enhanced 3D MRA using SENSE. Journal of magnetic resonance imaging; 12:671-677.
- 121. Wetzel S.G., Bilcer, D. and Lyrer P. (2000):** Cerebral dural arteriovenous fistulae: Detection by dynamic MR projection angiography. AJR; 174:1293-1295.
- 122. Willinek W.A., Falkenhausen M. and Born M. (2002):** Non invasive detection of steno-occlusive disease of the supraaortic arteries with three dimensional contrast enhanced magnetic resonance angiography - A prospective, comparative analysis with digital subtraction angiography. St 36:38-43.
- 123. Willinek W.A., Gieseke J. and Conrad R. (2005):** Randomly segmented central K-space ordering in high resolution, contrast enhanced MR angiography of the supraaortic arteries: Initial experience. Radiology; 225:583-588.
- 124. Wilms G., Bosnians H. and Beraaerel P. (2000):** Magnetic resonance angiography of intracranial vessels. Euro radiology; 38: 10-58.

125. **Winn W.B., Schmeidl U.P. and Reichcnbach D.D. (1998):** Detection and characterization of atherosclerotic IT caps with T2 weighted MR. *AJNR.*, 19: 129-134.
126. **Wutke R., Lang W. and Fellner C. (2002):** high resolution contrast-enhanced magnetic resonance angiography with elliptic centric k-space ordering of supra-aortic at compared with selective X-ray angiography. *Stroke*; 33: 1529.
127. **Yamaura A., Ono J. and Hiari S. (2000):** Clinical picture of intracranial non-traumatic dissecting aneurysm. *Neuropathology*; 20:85-90.
128. **Yang J.J., Hill M.D. and Morrish W.F. (2002):** Comparison of pre and post-contrast 3D time-of-flight MR angiography for the evaluation of distal intracranial branch occlusion in acute ischemic stroke. *AJNR*; 23: 557-567.
129. **Yano T., Kodama T. and Suzuki Y., et al., (1997):** Gadolinium enhanced magnetic resonance angiography. *Acta radiologica*; 38: 47-54.
130. **Younger D.S. (2004):** Vacuities of the nervous system. *Current opinion in neurology*; 17(3): 317-336.
131. **Zhu W., Feng D. and Qi J. (2004):** Evaluation of large intracranial aneurysm with cine MRA and 3D

contrast enhanced MRA. Journal of Huazhong University of science technology and medical sciences; 24: 95-98.

132. **Ziyeh S., Strecker R. and Berlis A. (2005):** Dynamic 3DMR Angiography of intra and extra cranial vascular malformations at 3T: A technical note. AJNR; 26: 630-634.