

REFERENCES

- *Abbot R, Yin-yin M, Reed D, Yan K. Risk of stroke in male cigarette smokers. N Eng J Med 2004;*
- *Albers GW: diffusion-weighted MRI for evaluation of acute stroke. Neurology 51(supplement 3):s47-s49,1998*
- *Alter M, Lai S, Friday G, stroke recurrence in diabetics. Does control of blood glucose reduce risk ? stroke 2005;*
- *American heart association 2002 heart and stroke stat update. Dallas , Texas: American heart association 2002*
- *Bammer R, Skare S, Newbould R, Liu C, Thijis V, Ropele S, Clayton DB, Krueger G, Moseley ME, and Glover GH, foundation of advanced magnetic resonance imaging. NeuroTX 2005 vol 2, 167-196.*
- *Barbier el, Lamalle L, Decorps M, 2002 methodology of brain perfusion imaging, 13:496-526*
- *Barber PA, Darby DG, Desmond PM et al. Prediction of stroke outcome with echoplanar perfusion- and diffusion-weighted MRI. Neurology 2005; 51: 418–426.*
- *Barone JC, (2001) mapping the ischemic penumbra with new approaches.*
- *Basser PJ, Pierpaoli C (2003) Microstructural and physiological features of tissues elucidated by quantitative-diffusion- tensor MRI. J Magn Reson B 111(3):209–219*
- *Beauchamp NJ, Barker P B & Wang P Y: imaging of acute cerebral ischemia radiology 2004; 212:307-324.*

- Calamante F, Gadian DG, Connelly A (2005) Delay and dispersion effects in dynamic susceptibility contrast MRI: simulations using singular value decomposition. *Magn Reson Med* 44:466–473
- Cipolla Mj, Porter Jm, Osol G. high glucose concentration dilate cerebral arteries and diminish myogenic tone through an endothelial mechanism. *Stroke* 2003;
- Copen WA et al (2003) Ischemic stroke: effects of etiology and patient age on the time course of the core apparent diffusion coefficient. *Radiology* 221(1):27–44
- Degraba TJ, Siren Al, Penix L, Mccarron Rm, Hargraves R, Sood S, Pettigrew Kd, Hallenbeck Jm. Increased endothelial expression of intracellular adhesion molecule-1 in symptomatic versus asymptomatic human carotid atherosclerosis plaque stroke. *Stroke* 2004
- De Myer W: Blood supply of the central nervous system: In *Neuroanatomy* De Myer W ed. John Wiley & Sons, Inc.; 2002, 335-353.
- Deok HL, Dong-Wha K, Ahn JS, Choi CG, Kim SJ, et al., imaging of the ischemic penumbra in acute stroke. *Korean journal of radiology*; 2005;
- Dirnag U, Iadecola C, Moskowitz Ma. Pathology of ischemic stroke: an integrated view. *Trends Neurosci* 2006.
- Di Salle R, Duvernoy H., Rabischong R, Scarabino T., Salvolini U., *Atlas of Morphology and Functional Anatomy of the Brain*. Springer-Verlag, Berlin, 2005.
- Ditullio M, Sacco Rl, Homma S, et al. atrial size and the risk of ischemic stroke in an ethnically mixed population. *Stroke* 2003;

- *Duvernoy H.M., The Human Brain: Surface, Blood Supply, and Three Dimensional Sectional Anatomy, 2nd edition. Springer, Vienna, 2001.*
- *Engelter S et al (2003) Infarct volume on apparent diffusion coefficient maps correlates with length of stay and outcome after middle cerebral artery stroke. Cerebrovasc Dis 15(3):188–191*
- *Engelter S et al (2004) The clinical significance of diffusion-weighted MR imaging in infratentorial strokes. Neurology 62(4):474–480*
- *Fiehler J et al (2002) Severe ADC decreases do not predict irreversible tissue damage in humans. Stroke 33(1):79–86*
- *Fiehler J et al (2001) Apparent diffusion coefficient decreases and magnetic resonance imaging perfusion parameters are associated in ischemic tissue of acute stroke patients. J Cereb Blood Flow Metab 21(5):577–584*
- *Folsam A, Prineas R, Kaye S, Munger R. incidence of hypertension and stroke in relation to body fat and other risk factor in older women. Stroke 2004;*
- *Gallucci M., Sparvoli L., Patologia delVosso temporale. In: Dal Pozzo G., Compendio di Risonanza Magnetica, Cranio e Rachide. UTET, Torino, 2001:807-836.*
- *Gorelick P. the epidemiology and trials. Brain ischemia basic concept and clinical releavance 2005,;*
- *Grandin CB, Duprez TP, Smith AM, Oppenheim C, Peeters A, Robert AR, Cosnard G (2002) Which MR-derived perfusion parameters are the best predictors of infarct growth in hyperacute stroke? Comparative study between relative and quantitative measurements. Radiology 223:361–370*

- *Grant PE et al (2001) Frequency and clinical context of decreased apparent diffusion coefficient reversal in the human brain. Radiology 221(1):43–50*
- *Heiss Wd, ischemic penumbra: evidence of functional imaging in man. cerebral blood flow 2005*
- *Herrero, Barcia C., Navarro J.M., Functional anatomy of thalamus and basal ganglia. Child's Nerv. Syst. 18:386-404, 2002.*
- *Hossmann Ka. Viability threshold and penumbra focal ischemia. Ann neurol 2004*
- *Karonen JO, Liu Y, Vanninen RL, Ostergaard L, Kaarina Partanen PL, Vainio PA, Vanninen EJ, Nuutinen J, Roivainen R, Soimakallio S, Kuikka JT, Aronen HJ (2000) Combined perfusion- and diffusion-weighted MR imaging in acute ischemic stroke during the 1st week: a longitudinal study Radiology 217:886–894*
- *Karonen JO, Nuutinen J, Kuikka JT. SPECT and diffusion-weighted MRI as a principle of assessment of infarct growth in acute stroke. 2005; 41:788-794.*
- *Kaufmann Am, Firlik AD, Fukui Mb, ischemic core and penumbra in human stroke 2006*
- *Kogure K, and Kato H. pathophysiology, diagnosis and management of acute cerebral stroke 2003*
- *Kuusisto J, Pyroala K, non insulin dependent diabetes and its metabolic control are important predictors of stroke in elderly subjects stroke 2004;*
- *Lanfermann H, Kugel H, Heindel W, Herholz K, Lackner K, metabolic changes in acute and subacute cerebral infarction 2005.*

- *Le Bihan D et al (2001) Diffusion tensor imaging: concepts and applications. J Magn Reson Imaging 13(4):534–546*
- *Liao D, Myers R, Hunt S, Shahar E .familial history of stroke. Stroke 1997;*
- *Liu y, Karonen JO, Vanninen RL Soimakallio S and Aronen HJ: Acute ischemic stroke: predictive value of 2D phase contrast angiography-serial study with combined diffusion perfusion MR imaging radiology 2004; 231:517:527.*
- *Marks MP et al (2002) Evaluation of early reperfusion and i.v. tPA therapy using diffusion- and perfusion-weighted MRI. Neurology 52(9):1792–1798*
- *Mullins ME et al (2002) CT and conventional and diffusion-weighted MR imaging in acute stroke. Radiology 224(2):353–360*
- *Nagesh V et al (2002) Time course of ADCw changes in ischemic stroke: beyond the human eye! Stroke 29(9):1778– 1782*
- *National institute of neurological disorders and stroke using t-pa study group; t-pa therapy for ischemic stroke 28;2109-2118, 2002*
- *Neumann-Haefelin T, Wittsack HJ, Wenserski F et al. Diffusion- and perfusion-weighted MRI. The DWI/PWI mismatch region in acute stroke. Stroke 2004; 30: 1591–1597.*
- *Niendorf T et al (2006) Biexponential diffusion attenuation in various states of brain tissue: implications for diffusion weighted imaging. Magn Reson Med 36(6):847–857*
- *Nolte J., The Human Brain. An introduction to its functional anatomy, 4th edition. Mosby, St. Louis, MO, 2004.*
- *O'Leary D, Anderson K, Wolf P., cholesterol and carotid atherosclerosis in older persons : the Framingham study. AM J epidemiol 2002.*

- *Ostergaard L, Jhoannsen P, Vestegaard-Poulsen P, Asobe H, et al; cerebral blood flow measurements by magnetic resonance imaging bolus tracking: 2004 18:935-940*
- *Powers WJ (2001) Cerebral hemodynamics in ischemic cerebrovascular disease. Ann Neurol 29:231–240*
- *Reed Sj, Hirano T, Abbott Df, Markus , Sachinidis Ji, Tochon-Danguy Hj. Et al., the fate of hypoxic tissue on 18f-fluoromisonidazole positron emission tomography after ischemic stroke, Ann Neurol 2002.*
- *Robert L, Herting J, and Nora R. Frohberg. Department of family medicine university of Iowa family practice handbook, 3rd edition, chapter 14, 2001.*
- *Rohl L et al (2003) Correlation between diffusion- and perfusion-weighted MRI and neurological deficit. Cerebrovasc Dis 12(3):203–213*
- *Rordorf G, Koroshetz WJ, Copen WA et al. Regional ischemia and ischemic injury in patients with acute stroke as defined by early diffusion-weighted and perfusion-weighted MRI. Stroke 2001; 29: 939–943.*
- *Rowely Ha, noninvasive imaging of the ischemic penumbra. Ann Neurol 2002*
- *Ryan SP, McNicholas MJ: The Central Nervous System: In Anatomy for diagnostic imaging. Ryan SP, McNicholas MJ, ed. WB Saunders Company Ltd; 2004: 43-84.*
- *Sa de Camagro EC, Koroshetz WJ. Neuroimaging of ischemia and infarction. NeuroRX.2005;2:256-276.*

- *Sacco R., identifying patient populating at high risk for stroke. Neurology 2002;*
- *Sette G. pathophysiology of ischemic stroke. In: Castillo J, Davalos A, Toni D., management of acute ischemic stroke 2004.*
- *Schaefer PW, Grant PE, Gonzalez RG (2002) Diffusion weighted MR imaging of the brain. Radiology 217(2): 331–345*
- *Schaefer PW et al (2003) Predicting cerebral ischemic infarct volume with diffusion and perfusion MR imaging. Am J Neuroradiol 23(10):1785–1794*
- *Schaefer PW, et al. The ischemic penumbra: operationally defined by diffusion and perfusion MRI. Neurology 53:1528–1537*
- *Schlaug G et al (2001) Time course of the apparent diffusion coefficient (ADC) abnormality in human stroke. Neurology 49(1):113–119*
- *Shih LC, Saver JL, Alger JR, Starkman S, Leary MC, Vinuela F, Duckwiler G, Gobin YP, Jahan R, Villablanca JP, Vespa PM, Kidwell CS (2006) Perfusion-weighted magnetic resonance imaging thresholds identifying core, irreversibly infarcted tissue. Stroke 34:1425–1430*
- *Shih LC, Saver JL, Duckwiler G, Gobin YP, Jahan R, Villablanca JP, Vespa PM, (2005) Diffusion weighted MR imaging of the brain. Stroke 22:998–1050*
- *Singer MB et al (2004) Diffusion-weighted MRI in acute subcortical infarction. Stroke 29(1):133–136*

- *Sorensen AG (2001) Predicting tissue outcome in acute human cerebral ischemia using combined diffusion- and perfusion-weighted MR imaging. Stroke 32:933–942*
- *Speck O, Chang L, DeSilva N, Ernst T (2004) Perfusion MRI of the human brain with dynamic susceptibility contrast. Gradient-echo versus spin-echo techniques. J Magn Reson Imaging 12:381–387*
- *Tamraz J.C, Comair Y.G., Atlas of Regional Anatomy of the Brain Using MRI with Functional Correlations. Springer- Verlag, Berlin, 2003.*
- *Thijs V, Somford D, Bammer R, Robberecht W, Moseley ME, Albers GW (2004) Influence of arterial input function on hypoperfusion volumes measured with perfusion-weighted imaging. Stroke 35:94–98*
- *Wei H, Chi O, Liu X, Weis H, oxygen balance in focal cerebral ischemia 2004.*
- *Werring DJ et al (2002) Diffusion tensor imaging can detect and quantify corticospinal tract degeneration after stroke. J Neurol Neurosurg Psychiatry 69(2):269–272*
- *Wittsack HJ, Ritzl A, Fink GR, Wenserski F, Siebler M, Seitz RJ, Modder U, Freund HJ (2002) MR imaging in acute stroke: diffusion-weighted and perfusion imaging parameters for predicting infarct size. Radiology 222:397–403*
- *Wolf P, Abbott R, Kannel W., independent risk factors for stroke 2002. stroke.*
- *Wu O, Koroshetz WJ, Ostergaard L, Buonanno FS, Copen WA, Gonzalez RG, Rordorf G, Rosen BR, Schwamm LH, Weisskoff RM,*
- *Yamada K, Wu O, Gonzalez RG, Bakker D, Ostergaard L, Copen WA, Weisskoff RM, Rosen BR, Yagi K, Nishimura T, Sorensen AG (2002) Magnetic resonance perfusion weighted imaging of acute*

cerebral infarction: effect of the calculation methods and underlying vasculopathy. Stroke 33:87–94

- *Yang Q et al (2004) Serial study of apparent diffusion coefficient and anisotropy in patients with acute stroke. Stroke 30(11):2382–2390*