

imaging results but high clinical suspicion of RVH, and patients with abnormal functional imaging results who have a contraindication to conventional angiography, such as renal failure or a history of allergy to iodinated contrast material.

REFERENCES

1. **Berkow R, Fletcher AJ:** The Merck Manual, Merck Research Laboratories, and 429- 430:1992
2. **Zierler RE, Bergelin Ro, Isaacson JA, et al:** Natural history of atherosclerotic renal artery stenosis: a prospective study with duplex · ultrasonography. J Vasc Surg; 19:250-258. 1994
3. **Decobelli F, Mellon R, Sailori M, et al:** Renal artery stenosis value of Screening with three-dimensional phases contrast MR angiography with phase array multicoil. Radiology.201:697-703, 1996
4. **Hunt JC, Sheps SG, Harrison EG, et al:** Functional characteristics of renovascular hypertension: a reasoned approach to diagnosis and management. Arch Intern Med 1974; 133:988-1999
5. **Safian RD, Textor SC:** Renal- artery stenosis. N Engl J Med 344:-422, 2001
6. **Bruce S Spinnowitz, Jonna Rodriguez MR, et al:** angiography of renal artery Stenosis; value of the combination of three dimensional time phase-contrast MR angiography sequence .AJR J Rontgenol 167(2): 489-494, 1996
7. **Dorros G, Jaff M, Matlniak L, et al:** Four-year follow-up of Palmaz-Schatz stent revascularization as treatment for atherosclerotic renal artery stenosis. Circulation; 98:642-647. 1998
8. **Krijnen P, van Jaarsveld BC, steyerberg EW,et al: and Man in Veld AJ, Schalel amp MA Jabbema JD:** A clinical

prediction-rule for renal artery stenosis Ann intern Med 129: 705-711, 1998.

9. Berland LL, Kuslin DB, Routh WD, et al: Keller FS. Renal artery stenosis: prospective evaluation of diagnosis with color duplex US compared with angiography. Radiology; 174:421-423. 1990

10. Miralles M, Cairols M, Cotillas J, et al: Gimenez A, Santiso A. Value of Doppler parameters in the diagnosis of renal artery stenosis. J Vasc Surg; 23:428-435. 1996

11. Journal of the American society of Nephrology volume 13 Number 11 November 2002 copy right © 2002 American society of Nephrology.

12. Beregi JP, Elkohen M, Deklunder G, et al: helical CT angiography compared with angiography in the detection of renal artery stenosis. AJR Am J Roentgenol 167:494-501, 1996.

13. Earls JP, Rofsky NM, DeCorato DR, et al: Breath-hold single dose Gd-enhanced three-dimensional MR aortography: usefulness of a timing examination and MR power injector. Radiology; 201:705-10. 1996

14- Van Jaarsveld BC, Pieterman H, Van Dijk LC ,et al: van Seijen AJ, Krinjen P, Derkx FH, Man in't Veld AJ, Schalekamp MA: Inter-observer variability in the angiographic assessment of renal artery stenosis 17: 1731-1736, 1999

15. Wasser MN, Wesenberg J, van der Hulst, et al: Hemodynamic significance of renal artery stenosis: digital subtraction versus systolically gated three-dimensional phase-contrast MR angiography. 1997

16. Dawson, Westenberg J, Aslmk Ragayan, et al: digital subtraction I Angiography, clinical radiology, 39, 5, 474-477, 1998

17. Taylor A, Nally J, Au-rell M, et al: Consensus report on ACE inhibitor renography for detecting renovascular hypertension. Nuclear Med; 37:1876-82. 1996

- 18. Taylor AT, Fletcher IW, Nally JR JV, et al:** Procedure guidelines for diagnosis of renovascular hypertension. *J Nucl Med*; 39:1297-1302, 1998
- 19. Snell Richard S:** clinical anatomy: 8th edition 13-135:453-4155, 2007
- 20. Resnick M1, Pounds DM, Boyce WH, et al:** Surgical anatomy of the human kidney and its applications. *Urology* 17:367, 1981.
- 21. Gray H, Pick TPB Howden R:** Gray's Anatomy, New York, Bounty Books, 556-557, 985-994. 1984
- 22. Lasts anatomy:**Regional and applied, 8 edition, 70-71,Churrchill Livingstone book company, U.K. 1991.
- 23. Sampaio FJ, Aragao AH;** Anatomical relationship between the intrarenal arteries and the kidney collecting system. *J Urol*, 1413:679, 1990.
- 24. Tanagho EA:** Anatomy of the lower urinary tract. InWalsh PC, Retik AB, and STOMEY TA, et al (Eds):Campbell's Urology. Philadelphia, WB Saunders, pp 40-69,1992.
- 25.Wilman AH, Riederer SJ, King BF, Dobbins .JP, Rossman PJ.EhmanRL:**f1uoroscopically triggered contrast-enhancedthree -dimensional MR angiography with elliptical centric view order: Application to the renal arteries.*Radiology* 205; 137 146, 1997.
- 26. Henry Gray:** Anatomy of the Human Body.1211-1222, Mosbey puplisher 1996
- 27. Walsh P, Rofsky NM,Krinsky GA, et al:**Asymmetric siognal intensity of renal artery stenosis follow administration of gadopentetate dimeglumine J; 20:812-814; 1996
- 28. Ashcraft KW:** pediatric Urology,W.B.Ssaunder,77-6-86:1999
- 29. Berkow R, Fletcher AJ:** The Merck Manual, Merck Research Laboratories, 429-430; 1992.
- 30. Rubin GD, VVa1ker PJ, Dake MD, et al:** Three-dimensional spiral computed tomographic angiography; an

alternative imaging modality for the abdominal aorta and its branches. *J Vase Surg* 18:656-665, 1993

31. Safian RD, Textor, or SC. Renal artery stenosis *Engl J Med*: 344:431- 42. 2001.

32. SOS, TA, Pickering, T.G, Sinderman, and K: Percutaneous transluminal renal angioplasty in Renal vascular hypertension due to atheroma or fibromuscular dysplasia. *N. Engl.j.Med* .309:274 1983.

33. Textor SC, Wilcox C Ischemic nephropathy/azotemic renovascular disease. *Sem Nephrol*, 20:489-502. 2000;

34. Lee VS, Rofsky NM, Krinsky GA, et al: Single dose breath-hold gadolinium-enhanced three-dimensional MR angiography of the renal arteries., 211:69-78., *Radiology* 1999 .

35. Walsh P, Rofsky NM, Krinsky GA, et al: Weinreb, Asymmetric signal intensity of the renal collecting system as a sign of unilateral renal, artery stenosis following administration of gadopentetate dimeglumine. *J.Assist Tomogr*; 20:812-814; 1996.

36. Anderson JR: Murray's textbook of pathology. London. Edward Arnold (publisher). 14, 1:14-36. 1995

37. Desantis RW, Doroghazi RM, Austin WG, et al: Aortic dissection, *N .Engl.J.Med.*:317:1060.1987

38. Hall S: Takayasu arteritis. A study of 32 North American patients. *Medicine*, 1985;64:89.

39. Schreiber MJ, Pohl MA, Novick AC: The natural history of atherosclerotic and fibrous renal artery disease. *Urol Clin NAm*, 11; 383-92. 1984.

40. Fry WJ, Ernest CB, Stanley JC: Renovascular hypertension in the pediatric patient. *Arch. Surg* 198;3:107,692-698.1983

51. Taylor Jr AT, Fletcher JW, Nally Jr JV, Blafox MD, Dubovsky EV, Fine EJ, et al.; Procedure guideline for diagnosis of renovascular hypertension. *Society of Nuclear Medicine Nucl Med*, 39;1297-302 1998

52. **Nada NC:** History of echocardiographic contrast agents
clin cardiol,20:17-11;1997.
53. **Burns physical principles of Doppler and spectral A:**
analysis of JCU; 15:567-590; 1995
54. **Olin JW:** Role of duplex ultrasonography in screening for
significant renal artery disease. Urol Clin North Am.; 0 21:215-
26; 1994.
55. **Barnes RW,Bone GE and Reinterson J:** Non invasive US
carotid angiography : prospective validation by
contrastangiography.surgery 1979;80382. 1979
56. **Francesco De Cobelli, , Massimo Venturini, , Angelo
Vanzulli, , Sandro Sironi, , Marco Salvioni, , Enzo Angeli,**
Renal Arterial Stenosis: Prospective Comparison of` Color
Doppler US and Breath-hold, Three-dimensional, Dynamic
Gadolinium-enhanced. MR. Angiography Radiology.
214:373-380, 2000.
57. **Krebs CA, Giyanni VZ, Eisenberg RL,et al:** Ultrasound
atlas of1 J vascular diseases. Appelton and Lange.; 184-187.
1999
58. **Hahn U, Miller S, Nagele T, et al:** Renal MR angiography
at 1.0 T: three-dimensional (3D) phase-contrast techniques
versus gadolinium-enhanced 3D fast low-angle shot breath-hold
imaging. AJR; 172:1501-1508.1999
59. **Duda SH, Schick F, Teufl F, et al.** Phase-contrast MR
angiography for detection of arteriosclerotic renal artery
stenosis. Acta Radiol; 38:287-291. 1997
60. **Rosner MH:** Renovascular hypertension: can we identify
apopulation at high risk,South Med J, 94:1058-1064; 2001.
61. **Nazzal M·MS, Hoballah JJ, Miller EV, et al:** Renal hilar
Doppler analysis is of value in the management of patients with
renovascular disease. Am J Surg 174:164-168, 1997.
62. **Melany ML, Grant EG, Duerinckx AJ, Watts TM,**
Levine0 BS: Ability of a phase shift US contrast agent to
improve imaging of` the main renal arteries, Radiology;
205:147-152; 1997.

- 63. Missouriis CG, Allen:CM, Balen FG, Buckenham T, LeesWR, MacGregor GA:** Non-invasive screening for renal artery5 stenosis with ultrasound contrast enhancement J Hypertens, 14:519-524;1996
- 64. Porter TR, Li S, Kilzer K:** Smaller intravenous perfluorocarbon-containing micro bubbles produce greater myocardial contrast with intermittent harmonic imaging and better delineation of risk area during acute myocardial ischemia J Am Soc Echocardiogr, 10:792-797; 1997.
- 65. Helenon O, El Rody F, Correias JM, Melki P, Chauveau D,Chretien Y, Moreau IF:** Color Doppler US of renovascular disease in native kidneys. Radiographics, 15:833-54; 1995
- 66. Souza de Oliveira IR, Widman A, Molnar LJ, JT,Praxedes_JN, Cerri GG:** Colour Doppler ultrasound: a new index improves the diagnosis of renal artery stenosisUltrasound Med Biol, 26:41-47; 2000.
- 67. Claudon M, Plouin PF, Baxter GM, Rohban T, Devos DM:** Renal arteries in patients at risk of renal arterial stenosis; multicenter evaluation of the echo-enhancer Renal Artery Study Group Radiology, 214:739-746; 2000.
- 68. Teixeira, Odila UN, Luiz A Bortolotto** and Helio Bernardes Silva. The contrast-enhanced Doppler ultrasound with perfluorocarhon exposed sonicated albumin does not improve the diagnosis of renal artery stenosis compared with angiography. Journal of Negative Results inBiomedicine, .11861477-5751-3-3; 2004.
- 69. Keller PJ, Saloner D, Potchen EJ, Siebert JE, and Haacke E.M:** Time of flight imaging in Magnetic Resonance Angiography, concepts and applications, St. Louis, Mosby ,:146-159. 1993
- 70. Schoenberg SO, Bock M, Knopp MV, et al:** Renal arteries: optimization of three-dimensional gadolinium-enhanced MR angiography with bolus-timing-independent last

multiphase acquisition in a single breath hold. Radiology,21 1:667-79. 1999.

71. Maki JH, Prince MR, Londy FJ, et al: The effects of time varying intravascular signal intensity and k-space acquisition order on three-dimensional MR angiography image quality. J Magn Reson Imaging, 6; 642-51. 1996.

72. Marks B, Mitchell DG, Simelaro JP, et al: Breath-holding in healthy and pulmonary-compromised populations: effects of hyperventilation and oxygen inspiration. JMRI; 7:595-597; 1997.

73. Gilfeather M., Yoon HC, Siegelman ES, et al:Renal artery stenosis: evaluation, With conventional angiography

versus gadolinium-enhanced MR angiography. Radiology; 210:367-9 372; 1999.

74. Prince MR, Yucel EK, Kaufman JA, et al: Dynamic gadolinium-enhanced three-dimensional abdominal MR arteriography. J Magn Reson Imaging; 3; 877-81+1998.

75. Schoenberg SO, Knopp MV, Londy F et al: Morphologic and functional magnetic resonance imaging of the renal artery stenosis: a multi-reader tri-center study. J Am Soc Nephrol; 13: 158-169; 2002

76. Schoenberg SO, Knopp MV, Bock M et al: Renal artery stenosis; grading of hemodynamic changes with cine phase-contrast MR blood flow measurements. Radiology; 11:203: 45-53; 1997.

77. Maki JH, Chenevert TL, Prince MR: Contrast-enhanced MRangiography. Abdomen imaging; .23:469-484; 1998.

78. Weiger M, Pruessmann KP, Kassner A et al: Contrast-enhanced 3D MRA using SENSE. J Magn Reson Imaging; 12: 671-677; 2000.

79. Wedeen VJ, Melni RA, Edelman RR Kassner A, et al. Contrast-enhanced 3D MRA using SENSE. J Magn Reson imaging; 12:671-7. 2000

- 80. Ruehm SG, Goye Barkhausen J et al:** Rapid magnetic resonance angiography for detection of atherosclerosis. *Lancet*; 357: 1086-1091; 2001.
- 81. Levy BI, Duriez M, Samuel JL.** Coronary micro vasculature alteration in hypertensive rats. Effect of treatment with diuretic and an ACE inhibitor. *Am J Hypertens*; 14: 7-13; 2001.
- 82. Dietrich O, Nikolou K, Wintersperger B et al:** Integrated parallel acquisition techniques (iPAT): applications for fast and cardiovascular MR imaging. *Electromedia*; 70: 133-145; 2002
- 83. Schoenberg SO, Bock M, Kallinowski F, Just A.** Correlation of hemodynamic impact and morphologic degree of renal artery stenosis in a canine model. *J Am Soc Nephrol*; 11:2190-2198; 2000.
- 84. Aumann S, Schoenberg SO, Just A, et al:** Quantification of renal perfusion using an intravascular contrast agent (part 1): results in a canine model. *Magn Resonance Med* 2003; 49:276–287
- 85. Buckley DL, Shurraf AE, Jones AP, et al:** Kilgallon JE, Mamtola H, Kalra PA. Quantitative assessment of renal function using dynamic Gd-DTPA-enhanced MRI. *Proceeding of the Tenth Scientific Meeting and Exhibition. International Society of Magnetic Resonance in Medicine*; 811; 2002.
- 86. FOX, S. H, Tanenbaum, L. N., Ackelsberg, S., He, H. D, Hsieh, J., Hu, and H.:** Future directions in CT technology. *Neuro imaging clinics of North America*; 8:497-513; 1998.
- 87. Brink JA, Davros WJ, Zeman RK, Costello P (edn):** Helical Spiral CT. a practical approach. New York, McGrawHill, 1-26; 1995.
- 88. Foo TK, saranathan M, Prince MR, et al:** automated detection of bolus arrival and initiation of data acquisition in fast, three-dimensional gadolinium-enhanced MR angiography. *Radio1ogy*; 203:275-80, 1997

- 89. Kaplan-Parloveie S, Nadja C:** Captopril renography and duplex Doppler sonography in the diagnosis of renovascular hypertension. *Nephrol Dial Transplant* 13:313-317, 1998.
- 90. Kopka L, Vosshenrich R, Rodenwaldt J, et al:** Differences in injection rates on contrast-enhanced breath-hold three-dimensional MR angiography. *AJR AmJ Rontgenol* 170:345-8. 1998.
- 91. Kuszyk BS, Fishman EK.** Technical aspects of CT angiography, *Semin Ultrasound MR*; 19:383-393; 1998.
- 92. Chapman S, Nakielny R:** A guide to radiological S procedures. *Baillire Tindal* 1, 201-241; 1997.
- 93. Uday D. Patil, Ashok Ragavan, Nadaraj,** Keshava M , Ravi Shankar, Bahar Bastani and Sudarshan H. Ballal, PJ; Renal artery stenosis: CT angiography-comparison of real-time volume rendering and intensity projectional algorithms *Nephrol Dial Transplant*, 16:1900-1904;2001
- 94. Prigent A. and Nally JV,** the diagnosis of renovascular hypertension: the role of captopril renal scintigraphy and related issues. *Eur J Nucl Med*; 20:625-644; 1998.
- 95. Aitkison F, Page A:** Diagnostic imaging of renal artery stenosis. *J Hum Hypertension*; 1999;13:595-603.
- 96. Boos M, Seheffler K, Haselhorst R, et al.** Arterial first pass gadolinium-CM dynamics as a function of several intravenous saline flush and Gd volumes. . *J Magn Reson Imaging* 13:568-76,2001.
- 97 . Rimmer J, Gennari FJ.** Atherosclerotic renovascular disease and progressive renal failure. *Ann Intern Med*, 118:712-9. 1993.
- 98. Handa N, Komada T.** Efficacy of echo-Doppler for the valuation of renovascular disease. *Ultrasound Med Biol*; 14:1-5.1998
- 99. Patriquin H, Lafortune M, Jéquier JC, et al:** Stenosis of the renal artery: assessment of slowed systole in the down stream circulation with Doppler sonography. *Radiology* 1992; 184:479-485.

- 100. Halpern EJ, Needleman L, Nack TL, East SA.** Renal artery stenosis: should we study the main renal artery or segmental vessels. *Radiology* 1995; 195:799-804
- 101. Prince MR,** Gadolinium-enhanced MR aortography. *Radiology* 1994; 191:155-164.
- 102. Bakker J, Beek FJ, Beutler JJ, et al.** Renal artery stenosis and accessory renal arteries: accuracy of detection and visualization with gadolinium-enhanced breath-hold MR angiography. *Radiology* 1998; 207:497-504
- 103 . Halpern EJ, Mitchell DG, Wechsler RJ, et al:** Preoperative evaluation of living renal donors: comparison of CT angiography and MR angiography. *Radiology* 2000; 216:434-439
- 104. Bruce A. Urban, Lloyd E. Ratner, and Elliot K. Fishman;** Three-dimensional Volume rendered CT Angiography of the Renal Arteries and Veins: Normal Anatomy, Variants, and Clinical Applications I Radiographics. 21:373-386; 2001.
- 105. Barrett BJ. And Halpern EJ,** Contrast nephrotoxicity. *J Am s Soc Nephrol*; 5: 125-137; 1994.
- 106. Siegel man ES, Gilgeather M, Holland GA,**et al: Breath-holdultrafast three-dimensional gadolinium enhanced MRangiography of Reno vascular system. *AM Rontgenol*; 168:p 1035-1040; 1997.
- 107. Gross CM, Kramer J, Weingartner O. et al:** Determination of renal arterial stenosis severity: comparison of pressure gradient and vessel diameter. *Radiology*. 2001; 220: 751-64].
- 108. Tegtmeier CJ, Peter B. Sachs** Renal Angioplasty,In:Abram's Angiography, Little Brown, RA: 294-325; 1999.
- 109. Mittai BR, Kumar P, Arora P, et al:** Role of captopril renography in the diagnosis of renovascular hypertension. *Am J Kidney Dis* 1996; 28:209-213

- 110. Prigent A.** The diagnosis of renovascular hypertension: the role of captopril renal scintigraphy and related issues. *Eur J Nucl Med* 1993; 20:625-644.
- 111. Harding MB, Smith LR, Hammerstein SI, et al:** Renal artery stenosis; prevalence and associated risk factors in patients undergoing routine cardiac catheterization.
- 112. Carman T, Olin JW, Czum J** Noninvasive imaging of renal arteries. *Urol Clin N am*: 28:815-26. 2001
- 113. Platt JF;** Doppler ultrasound of the kidney. *Semin Ultrasound CT MR* 18:22-32, 1997.
- 114. Malatino LS, Polizzi G, Garozzo M, et al:** Diagnosis of renovascular disease by extra- and intrarenal Doppler parameters. *Angiology* 49:707-721, 1998.
- 115. Isaacson J, Neumyer MM:** Direct and indirect renal arterial duplex and Doppler color flow evaluations. *Journal o Vascular Technology* 19(5-6): 309-316, 1995
- 116. Radermacher J, Chavan A, Bleek J, et al:** Use of Doppler ultrasonography to predict the outcome of therapy for renal artery stenosis. *N Engl J Med* 344:410-7. 2001
- 117. Miralles M, Santiso A, Gimenez A, et al:** Renal duplex scanning; Correlation with angiography and isotopic renography., *Eur J Vasc Surg* 7:188-194, 1993
- 118. Helenon O, Meki P, Correas J-M, Boyer J-c, Moreau J-F** Renovascular diseases: Doppler ultrasound. 18:136-146, 1997.
- 119. Leung DA, Hoffman U, Pfammatter T, et al:** Magnetic resonance angiography versus duplex sonography for L diagnosing renovascular disease. *Hypertension* 3:726-731, 1999
- 120 . Pedersen EB:** New tools in diagnosing renal artery stenosis. *Kidney Int* 57:2657-2677, 2000
- 121. Kaufman JA, McCarter D, Geller SC, et al:** Two-dimensional time-flight MR angiography of the lower extremities: artifacts and pitfalls. *AJR Am J Rontgenol*: 171:129-35. 1998

- 122. Saloner D:** Determinants of image appearance in contrast-enhanced magnetic resonance angiography; A review Invest Radiol 33:4188-4195, 1998
- 123. Hahn U, Miller S, Ngele T, et al;** Renal MR angiography at 1.0 T: Three-dimensional (3D) phase- contrast techniques versus gadolinium-enhanced 3D fast low angle .et.1 short breath-hold imaging. AJR Am J Roentge172: 11508, 1999
- 124. Joseph P. Hornak, Ph.D.** Copyright J.P. Hornak .AllRights Reserved. 1996-2002
- 125. Paul JF, Cherrak I, Jaulent MC,** Chatellier G, Plouin PF, Degoulet P, Gaux JC: Interobserver variability in the interpretation of renal digital subtraction angiography. Am J Rontgenol 173: 12854288, 1999
- 126. Teresa L.** Carman MD Jeffrey Olin DO Julianna Czum MD Urologic Clinics of North AmericaVo1ume 28 Number 4 Department of Vascular Medieine.2001

- 1.Berkow R, Fletcher AJ:** The Merck Manual, Merck Research Laboratories, and 429- 430:1992
- 2.Zierler RE, Bergelin Ro, Isaacson JA, et al:** Natural history of atherosclerotic renal artery stenosis: a prospective study with duplex · ultrasonography. J Vasc Surg; 19:250-258. 1994
- 3.Decobelli F, Mellon R, Sailori M, et al:** Renal artery stenosis value of Screening with three-dimensional phases contrast MR angiography with phase array multicoil. Radiology.201:697-703, 1996
- 4.Hunt JC, Sheps SG, Harrison EG, et al:** Functional characteristics of renovascular hypertension: a reasoned approach to diagnosis and management. Arch Intern Med 1974; 133:988-1999
- 5. Safian RD, Textor SC:** Renal- artery stenosis. N Engl J Med 344:-422, 2001
- 6. Bruce S Spinnowitz, Jonna Rodriguez MR, et al:** angiography of renal artery Stenosis; value of the combination of three dimensional time phase-contrast MR angiography sequence .AJR J Rontgenol 167(2): 489-494, 1996
- 7.Dorros G, Jaff M, Matlniak L, et al:** Four-year follow-up of Palmaz-Schatz stent revascularization as treatment for atherosclerotic renal artery stenosis. Circulation; 98:642-647. 1998
- 8.Krijnen P, van Jaarsveld BC, steyerberg EW, and Man in Veld AJ, Schalel amp MA Jabbema JD:** A clinical prediction-rule for renal artery stenosis Ann intern Med 129: 705-711, 1998.
- 9.Berland LL, Kuslin DB, Routh WD, et al:** Keller FS. Renal artery stenosis: prospective evaluation of diagnosis with color duplex US compared with angiography. Radiology; 174:421-423. 1990
- 10.Miralles M, Cairols M, Cotillas J, et al:** Gimenez A, Santiso A. Value of Doppler parameters in the diagnosis of renal arte stenosis. J Vasc Surg 1996; 23:428-435.
- 11. Journal of the American society of Nephrology** volume 13 Number 11 November2002 copy ight@2002 American society of Nephrology.
- 12.Beregi JP, Elkohen M, Deklunder G, et al;**helical CT angiography compared with angiography in the detection of renal artery stenosis. AJR Am j Roentgenol 167:4%-501, 1996.
- 13. Earls JP, Rofsky NM, DeCorato DR, et al:**Breath-hold single dose Gd-enhanced three-dimensional MR aortography: usefulness of a timing examination and MR power injector.Radio1ogy; 201:705-10. 1996
- 14- Van Jaarsveld BC, Pieterman H, Van Dijk LC ,et al:**alzvan Seijen AJ, Krinjen P Derkx FH, Man intVeld AJ, Schaleltamp MA: Inter-observer variability in the angiographic assessment of renal artery stenosis 17: 1731-1736,1999.

15. **Wasser MN, Wesenberg J, van der Hulst, et al:** Homodynamic significance of renal artery stenosis digital subtraction versus systolically gated through dimensional phase-contrast MR angiography. 1997
16. **Dawson, Westenberg J, Aslmk Ragayan et al:** digital subtraction I Angiography, clinical radiology, 39, 5, 474-477, 1998
17. **Taylor A, Nally J, Au rell M, et al:** Consensus report on ACE inhibitor renography for detecting of renovascular hypertension. Nuclear Med; 37:1876- 82. 1996
18. **Taylor AT, Fletcher IW, Nally JR JV, et al:** Procedure guidelines for diagnosis of renovascular hypertension. J Nucl Med; 39:1297-1302, 1998
19. **Snell Richard S:** clinical anatomy: 8th edition 13-135:453-4155, 2007
20. **Resnick M1, Pounds DM, Boyce WH:** Surgical anatomy of the human kidney and its applications. Urology 17:367, 1981.
21. **Gray H, Pick TPB Howden R:** Gray's Anatomy, New York, Bounty Books 1984, 556-557, 985-994.
22. **Lasts anatomy:** Regional and applied, 8 edition, 70-71, Churrchill Livingstone book company, U.K. 1991.
23. **Sampaio FJ, Aragao AH;** Anatomical relationship between the intrarenal arteries and the kidney collecting system. J Urol, 1413:679, 1990.
24. **Tanagho EA:** Anatomy of the lower urinary tract. In Walsh PC, Retik AB, and STOMEY TA, et al (Eds): Campbell's Urology. Philadelphia, WB Saunders, pp 40-69, 1992.
25. **Wilman AH, Riederer SJ, King BF, Dobbins .JP, Rossman PJ.** Ehman RL: fluoroscopically triggered contrast-enhanced three -dimensional MR angiography with elliptical centric view order: Application to the renal arteries. Radiology 205; 137 146, 1997.
26. **Henry Gray:** Anatomy of the Human Body. 1211-1222, Mosbey puplisher 1996
27. **Walsh, et al:** imaging of the renal arteries: value of MR angiography. AJR AM J Rontgenol 157:981-90. 1985
28. **Ashcraft, Pick TPB Howden R:** Gray's Anatomy, New York, Bounty Books 1984, 556-557, 985-994.
29. **Berkow R, Fletcher AJ:** The Merck Manual, Merck Research Laboratories, 429-430; 1992.
30. **Rubin GD, VVa1ker PJ, Dake MD, et al:** Three-dimensional spiral computed tomographic angiography; an alternative imaging modality for the abdominal aorta and its branches. J Vase Surg 18656-665, 1993, Spiral CT of renal artery stenosis: Comparison of three-dimensional rendering techniques .Radiology 190.181-189, 1994

31. **Safian RD, Textor, or SC.** Renal artery stenosis. *N Engl J Med*: 344:431- 42. 2001.
32. **SOS, TA, Pickering, T.G, Sinderman, and K:** Percutaneous transluminal renal angioplasty in Renal vascular hypertension due to atheroma or fibromuscular dysplasia. *N. Engl.j.Med* .309:274 1983.
33. **Textor SC, Wilcox C** Ischemic nephropathy/azotemic renovascular disease. *Sem Nephrol*, 20:489-502. 2000;
34. **Lee VS, Rofsky NM, Krinsky GA.** Single dose breath-hold gadolinium-enhanced three-dimensional MR angiography of the renal arteries., 211:69-78., *Radiology* 1999 .
35. **Walsh P, Rofsky NM, Krinsky GA.** Weinreb, Asymmetric signal intensity of the renal collecting system as a sign of unilateral renal artery stenosis following administration of gadopentetate dimeglumine. *J. Assist Tomogr*; 20:812-814; 1996.
36. **Anderson JR:** Murray's textbook of pathology. London. Edward Arnold (publisher). 14, 1:14-36. 1995
37. **Desantis RW, Doroghazi RM, Austin WG, et al.:** Aortic dissection, *N Engl.J.Med.*:317:1060.1987
38. **Hall S:** Takayasu arteritis. A study of 32 North American patients. *Medicine*, 1985;64:89.
39. **Schreiber MJ, Pohl MA, Novick AC:** The natural history of atherosclerotic and fibrous renal artery disease. *Urol Clin NAm*, 11; 383-92. 1984.
40. **Fry WJ, Ernest CB, Stanley JC:** Renovascular hypertension in the pediatric patient. *Arch. Surg* 1983;117:692-698. 1983
51. **Taylor Jr AT, Fletcher JW, Nally Jr JV, Blafox MD, Dubovsky EV, Fine EJ, et al.;** Procedure guideline for diagnosis of renovascular hypertension. *Society of Nuclear Medicine Nucl Med*, 39;1297-302 1998
52. **Nada NC:** History of echocardiographic contrast agents *clin cardiol*, 20:17-11; 1997.
53. **Burns physical principles of Doppler and spectral A:** analysis of JCU; 15:567-590; 1995
54. **Olin JW:** Role of duplex ultrasonography in screening for significant renal artery disease. *Urol Clin North Am.*; 21:215-26; 1994.
55. **Barnes RW, Bone GE and Reinterson J:** Non invasive US carotid angiography : prospective validation by contrast angiography. *surgery* 1979;80:382.
56. **Francesco De Cobelli, , Massimo Venturini, , Angelo Vanzulli, , Sandro Sironi, , Marco Salvioni, , Enzo Angeli, ,** Renal Arterial Stenosis: Prospective Comparison of Color Doppler US and Breath-hold, Three-dimensional, Dynamic, Gadolinium-enhanced. MR. Angiography *Radiology*. 214:373-380, 2000.

57. Krebs CA, Giyanni VZ, Eisenberg RL: Ultrasound atlas of 1 J vascular diseases. Appelson and Lange. 1999; 184-187.

58. Hahn U, Miller S, Nagele T, et al. Renal MR angiography at 1.0 T: three-dimensional (3D) phase-contrast techniques versus gadolinium-enhanced 3D fast low-angle shot breath-hold imaging. AJR; 172:1501-1508.1999

59. Duda SH, Schick F, Teufl F, et al. Phase-contrast MR angiography for detection of arteriosclerotic renal artery stenosis. Acta Radiol 1997; 38:287-291.

60. Rosner MH: Renovascularhypertension: can we identify a population at high risk? South Med J, 94:1058-1064; 2001.

61. Nazzari M-MS, Hoballah JJ, Miller EV, et al: Renal hilar – Doppler analysis is of value in the management of patients with renovascular disease. Am J Surg 174:164-168, 1997.

62. Melnyk ML, Grant EG, Duerinckx AJ, Watts TM, Levine BS: Ability of a phase shift US contrast agent to improve imaging of the main renal arteries, Radiology; 205:147-152; 1997.

63. Missouris CG, Allen CM, Balen FG, Buckenham T, Lees WR, MacGregor GA: Non-invasive screening for renal artery stenosis with ultrasound contrast enhancement J Hypertens, 14:519-524;1996

64. Porter TR, Li S, Kilzer K: Smaller intravenous perfluorocarbon-containing micro bubbles produce greater myocardial contrast with intermittent harmonic imaging and better delineation of risk area during acute myocardial ischemia J Am Soc Echocardiogr, 10:792-797; 1997.

65. Helenon O, El Rody F, Correias JM, Melki P, Chauveau D, Chretien Y, Moreau IF: Color Doppler US of renovascular disease in native kidneys. Radiographics, 15:833-54; 1995

66. Souza de Oliveira IR, Widman A, Molnar LJ, JT, Praxedes JN, Cerri GG: Colour Doppler ultrasound: a new index improves the diagnosis of renal artery stenosis Ultrasound Med Biol, 26:41-47; 2000.

67. Claudon M, Plouin PF, Baxter GM, Rohban T, Devos DM: Renal arteries in patients at risk of renal arterial stenosis; multicenter evaluation of the echo-enhancer Renal Artery Study Group Radiology, 214:739-746; 2000.

68. Teixeira, Odila UN, Luiz A Bortolotto and Helio Bernardes Silva. The contrast-enhanced Doppler ultrasound with perfluorocarbon exposed sonicated albumin does not improve the diagnosis of renal artery stenosis compared with angiography. Journal of Negative Results in Biomedicine, 11:861-877; 2004.

69. **Keller PJ, Saloner D, Potchen EJ, Siebert JE, and Haacke E.M:** Time of flight imaging in Magnetic Resonance Angiography, concepts and applications, St. Louis, Mosby ,:146-159. 1993
70. **Schoenberg SO, Bock M, Knopp MV, et al:** Renal arteries: optimization of three-dimensional gadolinium- enhanced MR angiography with bolus-timing-independent last multiphase acquisition in a single breath hold. *Radiology*, 211:667-79. 1999.
71. **Maki JH, Prince MR, Londy FJ, et al:** The effects of time varying intravascular signal intensity and k-space acquisition order on three-dimensional MR angiography image quality. *J Magn Reson Imaging*, 6; 642-51. 1996.
72. **Marks B, Mitchell DG, Simelaro JP.** Breath-holding in healthy and pulmonary—compromised populations: effects of hyperventilation and oxygen inspiration. *JMRI*; 7:595-597; 1997.
73. **Gilfeather M., Yoon HC, Siegelman ES, et al:** Renal artery stenosis: evaluation, With conventional angiography versus gadolinium-enhanced MR angiography. *Radiology*; 210:367-9 372; 1999.
74. **Prince MR, Yucel EK, Kaufman JA, et al:** Dynamic gadolinium-enhanced three-dimensional abdominal MR arteriography. *J Magn Reson Imaging*; 3; 877-81+1998.
75. **Schoenberg SO, Knopp MV, Londy F et al:** Morphologic and functional magnetic resonance imaging of the renal artery stenosis: a multireader tricenter study. *J Am Soc Nephrol*; 13: 158-169; 2002
76. **Schoenberg SO, Knopp MV, Bock M et al:** Renal artery stenosis; grading of hemodynamic changes with cine phase-contrast MR blood flow measurements. *Radiology*; 11:203: 45-53; 1997.
77. **Maki JH, Chenevert TL, Prince MR:** Contrast-enhanced MR angiography. *Abdomen Imaging*; 23:469-484; 1998.
78. **Weiger M, Pruessmann KP, Kassner A et al:** Contrast-enhanced 3D MRA using SENSE. *J Magn Reson Imaging*; 12: 671-677; 2000.
79. **Wedeen VJ, Melni RA, Edelman RR, Kassner A, et al.** Contrast-enhanced 3D MRA using SENSE. *J Magn Reson imaging*; 12:671-7. 2000
80. **Ruehm SG, Goye Barkhausen J et al:** Rapid magnetic resonance angiography for detection of atherosclerosis. *Lancet*; 357: 1086-1091; 2001.
81. **Levy BI, Duriez M, Samuel JL.** Coronary microvasculature alteration in hypertensive rats. Effect of treatment with diuretic and an ACE inhibitor. *Am J Hypertens*; 14: 7-13; 2001.
82. **Dietrich O, Nikolou K, Wintersperger B et al:** Integrated parallel acquisition techniques (iPAT): applications for fast and cardiovascular MR imaging. *Electromedia*; 70: 133-145; 2002

- 83. Schoenberg SO, Bock M, Kallinowski F, Just A.** Correlation of hemodynamic impact and morphologic degree of renal artery stenosis in a canine model. *J Am Soc Nephrol*; 11:2190-2198; 2000.
- 84. Aumann S, Schoenberg SO, Just A et al:** Quantification of renal perfusion using an intravascular contrast agent (part 1): results in a canine model. *Magn Resonance Med* 2003; 49:276–287
- 85. Buckley DL, Shurraf AE, Jones AP, Kilgallon JE, Mamtola H, Kalra PA.** Quantitative assessment of renal function using dynamic Gd-DTPA-enhanced MRI. *Proceeding of the Tenth Scientific Meeting and Exhibition. International Society of Magnetic Resonance in Medicine*: 811; 2002.
- 86. FOX, S. H, Tanenbaum, L. N., Ackelsberg, S., He, H. D., Hsieh, J., Hu, and H.:** Future directions in CT technology. *Neuroimaging clinics of North America*; 8:497-513; 1998.
- 87. Brink JA, Davros WJ, Zeman RK, Costello P (edn):** Helical/Spiral CT. a practical approach. New York, McGrawHill, 1-26; 1995.
- 88. Foo TK, saranathan M, Prince MR, et al** automated detection of bolus arrival and initiation of data acquisition in fast, three-dimensional gadolinium-enhanced MR angiography. *Radio1ogy*; 203:275-80, 1997
- 89. Kaplan-Parloveie S, Nadja C:** Captopril renography and duplex Doppler sonography in the diagnosis of renovascular hypertension. *Nephrol Dial Transplant* 13:313-317, 1998.
- 90. Kopka L, Vosschenrich R, Rodenwaldt J, et al:** Differences in injection rates on contrast-enhanced breath-hold three-dimensional MR angiography. *AJR AmJ Rontgenol* 170:345-8. 1998.
- 91. Kuszyk BS, Fishman EK.** Technical aspects of CT angiography, *Semin Ultrasound MR*; 19:383-393; 1998.
- 92. Chapman S, Nakielny R:** A guide to radiological procedures. Baillire Tindall, 201-241; 1997.
- 93. Uday D. Patil, Ashok Ragavan, Nadaraj,** Keshava M , Ravi Shankar, Bahar Bastani and Sudarshan H. Ballal, PJ; Renal artery stenosis: CT angiography-comparison of real-time volume rendering and intensity projection algorithms *Nephrol Dial Transplant*, 16:1900-1904; 2001
- 94. Prigent A. and Nally JV,** the diagnosis of renovascular hypertension: the role of captopril renal scintigraphy and related issues. *Eur J Nucl Med*; 20:625-644; 1998.
- 95. Aitkison F, Page A:** Diagnostic imaging of renal artery stenosis. *J Hum Hypertension*; 1999;13:595-603.
- 96. Boos M, Seheffler K, Haselhorst R, et al.** Arterial first pass gadolinium-CM dynamics as a function of several intravenous saline flush and Gd volumes. *J Magn Reson Imaging* 13:568-76, 2001.

- 97 . Rimmer J, Gennari FJ.** Atherosclerotic renovascular disease and progressive renal failure. *Ann InternMed*, 118:712-9. 1993.
- 98. Handa N, Komada T.** Efficacy of echo-Doppler examination for the valuation of renovascular disease. *Ultrasound Med Biol* 1988; 14:1-5.
- 99. Patriquin H, Lafortune M, Jéquier JC, et al:** Stenosis of the renal artery: assessment of slowed systole in the downstream circulation with Doppler sonography. *Radiology* 1992; 184:479-485.
- 100. Halpern EJ, Needleman L, Nack TL, East SA.** Renal artery stenosis: should we study the main renal artery or segmental vessels? *Radiology* 1995; 195:799-804
- 101. Prince MR,** Gadolinium-enhanced MR aortography. *Radiology* 1994; 191:155-164.
- 102. Bakker J, Beek FJ, Beutler JJ, et al.** Renal artery stenosis and accessory renal arteries: accuracy of detection and visualization with gadolinium-enhanced breath-hold MR angiography. *Radiology* 1998; 207:497-504
- 103 . Halpern EJ, Mitchell DG, Wechsler RJ, et al:** Preoperative evaluation of living renal donors: comparison of CT angiography and MR angiography. *Radiology* 2000; 216:434-439
- 104. Bruce A. Urban, Lloyd E. Ratner, and Elliot K. Fishman;** Three-dimensional Volume rendered CT Angiography of the Renal Arteries and Veins: Normal Anatomy, Variants, and Clinical Applications I *Radiographics*. 21:373-386; 2001.
- 105. Barrett BJ. And Halpern EJ,** Contrast nephrotoxicity. *J Am s Soc Nephrol*; 5: 125-137; 1994.
- 106. Siegel man ES, Gilgeather M, Holland GA.** Breath-holdultrafast three-dimensional gadolinium enhanced MRangiography of Reno vascular system. *AM Rontgenol*; 168·p 1035-1040; 1997.
- 107. Gross CM, Kramer J, Weingartner O. et al:** Determination of renal arterial stenosis severity: comparison of pressure gradient and vessel diameter. *Radiology*. 2001; 220: 751–64].
- 108. Tegtmeier CJ, Peter B. Sachs** Renal Angioplasty, In: *Abram's Angiography*, Little Brown, RA: 294-325; 1999.
- 109. Mittai BR, Kumar P, Arora P, et al:** Role of captopril renography in the diagnosis of renovascular hypertension. *Am J Kidney Dis* 1996; 28:209-213
- 110. Prigent A.** The diagnosis of renovascular hypertension: the role of captopril renal scintigraphy and related issues. *Eur J Nucl Med* 1993; 20:625-644.
- 111. Harding MB, Smith LR, Hammerstein SI, et al:** Renal artery stenosis; prevalence and associated risk factors in patients undergoing routine cardiac catheterization.

- 112. Carman T, Olin JW, Czum J** Noninvasive imaging of renal arteries. *Urol Clin N Am*: 28:815-26. 2001
- 113. Platt JF;** Doppler ultrasound of the kidney. *Semin Ultrasound CT MR* 18:22-32, 1997.
- 114. Malatino LS, Polizzi G, Garozzo M, et al:** Diagnosis of renovascular disease by extra- and intrarenal Doppler parameters. *Angiology* 49:707-721, 1998.
- 115. Isaacson J, Neumyer MM:** Direct and indirect renal arterial duplex and Doppler color flow evaluations. *Journal of Vascular Technology* 19(5-6): 309-316, 1995
- 116. Radermacher J, Chavan A, Bleek J, et al:** Use of Doppler ultrasonography to predict the outcome of therapy for renal artery stenosis. *N Engl J Med* 344:410-7. 2001.
- 117. Miralles M, Santiso A, Gimenez A, et al:** Renal duplex scanning; Correlation with angiography and isotopic renography., *Eur J Vasc Surg* 7:188-194, 1993.
- 118. Helenon O, Meki P, Correias J-M, Boyer J-c, Moreau J-F** Renovascular diseases: Doppler ultrasound. 18:136-146, 1997.
- 119. Leung DA, Hoffman U, Pfammatter T, et al:** Magnetic resonance angiography versus duplex sonography for diagnosing renovascular disease. *Hypertension* 3:726-731, 1999.
- 120 . Pedersen EB:** New tools in diagnosing renal artery stenosis. *Kidney Int* 57:2657-2677, 2000 .
- 121. Kaufman JA, McCarter D, Geller SC, et al:** Two- dimensional time-flight MR angiography of the lower extremities: artifacts and pitfalls. *AJR Am J Roentgenol*: 171:129-35. 1998.
- 122. Saloner D:** Determinants of image appearance in contrast- enhanced magnetic resonance angiography; A review *Invest Radiol* 33:4188-4195, 1998
- 123. Hahn U, Miller S, Ngele T, et al;** Renal MR angiography at 1.0 T: Three-dimensional (3D) phase- contrast techniques versus gadolinium-enhanced 3D fast low angle .et.1 short breath-hold imaging. *AJR Am J Roentgenol* 172: 11508, 1999,
- 124. Joseph P. Hornak, Ph.D.** Copyright J.P. Hornak .AllRights Reserved. 1996-2002.
- 125. Paul JF, Cherrak I, Jaulent MC,** Chatellier G, Plouin PF, Degoulet P, Gaux JC: Interobserver variability in the interpretation of renal digital subtraction angiography. *Am J Roentgenol* 173: 1285-1288, 1999 .
- 126. Teresa L. Carman MD Jeffrey Olin DO Julianna Czum MD** *Urologic Clinics of North America* Volume 28 Number 4 Department of Vascular Medicine. 2001.