

References

1. Arabshahi B, Cron RQ (2006) Temporomandibular joint arthritis in juvenile idiopathic arthritis: the forgotten joint. *Curr Opin Rheumatol* 18:490–495} .
2. Adame CG, Mange F, Munoz M, Granizo RM (1998): Effusion in magnetic resonance imaging of the temporomandibular joint. A study of 123 joints. *J. Oral Maxillofac Surg* ; 56: 314.
3. Arnett GW, Milam SB, Gottesman L (1996a,b). Progressive mandibular retrusion—idiopathic condylar resorption. Part I. *Am J Orthod Dentofacial Orthop* 110:8–15.
4. Arslan A, Orhan K, Paksoy CS, Uçok O, Ozbek M, Dura SI, et al. 2009 . MRI evaluation of the classification, frequency, and disc morphology of temporomandibular joint disc displacements: a multicenter retrospective study in a Turkish population. *Oral Radiol.*;1:14–21.
5. Barclay P, Hollender LG, Maravilla KR, Truelove EL (1999): Comparison of clinical and magnetic resonance imaging diagnoses in patients with disk displacement in the temporomandibular joint. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*; 88:37.
6. Beatty MW, Bruno MJ, Iwasaki LR, Nickel JC (2001). Strain rate dependent orthotropic properties of pristine and impulsively loaded porcine temporomandibular joint disk. *J Biomed Mater Res* 57:25–34.
7. Beatty MW, Nickel JC, Iwasaki LR, Leiker M (2003). Mechanical response of the porcine temporomandibular joint disc to an impact event and repeated tensile loading. *J Orofac Pain* 17:160–166.
8. Beek M, Aarnts MP, Koolstra JH, Feilzer AJ, van Eijden TM. (2001) Dynamic properties of the human temporomandibular joint disc. *Journal of Dental Research* 80:876–880.
9. Bell WE 1990: Temporomandibular joint disorders. Classification, diagnosis and management. 3rd edn. Chicago: Yearbook Medical publications; 145–157.

10. Bergman AA, Heidger PM. 1996 . Histology. Iowa City, IA: W.B. Saunders Company .
11. Berquist T (2000). MRI of the Musculoskeletal System. 4th ed. Baltimore, MD: Lippincott Williams & Wilkins.
12. Bernhardt O, Biffar R , Kocher T , Meyer G (2007): Prevalence and clinical signs of degenerative temporomandibular joint changes validated by magnetic resonance imaging in a non-patient group. *Annals of anatomy*.
13. Bertram S, Rudisch A, Innerhofer K, Pümpel E, Grubwieser G, Emshoff R. 2001 . Diagnosing TMJ internal derangement and osteoarthritis with magnetic resonance imaging. *JADA*;132:753–61.
14. Bui-Mansfield LT, Rohini D, Bagg M. Tenosynovial chondromatosis of the ring finger. *AJR Am J Roentgenol* 2005;184:1223–1224, with the greatest number of cases occurring in patients in the fifth decade of life (mean age, 46 years) (19,20,36).
15. Butt SH, Muthukumar T, Cassar-Pullicino VN, Mangham DC. Primary synovial osteochondromatosis presenting as constrictive capsulitis. *Skeletal Radiol* 2005;34:707–713. 4- Benjamin M, Ralphs JR. Biology of fibrocartilage cells. *Int Rev Cytol* 2004; 233:1–45.
16. Carvalho RS, Yen EH, Suga DM (1995): Glycosaminoglycan synthesis in the rat articular disk in response to mechanical stress. *Am J Orthod Dentofacial Orthop*; 107:401-410.
17. Chossegros C, Cheynet F, Guyot L, Bellot-Samson V, Blanc JL (2001): Posterior disk displacement of the TMJ: MRI evidence in two cases. *Cranio*;19: 289–293.
18. Cupp JS, Miller MA, Montgomery KD, et al. (2007) Translocation and expression of CSF1 in pigmented villonodular synovitis, tenosynovial giant cell tumor, rheumatoid arthritis and other reactive synovitides. *Am J Surg Pathol* 31:970–976.

19. Dalkiz M, Pakdemürlü E, Beydemür B (2001): Evaluation of Temporomandibular Joint Dysfunction by Magnetic Resonance Imaging. *Turk J Med Sci*; 31: 337-343.
20. Da Silva AF, Shaefer J, Keith DA. 2003 The temporomandibular joint: clinical and surgical aspects. *Neuroimaging Clin N Am*;13:573–582.
21. Davis RI, Hamilton A, Biggart JD. 1998 . Primary synovial chondromatosis: a clinicopathologic review and assessment of malignant potential. *Hum Pathol*;29:683–688.
22. Del Pozo R, Tanaka E, Tanaka M, et al (2002): The regional difference of viscoelastic property of bovine temporomandibular joint disc in compressive stress-relaxation. *Med Eng Phys*; 24:165-71.
23. De Leeuw R. 2008 Internal derangements of the temporomandibular joint. *Oral Maxillofac Surg Clin North Am*;20:159–68.}.
24. Debrul E.L. (1980): Sicher's Oral anatomy, St.Louis, Masby; 174-20.
25. Dimitroulis G (2005): The prevalence of osteoarthritis in cases of advanced internal derangement of the temporomandibular joint: a clinical, surgical and histological study. *Int J Oral Maxillofac Surg*; 34:345-349.
26. Drace J.E., and Enzman D.R. (1990): Defining the normal TMJ, closed, partially open and open mouth MR imaging of asymptomatic subjects. *Radiology*; 177:67-71.
27. Drace J.E., Young S., Enzman D. (1990) : TMJ meniscus and bilaminar zone MR imaging of substructure-diagnostic land marks and pitfalls of interpretation. *Radiology*; 177:73-76.
28. Drake R, Vogel W, Mitchell A(2005):GRAY'S Anatomy for students. Philadelphia; PP 847-875.
29. Eberhard D, Bantleon HP, Steger W (2002): The efficacy of anterior repositioning splint therapy studied by magnetic resonance imaging. *Eur J Orthod*; 24:343-352.

30. Edelman R and Hesselink J (1990) : Clinical Magnetic Resonance Imaging, Philadelphia, WB Saunders; 250-266.
31. Ekberg E, Vallon D, Nilner M (2002): Treatment outcome of headache after occlusal appliance therapy in a randomised controlled trial among patients with temporomandibular disorders of mainly arthrogenous origin. *Swed Dent J*; 26:115-124.
32. Emshoff R, Brandlmaier I, Schmid C, Bertram S, Rudisch A (2003): Bone marrow edema of the mandibular condyle related to internal derangement, osteoarthritis, and joint effusion. *J Oral Maxillofac Surg* ; 61: 35–40.
33. Emshoff R, Gerhard S, Ennemoser T, et al (2006): Magnetic resonance imaging findings of internal derangement, osteoarthritis, effusion, and bone marrow edema before and after performance of arthrocentesis and hydraulic distension of the temporomandibular joint. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*; 101:784-790.
34. Fabian FM, Mumghamba EG. 2008 Risk factors for signs and symptoms of TMD in a rural southeast Tanzanian population *Cranio*.;26:44–9 .
35. Fanghanel J, Gedrange T (2007) On the development, morphology and function of the temporomandibular joint in the light of the orofacial system. *Ann Anat* 189:314–319 .
36. Fetsch JF, Vinh TN, Remotti F, Walker EA, Murphey MD, Sweet DE. 2003 Tenosynovial (extraarticular) chondromatosis: an analysis of 37 cases of an underrecognized clinicopathologic entity with a strong predilection for the hands and feet and a high local recurrence rate. *Am J Surg Pathol*; 27:1260–1268 .
37. Forster H, Fisher J (1996): The influence of loading time and lubricant on the friction of articular cartilage. *Proc Inst Mech Eng [H]*; 210:109-119.
38. Gameiro GH, da Silva Andrade A, Nouer DF, Ferraz de Arruda Veiga MC. 2006 . How may stressful experiences contribute to the development of temporomandibular disorders? *Clin Oral Investig*.;10:261–8.}.

39. Gerhard S, Ennemoser T, Rudisch A, Emshoff R (2007) : Condylar injury: magnetic resonance imaging findings of temporomandibular joint soft-tissue changes. *Int. J. Oral Maxillofac. Surg.*; 36: 214–218.
40. Glaser C, Horng A, Mendilik T, Weckbach S, Hoffmann RT, Wagner S, Raya JG, Horger W, Reiser M (2007) T2 relaxation time in patellar cartilage: global and regional reproducibility at 1.5 Tesla and 3 Tesla. *Rofo* 179:146– 152 .
41. Greene, C. (2006). Concepts of TMD Etiology: Effects on Diagnosis and treatment. *TMDs An Evidence- Based Approach to Diagnosis and Treatment*. G. C. Laskin DM, and Hylander WL Hanover Park, Quintessence: 35- 52.
42. Greess H, Anders K. 2005 . Indications for validity of computed tomography and magnetic resonance imaging of the temporomandibular joint]. *Rontgenpraxis*; 56:1–11.
43. Haley, D.P., Schiffman, E.L., Lindgren, B.R., Anderson, Q., Andreasen, K., 2005 . The relationship between clinical and MRI findings in patients with unilateral temporomandibular joint pain. *J. Am. Dent. Assoc.* 132, 476–481.
44. Hayt MW, Abrahams JJ, Blair J (2000): Magnetic resonance imaging of the temporomandibular joint. *Top Magn Reson Imaging*; 11:138-146.
45. Helenius LMJ, Hietanen J, Helenius I, Kautiainen H, Piirainen H, Paimela L, Lappalainen M, Leirisalo-Repo M (2001): Focal sialadenitis in patients with ankylosing spondylitis and spondyloarthropathy: a comparison with patients with rheumatoid arthritis or mixed connective tissue disease. *Ann Rheum Dis.*; 60: 744–749.
46. Helenius L. M. J., Tervahartiala P., Helenius I., Al-Sukhun J., Kivisaari L., Suuronen R., Kautiainen H., Hallikainen D., Lindqvist C., Leirisalo-Repo M. (2006): Clinical, radiographic and MRI findings of the temporomandibular joint in patients with different rheumatic diseases. *Int. J. Oral Maxillofac. Surg.*; 35: 983–989.

47. Helms C.A., (1990): The TMJ in MRI of the musculoskeletal system, 2nd edition, edited by T.H.Berquist, Raven press Ltd. New York; chapter 4: 75-89.
48. Helms C.A., (1997): MRI of the body, 3rd edition, Lippincott-raven press New York, chapter 47: 1257-1266.
49. Heylings DJ, Nielsen IL, McNeill C (1995): Lateral pterygoid muscle and the temporomandibular disc. J Orofac Pain; 9: 9–16.
50. Hiraba K, Hibino K, Hiranuma K, Negoro T (2000). EMG activities of two heads of the human lateral pterygoid muscle in relation to mandibular condylar movement and biting force. J Neurophysiol 83:2120–2137.
51. Honda K, Bjørnland T. (2006) , Image-guided puncture technique for the superior temporomandibular joint space: value of cone beam computed tomography (CBCT). Oral Surg Oral Med Oral Pathol Oral Radiol Endod. ;102:281–6.
52. Hopyan S, Nadesan P, Yu C, Wunder J, Alman BA. (2005) Dysregulation of hedgehog signalling predisposes to synovial chondromatosis. J Pathol ; 206:143–150 .
53. Jensen U, Ruf S. (2007) Longitudinal changes in temporomandibular disorders in young adults: indication for systematic temporomandibular joint screening. J Orofac Orthop. ;68:501–9.
54. Kaneyama K, Segami N, Shin-Ichi T, Fujimura K, Sato J, Nagao T (2006): Anchored disc phenomenon with a normally positioned disc in the temporomandibular joint: Characteristics and behaviour. British Journal of Oral and Maxillofacial Surgery; 45 :279–283.
55. Kaneyama K, Segami N, Shin-Ichi T, Fujimura K, Sato J, Nagao T (2007): Anchored disc phenomenon with a normally positioned disc in the temporomandibular joint: Characteristics and behaviour. British Journal of Oral and Maxillofacial Surgery; 45: 279–283.
56. Katezberg R.W., Westesson P.L (1994): Normal anatomy. Diagnosis of the TMJ. Philadelphia : W.B. Sanders; 149-158.

57. Katzberg RW, Westesson PL, Tallents RH (1996): Anatomic disorders of the temporomandibular joint disc in asymptomatic subjects. *J Oral Maxillofac Surg* 54:147-153.
58. Kirc WS (1994): Sagittal Magnetic Resonance Image Characteristics and Surgical Findings of Mandibular Condyle Surface Disease In Staged Internal Derangement. *J Oral Maxillofacial Surg*; 52: 64.
59. Kircos L.T., Ortendahl, Mark L.s (1987): Magnetic resonance imaging of the TMJ disc in asymptomatic volunteer. *J Oral Maxillofac surg*; 45:852-854.
60. Klenoff JR, Lowlicht RA, Lesnik T, et al (2001): Mandibular and temporomandibular joint arthropathy in the differential diagnosis of the parotid mass. *Laryngoscope*; 111(12):2162-2165.
61. Kramer J, Recht M, Deely DM, et al. (1993) MR appearance of idiopathic synovial osteochondromatosis. *J Comput Assist Tomogr*;17:772–776.
62. Kransdorf MJ, Murphey MD. (2006) Synovial tumors. In: *Imaging of soft tissue tumors*. 2nd ed. Philadelphia, Pa: Lippincott Williams & Wilkins,; 412–436 .
63. Kurita H, Uehara S, Yokochi M (2006): A long-term follow-up study of radiographically evident degenerative changes in the temporomandibular joint with different conditions of disk displacement. *Int J Oral Maxillofac Surg*; 35(1):49-54.
64. Kursoglu P, Capa N. (2006) Elongated mandibular coronoid process as a cause of mandibular hypomobility. *Cranio*.;24:213–6.
65. Lafreniere CM, Lamontagne M, el-Sawy R (1997): The role of the lateral pterygoid muscles in TMJ disorders during static conditions. *Cranio*; 15:38–52.
66. Larheim TA, Westesson P-L, Hicks DG, et al (1999): Osteonecrosis of the temporomandibular joint: correlation of magnetic resonance imaging and histology. *J Oral Maxillofac Surg*; 57:888-898.

67. Larheim A, Katzberg W, Westesson L (2001): MR evidence of temporomandibular joint fluid and condyle marrow alterations: occurrence in asymptomatic volunteers and symptomatic patients. *Int J Oral Maxillofac Surg*; 30:113-117.
68. Larheim TA, Westesson PL, Sano T (2001): Temporomandibular joint disk displacement: comparison in asymptomatic volunteers and patients. *Radiology*; 218:428-432.
69. Larheim TA (2005): Role of magnetic resonance imaging in the clinical diagnosis of the temporomandibular joint. *Cells Tissues Organs*; 180(1):6- 21.
70. Laskin DM(1993). Diagnosis of pathology of the temporomandibular joint: Clinical and imaging perspectives. *Radiol Clin North Am.*; 31:135-147.
71. Laskin DM, Greene CS, Hylander WL(2006): Temporomandibular Disorders: An Evidence-Based Approach to diagnosis and treatment. Hanover Park, IL, Quintessence Publishing Co., Inc.
72. Levange K and Norkin C (1992): Joint structure and function. F.A. Davis company, Philadelphia.
73. Lieberman M, Hans G, Rozenzweig G(1992): MR imaging of the juvenile temporomandibular joint: preliminary report. *Radiology*; 182: 531-534.
74. Limchaichana N, Nilsson H, Ekberg EC, Nilner M, Petersson A (2007) Clinical diagnoses and MRI findings in patients with TMD pain. *J Oral Rehabil* 34:237–245 .
75. Liu ZJ, Yamagata K, Kuroe K, Suenaga S, Noikura T, Ito G (2000): Morphological and positional assessments of TMJ components and lateral pterygoid muscle in relation to symptoms and occlusion of patients with temporomandibular disorders. *J Oral Rehabil*; 27:860–874.
76. Major P, Ramos-Remus C, Suarez-Almazor ME, Hatcher D, Parfitt M, Russell AS (1999): Magnetic resonance imaging and clinical assessment of temporomandibular joint pathology in ankylosing spondylitis. *J Rheumatol*; 26: 616–621.

77. Manzione JV (1993): Head and neck radiology: RSNA meeting notes [abstr]. Radiology; 186(P):607.
78. Marguelles-Bonnet RE, Carpentier P, Yung JP, Defrennes D, Pharaboz C (1995): Clinical diagnosis compared with findings of magnetic resonance imaging in 242 patients with internal derangement of the TMJ. J Orofac Pain; 9:244–53.
79. Martini G, Bacciliero U, Tregnaghi A, Montesco MC, Zulian F. (2001) Isolated temporomandibular synovitis as unique presentation of juvenile idiopathic arthritis. J Rheumatol 28:1689–92.
80. Meikle MC. (2007) Remodeling the dentofacial skeleton: the biological basis of orthodontics and dentofacial orthopedics. J Dent Res; 86:12–24.
81. Mendenhall WM, Mendenhall CM, Reith JD, et al. (2006) Pigmented villonodular synovitis. Am J Clin Oncol . 29:548–550.
82. Milam B, Zardeneta G, Schmitz P (1998): Oxidative stress and degenerative temporomandibular joint disease: a proposed hypothesis. J Oral Maxillofac Surg; 56:214-223.
83. Milam SB. 2005 Pathogenesis of degenerative temporomandibular joint arthritides. Odontology; 93:7–15.
84. Minarelli AM, Del Santo M Jr, Liberti EA (1997): The structure of the human temporomandibular joint disc: a scanning electron microscopy study. J Orofac Pain; 11:95-100.
85. Mizoguchi I, Nakamura M, Takahashi I, Kagayama M, Mitani H. 1992 . A comparison of the immunohistochemical localization of type I and type II collagens in craniofacial cartilages of the rat. Acta Anat (Basel); 144:59–64.
86. Moffet B. (1968): The TMJ. In Sharry J.J. Complete Denture Prothodontics New York, Mc Grw. Hill; pp 56-104.
87. Molinari F ., Francesco P., Raffaelli L., Raffaelli R ., Pirroni T., Bonomo L (2007): Temporomandibular Joint Soft-Tissue Pathology, I: Disc Abnormalities. Semin Ultrasound CT MRI. Radiographics ; 28:192-204.

88. Murphy WA, Kaplan PA (1995): Temporomandibular joint. In: Resnick D, eds. *Diagnosis of bone and joint disorders*. Saunders; 1699-1754 .
89. Murray GM, Phanachet I, Uchida S, Whittle T (2004): The human lateral pterygoid muscle: a review of some experimental aspects and possible clinical relevance. *Aust Dent J*; 49:2–8.
90. Naidoo LC (1996): Lateral pterygoid muscle and its relationship to the meniscus of the temporomandibular joint. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*; 82:4–9.
91. Nitzan DW (2001): The process of lubrication impairment and its involvement in temporomandibular joint disc displacement: a theoretical concept. *J Oral Maxillofac Surg*; 59:36-45.
92. Ogura I. Magnetic resonance (2006) imaging characteristics of temporomandibular joint pain during opening and biting in patients with disc displacement. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*;102:669–72 .
93. Olin HB, Pedersen K, Francis D, et al (2001): A very rare benign tumour in the parotid region: calcium pyrophosphate dihydrate crystal deposition disease. *J Laryngol Otol*; 115(6):504-506.
94. Oliver J. S, Aigner F, Rudisch A, Gruber H, Fritsch H, Millesi W and Stiskal M (2003) : Cross-sectional and Functional Imaging of the Temporomandibular Joint: Radiology, Pathology, and Basic Biomechanics of the Jaw . *Radiographics*; 23:e14.
95. Orhan K, Nishiyama H, Tadashi S, Murakami S, Furukawa S. 2006 . Comparison of altered signal intensity, position, and morphology of the TMJ disc in MR images corrected for variations in surface coil sensitivity. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.*;101:515–22.
96. Pertes R.A., and Gross S.G. (1995): *Clinical management of TMJ disorders and orofacial pain*. Quintessence publishing Co, Inc. Chicago, Berlin, London, Tokyo, San Paulo, Moscow, Prague and Warsaw.

97. Phoebe A, Clyde A, Dussault R , Mark W and Nancy M (2001): Musculoskeletal MRI. Temporomandibular joint. Harcourt Health Sciences Company. Philadelphia. London. New York. St. Louis. Sydney. Toronto; 169-171.
98. Quartier P, Prieur AM (2007) [Juvenile idiopathic arthritis. (I) Clinical aspects] Rev Prat 57:1171–1178 } .
99. Rao VM (1995): Imaging of the temporomandibular joint. Semin Ultrasound CT MR; 16:513–526.
100. Raphael, K., Marbach, J.J. (1996) : Management of temporomandibular disorders. Journal of the American Dental Association; v127 n11 p1595.
101. Resnick D. (2002) Tumors and tumor-like lesions of soft tissues. In: Diagnosis of bone and joint disorders. 4th ed. Philadelphia, Pa: Saunders,; 4204–4273 .
102. Roth C ., Robert J ., Tsai S ., Zolotor W ., Tello R ., (2005): MR imaging of the TMJ: A pictorial essay. www.appliedradiology.com.
103. Rosa EL, Oleskovicz CF, Aragaõ BN. 2004 . Rapid prototyping in maxillofacial surgery and traumatology. Braz Dent J.; 15:243–7 .
104. Rudisch A, Innerhofer K, Bertram S, Emschhoff R (2001): Magnetic resonance imaging findings of internal derangement and effusion in patients with unilateral temporomandibular joint pain. Oral Surg Oral Med Oral Pathol Oral Radiol Endod; 92: 566–571.
105. Rutkiewicz T, Koˆnoˆnen M, Suominen-Taipale L, Nordblad A, Alanen P. (2006) Occurrence of clinical signs of temporomandibular disorders in adult Finns. J Orofac Pain.;20:208–17.
106. Sano T, Yamamoto M, Okano T (2003): Temporomandibular joint: MR imaging. Neuroimaging Clin N Am; 13:583-595.
107. Sano T, Westesson L, Yamamoto M (2004): Differences in temporomandibular joint pain and age distribution between marrow

edema and osteonecrosis in the mandibular condyle. *Cranio*; 22(4):283-288.

108. Sano T, Otonari-Yamamoto M, Otonari T, Yajima A (2007): Osseous Abnormalities Related to the Temporomandibular Joint. *Semin Ultrasound CT MRI* ; 28:213-221.
109. Saupe N, Pfirrmann CWA, Schmid MR, Schartler T, Manestar W, Weishaupt D (2007) MR imaging of cartilage in cadaveric wrists: comparison between imaging at 1.5 and 3.0 T and gross pathologic inspection. *Radiology* 243:180–187
110. Scapino R (1999): Morphology and mechanism of the jaw joint, in McNeill C (ed). *Science and Practice of Occlusion*. Chicago, IL, Quintessence Publishing Co., Inc.; pp 23-40.
111. Schneider A, Zahnert D, Klengel S, Loukota R and Eckelt U (2007): A comparison of MRI, radiographic and clinical findings of the position of the TMJ articular disc following open treatment of condylar neck fractures. *British Journal of Oral and Maxillofacial Surgery*; 74, D-01307.
112. Scolozzi P, Bosson G, Jaques B. (2005) Severe isolated temporomandibular joint involvement in juvenile idiopathic arthritis. *J Oral Maxillofac Surg* 63:1368–71.
113. Sezgin, O. S., P. Celenk, et al. (2007). "Mandibular asymmetry in different occlusion patterns." *Angle Orthod* 77(5): 803-7.)
114. Shahidi S, Haghnegahdar A, Falamaki FN, Khojastehpoor L. 2008 . Clinical evaluation of internal joint derangement using sonography. *Oral Radiol.*;24:34–8.
115. Shen G, Darendeliler MA. 2005 . The adaptive remodeling of condylar cartilage: a transition from chondrogenesis to osteogenesis. *J Dent Res*; 84:691–9.
116. Shibukawa Y, Young B, Wu C, Yamada S, Long F, Pacifici M, et al. 2007 . Temporomandibular joint formation and condyle growth require Indian hedgehog signaling. *Dev Dyn*; 236:426–34.

117. Smith HJ, Larheim TA, Aspestrand F (1992): Rheumatic and non rheumatic disease in the temporomandibular joint: gadolinium-enhanced MR imaging. *Radiology*; 185:229-234.
118. Sommer OJ, Aigner F, Rudisch A, et al (2003). Cross-sectional and functional imaging of the temporomandibular joint: Radiology, pathology, and basic biomechanics of the jaw. *RadioGraphics*.; 23:e14.
119. Stegenga B (2001): Osteoarthritis of the temporomandibular joint organ and its relationship to disc displacement. *J Orofac Pain* ;15:193-205.
120. Stehling C, Vieth V, Bachmann R, Nassenstein I, Kugel H, Kooijman H, Heindel W, Fischbach R (2007) Highresolution magnetic resonance imaging of the temporomandibular joint: image quality at 1.5 and 3.0 Tesla in volunteers. *Invest Radiol* 42:428–434)
121. Steven E., & Harms (1990): Clinical magnetic resonance imaging Edelman and Hesselink.; 1031:1056.
122. Styles C, Whyte A (2002): MRI in the assessment of internal derangement and pain within the temporomandibular joint: a pictorial essay. *Br J Oral Maxillofac Surg*.; 40: 220–228.
123. Suenaga S, Sonoda S, Oku T, Abeyama K, Noikura T. (1997) MRI of the temporomandibular joint disk and posterior disk attachment before and after nonsurgical treatment. *J Comput Assist Tomogr*;21:892–896.
124. Suenaga S, Abeyama K, Noikura T (1998): Gadolinium-Enhanced MR Imaging of Temporomandibular Disorders: Improved Lesion Detection of the Posterior Disk Attachment on TI-Weighted Images Obtained with Fat Suppression. *AJR*; 171:511-517.
125. Suenaga S, Ogura T, Matsuda T, Noikura T (2000): Severity of synovium and bone marrow abnormalities of the temporomandibular joint in early rheumatoid arthritis: Role of gadolinium-enhanced fat-suppressed T1-weighted spin echo MRI. *Comput Assist Tomogr*.; 24:461-465.

126. Svensson B, Adell R, Kopp S. (2000) . Temporomandibular disorders in juvenile chronic arthritis patients. A clinical study. *Swed Dent J* 24:83–92.
127. Takahashi T, Sasano T, Miyamoto Y (2006) Evaluation of the position, mobility and morphology of the disc by MRI before and after four different treatments for temporomandibular joint disorders. *Dentomaxillofac Radiol* 35:103–109 .
128. Tallents H, Macher J, Kyrkanides S, Katzberg W, Moss E (2002): Prevalence of missing posterior teeth and intraarticular temporomandibular disorders. *J Prosthet Dent*; 87: 45–50.
129. Tanaka E, Shibaguchi T, Tanaka M, et al (2000): Viscoelastic properties of the human temporomandibular joint disc in patients with internal derangement. *J Oral Maxillofac Surg*; 58:997-1002.
130. Tanaka E, Sasaki A, Tahmina K, et al (2001): Mechanical properties of human articular disk and its influence on TMJ loading studied with the finite element method. *J Oral Rehabil*; 28:273-279.
131. Tanaka E, van Eijden T (2003) : Biomechanical behavior of the temporomandibular joint disc. *Crit Rev Oral Biol Med*; 14:138-150.
132. Tasaki MM, Westesson PL (1993): Temporomandibular joint: diagnostic accuracy with sagittal and coronal MR imaging. *Radiology*; 189: 823–827.
133. Taskaya-Yilmaz N, Ceylan G, Incesu L(2005): A possible etiology of the internal derangement of the temporomandibular joint based on the MRI observations of the lateral pterygoid muscle. *Surg Radiol Anat*; 27:19-24.
134. Tins B, Cassar-Pullicino V. 2003 . Synovial osteochondromatosis in hereditary arthro-ophthalmopathy (Wagner-Stickler syndrome). *Skeletal Radiol*;32:302–305.
135. Tomas X. (1999) Estudio por resonancia magnética, mediante secuencias GE T2 (flash 2D) y SE T1, de la detección de patología disfuncional a nivel de la articulación

temporomandibular [doctoral thesis]. Barcelona, Spain: University of Barcelona,

136. Tomas X (2000): Estudio por resonancia magnética, mediante secuencias GE T2 (flash 2D) y SE T1, de la detección de patología disfuncional a nivel de la articulación temporomandibular. Doctoral Thesis, University of Barcelona; Code 3790.
137. Tomas X, Pomes J (2004): Articulación temporomandibular. In: Mercader JM, Viñuela F, eds. Neuror-radiología diagnóstica y terapéutica. Barcelona, Spain: Masson; 403–408 .
138. Tomas X., Pomes J., Berenguer J., et al. (2006) : MR Imaging of Temporomandibular Joint Dysfunction: A Pictorial Review. *RadioGraphics* 2006;26:765-781
139. Tomas X ., Pomes J., Berenguer J., Maria J, Pons F ., Donoso L (2007) : Temporomandibular Joint Soft-Tissue Pathology, II: Nondisc Abnormalities. *Semin Ultrasound CT MRI . Radiographics*; 28:205-212.
140. Toyama M, Kurita K, Rivera G (2000): Magnetic resonance arthrography of the temporomandibular joint. *J Oral Maxillofac Surg*; 58:978-983;
141. Trattnig S, Pinker K, Ba-Ssalamah A, Nöbauer-Huhmann IM (2006) The optimal contrast agents at high field MRI . *Eur Radiol* 16:1280–1287.
142. Twilt M, Mobers SM, Arends LR, ten Cate R, van Suijlekom-Smit L. (2004) . Temporomandibular involvement in juvenile idiopathic arthritis. *J Rheumatol* 31:1418–22.
143. Twilt M, Schulten AJ, Verschure F, Wisse L, Prahl-Andersen B, van Suijlekom-Smit LW (2008) Long-term followup of temporomandibular joint involvement in juvenile idiopathic arthritis. *Arthritis Rheum* 59:546–552 }
144. Tymofiyeva O, Proff P, Richter E, Jakob P, Nel J, Gedrange T, Rottner K (2007) : Correlation of MRT imaging with real-time axiography of TMJ clicks. Greifswald, Rotgerberstraße Greifswald, Germany; 8, D-17487.

145. Unni KK, Inwards CY, Bridge JA, Kindblom LG, Wold LE. 2005 . Synovial tumors. In: Tumors of the bone and joints. 4th ed. Silver Spring, Md: ARP Press;; 386–432)
146. USUMEZ S, OZ F and GURAY E (2004): Comparison of clinical and magnetic resonance imaging diagnoses in patients with TMD history. *Journal of Oral Rehabilitation*; 31: 52–56.
147. Vargas A, Teruel J, Trull J, et al (1997): Calcium pyrophosphate dehydrate crystal deposition disease presenting as a pseudotumor of the temporomandibular joint. *Eur Radiol*; 7(9):1452-1453.
148. Vogl TJ, Abolmaali N (2001): MRI of the temporomandibular joint: technique, results, indications. *Rofo Fortschr Geb Roentgenstr Neuen Bildgeb Verfahren*; 173:969-979.
149. Watabe T, Adachi M, Watabe T, Heshiki A, Mizuno H, Mizuno H (1995): MR imaging of temporomandibular joint (TMJ): mandibular fracture and traumatic disk injury. *Nippon Igaku Hoshasen Gakkai Zasshi*; 55:456–459.
150. Weiss PF, Arabshahi B, Johnson A et al (2008) High prevalence of temporomandibular joint arthritis at disease onset in children with juvenile idiopathic arthritis, as detected by magnetic resonance imaging but not by ultrasound. *Arthritis Rheum* 58:1189–1196 .
151. Westesson P.L., Tasaki M. (1993): TMJ Diagnostic Accuracy with sagittal and coronal MR imaging. *Radiology*; 186:723-729.
152. Westesson PL, Larheim TA, Tanaka H (1998): Posterior disc displacement in the temporomandibular joint. *J Oral Maxillofac. Surg.*; 56:1266-1273; discussion 1273-1274.
153. Whyte M, McNamara D, Rosenberg I, Whyte A (2006): Magnetic resonance imaging in the evaluation of temporomandibular joint disc displacement—a review of 144 cases. *Int. J. Oral Maxillofac. Surg.*; 35: 696–703.

154. William and Wilkin (1995): Imaging atlas of the TMJ. Baltimore. Philadelphia. Hong Kong. London. Munich. Sydney. Tokyo Awavery Co.
155. Wongwatana S, Kronman JH, Clark RE, Kabani S, Mehta N (1994): Anatomic basis for disk displacement in temporomandibular joint (TMJ) dysfunction. Am J Orthod Dentofacial Orthop; 105:257–264.
156. Yang X, Pemu H, Pyhtinen J (2001): MRI findings concerning the lateral pterygoid muscle in patients with symptomatic TMJ hypermobility. Cranio; 19:260-268.
157. Yang X, Pernu H, Pyhtinen J, Tiilikainen PA, Oikarinen KS, Raustia AM (2002): MR abnormalities of the lateral pterygoid muscle in patients with non-reducing disk displacement of the TMJ. Cranio; 20:209–221.
158. Yang C, Zhang SY, Wang XD, Fan XD. 2005. Magnetic resonance arthrography applied to the diagnosis of intraarticular adhesions of the temporomandibular joint. Int J Oral Maxillofac Surg.;34:733–8.
159. Yengin E, Evliyaoğlu G (1996): The Evaluation of TMJ by Magnetic Resonance Imaging. J Üstambul University Dental Faculty; 30: 37.