

REFERENCES

REFERENCES

- Abehsera M, Vilgrain V, Belghiti J, *et al.* Inflammatory pseudotumor of the liver: Radiologic-pathologic correlation. *Comput Assist Tomogr* 1995; 19: 80-90.
- Acunas B, Rozanes I, Celik L, *et al.* Purely hydatid disease of the liver. Treatment with percutaneous aspiration and injection of hypertonic saline. *Radiology* 1992; 182: 541-543.
- Adam A, Gibson RN, Soreide O, *et al.* The radiology of the fibrolamellar hepatoma. *Clin Radiology* 1986; 37: 355-358.
- Adjei ON, Tamura S, Sugimara H, *et al.* Contrast enhanced MR imaging of intrahepatic cholangiocarcinoma. *Clinical Radiology* 1995; 50: 6-10.
- Akhan O, Ozmen MN, Dincer A, *et al.* Liver hydatid disease long-term results of percutaneous treatment. *Radiology* 1996; 198: 259-264.
- Anthony Blakeborough, Janice Eard, Daniel Wilsons, Yoshiki Kajiga, *et al.* Hepatic lesion detection at MR imaging: A comparative study with four sequences. *Radiology* 1997; 203:759-765.
- Arrive L, Flejoue LF, Vilgrain V, *et al.* Hepatic adenoma: MR findings in 51 pathologically proved lesions. *Radiology* 1994; 193: 193-199.

Arrive L, Hricak H, Goldberg HI, et al. MR appearance of the liver after partial hepatectomy. *AJR* 1989; 152: 1215-1220.

Elman Abdel Rahman Saker MD degree thesis , Alexandria University, Magnetic Resonance Imaging and Color Doppler of focal hepatic lesions in patients with cirrhosis , 1999 .

Barakos J , Goldberg H , Brown JJ , *et al* . Comparison of computed tomography and magnetic resonance imaging in the evaluation of focal hepatic lesions . *Radiology* 1990 ; 15:93-101 .

Barnes PA, Thomas JL, Bernardine ME. Pit falls in the diagnosis of hepatic cysts by computed tomography. *Radiology* 1981; 141: 129-133.

Berger JF , Laissy JP , Limoto , et al . Differentiation between multiple liver hemangioma and liver metastasis of gastrinomas : value of enhanced MRI . *Comput Assist Tomogr* 1996;29:349-355.

Bergeron P, Oliva VL, Lalonde L, *et al*. Liver angiomyolipoma: classic and unusual presentation. *Abdom Imaging* 1994; 10: 543-549.

Bernard E , Van Beers , Marc Lacrosse , Jacques Jamart Cecile Grandin , *et al* . Detection and segmental location of malignant hepatic tumors: comparison of ferumoxides-enhanced gradient-Echo and T2-weighted spine-Echo MR imaging . *AJR* 1997; 168:713-717.

- bernardino M , Erwin B and Steinberg H . Delayed hepatic CT scanning : increased confidence and improved detection of hepatic metastasis. Radiology 1986 ; 159:71.
- Bismuth H. Surgical anatomy and anatomical surgery of the liver. World J Surgery 1982; 6: 3-9.
- Brandt DJ, Johson CD, Stephens DH, *et al.* Imaging of fibrolamellar hepatocellular carcinoma. AJR 1988; 151: 295-299.
- Bradely G . Flow phenomena in MR imaging . AJR1998; 150:983-994.
- Bret PM, Fond A, Bretagnolle M, *et al.* Percutaneous aspiration and drainage of hydatid cysts in the liver. Radiology 1988; 168: 617-620.
- Bret PM, Labadie M, Bretagnolle M, *et al.* Hepatocellular carcinoma: diagnosis by percutaneous fine needle biopsy. radiology 1988; 13: 253-255.
- Bresslesr El , Alpren MB , Glazer GM , *et al.* Hyper vascular hepatic metastasis : CT evaluation , radiology 1987; 162:49-51 .
- Brick SH, Hill MC, Loude IM, *et al.* The mistaken or indeterminate CT: diagnosis of hepatic metastasis: the value of sonography. AJR 1987; 148: 725-726.

- rinbaum BA, Weinreb JC, Megibow AJ, *et al.* Definitive diagnosis of hepatic hemangiomas: MR imaging versus TC^{99m} labeled red blood cell SPECT. *Radiology* 1990; 176: 95-101.
- rinbaum BA, Weinreb JC, Meyebou AJ, *et al.* Blinded retrospective comparison of MR imaging and TC^{99m} labeled red blood cell SPECT for definitive diagnosis of hepatic tumors. *Radiology* 1989; 173: 270.
- Brown J, Naylor M, and Yegan N. Imaging of hepatic cirrhosis. *Radiology* 1997; 202: 1-16.
- Brown JJ, Lee JM, J.K.T., *et al.* Focal hepatic lesion: differentiation with MR imaging at 0.6 T. *Radiology* 1991; 179: 675-679.
- Buetow PC, Buck JL, Pantongrag-Brown L, *et al.* Biliary cystadenoma and cystadenocarcinoma: Clinical imaging pathologic correlation with emphasis on the importance of ovarian stroma. *Radiology* 1995; 196: 805-810.
- Buetow PC, Pantongrag-Brown L, Buck JL, *et al.* Focal nodular hyperplasia of the liver. Radiologic pathologic correlation. *Radiographics* 1996; 16: 364.
- Bydder G, Young I. NMR imaging clinical use of the inversion recovery sequence. *J Computed Assisted Tomography*. 1985; 914: 569-675.

- Cameron RG, Greig PD, Farber E. Small encapsulated hepatocellular carcinoma of the liver: Provisional analysis of pathogenic mechanism. *Cancer* 1993; 72: 2550-2559.
- Carmody E, Yeung E, McLoughlin M: Angiomyolipomas of the liver in tuberous sclerosis: *Abd Imaging* 1994; 19: 537-540.
- Carr DH, Brown J, Bydder GM. Gadolinium DTPA as a contrast agent in MRI: Initial clinical experience in 20 patients. *AJR* 1984; 143: 215-221.
- Catasca JV ,and Mirowitz SA. T₂-weighted MR imaging of the abdomen: fast spin-echo conventional spine-echo sequences. *AJR* 1994; 162: 61-67.
- Chang JC, Lee YW, Kim HJ: Preoperative diagnosis of angiomyolipoma of the liver. *Abdom Imaging* 1994; 19: 546-553.
- Chen DS , Sung JL , Sheu JC , *et al* . Serum Alpha-Feto protein in the early stage of human hepatocellular carcinoma .*Gastroentrology* 1984;86:1404-9.
- Cherqui D, Rahmouni A, Charlotte F, *et al*. Management of focal nodular hyperplasia and hepatocellular adenoma in young women. A series of 41 patients with clinical, radiological and pathological correlations. *Hepatology* 1995; 22: 1674-1680.

- Hezmar JL, Rumancik WM, Megibow AJ, *et al.* Liver and abdominal screening in patients with cancer: CT versus MR imaging. Radiology 1988; 168: 43-47.
- Choi BI, Han MC, Park JH, *et al.* Giant cavernous hemangioma of the liver: CT and MR imaging in 10 cases. AJR 1989; 152: 1221-1226.
- Choi BL, Lim JH, Han MC, *et al.* Biliary cystadenoma and cystadenocarcinoma: CT and sonographic findings. Radiology 1989; 171: 57-61.
- Choi BL, Han ME and Kim CW. Small hepatocellular carcinoma versus small cavernous hemangioma : differentiation with MR imaging at 1.5 T. Radiology 1990 ; 176:103-108 .
- Chung KY, Mayo-Smith WW, Saini S, *et al.* Hepatocellular adenoma: MR imaging features with pathologic correlation. AJR 1995; 155: 303-309.
- Cottone M, Marceno MP, and Maringhini A. Ultrasound in the diagnosis of hepatocellular carcinoma associated with cirrhosis. Radiology 1983 ; 147:517-519 .
- Couinaud C and Le Foie. Segmental hepatic anatomy divisions. Radiology 1957 ; 9-12.

- Craig JR, Peters RL, Edmundson HA, *et al.* Fibrolamellar carcinoma of the liver. A tumor of adolescents and young adults with distinct clinicopathologic features. *Cancer* 1980; 46: 372-379.
- Davolio Marani SA, Canossi GC, Nicoli FA, *et al.* Hydatid disease: MR imaging study. *Radiology* 1990; 175: 701-706.
- De Lange EE, Mugler JP III, Gay SB, *et al.* Focal liver disease: Comparison of breath-hold T1-weighted MP-GRE MR imaging and contrast-enhanced CT-lesion detection, localization, and characterization. *radiology* 1996; 200: 465-473.
- Demas BE, Hticak H, Moseley M, *et al.* gallbladder bile: an experimental study in dogs using MR imaging and proton MR spectroscopy. *Radiology* 1985; 157: 453-455.
- Didier Mathieu, Valerie Vilgrain, Ahmed-Emad Mahfouz, *et al.* Benign liver tumors. *MRI clinics of North America* 1997; Volume 5, Number 2: 255-265.
- Do H, Lambiase RE, Deyoe L, *et al.* Periotenous drainage of hepatic abscesses: comparison of results in abscesses with and without intrahepatic biliary communication. *AJR* 1991; 157: 1209-1212.

- Donald G Mitchell and Davis D Stark. Hepatobiliary MRI. A text-atlas at mid and high field. Third Edition, Philadelphia, Anne S Patterson and George S. 1992; 47-200.
- Dousset M, Weissleder R, Hendrick RE. Short T_1 inversion recovery imaging of the liver pulse sequence optimization and comparison with spin echo imaging. Radiology 1989; 171: 327-333.
- Duyer A, Frank J, Sank VJ. Short T_1 inversion recovery pulse sequence: Analysis and clinical experience in cancer imaging. Radiology 1988; 168: 827-832.
- Earl's JP , theise ND, and Weinreb JC . Dysplastic nodules and hepatocellular carcinoma : Thin section MR imaging of explanted cirrhotic liver with pathologic correlation . Radiology 1996 ; 201:207 .
- Ebara M and Wantanabe Y. Diagnosis of small hepatocellular carcinoma: Correlation of MR imaging and tumor histologic studies. Radiology 1996; 159: 371-377.
- Ebara M, Otito M and Shinagawa T. Natural history of minute hepatocellular carcinoma smaller than three centimeters complicating cirrhosis. A study in 22 patients. Gastroenterology 1986; 90: 289-298.

- para M, Otito M and Watanabe Y. Diagnosis of small hepatocellular carcinoma: Correlation with MR imaging and tumor histologic studies. *Radiology* 1986; 159: 371-377.
- delman R, Hahn P, Buxton R. Rapid MR imaging with suspended respiration: clinical application in the liver. *Radiology* 1986; 161: 125-131.
- delman RR, Siegel JB, Singer A, *et al.* Dynamic MR imaging of the liver with Gd-DTPA: initial clinical results. *AJR* 1989; 153: 1213-1219.
- gglin TK, Rummeny E, Stark DD, *et al.* Hepatic tumors: quantitative tissue characterization with MR imaging. *Radiology* 1990; 176: 107-110.
- hman RL and Felmle JP. Adaptive techniques of high definition MR imaging of moving structures. *Radiology* 1989; 173: 225-263.
- hman RL, N Namara MT, Pallack M. Magnetic resonance imaging with respiratory gating: techniques and advantages. *AJR* 1984; 143: 1175-1182.
- lizondo G, Weissleder R, Stark D, *et al.* Amebic liver abscess. Diagnosis and treatment, evaluation with MR imaging. *Radiology* 1987; 165: 795-800.

- Evans SE, Harris I, Williams CE. Benign cholangioma of the liver. Clin radiology 1989; 40: 427-429.
- Helmlé JP, Ehman RL. Spatial presaturation; a method of suppressing flow artifacts and improving depiction of vascular anatomy in MR imaging. Radiology 1987; 64: 559-564.
- Merrel L, Wright T, Lake J, Roberts J, Ascher N. Incidence and diagnostic features of macroregenerative nodules Vs small hepatocellular carcinoma in cirrhotic liver. Hepatology 1992; 16: 1372-1381.
- Hertz CJ, Elizondo G, and Wessleder R. Superparamagnetic iron oxide-enhanced MR imaging: pulse sequence optimization for detection of liver cancer. Radiology 1989; 172: 393-397.
- Fisher MR, Wall SD, Hricak *et al.* Hepatic vascular anatomy on magnetic resonance imaging. AJR 1985; 144: 739-742.
- Flejou JF, Barge J, Menu Y, *et al.* Liver adenomatosis an entity distinct from liver adenoma. Gastroenterology 1985; 89: 1132-1135.
- Flisak ME, Budris DM, Olson MC, *et al.* Inflammatory pseudotumor of the liver. Appearance on MRI. Clin Imaging 1994; 18: 1.
- Foley WD, Berland LL, Lawson TL, *et al.* Contrast enhancement technique for dynamic hepatic computed tomographic scanning. radiology 1983; 147: 797-803.

- Boyle WD, Keeland B, Cates JD. Contrast optimization for the detection of focal hepatic lesions by MR imaging at 1.5 T. *AJR* 1987;149: 1155-1160.
- Greeny PC, Marks WM. Hepatic hemangioma: dynamic bolus CT. *AJR* 1986; 147: 711-719.
- Greeny P, Baron R and Teefy S. Hepato-cellular carcinoma reduced frequency of typical findings with dynamic contrast enhanced CT in non Asian population. *Radiology* 1992 ;182:143-148.
- Grenanderz M and Redvanly R. Primary malignant neoplasm in radiologic clinics of North America 1998 ; 36(2): 333-348.
- Friedman AC, Lichtenstein JE, Goodman C, *et al.* Fibrolamellar hepatocellular carcinoma. *Radiology* 1985; 157: 583.
- Furul S, Itai Y, Ohtomo K, *et al.* Hepatic epithelioid hemangioendothelioma: Report of five cases. *Radiology* 1989; 171: 63-68.
- Gabata T, Matsui O, Kadoya M, *et al.* MR imaging of hepatic adenoma. *AJR* 1990; 155: 1009-1011.

- lines PA, and Sampson MA. The prevalence and characterization of simple hepatic cysts by ultrasound examination. *Radiology* 1984; 62: 335-339.
- allego JC, Suarezl, Soler R. Multiple bile duct hamartomas. US, CT, and MR findings. A case report. *Acta radiology* 1995; 36: 273-290.
- ates LK Jr, Cameron AJ, Nagorney DM, *et al.* Primary leiomyosarcoma of the liver mimicking liver abscess. *Am J Gastroentrol* 1995; 90: 649-652.
- orgio A, Tarantino L, Mariniello N, *et al.* Pyogenic liver abscess: 13 years of experience in percutaneous needle aspiration with US guidance. *Radiology* 1995; 195: 122-124.
- oldsmith NA and Woodburne RT. The surgical anatomy pertaining to liver resection. *Surg Gynecol Obstet* 1957; 310: 195-202.
- oman JJ , Epsanza AR , Ridlin MS and Paoletta LP . Cavernous hemangioma of the liver : role of percutaneous biopsy . *Radiology* 1988 ; 166:135-138 .
- acke EM, Tkach JA. Fast MR imaging: techniques and clinical application. *Radiology* 1990; 95: 155-158.
- acke EM, Lenz GW. Improving MR image quality in the presence of motion by using rephasing gradients. *AJR* 1987; 148: 1251-1258.

References

- Huber M, Reuben A, Burrell M, *et al.* Multiple focal nodular hyperplasia of the liver associated with hemihypertrophy and vascular malformations. *Gastroenterology* 1995; 108: 1265.
- John PF, Stark DD, Saini S, *et al.* The differential diagnosis of ringed hepatic lesions in MR imaging. *AJR* 1990; 154: 287-290.
- John PF, Stark DD, Weissleder R. Clinical application for superparamagnetic iron oxide to MR imaging of tissue perfusion in vascular liver tumors. *Radiology* 1990; 174: 361-366.
- Mamm B, Vogl TJ, Branding G, *et al.* Focal liver lesions: MR imaging with Mn-DPDP. Initial clinical results in 40 patients. *Radiology* 1992; 182: 167.
- Mamm B, Wolf KJ, and Felix R. Conventional and rapid MR imaging of the liver with gadolinium-DTPA in clinical use. *Radiology* 1987; 164: 313-320.
- Hamrick-Turner JE, Abbitt PL, Ros PR. Intra-hepatic cholangiocarcinoma: MR appearance. *AJR* 1992; 158: 77-79.
- Hanafusa K, Ohashi L, Himeno Y, *et al.* Hepatic hemangiomas: Findings with two phase CT. *Radiology* 1995; 196: 465-469.
- Heiken JP, Weyman PJ, Lee JKT, *et al.*, . Detection of focal hepatic masses : prospective evaluation with CT, delayed CT, CT

References

- Haruchi R, Uchida T, Kojima T, *et al.* Inflammatory pseudotumor of the liver. Clinicopathologic study and review of the literature. *Cancer* 1990; 65: 1583-1590.
- Ab M. Reda , MD degree thesis , Alexandria University , Ultrasonography , Angiography , Computed Tomography and Magnetic Resonance Imaging in cases of hepatic masses: comparative study , 1992 .
- Osaki T, Chatani M, and Moris S. Dynamic computed tomography of hepatocellular carcinoma. *AJR* 1982; 139: 1099-1106.
- Shazato M, Isobe Y, Ueno E. Inflammatory pseudotumor of the liver. *AJR* 1990; 154: 201-211.
- Oh K , Nishumira K , Toaushik , Fujisawat , Noura S , Ozawa K , and Torizoka K . Hepatocellular carcinoma : MR imaging . *radiology* 1987 ; 164:21-25 .
- Iai Y, Ohtomo K, Furui S, *et al.* Non invasive diagnosis of small cavernous hemangioma of the liver: Advantage of MRI. *AJR* 1985; 145: 1195-1199.
- Iai Y, Terako T: Angiosarcoma of the liver mimicking cavernous hemangioma on dynamic CT. *J Comput Assist Tomogr* 1989; 13: 910-912.

References

- Ishii K, Nishimura K, Togashi K, *et al.* Hepatocellular carcinoma. MR Imaging, Radiology 1987; 164: 21-25.
- Ishii K, Saini S, Hahn PF, *et al.* Differentiation between small hepatic metastasis on MR images. Importance of size-specific quantitative criteria. AJR 1990; 155: 61-66.
- Wittenberg, David D. Strak Bruce H. Forman, *et al.* Differentiation of hepatic metastases from hepatic hemangiomas and cysts by using MR imaging. AJR 1988; 151:79-84.
- Ward, Fengche, J.Ashley Cuthrie, Danile Wilson, J.Peter, *et al.* Hepatic lesion detection after superamagnetic iron oxide enhancement : comparison of five T2 weighted sequences at 1.0 T by using alternative - free response operating characteristic analysis. Radiology 2000, 214:159-166.
- Brown, Jeffrey, Jae Mun Lee, Joseph K. Focal hepatic lesion: Differentiation with MR imaging at 0.5 T. Radiology 1991;179:675-679
- Johnson RD, Mueller PR, Ferrucci JT, *et al.* Percutaneous drainage of pyogenic liver abscesses. AJR 1985; 144: 463-467.
- Jones EC, Chezmar JL, Nelson RC, *et al.* Variability of hepatic enhancement with regard to imaging time and location with

References

- gadolinium-enhanced MR imaging. *Radiology* 1989; 3: 637-646.
- John N Rydberg , David J. Lomas , Kevin J . Coakley , David M , Richard L. Comparison of breath -Hold fast spine echo and conventional spin echo pulse sequences for T2-weighted MR imaging of liver lesions . *Radiology* 1995;194:431-437.
- Kudo M, Matsui O, Nakanuma Y, *et al.* Ciliated hepatic foregut cyst. Radiographic features. *Radiology* 1990; 175: 475-480.
- Kudo M , Matsui O, Takashima T , and Nnamwat . Hepatocellular carcinoma : correlation of MR imaging and histopathologic findings . *Radiology* 1992 ; 183: 819-825 .
- Mitsuyoshi Ito, Donald G, Mitchell, Erik K, Outwater *et al.* Hepatic lesions: Discrimination of non solid, benign lesions from solid malignant lesions with heavily T₂-weighted fast spin-echo MR imaging *radiology* 1997; 204: 729-737.
- Heiken JP, Weyman PJ, Lee JKT, *et al.* Detection of focal hepatic masses: Prospective evaluation with CT, delayed CT, CT during arterial portography, and MR imaging. *Radiology* 1989; 171: 47-51.
- Ken JC, Vansonnenberg E, Casola G, *et al.* Perforated amebic liver abscesses. Successful percutaneous treatment. *Radiology* 1989; 164: 145-197.

References

- Harlin P, Davis GL, McGill DB, *et al.* Hepatic adenoma and focal nodular hyperplasia: clinical, pathologic and radiologic features. *Gastroenterology* 1983; 84: 994-1002.
- Shuroo MS, Zargar SA, Mahajan R, *et al.* Echinococcus granulosus cysts in the liver, management with percutaneous drainage. *Radiology* 1991; 180: 141-145.
- Kim TK, Lee HJ, Tang HJ, Kim AY, Han JK. T2 weighted breath hold MRI of the liver at 1.0 T. Comparison of turbo-spin echo and HASTE sequences with and without fat suppression. *J Magn Reson Imaging* 1998; (6): 1213-8.
- Matsumura S, Lishi H, Tatsuta M, and Ishikawa H. Liver with hypoechoic nodular pattern as a risk factor for hepatocellular carcinoma. *Gastroenterology* 1995; 108: 1778-1784.
- Condo Y, Niwa and Akikusa B. A histopathologic study of early hepatocellular carcinoma. *Cancer* 1983; 52: 684-692.
- Korobkin M, Stephens D, Lee J, *et al.* Biliary cystadenoma and cystadenocarcinoma: CT and sonographic findings. *AJR* 1989; 153: 507-511.
- Kristin H, Lewis and Judith L, Chezmar. Hepatic metastases. *MRI Clinics of North America* 1997; 5: 319-329.

References

- L... MJ, Saini S, Hamm B, *et al.* Focal nodular hyperplasia of the liver. MR findings in 35 proved cases. AJR 1991; 156: 317-320.
- L... lie DF, Johnson CD, MacCarty RL, *et al.* Single pass CT of hepatic tumors: Value of globular enhancement in distinguishing hemangiomas from hypervascular metastasis. AJR 1995; 165: 1403-1406.
- vine E, Cook LT, Grantham JJ. Liver cysts in autosomal-dominant polycystic kidney disease. Clinical and computed tomographic study. AJR 1985; 145: 229-335.
- ev-Touff AS, Bach AM., Wechster RJ, *et al.* The radiologic and pathologic of biliary hamartomas. AJR Am J Roentgenol 1995; 165: 309-320.
- evy HM ,and Newhouse JR . imaging of portal vein thrombosis . AJR 1988; 151:283-286 .
- Lewis WH and Grey H. Anatomy of human body. 21st edition, Philadelphia, Lea and Febiger 1924; 1198-1202.
- Like C Glazer GM , Quint LE , *et al.* Distinction of hepatic cavernous hemangioma from hepatic metastasis with MR imaging . Radiology 1988; 169:409-415.
- Lim KO, Stark DD, and Lesse RT. MR hepatoability imaging: first human experience with Mn-DPDP. Radiology 1991; 178: 79-82.

- Livraghi T, Sangalli G and Vettori C. Adenomatous hyperplastic nodules in cirrhotic liver : a therapeutic approach .Radiology 1989;170:155-157.
- Lubin TG, Simeone JF, Mueller PR, *et al.* Gallbladder bile in cholecystitis: *in vitro* MR evaluation. Radiology 1985; 157: 457-459.
- Lombardo DM, Baker ME, Spritzer C, *et al.* Hepatic hemangiomas Vs metastasis: MR differentiation at 1.5 T. AJR 1990; 155: 55-59.
- Lorigan JC , Charnsangavi J , Carrasco CH , *et al* . Atrophy with compensatory hypertrophy of the liver in hepatic neoplasm : radiological finding . AJR 1988; 150:1291-1295.
- Robbers PR, Ros PR, Goodman ZD, *et al.* Accumulation of Technetium^{99m} sulfur colloid by hepatocellular adenoma: scintigraphic-pathologic correlation. AJR 1990; 148: 105-108.
- Sligowaska E, Connors SK, Shapiro JH. Liver abscess: sonography in diagnosis and treatment. AJR 1982; 138: 253-257.
- Upetin AR and Dash N. Intrahepatic rupture of hydatid cyst: MR finding. AJR 1988; 154 491-442.
- Mahfous A-E, Hamm B, Taupitz M, *et al.* Hypervascular liver lesion: Differentiation of focal nodular hyperplasia from malignant tumors with dynamic gadolinium-enhanced MR imaging. Radiology 1993; 186: 133.

- Miyahara J, Yoshida H, Nakabayashi K, *et al*. Fast spin echo imaging with suspended respiration : gadolinium enhanced MR imaging of the liver tumors . Jcomputed Assist. Tomogr.1987;11:73-80.
- Marchal G, Demaerel P, Decrop E, *et al*. Gadolinium-DDTA enhanced fast imaging of the liver tumors at 1.5 T. J Comput Assist Tomogr 1998; 14: 217-222. (Coated)
- Marago P and Rose P. Imaging of diffuse liver disease in radiologic clinics of North America 1998;vol.36-2:365-375 .
- Marchal G, Van HP, Demaerel P. Detection of liver metastases with superparamagnetic iron oxide in 15 patients. Results of MR imaging at 1.5 T. AJR 1989; 152: 771-775.
- Mark R, Paley and Pablo R Ros. MR imaging of the liver a practical approach MRI Clinics of North America 1997; 5: 415-422.
- Marthinoli C, Cittadini GJR, Rollandi GA, *et al*. Case report: Imaging of bile duct hamartomas. Clinical Radiology 1992; 45: 203-210.
- Marzi-Bonmati L, Menor F, Vizcaino I, *et al*. Lipoma of the liver. US, CT and MRI appearance. Gastrointest Radiology 1989; 141: 455-456.
- Martin D, Sentis M, and Zidan A. Fatty metamorphosis of hepatocellular carcinoma . Detection with chemical shift gradient echo . MR imaging . Radiology 1995 ; 195:125-129 .

References

- Masayuki Kanematsu , Hiroaki Hushi , Takomichi Murakami , Yoshitaka Inaba , *et al* .Focal hepatic lesion detection : Comparison of four T2 weighted MR imaging pulse sequences . Radiology 1998;206:167-175.
- Masayuki Kanematsu , Hiroaki Hushi , Kyoitoh , Takamichi Murakami , *et al* .Focal hepatic lesion detection : Comparison of four fat – suppressed T2 weighted MR imaging pulse sequences . Radiology 1999;211(2) :363-371.
- Mathieu D, Rahmouni A, Anglade MC, *et al*. Focal nodular hyperplasia of the liver: assessment with contrast-enhanced turbo-FLASH MR imaging. Radiology 1991; 180: 25-30.
- Matsui O, Kadoya M and Kameyama T. Adenomatous hyperplastic nodules in the cirrhotic liver differentiation from hepatocellular carcinoma with MR imaging. Radiology 1989; 173: 123-126.
- Matsui O, Kadoya M and Kameyama T. Benign and malignant nodules in cirrhotic liver : distinction based on blood supply . Radiology 1989; 171:335-338 .
- Matsui O, Kadoya M , and Kameyama T . Benign and malignant nodules in cirrhotic liver: distinction based on blood supply . Radiology 1991 ; 178:493-497 .

References

- Mattison GR, Glazer GM, Quint LE, *et al.* MR imaging of hepatic focal nodular hyperplasia: characterization and distinction from primary malignant hepatic tumors. *AJR* 1987; 148: 711.
- Murica N , Ralls P , Jeffeey R, and Johnson M . Color flow Doppler chracterization of focal hepatic lesions . *AJR* 1992 ; 159:1155-1179 .
- McFarland EG, Mayo SW, Saini S, *et al.* Hepatic hemangiomas and malignant tumors: improved differentiation with heavily T₂-weighted conventional spin-echo MR imaging. *Radiology* 1994; 193: 43-47.
- Hendenz RJ, Schiebler ML, Outwater, *et al.* Hepatic abscesses: MR imaging findings. *Radiology* 1944; 190: 431-436.
- Milan E , Martin N , ELS Van Persijn , Cornelis JH ,Jo Hermans , *et al.* Comparison of inversion-recovery gradient and spin echo and fast spin echo techniques in detection and characterization of liver lesions .*Radiology* 1998,290:427-434.
- Lirowitz SA, Lee JKT, Gutierrez E, *et al.* Dynamic gadolinium-enhanced rapid acquisition spine-echo MR imaging of the liver. *Radiology* 1991; 179: 371-376.
- Mitchell DG, Burk DL, Vintiski S. The biophysical basis of tissue contrast in extracranial MR imaging. *AJR* 1987; 149: 837-842.

References

- Mitchell DG, Vinitzki S, Bark DL. Motion artifacts reduction in MR imaging of abdomen: gradient moment nulling versus respiratory sorted phase encoding. *Radiology* 1988; 169: 155-160.
- Mitchell DG, Vinitzki S, Burk DL, *et al.* Variable-flip-angle spin-echo MR imaging of the pelvis: more versatile T₂-weighted images. *Radiology* 1984; 171: 525-529.
- Mitchell DG, Vinitzki S, Refkin. Sampling band width and fat suppression: effects of long TR/TE/MR imaging on the abdomen and pelvis at 1.5 T. *AJR* 1989; 153: 419-425.
- Mitchell DG, Vinitzki S, Sponara S. The liver and pancreas: improved spin-echo T₁ contrast by shorten TE and fat suppression at 1.5 Tesla. *Radiology* 1991; 178: 67-71.
- Mitchell DG, Rubin R and Siegelman. Hepatocellular carcinoma within siderotic regenerative nodules (the nodule within nodule sign) on MR imaging . *Radiology* 1991 ; 178:101-103 .
- Mitchell DG, Saini S, Weinreb J, *et al.* Hepatic metastasis and cavernous hemangiomas : distinction with standard and triple dose gadolinium – enhanced MR imaging . *Radiology* 1994 ;193:49-57.

References

- Moinuddin M, Auison JR, and Montgomery JLL. Sonographic diagnosis of hepatic hemangioma : its role in the management of hepatic mass lesion . A JR 1985 ; 145:223-228 .
- Uchida F . Doppler ultrasound in diagnosis of liver tumor and portal hypertension . Radiology 1993 ; 85:44-55 .
- Uchida JK, Stack C, Turner DA. Imaging of surgically relevant hepatic vascular and segmental anatomy. 1. Normal anatomy. AJR 1987; 149: 287-292.
- Uchida JK, Stack C, Turner DA. Imaging of surgically relevant hepatic vascular and segmental anatomy: 11. Extent and respectability of hepatic neoplasm. AJR 1987; 1987: 293-298.
- Ungovan JA, Cronan JJ, Vacarro J: Hepatic cavernous hemangiomas: Lack of enlargement over time. Radiology 1994; 191: 111-113.
- Nelson RC, Chezmar JL. Diagnostic approach to hepatic hemangiomas. Radiology 1990; 176: 11-13.
- Nelson RC, Lovettke, Chezmar JL, *et al* .Comparison of pulsed Doppler sonography and angiography in patients with portal hypertension . AJR 1987 ; 149:77-81 .

- Nelson RC , Chezmar JL , Sugarbaker PH , *et al* . Hepatic tumors : comprison of CT and MR imaging for preoperative evaluation . Radiology 1989 ; 172:27-34 .
- Omishi KO and Nomurra F. Ultrasonic Doppler studies of hepatocellular carcinoma and comparison with other hepatic focal lesions . Gastroentrology 1989 ; 97:1429-1435.
- Omomo K, Baron RL and Dodd III GD. Confluent hepatic fibrosis in advanced cirrhosis: appearance at CT. Radiology 1993; 188: 31-35.
- Omomo K, Haii Y, and Yoshikaw K. Hepatic tumors : dynamic MR imaging. Radiology 1987; 163: 27-32.
- Omomo K, Itali Y, Yoshikwa K, *et al*. Hepatocellular carcinoma and cavernous hemangioma: differentiation with MR imaging. Radiology 1988; 168: 621-623.
- Omomo K, Tsutomu A, Ital Y, *et al*. MR imaging of malignant mesenchymal tumors of the liver. Gastrointest Radiology 1992; 17: 58-62.
- Over J and Baron R . Helical biphasic contrast enhanced CT of the liver technique , indications , interpretation and pit falls . Radiology 1996 ; 201:1-14 .

- Overwater EK, Mitchell DG, Vinitki S. Abdominal MR imaging: Evaluation at a fast spine-echo sequence,. Radiology 1994; 190: 425-429.
- Tagani JJ. Intrahepatic vascular territories shown by computed tomography (CT). Radiology 1983; 147: 173-179.
- Yang MR, Abbit PL, Mugler JJ. Liver metastasis: optimization of MR imaging pulse sequences at 1 T. Radiology 1988; 167: 659-699.
- Patricia J, Mergo and Pablo R, Ros. MR imaging of inflammatory disease of the liver. MRI Clinics of North America 1997; (5) 2: 367-370.
- Paulson EK, McClellan JS, Washington K, *et al.* Hepatic adenoma: MR characteristics and correlation with pathologic findings. AJR 1994; 163: 113-116.
- Peter C, Buetow, Russell B and Midkiff. Primary malignant neoplasms in the adult. MRI Clinics of North America 1997; 5: 289-293.
- Philippe Soyer, David A, Bluemke and Roland Rymer. MR imaging of the liver. MRI Clinics of North America 1997; 5: 205-221.
- Philippe Soyer, M'Hamed Gouhiri, Yves Rondeau, *et al.* Non-breath-hold fast spin-echo versus breath-hold fast-echo and spoiled gradient-recalled echo MR imaging in the detection of hepatic

tumors: correlation with surgical findings. *AJR* 1997; 168: 1199-1204.

Philippe Soyer, Sandrin Le Normard , Sebastien Clement , *et al* . T2 weighted spin echo MR imaging of the liver : Breath hold fast spine -echo versus non Breath Hold fast spine -echo imaging with and without fat suppression .*AJR*1996;166:593-597.

Flowers C, Ros PR, Stoupis C, *et al*. Primary liver neoplasms. MR imaging with pathologic correlation. *Radiology* 1994; 14: 549-482.

O'Leary SF, Benjamin GG. Hepatic hemangiomas: Simple diagnostic sign with dynamic bolus CT. *Radiology* 1992; 182: 545-548.

Madan DR, Cruik JR, Colletti PM, *et al*. Hepatic epithelioid hemangioendothelioma. *Radiology* 1988; 169: 145-148.

Hills PW, Henley DS, Colletti PM, *et al*. Amebic liver abscess, MR imaging. *Radiology* 1987; 165: 801-804.

Winertson TE, Fortune JB, Peters JC, *et al*. Primary leiomyoma of the liver. A case report and review of the literature. *Dig Dis Sci* 1992; 4: 622 (coated).

Reining JW, Dwyer AJ, Miller DL, Frank JA, *et al*. Liver metastasis: detection with MR imaging at 0.5 and 1.5 T. *Radiology* 1989; 170: 149-153.

References

- Reining JW. Breath-hold fast spin-echo MR imaging of the liver: a technique for high-quality T₂-weighted images. *Radiology* 1995; 194: 303-304.
- Richard C, Semelka, Carolyn M. Hepatic hemangiomas. *MRI clinic of North America* 1997; 5: 241-242.
- Richard F, Marton J and Richard Hebel (1984) : Correlation : In a study guide to epidemiology and Biostastics , Text book , 2nd (ed.), P(81-94).
- Rogers WF, Rall PW, Boswell WP, *et al.* Amebiasis: unusual radiographic manifestation. *AJR* 1980; 135: 1253-1257.
- Ros PR, Lubbers PR, Olsted WW, *et al.* Hemangioma of the liver: heterogeneous appearance of T₂-weighted images. *AJR* 1987; 149: 1167-1170.
- Rummeny E, Saini S, Wittenberg J, *et al.* MR imaging of the liver neoplasms. *AJR* 1989; 152: 443-499.
- Rummeny E, Weissleder R and Stark D. Primary liver tumor: Diagnosis by MR imaging. *AJR* 1989; 152: 63-72.
- Rummeny E, Weissleder R, Sironi S, *et al.* Central scars in primary liver tumors. MR features, specificity, and pathologic correlation. *Radiology* 1989; 171: 323-326.

References

- Rummeny EJ, Wernecke K, Saini S, *et al*: Comparison between high-field-strength MR imaging and CT for screening of hepatic metastases. A receiver operating characteristic analysis. *Radiology* 1992; 182: 879-886.
- Runge VM, Clanton JA, Patain CL. Respiratory gating in magnetic resonance imaging at 0.5 tesla. *Radiology* 1984; 151: 521-254.
- Rydberg JN, Lomas DJ, Coakley KJ, *et al*. Comparison of breath hold fast spin-echo and conventional spine-echo pulse sequences for T₂-weighted MR imaging of liver lesions. *Radiology* 1995; 144: 431-437.
- Saini S, Edelman R, Sharma P and Li W. Blood-pool MR contrast material for detection and characterization of focal hepatic lesion: Initial clinical experience with ultra small super paramagnetic iron oxide (Ami-227). *AJR* 1995; 164: 1147-1152.
- Saini S, Stark DD, Brady TJ, *et al*. Dynamic spine-echo MRI of liver cancer using gadolinium-DTPA: animal investigation. *AJR* 1986; 147: 357-362.
- Saini S, Stark DD, Hahn PP. Ferrite particle a superparamagnetic MR contrast agent for the reticuloendothelial system. *Radiology* 1987; 162: 211-216.

- Saini S, Stark DD, Hahn PP. Ferrite particles: a superparamagnetic MR contrast agent for enhanced detection of liver carcinoma. *Radiology* 1987; 162: 217-222.
- Saini S, Stark DD, Wittenberg J, Dynamic gadolinium DTPA imaging of liver cancer, animal investigation. *AJR* 1986; 147: 357-363.
- Semelka RC, Sijhoenut JP, Kroeker MA, *et al.* Focal liver disease: comparison of dynamic contrast-enhanced CT and T2-weighted fat suppressed, FLASH, and dynamic gadolinium-enhanced MR imaging at 1.5 T. *Radiology* 1992; 184: 687-694.
- Semalk RC , Brown ED , Ascher SM, *et al* . Hepatic hemangiomas : a multi – institutional study of appearance on T2 weighted and serial gadolinium – enhanced gradient echo MR images . *Radiology* 1994; 192:401-406 .
- Sexton CC, Zeman RK. Correlation of Computed tomography, Sonography and Gross anatomy of the liver. *AJR* 1983; 141: 711-715.
- Schwartz L , Seltzers , Tempany C , *et al* . Prospective comparison of T2 – weighted fast –spine echo , with and without fat suppression and conventional spin- echo pulse sequences in the upper abdomen. *Radiology* 1993 ; 189:411-416 .

References

- Shimamoto K , Sakuma S , Ishigaski T , Ischiguch T , Itoh S , and Fukatsu H . Hepatocellular carcinoma : evaluation with color Doppler US and MR imaging . Radiology 1992 ; 182: 149-153.
- Shoenut JP, Semelka RC, Levi C, *et al.* Ciliated hepatic foregut cysts: US, CT and contrast-enhanced MR imaging. Abdom Imaging 1994; 19: 150-160.
- Shuman W, Baron R, Peters M, Taziodi P. Comparison of STIR and spin-echo MR imaging at 1.5 T in 90 lesions of the chest, liver and pelvis. AJR 1989; 152: 853-857.
- Siskin GP, Haller JO, Miller S, *et al.* AIDS-related lymphoma, radiologic features in pediatric patients. Radiology 1995; 196: 63-66.
- Slone HW, Bennett WF, Bova JG. MR findings of multiple biliary hamartomas. AJR Am J Roentgenol 1993; 101: 581-590.
- Spiegel RM, King DL, Green WM. Ultrasonography of primary cysts of the liver. AJR 1978; 131: 235-238.
- Spritzer C, Kressel MY, Mitchell D, *et al.* MR imaging of normal extrahepatic bile ducts. J Comput Assist Tomography 1987; 11: 248-252.
- Stark DD, Felder RC, Wittenberg J, *et al.* Magnetic resonance imaging of hepatic cavernous hemangioma of liver: tissue specific characterization. AJR 1985; 145: 213-220.

References

- Stark DD , Hahn PF , Felder RC and Wittenberg . Magnetic resonance imaging of cavernous hemangioma of the liver . Tissue specific characterization . AJR 1985 ; 145:213-220 .
- Stark DD, Hendrick RE, Hahn PF, *et al.* Motion artifact suppression by fast spin-echo imaging. Radiology 1987; 164: 183-191.
- Stark DD, Wittenberg J, Butch RJ, Ferrucci JP. Hepatic metastasis randomized controlled comparison of detection with MR imaging and CT. Radiology 1987; 165: 399-406.
- Stark DD, Wittenberg J, Edelman R. Detection of hepatic metastasis: analysis of pulse sequence performance in MR imaging. Radiology 1986; 159: 365-370.
- Stephens DH, Johnson DC. Primary malignant neoplasm of the liver. Alimentary Tract Radiology 1994; 166-1687.
- Sugarbaker PH. Surgical decision making for large bowel cancer metastatic to the liver. Radiology 1990; 103: 278-288.
- Takayama T, Makuuchi M, Hirohashi S. Malignant transformation of adenomatous hyperplasia to hepatocellular carcinoma. Lancet 1990; 336: 1150-1153.
- Takayasu K, Moriyama N, Shima Y, *et al.* A typical radiologic findings in hepatic cavernous hemangioma: correlation with histologic features. AJR 1986; 146: 1149-1153.

References

- Takayasu K , Furukawa H , Wakao F , Muramatsu Y , Abe H , Terachi T
Winter T , *et al* . CT diagnosis of early hepatocellular
carcinoma :Sensitivity , findings, and CT pathologic
correlation . AJR1995;164:885-890.
- Tanaka S, Kitamura Imoka S, and Sasaki Y. Hepatocellular carcinoma:
sonographic and histologic correlation. AJR 1982; 140: 701-
707.
- Tanaka S , Kitamura T, Nakahish K and Okuda S. Color Doppler flow
imaging of liver tumors . AJR1990;154:509-414.
- Tanaka S , Kitamura T , Fujita , Kasugai Hand Ishiguro S. Small
hepatocellular carcinoma . Differentiation from adenomatous
hyperplastic nodule with color Doppler flow imaging .
Radiology 1992;182:161-165.
- Tanaka K , Numata K , Okazaki H , Nakamura S , Inoue S , and Takamura
Y . Diagnosis of portal vein thrombosis in pateints with
hepatocellular carcinoma. Efficiency of color Doppler
sonography compared with angiography . AJR 1993 ;
160:1279-1283 .
- Tangy , Yamashita Y , Namimoto T . Characterization of focal liver lesions
with half - fourier acquisition single shot turbo - spine- echo
(HASTE) and Inversion Recovery (IR) -(HASTE) sequences
. J Magn Reson Imaging 1998 ; (2):348-45.

- Taupitz M, Speidel A, Hamm B, *et al.* T₂ weighted breath-hold MR imaging of the liver at 1.5 T: results with a three-dimensional steady-state free precession sequence in 87 patients. *Radiology* 1995; 194: 439-446.
- Terda T and Nakanuma Y. Survey of iron accumulative macroregenerative nodules in cirrhotic livers. *Hepatology* 1989; 10 (5): 851-854.
- Tessler F , Gehring B , Gones A , Perella R , Ravendra N , busottil R , and Grount E . Diagnosis of portal vein thrombosis : value of color Doppler imaging . *AJR* 1991 ; 157:293-296 .
- Theoni RF, Luh N, Winkler M. Low-field MR imaging of the liver. *Radiology* 1986; 169: 64-59; 1988.
- Thomas JL, Bernardino ME, Vermess M, *et al.* EOE-13 in the detection of hepatosplenic lymphoma. *Radiology* 1982; 145: 629-634.
- Titelbam DS, Hatubu H, Scheiber ML, *et al.* Fibrolamellar hepatocellular carcinoma, MR appearance. *J Comput Assist Tomogr* 1988; 12: 588-591.
- Tomohiro Nammoto , Yasuyuki Yamashita , Seiya Sumi , Xi Tang , and Mutsusa Takahashi : Focal liver masses : characterization with Diffusion -weighted Echo-Planner imaging . *Radiology* 1997 ; 204: 739-744.

- Tung GA, Vaccaro JP, Cronan JJ, *et al.* Cavernous hemangioma of the liver: Pathologic correlation with high-field MR imaging. *AJR* 1994; 162: 1113-1117.
- Tylor KJW , Ramos T , Morese SS , Fortune K , and Hammers L .Focal liver masses : differential diagnosis with pulsed Doppler US . *Radiology* 1987 ; 146:643-649 .
- Van Beers B, Demeure R, Pringot J. Dynamic Spin echo imaging with Gd-DTPA value in the differentiation of hepatic tumors. *AJR* 1990; 154: 515-519.
- Van Beers B, Materne R ,Lacrosse M . MR imaging of hypervascular liver tumors : Timing optimization during the arterial phase .*J Magn Rson Imaging* 1999;(4) : 562-7.
- Van GD , Avi E , Englhom L ,and Struyven J . Sonographic features of portal vein thrombosis . *AJR* 1985; 144:749-752.
- Vilgrain V, Flejou JF, Arrivé L, *et al.* Focal nodular hyperplasia of the liver. MR imaging and pathologic correlation in 37 patients. *radiology* 1992; 184: 699.
- Vilgrain V, Silbermann O, Benhamau JP, *et al.* MR imaging in intracystic hemorrhage of simple hepatic cysts. *Abdom imaging* 1993; 18: 164-169.

- Vogl TJ, Kommel S, Hammersting R, *et al.* Liver tumors: Comparison of MR imaging with Gd-EoB-DTPA and Gd-DTPA. *Radiology* 1996; 200: 59.
- Wanless IR, Mawdsley C, Adams R. On the pathogenesis of focal nodular hyperplasia of the liver. *Hepatology* 1985; 5: 1194.
- Wehrli FW, Macfall JR, Glover GH, *et al.* The dependence of nuclear magnetic resonance (NMR) image contrast on intrinsic and pulse sequence timing parameters. *Magnetic Reson Imaging* 1984; 2: 3-16.
- Weinreb JC, Brateman L, and Maravilla KR. Magnetic resonance imaging of hepatic lymphoma. *AJR* 1984; 143: 1211-1214.
- Welch IJ, Sheedy PF, Johnson CM, Stephens DH, May GR, and Adson MA. Focal nodular hyperplasia: comparison of angiography, CT, US and scintigraphy. *Radiology* 1985; 156: 593-595.
- Werencke K, Rummary E, Bongart G, *et al.* Detection of hepatic masses in patients with carcinoma: comparative sensitivities with sonography, CT MR imaging. *AJR* 1991; 157: 731-740.
- Whitney WS, Herfkens RJ, Jeffrey RB, *et al.* Dynamic breath-hold multiplanar spoiled gradient-recalled MR imaging with gadolinium enhancement for differentiating hepatic hemangiomas from malignancies at 1.5 T. *Radiology* 1993;

- 189: 863-870. Adson MA. Surgery symposium: mass lesions of the liver. Mayo Clin Proc 1986; 61: 362-368.
- Wilcox DM, Weinreb JC, Lesh P. MR imaging of a hemorrhagic hepatic cyst in a patient with polycystic liver disease. J Comput Assist Tomogr 1985; 9: 183-189.
- Winkler M, Theoni R, Luh N. Hepatic neoplasia: Breath hold MR imaging. Radiology 1989; 170: 801-806.
- Wittenberg J, Stark DD, Forman BH, *et al*. Differentiation of hepatic masses from hepatic hemangiomas and cysts by using MR imaging. AJR 1988; 151: 79-84.
- Wittenberg J, Tosteson AA, Karstaedt N, *et al*. MR imaging versus CT : a multi-institutional comparison of hepatic metastatic tumors detection accuracy . Radiology 1988 ; 169(P) 63 .
- Wolfgang Schima , Sanjay Saini , Jose A, Echeverri , *et al* . Focal liver lesions : characterization with conventional spin-echo versus fast spine-echo T2 weighted MR imaging . Radiology 1997 ; 202:369-393.
- Yamakado K , Sakuma H , Murashima S, Nakatsuka A. Comparison of breath – hold multishot echoplaner and respiratory triggered fast –spin echo- sequences for T2 weighted MRI of the liver lesions . J Magn Reson Imaging 1998; (2) : 432-7.

References

- Yamashita Y, Mitsuzaki K, Tang Y, Ogata T, *et al*. Small hepatocellular carcinoma in patients with chronic liver damage: prospective of detection with dynamic MR imaging and helical CT of the whole liver. 1996; 200: 79-84.
- Yoshikawa J, Matsui O , Takashima T , *et al* . Fatty metamorphosis in hepatocellular carcinoma . Radiographic features in 10 cases . AJR 1988; 151: 717-720.
- Yu Js , Kim KW , Kim YH , Jeong EK . Comparison of multishot turbo spin echo and HASTE sequences for T2 weighted MRI of liver lesions . J Magn. Reson Imaging , 1998 ; 8(5) : 1079-84.
- Zaki M and Osman M . Evaluation of the role of color Doppler sonography in diagnosis of diffuse H.C.C in cirrhotic liver . The Egyptian Journal of radiology & Nuc.Med. 1997 ; 28(2); 283-293.
- Zerhouni EA, Rutter C, Hamilton SR, *et al*. CT and MR imaging in the staging of colorectal carcinoma: Report of the radiology diagnostic oncology group II. Radiology 1996; 200: 443-451.
- Zoranza J and Ginaldi S. Computed tomography in hepatic lymphoma. Radiology 1981; 138: 405-410.