

References

- Arenault TM, King PF, Marsh JW Jr et al:** Systemic gadolinium toxicity in patients with renal insufficiency and renal failure: retrospective analysis of an initial experience. *MayoClin. Proc.*, 71: 1150-1154, (1996).
- Allkemper T, Tombach B and Heindel W:** Gd-BOPTA enhanced excretory MR urography without administration of diuretics. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr.*, 173 (2): 115-120, (2001).
- Anson BJ and McVay CB:** *Surgical anatomy*, 6th ed. Philadelphia, WB Saunders, pp: 833, (1984).
- Beer A, Saar B, Link TM et al:** Virtual endoscopy of the urinary tract from T(2)-weighted and gadolinium-enhanced T(1)-weighted MR urographic images. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr.*, 173 (11): 997-1005, (2001).
- Bigongiari LR, Davis RM, Novak WG et al:** Visualization of the medullary rays on excretory urography in experimental ureteric obstruction. *AJR*, 129: 89-93, (1977). Quoted from **Talner LB:** Urinary Obstruction. In Grainger RG and Allison D. *Grainger & Allison's Diagnostic Radiology*. 3rd ed. New York, Churchill Livingstone, (1997).
- Blandino A, Gaeta M, Minutoli F et al:** MR pyelography in 115 patients with a dilated renal collecting system. *Acta Radiol.*, 42 (5): 532-536, (2001).
- Blandino A, Minutoli F, Gaeta M et al:** MR pyelography in the assessment of hydroureteronephrosis: single-shot thick-slab RARE versus multislice HASTE sequences., *Abdom Imaging.*, 28 (3): 433-439, (2003).
- Blandino A, Scribano E, Gaeta M et al:** Magnetic resonance pyelography: clinical use of ultrafast breath-hold sequences in obstructive uropathy. Personal experience. *Radiol Med (Torino)*., 96 (4): 353-359, (1998).
- Boridy IC, Nikolaidis P, Kawashima A et al:** Noncontrast helical CT for ureteral stones. *World J Urol.*, 16: 18-21, (1998).

Bosniak MA, Megibow AJ, Ambos MA et al: Computed tomography of ureteral obstruction. *AJR.*, 138: 1107-1113, (1982).

Boulay I, Holtz P, Foley WD et al: Ureteral calculi: diagnostic efficacy of helical CT and implications for treatment of patients. *AJR Am J Roentgenol.*, 172: 1485-1490, (1999).

Brown SCW, Upsdell SM and O'Reilly PH: The importance of renal function in the interpretation of diuresis renography. *Br J Urol.*, 69: 121-125, (1992).

Budau M, Onu M, Jinga V et al: US and MRI in renal obstruction evaluation. *Arch Ital Urol Androl.*, 74 (2): 59-60, (2002).

Burge HJ, Middleton WD, McClennan BL, Hildebolt CF: Ureteral jets in healthy subjects and in patients with unilateral ureteral calculi: comparison with color Doppler US. *Radiology*, 180: 437-442, (1991).

Caoili EM, Cohan RH, Korobkin M et al: Urinary Tract Abnormalities: Initial Experience with Multi-Detector Row CT Urography. *Radiology*, 222:353-360, (2002).

Catalano C, Pavone P, Laghi A et al: MR pyelography and conventional MR imaging in urinary tract obstruction. *Acta Radiol*, 40: 198-202, (1999).

Chen MY, Zagoria RJ and Dyer RB: Radiologic findings in acute urinary tract obstruction. *J Emerg Med.*, 15(3): 339-343, (1997).

Chibber GD, Hargrave TB and Merrick MV: 99m Technetium DMSA and the prediction of recovery in obstructive uropathy. *Br J Urol*, 53: 492-495, (1981).

Cohnen M, Jung G, Fritz B et al: Modern diagnostic imaging: MR urography. *Urologe A.*, 41 (6): 542-547, (2002).

Conway JJ: "Well-tempered" diuresis renography: Its historical development, physiological and technical pitfalls, and standardized technique protocol. *Semin Nucl Med*, 22: 74-84, (1992).

Coroneos E, Assouad M, Krishnan B and Troung LD: Urinary obstruction causes irreversible renal failure by inducing chronic tubulointerstitial nephritis. *Clinical Nephrology*, 48(2), 125 – 128, (1997).

Cronan JJ and Tublin ME: Role of the resistive index in the evaluation of acute renal obstruction. *AJR*, 164: 377–378, (1995).

Dal Canton A, Corradi A, Stanziale R et al: Glomerular hemodynamics before and after release of 24-hour bilateral ureteral obstruction. *Kidney Int*, 17: 491–496, (1980).

Dalla Palma L, Stacul F, Bazzocchi M et al: Ultrasonography and plain film versus intravenous urography in ureteric colic. *Clin Radiol*, 47: 333–336, (1993).

Deyoe LA, Cronan JJ, Breslaw BH and Ridlen MS: New techniques of ultrasound and color Doppler in the prospective evaluation of acute renal obstruction. Do they replace the intravenous urogram? *Abdom Imaging*, 20: 58–63 (1995).

Di Girolamo M, Pirillo S, Laghi A et al: Urography with magnetic resonance: a new method for study of the renal collecting system in patients without obstructive uropathy. *Radiologia Med.*, 92: 758–764 (1996).

Diamond JR, Richardo SD and Klahr S: Mechanisms of interstitial fibrosis in obstructive nephropathy. *Seminars in Nephrology*, 18 (6): 594 – 602, (1998).

Diamond JR: Macrophages and progressive renal disease in experimental hydronephrosis. *American Journal of Kidney Diseases*, 26 (1): 133 – 140, (1995).

Dretler SP: Ureteral stone disease: Options for management. *Urology Clinics of North America*, 17: 217- 230 (1990).

El-Diasty T, Mansour O and Farouk A: Diuretic contrast-enhanced magnetic resonance urography versus intravenous urography for depiction of nondilated urinary tracts. *Abdom Imaging.*, 28(1): 135-145, (2003).

Eleman RR, Wielopolski P and Schmitt F: Echo-planar MR imaging. *Radiology*. 192: 600-612, (1994).

Elkin M: Obstruction and dilatation of the urinary tract. In: Elkin M (ed) *Radiology of the urinary system*. Little Brown, Boston, Ch 8, (1980).

Elkin M: Radiology of the urinary tract: some physiological considerations. *Radiology*, 116: 259–270 (1975). Quoted from **Talner LB:** Urinary Obstruction. In Grainger RG and Allison D. *Grainger & Allison's Diagnostic Radiology*. 3rd ed. New York, Churchill Livingstone, (1997).

Eshima D and Taylor A Jr: Technetium-99m(^{99m}Tc) mercaptoacetyltriglycine: Update on the new ^{99m}Tc renal tubular function agent. *Semin Nucl Med*, 22: 61–73 (1992).

Farres MT, Gattegno B, Ronco P et al: Nonnephrotoxic, dynamic, contrast enhanced magnetic resonance urography: use in nephrology and urology. *J Urol*, 163 (4): 1191-6, (2000).

Fielding JR, Fox LA, Heller H et al: Spiral CT in the evaluation of flank pain: overall accuracy and feature analysis. *J Comput Assist Tomogr*, 21 (4): 635-638, (1997).

Fine EJ: Interventions in renal scintigraphy. *Semin Nucl Med*, 21: 116–127, (1991).

Fredrick A, Gulmi, Felsen D and Vaughan D: Pathophysiology of urinary tract obstruction In: Walsh PC, Vaughan ED Jr, Wein AJ, (eds). *Campbell's Urology*. 8th ed. Philadelphia, Pa: WB Saunders, Vol 1 pp: 411 – 462, (2002).

Friedburg H, Hennig J and Frankenschmidt: A. RARE-MR urography: a fast new tomographic imaging procedure for demonstrating the efferent urinary pathways using nuclear magnetic resonance. *Radiology*, 27: 45-47, (1987).

Garcia-Pena BM, Keller MS, Schwartz DS et al: The ultrasonographic differentiation of obstructive versus nonobstructive hydronephrosis in children: a multivariate scoring system. *J Urol* 158: 560–565, (1997).

Gulmi FA, Matthews GJ and Marion D et al: Volume expansion enhances the recovery of renal function and prolongs the diuresis and natriuresis after release of

bilateral ureteral obstruction: A possible role for atrial natriuretic peptide. *J Urol*, 153: 1276–1283, (1995).

Hasegawa N, Miki K, Kato N and Ohishi Y: Evaluation of clinical usefulness of MR urography. *Nippon Hinyokika Gakkai Zasshi.*, 90 (8): 706-712, (1999).

Henning J, Nauwerth A and Friedberg H: RARE imaging a fast imaging method for clinical MR. *Magn Reson Med*, 3: 391–395, (1986).

Hinman F Jr: *Atlas of Urosurgical Anatomy*. Philadelphia, WB Saunders, pp: 356, (1993).

Huang A, Palmer LS, Hom D et al: The role of nitric oxide in obstructive nephropathy. *The Journal of Urology*, 163(4): 1276 – 1281, (2000).

Hughes J, Jan W, Goodie J et al: MR urography: evaluation of different techniques in non-dilated tracts. *Clin Radiol.*, 57 (11): 989-994,(2002).

Hussain S, O'Malley M, Jara H et al: MR urography. *Magn Reson Imaging Clin N Am.*, 5 (1): 95-106, (1997).

Hwang SI, Kim SH, Kim YJ et al: Effectiveness of MR urography in the evaluation of kidney which failed to opacify during excretory urography: comparison with ultrasonography. *Korean J Radiol.*, 1(3):152-158, (2000).

Jackson EK: Diuretics. In Hardman JG, Limbird LE, Molinoff PB, Goodman & Gilman's A (eds) *The pharmacological bases of therapeutics 9th ed.* McGraw-Hill, New York, pp: 691.692.697-701, (1996).

James D and Brooks: Anatomy of the lower urinary tract and male genitalia. In: Walsh PC, Vaughan ED Jr, Wein AJ, (eds). *Campbell's Urology. 8th ed.* Philadelphia, Pa: WB Saunders, Vol 1 pp: 41 – 81, (2002).

Jones DA, Lupton EW and George NJR: Effect of bladder filling on upper tract urodynamics in man. *Br J Urol.*, 65: 492–496, (1990).

Jung P, Brauersl A, Nolte-Ernsting CA et al: Magnetic resonance urography enhanced by gadolinium and diuretics: a comparison with conventional urography in diagnosing the cause of ureteric obstruction. *BJU International*, 86: 960 – 965 (2000).

Kabalin JN: Surgical anatomy of the retroperitoneum, kidneys and ureters. In: Walsh PC, Vaughan ED Jr, Wein AJ, (eds). *Campbell's Urology*. 8th ed. Philadelphia, Pa: WB Saunders, Vol 1 pp: 3 – 40, (2002).

Kamholtz RG, Cronan JJ and Dorman GS: Obstruction and the minimally dilated renal collecting system: US evaluation. *Radiology*, 170: 51–53, (1989).

Kass EJ, Majd M and Belman AB: Comparison of the diuretic renogram and the pressure perfusion study in children. *J Urol.*, 134: 92–96, (1985).

Katz DS, Hines J, Rausch DR et al: Unenhanced helical CT for suspected renal colic. *AJR Am J Roentgenol.*, 173:425–430, (1999).

Katzberg RW, Buonocore MH, Ivanovic M et al: Functional, dynamic, and anatomic MR urography: feasibility and preliminary findings. *Acad Radiol.*, 8(11): 1083–1099 (2001).

Kessel RG, Kardon RH: Tissues and Organs: A Text-Atlas of Scanning Electron Microscopy, (1979). Quoted from anatomy of the retroperitoneum, kidneys and ureters. In: Walsh PC, Vaughan ED Jr, Wein AJ, (eds). *Campbell's Urology*. 8th ed. Philadelphia, Pa: WB Saunders, Vol 1 pp: 411 – 462, (2002).

Klahr S and Morrissey J: Obstructive nephropathy and renal fibrosis. *Am J Physiol Renal Physiol* 283(5): 861–875, (2002).

Klahr S: Obstructive nephropathy. *Kidney International*, 54 (1): 286 –300, (1998a).

Klahr S, Bueskert J and Morrison A: Urinary tract obstruction. In Brenner BM and Rector FC (eds), *The kidney 3th ed*. Philadelphia: W.B. Saunders. Pp: 1443 – 1490, (1986).

Klahr S: Obstructive uropathy. In R.J. Glasscock (Ed.), *Current therapy in nephrology and hypertension*, 4th ed. St. Louis, MO: Mosby. pp: 69 –74, (1998b).

-
- Klahr S:** Pathophysiology of obstructive nephropathy. *Kidney International*, 23: 414 – 426, (1983).
- Klein LT, Frager D, Subramaniam A, Lowe FC:** Use of magnetic resonance urography. *Urology*, 52: 602-608, (1998).
- Knesplova L and Krestin GP:** Magnetic resonance in the assessment of renal function. *Eur Radiol*, 8: 201-211, (1998).
- Knopp MV, Dorsam J, Oesingmann N et al:** Functional MR urography in patients with kidney transplantation. *Radiologe.*, 37(3): 233-238, (1997).
- Koelliker SL and Cronan JJ:** Acute urinary tract obstruction: Imaging update. *Urologic Clinics of North America*, 24(3): 571 – 582, (1997).
- Kramer LA:** Magnetic resonance imaging of renal masses. *World J Urol* 16: 22–28, (1998).
- Krestin GP:** Genitourinary MR kidneys and adrenal glands. *Eur Radiol*, 9: 1705-1714, (1999).
- Kuuliala IK, Niemi LK and Ala-Opas MY:** Ultrasonography for diagnosis of obstructing ureteral calculus. *Scand J Urol Nephrol.*, 22 (4): 275-277, (1988).
- Laing FC, Jeffrey RB Jr and Wing VW:** Ultrasound versus excretory urography in evaluating acute flank pain. *Radiology*, 154: 613–616, (1985).
- Lemaitre L, Claudon M, Fauquet I et al:** Imaging of chronic and intermittent adult upper urinary tract obstruction. *J Radiol.*, 85 (2 Pt 2): 197-216, (2004).
- Leppert A, Nadalin S, Schirg E et al:** Impact of magnetic resonance urography on preoperative diagnostic workup in children affected by hydronephrosis: should IVU be replaced? *J Pediatr Surg.*, 37 (10): 1441-1445, (2002).
- Li W, Chavez D, Edelman RR and Prasad PV:** Magnetic resonance urography by breath-hold contrast-enhanced three-dimensional FISP. *J Magn Reson Imaging.*, 7: 309-311, (1997).
-

Li X, Han Z, Wang W et al: Magnetic resonance urography in the diagnosis of urinary tract obstruction after renal transplantation. *Chin Med J (Engl)*., 115(4):540-542, (2002).

Louca G, Liberopoulos K, Fidas A et al: MR urography in the diagnosis of urinary tract obstruction. *Eur Urol.*, 35 (2): 102-108, (1999).

Low RN, Martinez AG, Steinberg SM et al: Potential renal transplant donors: evaluation with gadolinium-enhanced MR angiography and MR urography. *Radiology*, 207:165-172, (1998).

Low RN, Martinez AG, Steinberg SM et al: Potential renal transplant donors: evaluation with gadolinium-enhanced MR angiography and MR urography. *Radiology*, 207:165-172, (1998).

Lu Y, Hong W and Lu L: Diagnostic evaluation of magnetic resonance urography in urinary tract dilatation and obstruction. *Zhonghua Yi Xue Za Zhi.*, 78 (9): 670-673, (1998). Quoted - ?!

Maizels M, Zaontz MR and Firlit CF: Role of in-office ultrasonography in screening infants and children for urinary obstruction. *Urol Clin North Am.*, 17 (2): 429-435, (1990).

Massague JJ: The transforming growth factor-b family. *Annals of Biochemistry*, 6: 597- 641, (1990).

Miller OF, Rineer SK, Reichard SR et al: Prospective comparison of unenhanced spiral computed tomography and intravenous urogram in the evaluation of acute flank pain. *Urology*, 52: 982-987, (1998).

Moody TE, Vaughan ED Jr and Gillenwater JY: Comparison of the renal hemodynamic response to unilateral and bilateral ureteral occlusion. *Investig Urol*, 14: 455-459, (1977). Quoted from Pathophysiology of urinary tract obstruction In: Walsh PC, Vaughan ED Jr, Wein AJ, (eds). *Campbell's Urology*. 8th ed. Philadelphia, Pa: WB Saunders, Vol 1 pp: 411 - 462, (2002).

Moody TE, Vaughn ED Jr and Gillenwater JY: Relationship between renal blood flow and ureteral pressure during 18 hours of total unilateral ureteral occlusion. Implications for changing sites of increased renal resistance. *Investig Urol*, 13: 246–251, (1975). Quoted from Pathophysiology of urinary tract obstruction In: Walsh PC, Vaughan ED Jr, Wein AJ, (eds). *Campbell's Urology*. 8th ed. Philadelphia, Pa: WB Saunders, Vol 1 pp: 411 – 462, (2002).

Murano F: Ultrasonic evaluation of hydronephrosis during pregnancy and puerperium. *Gynecological Obstetrical Investigator*, 35: 94 – 98, (1993).

Mustonen S, Ala-Houhala I and Tammela TLJ: Proteinuria and renal function during and after acute urinary retention. *The Journal of Urology*, 161(6): 1781 – 1784, (1999).

Nolte-Ernsting C, Adam G, Bucker A et al: Contrast-enhanced magnetic resonance urography. First experimental results with a polymeric gadolinium bloodpool agent. *Invest Radiol.*, 32: 418-423, (1997).

Nolte-Ernsting C, Staatz G, Wildberger J and Adam G: MR-urography and CT-urography: principles, examination techniques, applications. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr.*, 175 (2): 211-222, (2003). *Quotation*.

Nolte-Ernsting CC, Adam GB and Gunther RW: MR urography: examination techniques and clinical applications. *Eur Radiol.*, 11(3): 355-372, (2001).

Nolte-Ernsting CC, Bucker A, Adam GB et al: Gadolinium-enhanced excretory MR urography after low-dose diuretic injection: comparison with conventional excretory urography. *Radiology*, 209: 147-157 (1998).

Nolte-Ernsting CC, Bucker A, Neuerburg JM et al: MR imaging – guided percutaneous nephrostomy and use of MR compatible catheters in non dilated porcine urinary tract. *J Vas Intervent Radiol.*, 10: 1305 – 1314, (1999).

Nolte-Ernsting CC, Krombach G, Staatz G et al: Virtual endoscopy of the upper urinary tract based on contrast-enhanced MR urography data sets. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr.*, 170 (6): 550-556, (1999).

- Rosi P, Virgili G, Distasi SM et al:** Diuretic ultrasound. A non-invasive technique for the assessment of upper tract obstruction. *Br J Urol* 65: 566-569, (1990).
- Rothepearl A, Frager D, Subramanian A et al:** MR urography: technique and application. *Radiology*, 194: 125-130, (1995).
- Roy C, Saussine C, Jahn C et al:** Evaluation of RARE-MR urography in the assessment of ureterohydronephrosis. *J Comput Assist Tomogr.*, 18 (4): 601-608, (1994).
- Salinas SAS, Guardiola MA, Moreno AJ and Server FG:** Puncture of urinary tract for the diagnosis of its obstruction. Review of 60 cases. *Arch Esp Urol.*, 42 (1):43-46, (1989).
- Schad LR, Semmler W, Knopp MV et al:** Preliminary evaluation: Magnetic resonance urography using a saturation inversion projection spin echo sequence. *Magn Reson Imaging*, 11: 319 – 327, (1993).
- Schubert RA, Gockeritz S, Mentzel HJ et al:** Imaging in ureteral complications of renal transplantation: value of static fluid MR urography. *Eur Radiol.*, 10 (7): 1152-1157, (2000).
- Schuhmann-Giampieri G and Krestin GP:** Pharmacokinetics of Gd-DTPA in patients with chronic renal failure. *Invest. Radiol*, 26: 975-979, (1991).
- Schulam PG, Kawashima A, Sandler C et al:** Urinary tract imaging-Basic principles In: Walsh PC, Vaughan ED Jr, Wein AJ, (eds). *Campbell's Urology*. 8th ed. Philadelphia, Pa: WB Saunders, Vol 1 pp: 122 – 166, (2002).
- Shokeir AA, El-Diasty T, Eassa W et al:** Diagnosis of noncalcareous hydronephrosis: role of magnetic resonance urography and noncontrast computed tomography. *Urology*. 63 (2): 225-229, (2004).
- Shokeir AA:** The diagnosis of upper urinary tract obstruction. *BJU International.*, 83: 893 – 901, (1999).
- Sigmund G, Stoeve B, Zimmerhackl LB et al:** RARE-MR-urography in the diagnosis of upper urinary tract abnormalities in children. *Pediatr Radiol.*, 21 (6): 416-420, (1991).

-
- Sigmund G, Stover B, Zimmerhackl LB et al:** RARE-MR urography in the diagnosis of upper urinary tract abnormalities in children. *Pediatr Radiol*, 21: 416-420, (1991).
- Smith RC and Coll DM:** Helical computed tomography in the diagnosis of ureteric colic. *BJU Int*, 86 (1): 33-41, (2000).
- Smith RC, Rosenfield AT, Choe KA et al:** Acute flank pain: comparison of non-contrast-enhanced CT and intravenous urography. *Radiology*, 194: 789-794, (1995).
- Smith RC, Verga M, McCarthy S and Rosenfield AT:** Diagnosis of acute flank pain: value of unenhanced helical CT. *AJR Am J Roentgenol*, 166: 97-101 (1996).
- Spencer JA, Chahal R, Kelly A et al:** Evaluation of painful hydronephrosis in pregnancy: magnetic resonance urographic patterns in physiological dilatation versus calculous obstruction. *J Urol*, 171 (1): 256-260, (2004).
- Staatz G, Nolte-Ernsting CC, Adam GB et al:** Feasibility and utility of respiratory-gated, gadolinium-enhanced T1-weighted magnetic resonance urography in children. *Invest Radiol*, 35: 504-512, (2000).
- Staatz G, Nolte-Ernsting CC, Haage P et al:** Gadolinium-enhanced T(1)-weighted MR urography versus T(2)-weighted (HASTE) MR urography in children. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr*. 173 (11): 991-996, (2001).
- Sudah M, Vanninen R, Partanen K et al:** MR urography in evaluation of acute flank pain: T2-weighted sequences and gadolinium-enhanced three-dimensional FLASH compared with urography. Fast low-angle shot. *AJR Am J Roentgenol*, 176(1): 105-112 (2001).
- Sudah M, Vanninen RL, Partanen K et al:** Patients with Acute Flank Pain: Comparison of MR Urography with Unenhanced Helical CT. *Radiology*, 223: 98-105, (2002).
- Swan SK, Lambrecht LJ, Townsend R et al:** safety and pharmacokinetics profilr of gadolinium in subjects with renal impairment. *Ivtest. Radiol*, 34: 443-448 (1999).
-

Szopinski K, Szopinska M, Borowka A and Jakubowski W: Magnetic resonance urography: initial experience of a low-dose Gd-DTPA-enhanced technique. *Eur Radiol.*, 10 (7): 1158-1164, (2000).

Talner LB: Specific causes of obstruction. In Pollack HM ed: *Clinical Urography*, vol. 2. Philadelphia, WB Saunders, pp: 1629-1751, (1990a).

Talner LB: Urinary Obstruction. In Grainger RG and Allison D. *Grainger & Allison's Diagnostic Radiology*. 3rd ed. New York, Churchill Livingstone, (1997).

Talner LB: Urinary obstruction. In Pollack H ed: *Clinical Urology: An Atlas and Textbook of Urological Imaging*. Philadelphia, WB Saunders, pp: 1535-1628, (1990b).

Tang Y, Yamashita Y, Namimoto T et al: The value of MR urography that uses HASTE sequences to reveal urinary tract disorders. *AJR Am J Roentgenol*, 167: 1497-1502, (1996).

Taylor A Jr and Eshima D: Effects of altered physiologic states on clearance and biodistribution of technetium-99m MAG3, iodine-131 OIH, and iodine-125 iothalamate. *J Nucl Med*, 29: 669-675, (1988).

Taylor A Jr and Nally JV: Clinical applications of renal scintigraphy. *AJR Am J Roentgenol*, 164: 31-41, (1995).

Taylor AJr and Nally JV: Clinical application of renal scintigraphy. *Am J Roentgen.*, 164: 31-41, (1995).

Thoeny HC and Hoppe H: Unenhanced spiral CT in urolithiasis: indication, performance and interpretation. *Rofo Fortschr Geb Rontgenstr Neuen Bildgeb Verfahr.*, 175 (7): 904-910, (2003).

Thomsen HS and Dorph S: The upper urinary tract. Pyelography and interventional procedures. *Acta Radiol.*, 28 (2): 129-137, (1987).

Truong LD, Sheikh-Hamad D, Chakraborty S and Suki WN: Cell apoptosis and proliferation in obstructive uropathy. *Seminars in Nephrology*, 18(6): 641 - 651, (1998).

Tublin ME, Dodd GD III and Verdile VP: Acute renal colic: diagnosis with duplex Doppler US. *Radiology*, 193: 697-701, (1994).

United States Renal Data System (USRDS). Annual data report: Incidence and prevalence of ESRD. *American Journal of Kidney Disease*, 18 (S2): S34 - S47, (1996).

Upsdell SM, Leeson SM, Brooman PJ and O'Reilly: Diuretic-induced urinary flow rates at varying clearances and their relevance to the performance and interpretation of diuresis renography. *Br J Urol.*, 61: 14-18, (1988).

Upsdell SM, Testa MJ and Lawson RS: The F-15 diuresis renogram in suspected obstruction of the upper urinary tract. *Br J Urol.*, 69: 126-131, (1992).

Verswijvel GA, Oyen RH, Van Poppel HP et al: Magnetic resonance imaging in the assessment of urologic disease: an all-in-one approach. *Eur Radiol.*, 10: 1614-1619, (2000).

Vieweg J, Teh C, Freed K et al: Unenhanced helical computerized tomography for the evaluation of patients with acute flank pain. *J Urol.*, 160: 679-684, (1998).

Vinee P, Stover B, Sigmund G et al: MRI RARE urography: an alternative to IVU. *Ann Radiol* 36: 109-113, (1993).

Watson G: Problems with double-J stents and nephrostomy tubes. *J Endourol.*, 11 (6): 413-417, (1997).

Webb JAW, Reznick RH, White FE et al: Can ultrasound and computed tomography replace high-dose urography in patients with impaired renal function? *QJ Med*, 53: 411-425, (1984).

Whitaker R: Perfusion pressure flow studies. In O'Reilly P, George N, Weiss R (eds): *Diagnostic Techniques in Urology*. Philadelphia, WB Saunders. Pp: 135-141, (1990).

Whitaker RH and Buxton-Thomas MS: A comparison of pressure flow studies and renography in equivocal upper urinary tract obstruction. *J Urol.*, 131: 446–449, (1984).

Whitaker RH: Methods of assessing obstruction in dilated ureters. *Br J Urol.*, 45: 15–22, (1973). Quoted from **Shokeir AA:** The diagnosis of upper urinary tract obstruction. *BJU International*, 83: 893 – 901, (1999).

Whitfield HN, Britton LE, Hendry WF et al: The distinction between uropathy and nephropathy by radioisotope transit times. *Br J Urol.*, 50: 433–8, (1978). Quoted from **Shokeir AA:** The diagnosis of upper urinary tract obstruction. *BJU International*, 83: 893 – 901, (1999).

Wille S, von Knobloch R, Klose KJ et al: Magnetic resonance urography in pediatric urology. *Scand J Urol Nephrol.*, 37 (1): 16–21, (2003).

Williams PL, Bannister LH, Berry MM et al: *Gray's Anatomy*, 38th ed. New York, Churchill Livingstone, pp: 1385 – 1410, (1995).

Wilson D and Klahr S: Urinary tract obstruction. In Schrier R, Gottschalk C (ed): *Diseases of the Kidney*, 5th ed. Boston, Little, Brown, pp: 657–688, (1993).

Yarger WE, Schocken DD and Harris RH: Obstructive nephropathy in the rat: Possible roles for the renin-angiotensin system, prostaglandins, and thromboxanes in postobstructive renal function. *J Clin Invest*, 65: 400–412, (1980).

Yarger WE: Urinary tract obstruction. In Brenner BM and Rector FC (eds): *The kidney*, 4th ed. Philadelphia, Pa: WB Saunders, pp: 1772 – 1793, (1991).

Yilmaz S, Sindel T, Arslan G et al. Renal colic: comparison of spiral CT, US and IVU in the detection of ureteral calculi. *Eur Radiol.*, 8: 212–217, (1998).

Youssefzadeh D, Katz DS and Lummerman JH: Unenhanced helical CT in the evaluation of suspected renal colic. *AUA updates* 28: 203–207, (1999).

Zerhau P, Kubatova J, Skotakova J and Mach V: Magnetic resonance urography as an indicator for surgical treatment of anomalies of the upper urinary tract. *Rozhl Chir.*, 82 (12): 638–644, (2003).

References

Zielonko J, Studniarek M and Markuszewski M: MR urography of obstructive uropathy: diagnostic value of the method in selected clinical groups. *Eur Radiol.*, 13 (4): 802-809, (2003).