
REFERENCES

- Ahern D.J. and Maher J.F. (1992):* Heart failure as a complication of hemodialysis arteriovenous fistula. *Ann. Intern. Med.* 77: 201-204.
- Ahmad S.; Blagg C.R.; and Scribner B.H. (1988):* Center and home chronic hemodialysis. In schrier, R.W., C.W. Gottschalk (Eds.): *Diseases of the kidney*, 4th ed., Little, Brown, Boston. 3281-3322.
- Anderson C.B.; Etheridge E.E.; Harter H.R. et al. (1997):* Blood flow measurements in arteriovenous fistulas. *Surgery.* 136: 292-297.
- Applebaum H.; Shashikumar V.I.; Somers I.A. et al. (1990):* Improved hemodialysis vascular access in children. *J. Pediat. Surg.* 15: 764-769.
- Bacchini G.; La-Milia V.; Andruli S. et al. (2001):* Color Doppler ultrasonography imaging to guide transluminal angioplasty of venous stenosis. 58(4): 1870-3. *ang.*
- Baier R.E.; Akers C.K. and Perlmutter S. (1986):* Processed human umbilical veins for vascular reconstructive surgery. *Trans Am. Soc. Artif. Intern. Organs.* 22: 514-526.
- Baird D.E.; Omdal D.G.; Yuill S.C. et al. (1994):* Doppler analysis of hemodialysis grafts: resistive index as a predictor of stenosis. *J. Ultrasound Med.* 13: 791-796.
- Baker L.D.; Johnson J.D.M.; Goldfarb D. (1986):* Expanded polytetrafluoroethylene (PTFE) subcutaneous arteriovenous conduit: An improved vascular access for chronic haemodialysis. *Trans Am. Soc. Artif. Intern. Organs.* 22: 3872.

Bandyk D.F.; Calligaro K.D.; Andros G. et al. (1993): Selective use of duplex ultrasound to replace preoperative arteriography for failing arterial vein grafts (discussion). J. Vasc. Surg. 27: 94-95.

Bandyk D.F.; Calligaro K.D.; Andros G. et al. (1998): Selective use of duplex ultrasound to replace preoperative arteriography for failing arterial vein grafts (discussion). J. Vasc. Surg. 27: 94-95.

Barnes R.W. (1982): Doppler ultrasonic angiography. In Bernstein FE (ed): "Non-invasive techniques in vascular disease. CV Mosby Company. St. Louis P. 168.

Barril G.; Besada E.; Cirugeda A. et al. (2001): Hemodialysis vascular assessment by ultrasound dilatation method. 32(3): 459-82.

Basmajian J.V. and Slonecker C.E. (1993): Grants-method of anatomy 7th edition by Williams and Wilkins, Hong Kong, London. Sydney. P. 257.

Beach K.W. and Strandness D. Jr. (1985): Pulsed Doppler ultrasound for blood velocity measurements. In Bernstein E.F. (ed). Non-invasive techniques in vascular disease. CV Mosby Company, St. Louis. P. 25.

Beathard G.A. (1992): Percutaneous transvenous angioplasty in the treatment of vascular access stenosis. Kidney Int. 42: 1390-1397.

Beathard G.A. (1993): Gianturco self-expanding stent in the treatment of stenosis in dialysis access grafts. Kidney Int. 43: 872-877.

Beathard G.A. (1994): Percutanenous therapy of vascular access stenosis and thrombosis. Semin Dial. 7: 165-167.

- Beathard G.A. (1995):** Percutaneous angioplasty for the treatment of venous stenosis: a nephrologist, view-Semin. Dial. 8: 166-170.
- Bender M.H.M. Bruynincks C.M.A. and Gerlag P.G.G. (1994):** The brachiocephalic elbow fistula: A useful alternative angioaccess for permanent haemodialysis. J. Vasc. Surg. 20: 808-813.
- Bennion R.S. and Williams R.A. (1982):** The radiocephalic fistula. Contemp Dial. 3: 12-16.
- Besarab A.; Sullivan K.L.; Ross R.P. and Moritz M.J. (1995):** Utility of intra-access pressure monitoring in detecting and correcting venous outlet stenosis prior to thrombosis. Kind. Int. 1364-1373.
- Bhat D.J.; Tellis V.A.; Kohl W.I. et al. (1980):** Management of sepsis involving expanded polytetrafluoroethylene grafts for hemodialysis access. Surgery 87: 445-450.
- Bimble K.S.; Rabbat C.G.; Schiff D. et al., (2002):** The clinical utility of Doppler ultrasound prior to arteriovenous fistula creation. Intern. Med. 5: 314-7.
- Black well R. (1985):** General scanning. In Clinical diagnostic ultrasound. 1st edition. Barnett E. and Marley P. (eds). Blackwell scientific publications. Oxford, London, Edinburgh, Boston. P. 3.
- Blankestijn P.J.; Bosman P.J. and Boereboom F.T.J. (1996):** Haemodialysis access without thrombosis. Is it possible? Nephrol. Dial. Transplant. 11: 2394-2397.
- Bonahmi U.; Civalleri D.; Rovidia S.; Adami G.F.; Gianetta E. and Griffanti-Batoli F. (1982):** Nine years experience with end-to-end arteriovenous fistula at the anatomical snuffbox for maintenance haemodialysis. Br. J. Surg. 69: 486-488.

Bone G.E. and Pomajzl M.J. (1989): Management of dialysis fistula thrombosis. *Am. J. Surg.* 138: 901-906.

Bone G.E. and Pomajzl M.J. (1990): Prospective comparison of polytetrafluoroethylene and bovine grafts for dialysis. *J. Surg. Res.* 29: 223-227.

Bourquelot P.; Wolfscher I. and Lamy I. (1991): Microsurgery for haemodialysis distal arteriovenous fistulae in children weighing less than 10kg. *Proc. Eur. Dial. Transplant Assoc.* 18: 537-541.

Bouthier J.D.; Levenson J.A.; Simon A.C.; Bariety J.M.; Bourquelot P.E. and Safar M.E. (1990): A noninvasive determination of fistula blood flow in dialysis patients. *Artif. Org.* 404-409.

Bracewell R.N. (1989): The Fourier transform. *Scientific American.* 260:86.

Brescia M.J.; Cimino J.E.; Appel K. and Burwich B.J. (1966): Chronic hemodialysis using venipuncture and surgically created arteriovenous fistula. *New Engl. J. Med.* 275: 1089-1092.

Brescia M.J.; Cimino J.E.; Appel K. and Hurwich B.J. (1986): Chronic haemodialysis using vein-puncture and a surgically created arteriovenous fistula. *N. Engl. J. Med.* 275: 1089-1092.

Brooks J.L.; Sigby R.D.; May K.J. and Mack R.M. (1987): Transluminal angioplasty versus surgical repair for stenosis of hemodialysis grafts. *Am. J. Surg.* 153: 530-531.

Broviac J.W.; Cole J.J. and Scribner B.H. (1997): Silicone rubber atrial catheter for prolonged parenteral nutrition. *Surg. Gynecol. Obstet.* 136: 602.

Bulter H.G.; Baker L.D. and Johnson J.M. (1987): Vascular access for chronic haemodialysis. Polytetrafluoroethylene (PTFE) versus bovine heterograft. Am. J. Surg. 134: 791-793.

Burdick J.F.; Scott W. and Coslml A.B. (1988): Experience with darcon graft arteriovenous fistulas for dialysis access. Ann. Surg. 187: 262-266.

Burger H.; Zijlstra J.J.; Kluchert S.A. et al. (1990): Percutaneous transluminal angioplasty improves longevity in fistulae 1-shunts for haemodialysis: Nephrol. Dial. Transplant. 5: 608-611.

Burns P.N. (1995): The physical principles of Doppler and spectral analysis. JCU. 15: 567.

Bussel J.A.; Abbott J.A. and Lim R.C. (1996): A radial steal syndrome with arteriovenous fistula for hemodialysis. Ann. Intern. Med. 75: 387-394.

Bussel J.A.; Abott J.A. and Linn R.C. (1997): A radial steal syndrome with arteriovenous fistula for hemodialysis. Ann. Ihtern. Med. 75: 387-394.

Butt K.M.H. (1998): Blood access. Clin. Nephrol. 9: 138-143.

Butt K.M.H.; Friedman E.A. and Kountz S.A. (1996): Angio-access. Curr. Probl. Surg. 13: 1-67.

Calligaro K.D.; Syrek J.R.; Dougherty M.J. et al. (1998): Selective use of duplex ultrasound to replace preoperative arteriography for failing arterial vein grafts. J. Vasc. Surg. 27: 89-94.

Caps M.T.; Bergelin R.O.; Cantwell-Gab K. et al. (1995): Vein graft lesions: time of onset and rate of progression. J. Vasc. Surg. 22: 466-474.

-
- Carty G.A. and Davis V.H. (1990):** Mid-graft stenosis in expanded polytetrafluoroethylene hemodialysis conduits. *Dial. Transplant.* 19: 486-489.
- Cerilli J. and Haisch C. (1986):** Vascular access procedures for renal dialysis. In : Sabiston D.C., ed. *Textbook of surgery.* Philadelphia. Saunders, pp. 429-433.
- Cimochovski G.F.; Worley E.; Rutherford R.F. et al. (1990):** Superiority of the internal jugular over the subclavian access for temporary dialysis. *Nephron.* 54: 154.
- Coburn M.C. and Carney W.F. (1994):** Comparison of basilic vein and polytetrafluoroethylene for brachial arteriovenous fistula. *J. Vasc. Surg.* 20: 896-902.
- Compy C.M. and Shapiro F.L. (1988):** Cardiac complications of regular dialysis therapy. in Drukker W., F.m. Parsons, J.F., Maher (eds.): *Replacement of renal function by dialysis.* 2nd ed. Martinus Nijhoff, Boston. 595-601.
- Conly J.M.; Grieves K. and Peters B. (1989):** A prospective randomized study comparing transparent and dry gauze dressings for central venous catheters. *J. Infect. Dis.* 159: 310-319.
- Dapont O.; Feurstein M.; Rendl K.H. and Prenner K. (1987):** Transluminal angioplasty versus conventional operation in the treatment of haemodialysis fistula stenosis. Results from a 5-year study. *Br. J. Surg.* 74: 1005.
- Dardik H.; Ibrahim I.M. and Sussman B. (1989):** Gluteraldehyd-estabilised umbilical vein prosthesis for revascularization of the legs. *Am. J. Surg.* 138: 234-237.
- Davis D.; Petersen J.; Feldman R. et al. (1984):** Subclavian venous stenosis. A complication of subclavian dialysis. *JAMA.* 252: 3404-3406.
-

- Depner T.A. and Reasons A. (1996):** Longevity of peripheral AV grafts and fistulae for haemodialysis is related to access blood flow (Abst.) J. Am. Soc. Nephrol. 7: 1405.
- Dettester J.; Vanholder R. and Ringole S. (1992):** Factors affecting catheter and technique survival in permanent silicone single lumen dialysis catheters (Abstract). J. Am. Soc. Nephrol. 3: 361.
- Dobkin J.F.; Miller M.H. and Stelgblgel N.H. (1987):** Septicemia in patients on chronic hemodialysis. Ann. Intern. Med. 88: 28-33.
- Doken W.; Landwehr P.; Frank F.F. and Lackner K. (1996):** Calculation of flow velocities from the geometrical shape of contrast induced time density curves using DSA. Europ. J. Radiol. 208-213.
- Dolmatch B.L.; Gray R.J. and Horton K.M. (1994):** Will iatrogenic pulmonary embolization be our pulmonary embarrassment? Radiology 191: 615-617.
- Dousset V.; Grenier N.; Douws C. et al. (1999):** Hemodialysis grafts. color Doppler imaging correlated with digital subtraction angiography and functional status. Radiology. 181: 89-94.
- Duncan H.; Ferguson I. and Faris I. (1986):** Incidence of the radial steal syndrome in patients with Brescia fistula for hemodialysis. Its clinical significance. J. Vasc. Surg. 4: 144-147.
- Duncan H.; Ferguson L. and Faris I. (1996):** Incidence of the radial steal syndrome in patients with Brescia fistula for hemodialysis its clinical significance. J. Vasc. Surg. 4: 144-147.
- Enkina T.N.; Riabov S.L.; Lukichev et al. (2002):** Course of heart failure in patients with terminal chronic kidney failure on chronic hemodialysis. 73(6): 15-20.

- Fee H.J. and Goldberg A.L. (1996):** Lower extremity ischemia after femoral arteriovenous boving shunts. *Ann. Surg.* 183: 42-45.
- Finlay D.E.; Longley D.G.; Foshager M.C. et al. (1993):** Duplex and color Doppler sonography of hemodialysis arteriovenous fistulas and grafts. *Radiographics.* 13: 983-999.
- Flores L.; Dunn I.; Frumkin E. et al. (1993):** Dacron arteriovenous shunts for vascular access in haemodialysis. *Trans Am. Soc. Artif. Intern. Organs.* 19: 33-37.
- Fruhwald F. and Blackwell D.E. (1992):** CCDS physical principles and technical considerations. In "Atlas of Color-Coded Doppler Sonography" Fruhwald F and Blackwell D.E. (eds). Spring Verlag, Wien, New York, Chapter I.
- Gaylord G.M. and Taber T.E. (1993):** Long-term hernodialysis access salvage. Problems and challenges for nephrologists and interventional radiologists. *JVIR* 4: 103-107.
- Gibson S.P. and Mosquera D. (1991):** Five years experience with Quinton Perm Cath for vascular access. *Nephrol. Dial. Transplant.* 6: 269-274.
- Gibson T.C.; Dyer D.P.; Postlethwaite R.J. and Gough D.C.S. (1987):** Vascular access for acute hemodialysis. *Arch. Dis. Child.* 62: 141-145.
- Gilbert M.S.; Robinson A.; Baez A. et al. (1988):** Carpal tunnel syndrome in patients who are receiving long-term renal hemodialysis. *J. Bone. Joint. Surg. (am)* 70: 1145.
- Gilula L.A.; Staple T.W.; Anderson C.B. and Anderson L.S. (1995):** Venous angiography of hemodialysis fistulas. *Radiology* 115: 555-562.

- Gmelin E.; Wintherhoff R. and Rinast E. (1999):** Insufficient hemodialysis access fistulas. Late results of treatment with percutaneous balloon angioplasty. *Radiology* 171: 657-660.
- Goldstein A. (1981):** Range ambiguities in real-time ultrasound. *J. Clin. Ultrasound*. 9: 83.
- Gooding G.A. and Effeney D.J. (1980):** Ultrasound of femoral artery aneurysms. *AJR. Am. J. Roentgenol.* 134: 477-480.
- Grant E.G. (1985):** Duplex ultrasonography : Its expanding role in noninvasive vascular diagnosis. *Radiol. Clin. North Am.* 23: 563.
- Grant J.P. (1972):** Grant's atlas of anatomy, 7th edition by Grant, J.P., the Williams and Wilking Co. Baltimore. 220.
- Gravin P.J.; Casteneda M.A. and Codd J.E. (1982):** Etiology and management of bovine graft aneurysms. *Arch. Surg.* 117: 281-284.
- Haimov M.; Burrows L.; Schanzer H. et al. (1985):** Experience with arterial substitutes in the construction of vascular access for hemodialysis. *J. Cardiovasc. Surg.* 21: 149-154.
- Haisch C. and Cerilli E.J. (1991):** Vascular access produres for renal dialysis (including peritoneal dialysis).in : sabsiston. D.C. (ed.0: Textbook of surgery 14th ed. Saunders (Philadelphia). 393-401.
- Harder F. and Landmann J. (1984):** Trends in access surgery for haemodialysis. *Surg. Annu.* 16: 135-149.
- Hedrick W.R. and Hykes M.S. (1988):** Doppler physics and instrumentation: A review. *Journal of Diagnostic* 4th edition. McGraw-Hill Book Company. 923-934.

- Helvie M.A. and Rubin J. (1989):* Evaluation of traumatic groin arteriovenous fistulas with duplex Doppler sonography. *J. Ultrasound Med.* 8: 21-24.
- Hiatt J.R. and Busuttil R.W. (1993):* A method for vascular access in small children. *Surgery* 93: 343-344.
- Higgins R.M. (1989):* Infections in a renal unit. *QJM.* 70: 41-51.
- Ishihava A.M. and Meyers C.H. (1998):* Longevity of haemodialysis arteriovenous shunts for haemodialysis. *Ann. Surg.* 168: 281-286.
- Jagar K.A.; Ricketts H.J. and Strandess D.E. Jr. (1985):* Duplex scanning for the evaluation of lower limb arterial disease. In Bernstein EF (ed): *Non-invasive diagnostic techniques in vascular disease.* St-Louis CV, Mosby: 619-631.
- Jagar K.A.; Ricketts H.J. and Strandness D.E. (1985):* Duplex scanning for the evaluation of lower limb arterial disease, in Bernstein, EF (ed) : *Noninvasive diagnostic techniques in vascular disease.* St. Louis CV, Mosby,pp. 619-631.
- Jamil Z.; O'Donnell J.A.; Merk E.A.and Hobson R.W. (1989):* A comparison of knitted Dacron velour and bovine heterograft for haemodialysis access. *J. Surg. Res.* 26: 423-429.
- Jenkins A.M.; Buist T.A.S. and Glover S.D. (1980):* Meidum-turn follow up of forty autogenous vein and forty polytetrafluoroethylene (Gortex) grafts for vascular access. *Surgery*, 88: 667-672.
- Jensen J.A.; Coodson W.H.; Omachi R.S. et al. (1987):* Subcutaneous tissue oxygenation falls during haemodialysis. *Surgery.* 107: 416-421.

- Johnson G. and Blythe W.B. (1990):** Hemodynamic effects of arteriovenous shunts used for hemodialysis. *Ann. Surg.* 171: 715-721.
- Johnson G.; Dart C.H.; Peters R.M. and Steele F. (1996):** The importance of venous circulation in arteriovenous fistula. *Surg. Gynecol. Obstet.* 123: 995-1000.
- Kanterman R.Y.; Vesely T.M.; Pilgram T.K.; Guy B.W.; Windus D.W. and Picus D. (1995):** Dialysis access grafts: Anatomic location of venous stenosis and results of angioplasty. *Radiology.* 195: 135-139.
- Kaplowitz L.G.; Comstock J.A.; Landwehr D.M. et al. (1988):** A prospective study of infections in hemodialysis patients. Patient hygiene and other risk factors for infection. *Infect. Control Hosp. Epidemiol.* 9: 534-541.
- Kherlakian G.M.; Roedersheimer L.R.; Arbaugh J.J. et al. (1986):** Comparison of autogenous fistula versus expanded polytetrafluoroethylene graft fistula for angioaccess in hemodialysis. *Am.J. Surg.* 152: 235.
- Kiannaert P.; Janssen F. and Hall M. (1993):** Elbow arteriovenous fistula (EAVF) for chronic hemodialysis in small children. *J. Pediatr. Surg.* 18: 116-119.
- Kiannaert P.; Vereestraten P.; Toussaint C. and van Gertruyden J. (1997):** Nine years experience with internal arteriovenous fistulas for hemodialysis. A study of some factors influencing the results. *Br. J. Surg.* 64: 242-246.
- Knezvic W. and Mastaglia F.L. (1984):** Neuropathy associated with Brescia-Cimino arteriovenous fistulas. *Arch. Neurol.* 41: 1184-1186.

- Kohler T.R.; Nance D.R.; Cramer M.M. et al. (1997):** Duplex scanning for diagnosis of aortoiliac and femoro-popliteal disease. A prospective study. *Circulation* 76: 1074-1080.
- Kolff W.J. (1995):** The first clinical experience with the artificial. *Ann. International. Med.* 62: 609-619.
- Kottle S.P.; Gonzalez A.C.; Macon E.J. and Fellner S.K. (1998):** Ultrasonographic evaluation of vascular access complications. *Radiology.* 129: 751-754.
- Kremkau F.W. (1990):** Doppler principles. *Semin. In : Reontgenology.* 27(1): 6.
- Kremkau F.W. (1990):** Doppler ultrasound: principles and instruments. First edition (edited by Kremkau F.W.). W.B. Sanders Company. Philadelphia, USA.
- Lemoson M.S.; Tordoir J.H.; VanDet R.J. et al. (2002):** Effects of venous cuff at the venous anastomosis of polytetrafluoroethylene grafts for haemodialysis vascular access. *Vas. Calci. Surg. Dec.* 32(6): 1156-63.
- Levin A.; Mason A.J.; Jindal K.K. et al. (1991):** Prevention of hemodialysis subclavian vein catheter infection by topical providoneiodine. *Kidney Int.* 40: 934-938.
- Levowitz B.S.; Flores L.; Dunn I. and Frumkin E. (1986):** Prosthetic arteriovenous fistula for vascular access in haemodialysis. *Am. J.Surg.* 132: 368-372.
- Lilly I.; Negheim D.; Mendez-Picon G. and Lee H.M. (1980):** Comparison between bovine heterografts and expanded PTFE grafts for dialysis access. *Am. Surg.* 46: 694-696.

- Lumsden A.B.; McDonald M.I.; Isiklar H.; Martin L.G.; Kiker D.; Harker I.A. et al. (1997):** Central venous stenosis in the hemodialysis patients. Incidence and efficacy of endovascular treatment. *Cardiovasc. Surg.* 5(5): 504-509.
- Lund G.B.; Trerotola S.O.; Sheel P.F. et al. (1996):** Outcome of tunneled hemodialysis catheters placed by radiologists. *Radiology.* 198: 467-472.
- Mahmutyazicioglu K.; Kesenci M. et al. (2000):** Hemodynamic changes in the early phase of artificially created arteriovenous fistula. *Intern. Med.* Vol. 16. Issue. 12813.
- Mattos M.A.; vanBemmelen P.S.; Hodgson K.J. et al. (1993):** Does correction of stenoses identified with color duplex scanning improve infrainguinal graft patency? *J. Vasc. Surg.* 17: 54-66.
- May J.; Tiller D. and Johnson J. et al. (1989):** Saphenous vein arteriovenous fistula in regular dialysis treatment. *N. Engl. J. Med.* 280: 770.
- Mennes P.A.; Gilula L.A.; Anderso C.B.; Etheredge E.E.; Weerts C. and Harter H.R. (1998):** Complications associated with arteriovenous fistulas in patients undergoing chronic haemodialysis. *Arch. Intern. Med.* 138: 1117-1121.
- Middleton W.D.; Erickson S. and Melson G.L. (1999):** Perivascular color artifacts: Pathologic significance and appearance on color Doppler US images. *Radiology.* 171: 647-652.
- Middleton W.D.; Picus D.D.; Marx M.V. and Melson G.L. (1989):** Color Doppler sonography of hemodialysis vascular access. Comparison with angiography. *AJR.* 152: 633-639.
- Middleton W.D.; Picus D.D.; Mary M.V. and Melson G.L. (1994):** Color Doppler sonography of hemodiaylsis vascular access:

Comparison with angiography. Amer. J. Roentgenol. 152: 633-639.

Mills J.L.; Harris E.J.; Taylor L.M. et al. (1990): The importance of routine surveillance of distal bypass grafts with duplex scanning: a study of 379 reversed vein grafts. J. Vasc. Surg. 12: 379-389.

Miranda C.L. (1997): Ultrasound derived flow volumes in the evaluation of hemodialysis vascular accesses: variability due to measurement technique. J. Vasc. Technol. 21(3): 155-159.

Moran M.R.; Radriguez J.M.; Boyero M.R. et al. (1985): Flow of dialysis fistulas. Nephron 40: 63-66.

Morgan A.P.; Knight D.C.; Tilney N.L. and Lazarus J.M. (1980): Femoral triangle sepsis in dialysis patients. Ann. Surg. 191: 460-464.

Moss A.H.; Vasilis B.S. and Holley J.L. (1990): Use of a silicone dual-lumen catheter with a Dacron cuff as a long-term vascular access for haemodialysis patients. Am. J. Kidney Dis. 16: 211-215.

Munda R.; First M.R.; Alexander J.W. et al. (1983): Polytetrafluoroethylene graft survival in hemodialysis. JAMA. 249: 219-222.

Nelson T.R. and Pertorius D.B. (1988): The Doppler signal. Where does it come from and what does it mean? A.J.R. 151: 439.

Netter F.H. (2000): Atlas of human anatomy. Upper limb. Neurovasculature. 452-53.

Nevins T.E. and Kjellstrand C.M. (1983): Hemodialysis for children. A review, Int. J. Pediatric Nephrology. 4: 155-169.

- O'Brien T.F. (1982):** Infection in dialysis and transplant patients. In : Tilney, N.L., Lazarus, J.M., eds. *Surgical Care of the Patients with Renal Failure*. Philadelphia. W.B. Saunders. 67-97.
- Oakes D.D.; Spees E.K.; Light J.A. and Flye M.W. (1988):** A three years experience using modified bovine arterial heterografts for vascular access in patients requiring haemodialysis. *Am. Surg.* 187: 423-429.
- Odland M.D.; Kelly P.H.; Ney A.L. et al. (1991):** Management of dialysis-associated steal syndrome complicating upper extremity arteriovenous fistulas. Use of intraoperative digital photoplethysmography. *Surgery* 110: 664-670.
- Offner g.; Aschendorff C.; Hoyer P.F. et al. (1988):** End stage renal failure. 14 years experience of dialysis and renal transplantation. *Arch. Disc. Child.* 63: 120-126.
- Ohnishi K.; Saito M. and Sato S. (1986):** Clinical utility of pulsed Doppler flowmetry in patients with portal hypertension. *Am. J. Gastrointestinal.* 81(1): 1.
- Oudenhoven I.F.I.J.; Pattynama P.M.T.; DeRoos A. et al. (1994):** Magnetic resonance. A new method for measuring blood flow in hemodialysis fistulas. *Kidney Int.* 45: 884-889.
- Owens M.L.; Stabile B.F.; Gahr J.E. and Wilson S.E. (1989):** Vascular grafts for haemodialysis. An evaluation of sites and materials. *Dial. Transplant.* 8: 521-525.
- Pagano D.; Green M.A.; Henderson M.J. et al. (1994):** Surveillance policy for early detection of failing arteriovenous fistula for hemodialysis. *Nephrol. Dial. Transplant.* 9: 277-279.
- Pagano D.; Green M.A.; Henderson M.J.; Kmiot W.A. and Goldman M.D. (1994):** Surveillance policy for early detection of failing

arteriovenous fistulae for hemodialysis. *Nephrol. Dial. Transplant* 9: 277-279.

Palder S.B.; Kirkman R.I.; Wittemore A.D. et al. (1985): Vascular access for haemodialysis. Patency rates and results of revisions. *Ann. Surg.* 202: 235-239.

Pareek S.K. and Malhotra V. (2001): Angioaccess for haemodialysis. *Jul.* 99(7): 382-4.

Passman M.A.; Criado E.; Farber M.A.; Gooff L. et al. (1998): Efficiency of color flow duplex imaging for proximal upper extremity venous outflow obstruction in hemodialysis patients. *J. Vasc. Surg.* 28: 869-875.

Philips D.J. and Strandness D. Jr. (1985): Duplex scanning: Practical aspects of instrument performance. In Bernstein EF (ed) ; Non invasive diagnostic techniques in vascular disease. The CV mosby Company. Saint Louis. 397.

Philips D.J. et al. (1985): Detection of peripheral vascular disease using the Duplex Scanner III. *Ultrasound Med. Biol.* 63: 205.

Pietura R.; Zaluska W. Janicki K. et al. (2001): Chronic venous occlusion in hemodialysis- related fistula: diagnosis and endo vascular treatment. 58(6): 474-8.

Polak J.F. (1995a): Color flow and duplex sonography in lower extremity ischaemia. In : Clinical applications of Doppler ultrasound, Taylor, K.J.W.; Burns P.N. and Wells P.N.T. (eds. 2nd ed.), Raven Press Ltd. New York. 337-353.

Polak J.F. (1995b): Peripheral arterial disease evaluation with color flow and Doppler sonography. *Rad. Clin. North Am.* 33: 71-90.

Polak J.F.; Dobkin G.R.; O'Leary D.H. et al., (1989a): Internal carotid artery stenosis. Accuracy and reproductibility of color-Doppler-assisted duplex imaging. *Radiology.* 173: 793-798.

Polak J.F.; Donaldson M.C.; Dobkin G.R. et al. (1990a): Early detection of saphenous vein arterial bypass stenosis by color-assisted duplex sonography. A prospective study. *A.J.R. Am. J. Roentgenol.* 154: 857-861.

Polak J.F.; Donaldson M.C.; Whittemore A.D. et al. (1989b): Pulsatile masses surrounding vascular prostheses. Real-time US color flow imaging. *Radiology.* 170: 363-366.

Polak J.F.; Karmel M.I.; Mannick J.A. et al. (1990b): Determination of the extent of lower-extremity peripheral arterial disease with color-assisted duplex sonography. Comparison with angiography. *AJR. Am. J. Roentgenol* 155: 1085-1089.

Powid R.L. (1984): Color flow imaging : understanding its science and technology. *Journal of Diagnostic Medical Sonography.* 4: 236.

Quinton W.E.; Diller D. and Schriber B.H. (1990): Cannulation of blood vessels for prolonged haemodialysis. *Trans. Am. Soc. Artif. Intern. Organs.* 6: 104-113.

Rae R.E. (1995): Vascular sonography, principles and instrumentation. Chapter 19. In : *Textbook of diagnostic ultrasonography*, ed. By Sandera, L.H. Four edition, London, Philadelphia and Toronto.

Ranke C.; Creutzig A. and Alexander K. (1992): Duplex scanning of the peripheral arteries. Correlation of the peak systolic velocity ratio with angiographic diameter reduction. *Ultrasound in Med. Biol.* 18: 433-440.

- Reilly D.T.; Wood R.F.M. and Bell P.R.F. (1985):** Prospective study of dialysis fistulas. Problem patients and their treatment. *Br. J. Surg.* 69: 549-553.
- Rittgers S.E.; Garcia Valdez C.; McCormick I.T. and Posner M.P. (1986):** Noninvasive blood flow measurement in expanded polytetrafluoroethylene grafts for haemodialysis access. *J. Vasc. Surg.* 3: 633-642.
- Robin M.L.; Oser R.R.; Allon M.; Clements M.W. et al. (1998):** Hemodialysis access graft stenosis. US detection. *Radiology.* 208: 655-661.
- Rogers A.W. (1992):** Textbook of anatomy, Eighteen edition. Churchill. Livingstone, London. P. 619.
- Ronald O.; Bude M.D. and Rabin J.M. (1996):** Power Doppler sonography. *Radiology.* 200: 21-23.
- Roubidoux M.A.; Hertzberg B.S.; Carroll B.A. et al. (1990):** Color flow and image-directed Doppler ultrasound evaluation of iatrogenic arteriovenous fistulas in the groin. *J. Clin. Ultrasound* 18: 463-469.
- Rubin J.M.; Bude R.O.; Carson P.L.; Bree R.L. and Alder R.L. (1994):** Power Doppler ultrasonography a potentially useful alternative to mean ultrasonography frequency-based color Doppler ultrasonography. 553-856.
- Ryan S.P. and McNicholas M.M.J. (1994):** Anatomy for diagnostic imaging; W.B. Saunders Company Ltd. London. Philadelphia. P. 151-153.
- Sacks D. (1992):** Peripheral arterial duplex ultrasonography. *Seminars in Roentgenology.* Vol. XXVII, No. 1, 28: 38. January.

-
- Sacks D.; Robinson M.L.; Marinelli D.L. et al. (1992):** Peripheral arterial Doppler ultrasonography. Diagnostic criteria. J. Ultrasound Med. 11: 95-103.
- Safa A.A.; Val J.I.K.; Roberts A.C.; Ziegler T.W.; Hye R.J. and Oglevie S.B. (1996):** Detection and treatment of dysfunctional hemodialysis access grafts. Effect of a surveillance program on graft patency and the incidence of thrombosis. Radiology. 199: 653-657.
- Sands J.J. and Kapsick B. (1999):** Assessment of haemodialysis access performance by colour flow Doppler ultrasound. J. Biomaterials Applications. 13: 225-237.
- Saunders R.C. (1991):** Doppler principles. In Saunders RC (ed) "Clinical sonography a practical guide" 2nd edition. Little Brown Company, Boston, Toronto, London, p. 25.
- Schwab S.J. (1991):** Hemodialysis vascular access. In Jacobson, H.R.; Striker G.E. and Klahr S. eds. The principles and practice of Nephrology. Philadelphia. BC Decker 766-772.
- Schwab S.J. and Berkoben M.S. (1995):** Vascular access for hemodialysis. In : Nissenson, A.R.; Fine, R.N. and Gentile D.E., eds. Clinical dialysis. 3: 26-45.
- Schwab S.J.; Raymond J.R.; Saeed M. et al. (1989):** Prevention of hemodialysis fistula thrombosis: early detection of venous stenosis. Kidney Int. 36: 707-711.
- Schwab S.J.; Raymond J.R.; Saeed M. et al. (1989):** Prevention of hemodialysis fistula thrombosis. Early detection of venous stenosis. Kidney Int. 36: 707-711.
- Semer N.B.; Goldberg N.H. and Cuono C.B. (1989):** Upper extremity entrapment neuropathy and tourniquet use in patients undergoing hemodialysis. J. Hand Surg. (Am). 14: 897.
-

- Shakeleton C.R.; Taylor D.C.; Buckley A.R. et al. (1987):** Predicting failure in polytetrafluoroethylene vascular access grafts for hemodialysis. A pilot study. *Cand. J. Surg.* 30: 442-444.
- Shang K.K.; Cloutier G. and Lim C.C. (1992):** the effects of hematocrit, shear rate and turbulence on ultrasound Doppler spectrum from blood. *IEEE Trans Biomed Eng.* 39: 462-469.
- Sherman R.A. (1993):** The measurement of dialysis access recirculation. *Ann. Kidney Dis.* 22: 616-621.
- Shillinger F.; Schillinger D.; Montagnae R. and Milcent R. (1991):** Post-catheterization vein stenosis in hemodialysis comparative angiographic study of fifty subclavian and fifty internal jugular access. *Nephrol. Dial. Transplant.* 6: 722-724.
- Silva M.B.; Hobson R.W.; Pappas P.J. et al. (1997):** A strategy for increasing use of autogenous hemodialysis access procedures. Impact of preoperative noninvasive evaluation. *J. Vasc. Surg.* 27: 302-308.
- Spencer M.P. and Reid J.M. (1989):** Quantitation of carotid stenosis with continuous wave (C-W) Doppler ultrasound. *Stroke* 10: 326-330.
- Stehbens W.F. and Kanmody A.M. (1995):** Venous atherosclerosis associated with arteriovenous fistulas for hemodialysis. *Arch. Surg.* 110: 176-180.
- Storey B.G.; George C.R.P.; Stewart J.O.H. et al. (1989):** Embolic and ischemic complications after anastomosis of radial artery to cephalic vein. *Surg.* 66: 325-327.
- Strandnen Jr. (2001):** Vein graft surveillance. Duplex scanning in vascular disorders. 18: 296.

- Strandness D. Jr. (1985):** Doppler ultrasonic techniques. In : Bernstein E.F. (ed). Noninvasive diagnostic techniques in vascular disease. The CV Mosby Comp. St. Louis, Chapter 2: 18.
- Strandness D.E. (1992):** The role of non invasive procedures in the management of extremity arterial disease, Chapter 12. In : Introduction to vascular ultrasonography; ed. By Zwiebel, W.J. 3rd edition. W.B. Saunders Company. 176.
- Strandness D.Jr. (1985):** Non-invasive evaluation of arteriosclerosis: Comparison of methods. Arteriosclerosis. 3: 103-116.
- Suhocki P.; Conlon P.; Knelson M. et al. (1996):** Silastic cuffed catheters for hemodialysis vascular access. Thrombolytic and mechanical correction of malfunction. Am. J. Kidney Dis. 28: 379-386.
- Surratt R.S.; Picus D.D. and Hicks M.E. (1991):** The importance of preoperative evaluation of the subclavian vein in dialysis access planning AJR. 156: 623-625.
- Swedberg S.H.; Brown B.G.; Sigley R.T.; Wight T.U.; Gordon D. and Nicholls S.C. (1989):** Intimal fibromuscular hyperplasia at the venous anastomosis of PTEF grafts in hemodialysis patients. Circulation 80: 1726-1736.
- Taylor K.J.W. and Halland S. (1990):** Doppler U.S. part I. Basic principles, instrumentation and pitfalls. Radiology. 174: 297-307.
- Taylor K.J.W. and Holland S. (1990):** Doppler US part I. Basic principles, instrumentation and pitfalls. Radiology. 174: 297-307.
- Thomas S. and Zierler R.E. (1991):** Color Doppler characteristics in normal lower extremity arteries. Ultrasound, Med. Biol. 18(2): 167.

- Tordoir J.H.; deBruin H.G.; Hoeneveld H. et al. (1989):** Duplex ultrasound scanning in the assessment of arteriovenous fistulas created for haemodialysis access. Comparison with digital subtraction angiography. *J. Vasc. Surg.* 10: 122-128.
- Tortora G.I. (1989):** Principles of human anatomy 5th edition Harper and Publishers, New York. P. 380.
- Trerotola S.O.; Scheel P.J. and Powe N.R. et al. (1996):** Screening for access graft malfunction. Comparison of physical examination with US. *J. Vasc. Intern. Radiol.* 7: 15-20.
- Tullis M.J.; Primozich J.; Strandness D.E.Jr (1997):** "Detection of functional" valves in reversed saphenous vein bypass grafts: identification with duplex ultrasonography. *J. Vasc. Surg.* 25: 522-527.
- Turmel-Rodrigues I.; Pengloan J.; Blanchier D. et al. (1993):** Insufficient dialysis shunts. Improved long-term patency rates with close hemodynamic monitoring repeated percutaneous balloon angioplasty, and stent placement. *Radiology* 187: 273-278.
- Valji K.; Bookstein J.J.; Roberts A.C. and Davis G.B. (1991):** Pharmacomechanical thrombolysis and angioplasty in the management of clotted hemodialysis grafts. Early and late clinical results. *Radiology* 178: 243-247.
- VanStone J.C. and Daugirdas J.T. (1994):** Haemodialysis and physiologic principles, in *Handbook of Dialysis*. 2nd ed., Daugirdas, J.T. and Tod S. Ing. Little Brown and Company, Boston, New York, London.
- Vorwerk D.; Gunther R.W.; Mann H. et al. (1995):** Venous stenosis and occlusion in hemodialysis shunts. Follow-up results of stent placement in 65 patients. *Radiology* 195: 140-146.

- Wesley S. and Moore W.S. (1998):* Vascular surgery. A comprehensive review. 5th ed. W.B. Saunders Company. A Division of Harcourt Brace and Company. Philadelphia.
- Williams P.L.; Warwick R.; Dyson M. and Bannister L.H. (1989):* Gray's anatomy 37th edition. Churchill Livingstone, New York. P. 758.
- Wilson S.E.; Stabile B.E.; Williams R.A. and Owens M.L. (1982):* Current status of vascular access techniques. Surg. Clin. North Am. 62: 531-551.
- Winsett O.E. and Wolma F.J. (1985):* Complications of vascular access for hemodialysis. South Med. J. 78: 513-517.
- Woodcock J.P. and Baird R.N. (1983):* Upstream stenosis- its diagnosis by Doppler signal from the femoral artery. Arch. Surg. 115: 1316-1319.
- Zagzebski J.A. (1992):* Physics and instrumentation in Doppler and M. Mode ultrasonography. Chapter 2. In : Introduction to vascular ultrasonography; ed. By Zwiebel, W.J. 3rd edition. W. B. Sanders Company. 25-28.
- Zawada Edward T. Jr. (1994):* Indications for dialysis. Handbook of dialysis 2nd edition, Daugirdas, J.T. and Tod S. Ing. Little Brown and Company. Boston, New York, Toronto, London.
- Zwiebel W.J. (1992):* Spectrum analysis in Doppler vascular diagnosis. Chapter 3. In Introduction of vascular ultrasonography, ed. By : Zwiebel, W.J., 3rd ed. W.B. Saunders Company. 45-65.