
REFERENCES

Andrish, J.T. (1985):

Ligament injuries of the knee. Orthop. Clin. North Am., 16: 273.

Apley, A.G. and Solomon, L. (1982):

Fractures and joint injuries in Apley's system of Orthopaedics and fractures, ELBS 1 Butterworth, P. 435-478.

Applegate, G.R.; Flannigan, B.D.; Tolin, B.S. et al. (1993):

MR diagnosis of recurrent tears in the knee: value of intraarticular contrast material. AJR Am. J. Roentgenol., 161: 821.

Araki, Y.; Ootani, F.; Tsukaguchi, I. et al. (1992):

MR diagnosis of meniscal tears of the knee: Value of axial three-dimensional Fourier transformation GRASS images. AJR Am. J. Roentgenol., 158: 587.

Arnoczky, S.P. (1982):

Anatomy of the anterior cruciate ligament. Clin. Orthop., 172: 19.

Arnoczky, S.P. (1992):

Gross and vascular anatomy of the meniscus and its role in meniscal healing, regeneration, and repair. In Mow VC, Arnoczky SP, Jackson DW (eds): knee Meniscus: Basic and Clinical Foundations. New York, Raven Press.

Arnoczky, S.P.; Warren, R.F. (1994):

Anatomy of the cruciate ligaments. In Feagin JA (ed): The Crucial Ligaments. Diagnosis and Treatment of Ligamentous Injuries About the Knee, ed. 2. New York, Churchill Livingstone, p. 269.

Bach, B.A.; Warren, R.F. (1989):

"Empty wall" and "vertical strut" signs of ACL insufficiency. Arthroscopy, 5: 137

Baker, H.L.; Berquist, T.H.; Kispert, D.B. et al. (1985):
Magnetic resonance imaging on a routine clinical setting. Mayo Clin. Proc., 60: 75-75.

Barber, F.A. (1992):
What is the terrible triad? Arthroscopy 8:19.

Barnett, M.J. (1993):
MR diagnosis of internal derangements of the knee: Effect of field strength on efficacy. AJR, AM. J. Roentgenol., 161: 115.

Barry, K.P.; Mesagarzadeh, M.; Moyer, R. et al. (1991):
Pattern and accuracy of diagnosis of anterior cruciate ligament tears with MR imaging. Radiology, 181: 303.

Bennett, J.D.; Munk, P.L. (1992):
The patello-femoral joint and quadriceps mechanism In: Munk PL, Helmus CA and Vellet AD: MRI of the knee. An. Aspen. Publications, P. 41-55.

Berquist, T.H. (1988):
Magnetic resonance imaging of musculoskeletal system. In: Whrli FW, Show D and Kneeland J.B.: Biochemical Magnetic Resonance Imaging, Principles, Methodology and Applications, VCH, New York, p. 383-417.

Bessette, G.C. (1992):
The meniscus. Orthopedics 15:35.

Bianchi, M. (1983):
Acute tears of the PCL: Clinical study and results of operative treatment in 27 cases. Am. J. Sports. Med., 11: 308.

Bradley, W.G.; Kortman, K.E. and Grues, J.V. (1985):
Central nervous system. High-resolution magnetic resonance imaging: Effect of imaging spatial resolution on resolving power, Radiology, 156: 93-98.

Brown, J.J.; Malchow, S.C. and Totty, W.G. (1991):
MR examination of the knee: Interpretation with multiscreen digital workstation vs hardcopy format. AJR Am. J. Roentgenol., 157: 81.

-
- Brown, S.M.; Muroff, L.R. and Bradley, W.G. (1992):**
Ultrafast Kinematic MR imaging of the knee: increased sensitivity with a quadriceps Loading Device" in works in progress supplement to annual meeting program. Chicago, Society for Magnetic Resonance imaging, 29.
- Buckwalter, K.A.; Braunstein, E.M. and Janizek, D.B. (1993):**
MR imaging of meniscal tears: Narrow vs conventional window width photography. Radiology, 187: 827.
- Burk, D.L.; Mitchell, D.G.; Rifkin, M.D. et al. (1990):**
Recent advances in magnetic resonance imaging of the knee, Radiologic Clinics of North America, 28: 374-393.
- Burk, D.L.Jr.; Dalinka, M.K. and Kanal, E. (1988):**
Meniscal and ganglion cysts of the knee: MR evaluation. AJR Am. J. Roentgenol., 150: 331.
- Burk, D.L.; Kanal, E.; Brunberg, J.A. et al. (1986):**
1.5 T. surface coil MRI of the knee.
ARJ, 147: 293-300.
- Burks, R.T. (1990):**
Gross anatomy. In Daniel D, Akeson W, O'Connor J (eds): Knee Ligaments: Structure, Function, Injury, and Repair. New York, Raven Press, p. 59.
- Burnett, G.M.; Fowler, P.J. (1985):**
Reconstruction of the anterior cruciate ligament: Historic overview. Orthop. Clin. North Am., 16: 143.
- Burstein, D.B.; Fritts, H.M. and Fischer, D.A. (1991):**
Diagnosis of meniscal and cruciate ligament tears using MRI: Evaluation of two scanning techniques. Minn. Med., 74: 29.
- Chew, F.S. (1990):**
Medial meniscal flounce: Demonstration on MR imaging of the knee (letter). AJR. Am. J. Roentgenol., 155: 199.
- Cobby, M.J.; Schweitzer, M.E. and Resnick, D. (1992):**
The deep lateral femoral notch: An indirect sign of a torn anterior cruciate ligament. Radiology, 184: 855.
-

Crotty J.M., Monu M.B. and Pope T.L. (1996):

Magnetic Resonance Imaging of the Musculoskeletal System, part 4, the knee Clinical Orthop. Rad Res., 288-303.

Crues, J.V.III, Ryu, R.; Morgan, F.W. (1995):

Meniscal pathology: The expanding role of magnetic resonance imaging. Clin. Orthop., 252: 80.

Dandy, D.J. (1990):

The arthroscopic anatomy of symptomatic meniscal lesions. J. Bone, Joint Surg., 72B: 628.

De Smet, A.A.; Graf, B.K. (1994):

Meniscal tears missed on MR imaging: Relationship to meniscal tear patterns and anterior cruciate ligament tears. AJR, Am. J. Roentgenol., 162: 905.

De Smet, A.A.; Norris, M.A.; Yandow, D.R. (1993):

Diagnosis of meniscal tears of the knee with MR imaging: Effect of observer variation and sample size on sensitivity and specificity. AJR Am. J. Roentgenol., 160: 555.

De Smet, A.A.; Tuite, M.J. Norris, M.A. (1994):

MR diagnosis of meniscal tears: Analysis of causes of errors. AJR Am. J. Roentgenol., 163: 1419.

DeHaven, K.E. (1990):

Decision-making factors in the treatment of meniscus lesions. Clin. Orthop., 252: 50.

Delee, J.; Riely, M. and Rockwood, C. (1983):

Acute straight later instability of the knee. Am. J. Sports Med., 11: 404.

Deutsch, A.L. and Mink, J.H. (1992):

Magnetic resonance imaging of knee trauma. In: Bloem J.L. and Sartoris, D.J.: MRI and CT of the Musculoskeletal System. A text-Atlas, Williams and Wilkins, P. 421-441.

Deutsch, A.L.; Mink, J.H. and Fox, J.M. (1990):

Peripheral meniscal tears: MR findings after conservative treatment or arthroscopic repair. Radiology, 176: 485.

Disler, D.G.; Kattapuram, S.V. and Chew, F.S. (1993):

Meniscal tears of the knee: Preliminary comparison of three-dimensional MR reconstruction with two-dimensional MR imaging and arthroscopy. *AJR Am. J. Roentgenol*, 160: 343.

Duncan, J.B.; Hunter, R. and Purnell, M. (1995):

Meniscal injuries associated with acute anterior cruciate ligament tears in alpine skiers. *Am. J. Sports Med.*, 23: 170.

Evancho, A.M.; Fajman, W.A.; Stiles, R.G. (1990):

Magnetic resonance imaging of knee menisci: Comparison of spin echo pulse sequences. *Invest. Radiol.*, 2: 1224.

Farley, T.E.; Howell, S.M. and Love, K.F. (1991):

Meniscal tears: MR and arthrographic findings after arthroscopic repair. *Radiology*, 180: 517.

Finsterbush, A.; Frankl, V.; Matan, Y. et al. (1990):

Secondary damage to the knee after isolated injury of the anterior cruciate ligament. *Am. J. Sports Med.*, 18: 475.

Firooznia, H. (1992):

Knee in Firooznia H., Golimbu C.N., Rofii: MRI and CT of the Musculoskeletal System. Mosby Year Book, P. 661-797.

Fischer, S.P.; Fox, J.M. and Del Pizzo, W. (1991):

Accuracy of diagnosis from magnetic resonance imaging of the knee: A multi-center analysis of one thousand and fourteen patients. *J. Bone Joint. Surg.*, 73A:2.

Fisher, M.R.; Baker, B. and Amparo, E.G. (1985):

MR imaging using specialized coils. *Radiology*, 157: 443-447.

Fitzgerald, S.w.; Remer, E.M.; Friedman, H. et al. (1993):

MR evaluation of the anterior cruciate ligament: Value of supplementing sagittal images with coronal and axial images. *AJR Am. J. Roentgenol.*, 160: 1167.

Fitzgibbons, R.E.; Shelbourne, K.D. (1995):

"Aggressive" non-treatment of lateral meniscal tears seen during anterior cruciate ligament reconstruction. *Am. J. Sports Med.*, 23: 156.

Fowler, P.J.; Lubliner, J.A. (1989):

The predictive value of five clinical signs in the evaluation of meniscal pathology. *Arthroscopy*, 5: 184.

Fulkerson, J.P. and Hungerford, D.S. (1990):

Disorders of the patellofemoral joint, 2nd ed. Baltimore: Williams & Wilkins.

Garvin, G.J.; Munk, P.L. and Vellet, A.D. (1993):

Tears of the medial collateral ligament: Magnetic resonance imaging findings and associated injuries. *J. Can. Assoc. Radiol.*, 44: 199.

Gentili, A.; Seeger, L.L.; Yao, L. et al. (1994):

Anterior cruciate ligament tear: Indirect signs at MR imaging. *Radiology*, 193: 835.

Clashow J.L., Kat Z.R., Schneider M. et al., (1989):

Double blind assessment of the value of magnetic resonance imaging with diagnosis of anterior cruciate and mensicall lesions, *J. Bone-Joint surgery {Am.}* ; 71-A:113-119.

Grover, J.S.; Bassett, L.W.; Gross, M.L. et al. (1990):

Posterior cruciate ligament: MR imaging. *Radiology*, 174: 527.

Haggar, A.M.; Froelich, J.W.; Hearshen, D.O. et al. (1988):

Meniscal abnormalities of the knee: 3DFT fast-scan GRASS MR imaging. *AJR*, 150: 1341-1344.

Hajek, P.C.; Gylys-Morin, V.M. and Baker, L.L. (1987):

The high-signal intensity meniscus of the knee: Magnetic resonance evaluation and in vivo correlation. *Invest. Radiol.*, 22: 883.

Hall, F.M. (1991):

Buckled meniscus revisited (letter). *AJR Am. J. Roentgenol.*, 156: 200.

Haramatie, N.; Staron, R.B. and Rubin, S. (1993):

The flipped meniscus sign. *Skeletal Radiol.*, 22: 273.

Helms, C.A. (1995):

MR imaging of the knee in Helms CA: Fundamentals of skeletal radiology, W.B. Saunders Company, Philadelphia, P. 172-195.

Herman, L.J.; Beltran, J. (1988):

Pitfalls in MR imaging of the knee. Radiology, 167: 775.

Heron, C.W.; Calvert, P.T. (1992):

Three-dimensional gradient-echo MR imaging of the knee: Comparison with arthroscopy in 100 patients. Radiology, 183: 839.

Hodler, J.; Haghighi, P. and Pathria, M.N. (1992):

Meniscal changes in the elderly: Correlation of MR imaging and histologic findings. Radiology, 184: 221.

Hodler, J.; Haghighi, P.; Trudell, D. et al. (1992):

Thecruciate ligaments of the knee: Correlation between MR appearance and gross and histologic findings in cadaveric specimens. AJR Am. J. Roentgenol., 159: 357.

Howe, J.; Johnson, R.J. (1985):

Knee injuries in skiing. Orthop. Clin. North Am., 16: 303.

Howell, S.M.; Clark, J.A. (1992):

Tibial tunnel placement in anterior cruciate ligament reconstruction and graft impingement. Clin. Orthop., 283: 187.

Howell, S.M.; Knox, K.F.; Larley, I.E. et al. (1995):

Revascularization of a human anterior cruciate ligament graft during the first two years of implantation. Am. J. Sports. Med., 23: 42.

Hughston, J.C.; Andrew, J.R.; Cross, M.J. et al. (1976):

Classification of knee ligament instabilities. Part II. The lateral compartment. J. Bone Joint Surg., 58A: 173.

Hughston, J.C.; Andrews, J.R.; Cross, M.J. et al. (1976):

Classification of knee ligament instabilities. Part I. The medial compartment and cruciate ligaments. J. Bone Joint Surg., 58A: 159.

Irizarry, J.M.; Recht, M.P. (1997):

MR imaging of the knee ligaments and the postoperative knee, Radiologic clinics of North America. Volume 35, Number 1.

Johnsone R.J. (1988):

Prevention of cruciate ligament injuries In: Feagin JA: The cruciate ligaments: Diagnosis and treatment of ligamentous injuries about the knee, New Yourk, p. 189-222.

Jorgensen, U.; Sonne-Holm, S.; Lauridsen, F. et al. (1987):

Long-term follow up of meniscectomy in athletes: A prospective longitudinal study. J. Bone Joint Surg., 69B: 80.

Justice, W.W.; Quinn, S.F. (1995):

Error patterns in the MR imaging evaluation of menisci of the knee. Radiology, 196: 617.

Kannus, P. (1988):

Long term results of conservatively treated medial collateral ligament injuries of the knee. Clin. Orthop., 226: 103.

Kaplov S.R., Tersi L.M., Bradley W.G et al., (1993):

Bone contusions of the knee: Increased lesion detection with fast spin-echo MR imaging with spectroscopic fat saturation, Radiology, 189:901-904.

Kerlan, R.K. and Glosuman, R.E. (1988):

Tibial collateral ligament bursitis. Am. J. Sports Med., 16: 344.

Kimori, K.; Suzu, F. and Yamashita, F. (1989):

Evaluation of arthrography and arthroscopy for lesions of the posteromedial corner of the knee. Am. J. Sports Med., 17: 638.

Kneeland, J.B.; Hyde, J.S. (1989):

High-resolution MR imaging with local coils. Radiology, 171:1.

Kornick, J.; Trefelner, E. and McCarthy, S. (1990):

Meniscal abnormalities in the asymptomatic population at MR imaging. Radiology, 177: 463.

Langer J.E., Meyer S.J.F. and Dalinka M.K. (1990):

Imaging of the knee, Radiologic clinics of North America, vol. No. 5, p. 976-990.

Lee, J.K.; Yao, L.; Phelps, C.T. et al. (1988):

Anterior cruciate ligament tears: MR imaging compared with arthroscopy and clinical tests. Radiology, 166: 861.

Lindenbaum, B.L. (1981):

Complications of knee joint arthroscopy, Clin. Orthop. Rel. Res., 158: 162-170.

Lynch, T.C.P.; Crues, J.V.; Morgan, F.W. et al. (1989):

Bone abnormalities of the knee: Prevalence and significance at MR imaging. Radiology, 171: 761.

Mesgarzadeh, M.; Schneck, C.D. and Bondakdorpour, A. (1988):

Magnetic resonance imaging of the knee and correlation with normal anatomy. Radiographics, 8: 707-733.

Michell, M.R.; Tarr, R.W. and Conturo, T.E. (1986):

Spin-echo technique selection: basic principles for choosing MRI pulse sequence timing intervals, Radiographics, 6: 245-260.

Mink, J.H. (1993):

The cruciate and collateral ligaments. In Mink JH, Reicher MA, Crues JV (eds): Magnetic Resonance Imaging of the knee. New York, Raven Press.

Mink, J.H.; Levy, T. and Crues, J.V. III (1988):

Tears of the anterior cruciate ligament and menisci of the knee: MR imaging evaluation. Radiology, 167: 769.

Morgan, C.D.; Wojtys, E.M.; Casscells, C.D. et al. (1991):

Arthroscopic meniscal repair evaluated by second-look arthroscopy. Am. J. Sports Med., 19: 63.

Munk, P.L.; Helms, C.A. and Vellet, A.D. (1992):

The menisci. In: Munk PL, Helms CA and Vellet AD: MRI of the Knee. An Aspen Publications, p. 57-76.

-
- Munk, P.L.; Helms, C.A.; Genaut, H.K. et al. (1989):**
Magnetic resonance imaging of the knee: Current status, new directions, *Skeletal radiology*, 18: 569-577.
- Murphy, B.J.; Hechtman, K.S.; Uribe, J.W. et al. (1992):**
Hiolibial bands friction syndrome: MR imaging findings. *Radiology*, 185: 569.
- Newman, A.P.; Daniels, A.U. and Burks, R.T. (1993):**
Principles and decision making in mensical surgery. *Arthroscopy*, 9:33.
- Noyes, F.R.; Bassett, R.W.; Grood, E.S. et al. (1980):**
Arthroscopy of an acute traumatic hemarthrosis of the knee. *J. Bone Joint surg.*, 62A: 687.
- Oleaga, L.; Kressel, H. (1990):**
High signal intensity in the posterior cruciate ligament on sagittal gradient-echo images: Normal variant. *Radiology*, 177: 295.
- Peterfy, C.G.; Janzen, D.L.; Tirman, P.F.J. et al. (1994):**
"Magic angle" phenomenon: A cause of increased signal in the normal lateral meniscus on short-TE MR images of the knee. *AJR Am. J. Roentgenol.*, 163: 149.
- Potter, H.G.; Rodeo, S.A.; Wickiewicz, T.L. et al. (1996):**
MR imaging of meniscal allografts: Correlation with clinical and arthroscopic outcomes. *Aradiology*, 198: 509.
- Quinn, S.F.; Brown, T.F. (1991):**
Meniscal tears diagnosed with MR imaging versus arthroscopy: How reliable a standard is arthroscopy? *Radiology*, 181: 843.
- Recht, M.P.; Applegate, G.R.; Kaplan, P.A. et al. (1994):**
MR appearance of cruciate ganglion cysts: Report of 16 cases. *Skeletal Radiol.*, 23: 597.
- Reeder, J.D.; Matz, S.O.; Becker, L. et al. (1989):**
MR imaging of the knee in the sagittal projection; comparison of three dimensional gradient-echo and spin-echo sequences. *AJR*, 153: 537-550.
-

-
- Reicher, M.A.; Rauschnig, W.; Gold, R.H. et al. (1985) :**
High – resolution magnetic resonance imaging of the knee joint
: Normal anatomy, AJR, 145 : 895-901.
- Reiser, M.F.; Bongartz, G.; Erleman, R. et al. (1989):**
Magnetic resonance in cartilagenous lesions of the knee joint
with three dimensional echo-imaging, Skeletal radiology, 17:
465-471.
- Remer, E.M.; Fitzgerald, S.W.; Friedman, H. et al. (1992):**
Anterior cruciate ligament injury: MR imaging diagnosis and
patterns of injury. Radiographics, 12: 901-915.
- Renig, J.W.; McDevitt, E.R. and Ove, P.N. (1991):**
Progression of meniscal degenerative changes in college
football players: Evaluation with MR imaging. Radiology, 181:
255.
- Robertson, P.L.; Schweitzer, M.E.; Bartolozzi, A.R. et al. (1994):**
Anterior cruciate ligament tears: Evaluation of multiple signs
with MR imaging. Radiology, 193: 829.
- Rose, D.J.; Parisien, J.S. (1988):**
Popliteus tendon rupture: Case report and review of the
literature. Clin. Orthop., 226: 113.
- Rothschild, P.A.; Domesek, J.M.; Kaufman, L. et al. (1990):**
MR imaging of the knee with a 0.064-T permanent magnet.
Radiology, 175: 775.
- Rubin, D.A. (1992):**
MR imaging of the knee menisci. Radiologic Clinics of North
America, Volume 35, Number 1.
- Rubin, D.A.; Britton, C.A.; Towers, J.D. et al. (1994):**
Are MR imaging signs of meniscocapsular separation valid?
Radiology, in press.
- Rubin, D.A.; Kneeland, J.B.; Listerud, J. et al. (1994):**
MR diagnosis of meniscal tears of the knee: Value of fast spin-
echo vs conventional spin-echo pulse sequences. AJR Am. J.
Roentgenol., 162: 1131.
-

-
- Ruiz, M.E.; Erickson, S.J. (1994):**
Medial and lateral supporting structures of the knee. Normal MR imaging anatomy and pathologic findings. MRI clin. North Am., 2: 381.
- Ruwe, P.A.; Wright, J.; Randall, R.L. et al. (1992):**
MR imaging effectively replace diagnostic arthroscopy? Radiology, 183: 335.
- Schils, J.P.; Spindler, K.P.; Piraino, D.W. et al. (1992):**
Bone bruises and chondral lesions in anterior cruciate ligament tears: A prospective study. J. Magn Reson. Imaging, 2: 94.
- Schnarkowski, P.; Tirman, P.F.J.; Fuchigami, K.D. et al. (1995):**
Meniscal ossicle: Radiographic and MR imaging findings. Radiology, 196: 47.
- Schweitzer, M.E.; Tran, D.; Deely, D.M. et al. (1995):**
Medial collateral ligaments injuries: Evaluation of multiple signs, prevalence and location of associated bone bruises, and assessment with MR imaging. Radiology, 194: 825.
- Shankman, S.; Beltran, J. (1994):**
The knee. In: Haaga JR, Lanzieri CF, Sartoris DJ et al.: computed Tomography and Magnetic Resonance Imaging of the Whole Body, Mosby Year Book, p. 1477-1515.
- Shelbourne, K.D.; Nitz, P.A. (1991):**
The O'Donoghue triad revisited: Combined knee injuries involving the anterior cruciate and medial collateral ligament tears. Am. J. Sports, Med., 19: 474.
- Shelbourne, K.D.; Martini, D.J.; McCarroll, J.R. et al. (1995):**
Correlation of joint line tenderness and meniscal lesions in patients with acute anterior cruciate ligament tears. Am. J. Sports Med., 23: 166.
- Shellock, F.G.; Deutsch, A.L.; Mink, J.H. (1993):**
"Kinematic MR imaging of the Patellofemoral joint Basic and vanced Techniques", RSNA Categorical course in musculoskeletal Radiology, 173-88.

Shellock, F.G.; Mink, J.H. and Deutsch, A.L. (1993):

“Patellofemoral joint: Identification of abnormalities with active-movement, “unloaded” versus “loaded “Kinematic MR imaging techniques”, Radiology, 188: 2, 575-78.

Shellock, F.G.; Mink, J.H.; Deutsch, A. et al. (1990):

“Evaluation of patients with persistent symptoms following lateral retinacular release by kinematic MRI of the patellofemoral joint. Arthroscopy, 6: 226-34.

Shellock, F.G.; Mink, J.H. and Deutsch, A.L. (1994):

Effect of a patellar realignment brace on patellofemoral relationships: Evaluation with Kinematic MR imaging. JMR, 4: 590-94.

Sherman, O.; Fox, J.; Synder, S.J. et al. (1986):

Arthroscopy-no problem surgery: An analysis of 2640 cases. J. Bone Joint Surg., 69A: 256.

Shogry, M.E.C.; Pope, T.L.Jr. (1991):

Vacuum phenomenon simulating meniscal or cartilaginous injury of the knee at MR imaging. Radiology, 180: 513.

Silverman, J.M.; Mink, J.H.; Deutsch, A.L. (1989):

Discoid menisci of the knee: MR imaging appearance. Radiology, 173: 351.

Singson, R.D.; Feldman, F.; Staron, R. et al. (1991):

MR imaging of displaced bucket-handle tear of the medial meniscus. AJR Am. J. Roentgenol., 156: 121.

Sisk T.D. (1992):

Knee Injuries In: Crenshaw A.H.: Campbell's Operative Orthopaedics, vol. III, 8th edn., Mosby Year Book, p. 1487-1732.

Slocum D.B.; Larson R.L (1988):

Rotatory instability of the knee: Its pathogenesis and a clinical test to demonstrate its presence, J. Bone Joint surg. {Am.}, 50: 211-225.

-
- Shellock, F.G.; Mink, J.H. and Deutsch, A.L. (1993):**
“Patellofemoral joint: Identification of abnormalities with active-movement, “unloaded” versus “loaded “Kinematic MR imaging techniques”, Radiology, 188: 2, 575-78.
- Shellock, F.G.; Mink, J.H.; Deutsch, A. et al. (1990):**
“Evaluation of patients with persistent symptoms following lateral retinacular release by kinematic MRI of the patellofemoral joint. Arthroscopy, 6: 226-34.
- Shellock, F.G.; Mink, J.H. and Deutsch, A.L. (1994):**
Effect of a patellar realignment brace on patellofemoral relationships: Evaluation with Kinematic MR imaging. JMR, 4: 590-94.
- Sherman, O.; Fox, J.; Synder, S.J. et al. (1986):**
Arthroscopy-no problem surgery: An analysis of 2640 cases. J. Bone Joint Surg., 69A: 256.
- Shogry, M.E.C.; Pope, T.L.Jr. (1991):**
Vacuum phenomenon simulating meniscal or cartilaginous injury of the knee at MR imaging. Radiology, 180: 513.
- Silverman, J.M.; Mink, J.H.; Deutsch, A.L. (1989):**
Discoid menisci of the knee: MR imaging appearance. Radiology, 173: 351.
- Singson, R.D.; Feldman, F.; Staron, R. et al. (1991):**
MR imaging of displaced bucket-handle tear of the medial meniscus. AJR Am. J. Roentgenol., 156: 121.
- Sisk T.D. (1992):**
Knee Injuries In: Crenshaw A.H.: Campbell's Operative Orthopaedics, vol. III, 8th edn., Mosby Year Book, p. 1487-1732.
- Slocum D.B.; Larson R.L (1988):**
Rotatory instability of the knee: Its pathogenesis and a clinical test to demonstrate its presence, J. Bone Joint surg. {Am.}, 50: 211-225.

Smith, D.K.; Totty, W.G. (1990):

The knee after partial meniscectomy: MR imaging features. Radiology, 176: 141.

Sonin, A.H.; Fitzgerald, S.W.; Hoff, F.L. et al. (1995):

MR imaging of the posterior cruciate ligament: Normal, abnormal and associated injury patterns, Radiographics, 15: 551-561.

Sonin A.H.; Rogers L.F. (1997):

Skeletal Trauma In: Grainger R.G. and Allison D.: Grainger and Allison's Diagnostic Radiology: A Text book of Medical Imaging, 3th edn., vol. II, Churchill Livingstone, p. 1573-1626.

Sonin, A.H.; Fitzgerald, S.W.; Friedman, H. et al. (1994):

Posterior cruciate ligament injury: MR imaging diagnosis and patterns of injury. Radiology, 190: 455.

Snearly, W.N.; Kaplan, P.A.; Dussault, R.G. (1996):

Lateral compartment bone contusions in adolescents with intact anterior cruciate ligaments. Radiology, 198: 205.

Spiers, A.S.D.; Meagher, T.; Ostlere, S.J. et al. (1993):

Can MRI of the knee affect arthroscopic practice? J. Bone Joint Surg., 75B: 49-52.

Stoller, D.W.; Cannon, W.D. and Aderson, L.J. (1993):

The knee In: Stoller DW: Magnetic Resonance Imaging In Orthopaedics & Sports Medicine, J.B. Lippincott Company, Philadelphia, p. 139-372.

Thijn, C.J.P. (1982):

Accuracy of double contrast arthrography and arthroscopy of the knee, Skeletal Radiology, 8: 187-192.

Tolin, B.S.; Sapega, A.A. (1993):

Arthroscopic visual field mapping at the periphery of the medial meniscus: A comparison of different portal approaches. Arthroscopy, 9: 265.

Trichey, E.L. (1980):

Injuries to the posterior cruciate ligament: Diagnosis and treatment of early injuries and reconstruction of late instability. Clin. Orthop., 147: 76.

Trickey E.L. (1984):

Acute ligamentous injuries In: Jackson JP and Wough Weds.
Surgery of the Knee Joint, champman and Hall p. 323-401.

Tuckman, G.A.; Miller, W.J.; Remo, J.W. et al. (1994):

Radial tears of the menisci: MR findings. AJR Am. J.
Roentgenol., 163: 395.

Tung, G.A.; Davis, L.M.; Wiggins, M.E. et al. (1993):

Tears of the anterior cruciate ligament: Primary and secondary
signs at MR imaging. Radiology, 186: 661.

Turner, D.A.; Rapoport, M.I.; Erwin, W.D. et al. (1991):

Truncation artifact: A potential pitfall in MR imaging of the
menisci of the knee. Radiology, 179: 629.

Vahey, T.N.; Meyer, S.F.; Shelbourne, K.D. et al. (1994):

MR imaging of anterior cruciate ligament injuries. MRI Clin.
North Am., 2: 365.

Vallet, A.D.; Frad, D.; Marks, P.H. et al. (1991):

Occult post-traumatic osteochondral lesions of the knee:
prevalence, classification and short term sequelae evaluated
with MR imaging. Radiology, 178: 271-276.

Van Dommelen, B.; Fowler, P. (1989):

Anatomy of the posterior cruciate ligament. A review. Am. J.
Sports Med., 17: 24.

Veltri, D.M.; Warren, R.F. (1994):

Posterolateral instability of the knee. J. Bone Joint Surg., 76A:
460.

Walker, W.C.; Moore, T.E. (1997):

Imaging of skeletal and soft tissue injuries in and around the
knee. Radiologic. Clinics of North America, Vol. 35, No. 3, P.
631-653.

Watanabe, A.T.; Carter, B.C.; Teitelbaum, G.P. et al. (1989):

Normal variations in MR imaging of the knee: Appearance and
frequency. AJR Am. J. Roentgenol., 153: 341.

Weber, W.; Neumann, C.; Barakos, J. et al. (1991):

Lateral tibial rim (Segond) fractures: MR imaging characteristics. Radiology, 180: 731.

Weiss, K.L.; Morehouse, H.T.; Levy, I.M. (1991):

Sagittal MR images of the knee: A low-signal band parallel to the posterior cruciate ligament caused by a displaced bucket-handle tear. AJR Am. J. Roentgenol., 156: 117.

Winkler, M.L.; Orlendahl, D.A.; Mills, T.C. et al. (1988) :

Characteristics of partial flip angle and gradient reversal MR imaging, radiology, 166 : 17-26.

Wolf, G.L. (1989):

Current status of MR imaging contrast agents: Special report, Radiology, 172: 709-710.

Wright, D.H.; De Smet, A.A.; Norris, M. (1995):

Bucket-handle tears of the medial and lateral menisci of the knee: Value of MR imaging in detecting displaced fragments. AJR Am. J. Roentgenol., 165: 621.

Yerys, P. (1991):

Arthroscopic posterior cruciate ligament repair. Arthroscopy, 7: 111.

Yu, J.S.; Salonen, D.C.; Holder, J. et al. (1996):

Posterolateral aspect of the knee: Improved MR imaging with a coronal oblique technique. Radiology, 198: 199.

Zobel, M.S.; Borrello, J.A.; Siegel, M.J. et al. (1994):

Pediatric knee MR imaging: Pattern of injuries in the immature skeleton. Radiology, 190: 397.