

REFERENCES

Ahren B. and Pacini G. (2007):

Islet adaptation to insulin resistance: mechanisms and implications for intervention. *Diabetes Obes Metab.*; 1:2-8.

Arkan, M.C., Hevener, A.L., Greten, F.R., Maeda, S., Li, Z.W., Long, J.M., Wynshaw-Boris, A., Poli, G., Olefsky, J., and Karin, M. (2005):

IKK- β links inflammation to obesity-induced insulin resistance. *Nat. Med.*; 11: 191–198.

Aronne LJ and Segal KR. (2002):

Adiposity and fat distribution outcome measures: assessment and clinical implications. *Obes Res.*;10 (1):14S–21S.

Atsushi Tsuchida, Toshimasa Yamauchi, Yusuke Ito, Yusuke Hada, Toshiyuki Maki, Sato Takekawa, Junji Kamon, Masaki Kobayashi, Ryo Suzuki, Kazuo Hara and Naoto Kubota (2004):

Insulin/Foxo1 Pathway Regulates Expression Levels of Adiponectin Receptors and Adiponectin Sensitivity. *J. Biol. Chem.*; 279(29): 30817-30822

Bak, J.F., Moller, N., Schmitz, O., Saaek, A., and Pedersen, O. (1992):

In vivo insulin action and muscle glycogen synthase activity in type 2 (non-insulin-dependent) diabetes mellitus: Effects of diet treatment *Diabetologia.*; 35: 777–784.

Balon TW. and Nadler JL. (1997):

Evidence that nitric oxide increases glucose transport in skeletal muscle. *J Appl Physiol.*; 82:359–363.

Banerjee, R.R., Rangwala, S.M., Shapiro, J.S., Rich, A.S., Rhoades, B., Qi, Y., Wang, J., Rajala, M.W., Poci, A. and Scherer, P.E. (2004):

Regulation of fasted blood glucose by resistin. *Science*; 303: 1195–1198.

Barish GD, Narkar VA and Evans RM. (2006):

PPAR delta: a dagger in the heart of the metabolic syndrome. *J Clin Invest.* ;116(3):590-7.

Berg AH, Combs TP, Du X, Brownlee M and Scherer PE. (2001):

The adipocyte-secreted protein Acrp30 enhances hepatic insulin action. *Nat Med.*; 7:947–953.

Bergman, R.N., Kim, S.P., Hsu, I.R., Catalano, K.J., Chiu, J.D., Kabir, M., Richey, J.M., and Ader, M. (2007):

Abdominal obesity: Role in the pathophysiology of metabolic disease and cardiovascular risk. *Am. J. Med.*; 120 (1): S3– S8.

Bernard, C. (1854):

Leçons de physiologie expérimentale appliquées à la médecine
Baillere et fils, Paris.

Biddinger and Kahn. (2006):

Insulin resistance etiology and underlying causes *Annu. Rev.Physiol.*; 68:123-58.

Borghouts LB. and Keizer HA. (2000):

Exercise and insulin sensitivity: a review. *Int J Sports Med.*; 21:1–12.

Bruce D. Wingred and Geoffery stein (1988):

Rat dissection manual, third edition:45-51.

Cai, D., Yuan, M., Frantz, D.F., Melendez, P.A., Hansen, L., Lee, J., and Shoelson, S.E. (2005):

Local and systemic insulin resistance resulting from hepatic activation of IKK- β and NF- κ B. Nat. Med.; 11: 183–190.

Cefalu WT. (2001):

Insulin resistance: cellular and clinical concepts. Exp Biol Med.; 226:13–26.

Chandran, M., Phillips, S.A., Ciaraldi, T., and Henry, R.R. (2003):

Adiponectin: More than just another fat cell hormone? Diabetes Care.; 26: 2442–2450.

Chang SP, Chen YH, Chang WC, Liu IM and Cheng JT.(2006):

Increase of adiponectin receptor gene expression by physical exercise in soleus muscle of obese Zucker rats. Eur J Appl Physiol. :97(2):189- 95

Cheatham B. and Kahn CR. (1995):

Insulin action and the insulin signaling network. Endocr Rev.; 16:117–142.

Chiasson JL. And Rabasa-Lhoret R. (2004):

Prevention of type 2 diabetes: insulin resistance and beta-cell function. Diabetes.;53 Suppl 3:S34-8.

Chou CK, Dull TJ, Russell DS, Gherzi R, Lebowitz D, Ullrich A and Rosen OM. (2000):

Human insulin receptors mutated at the ATP-binding site lack protein tyrosine kinase activity and fail to mediate postreceptor effects of insulin. *J Biol Chem.*; 262:1842–1847.

Combs TP, Wagner JA, Berger J, Doebber T, Wang WJ, Zhang BB, Tanen M, Berg AH, O'Rahilly S, Savage DB, Chatterjee K, Weiss S, Larson PJ, Gottesdiener KM, Gertz BJ, Charron MJ, Scherer PE and Moller DE (2002):

Induction of adipocyte complement-related protein of 30 kilodaltons by PPAR gamma agonists: a potential mechanism of insulin sensitization. *Endocrinology.*; 143:998-1007, 2002.

Combs, TP, Pajvani, U.B., Berg, A.H., Lin, Y., Jelicks, L.A., Laplante, M., Nawrocki, A.R., Rajala, M.W., Parlow, A.F. and Cheeseboro, L. (2004):

A transgenic mouse with a deletion in the collagenous domain of adiponectin displays elevated circulating adiponectin and improved insulin sensitivity. *Endocrinology.*; 145: 367–383.

Craig B.W and Foley P.J (1984):

Effects of cell size and exercise on glucose uptake and metabolism in adipocytes of female rats. *J. appl. Physiology.*; 57:1120-1125.

Dandona, P., Aljada, A., and Bandyopadhyay, A. (2004):

Inflammation: The link between insulin resistance, obesity and diabetes. *Trends Immunol.*; 25: 4–7.

Dela, F., K. J. Mikines, J. J. Larsen, and H. Galbo (1994):

Training-induced enhancement of insulin action in human skeletal muscle: the influence of aging. *J. Gerontol.*; 51A: B-247-B252.

Devlin JT. (1992):

Effects of exercise on insulin sensitivity in humans. *Diabetes Care.*;15(11):1690-3

Devlin, J.T., and E. S. Horton(2005):

Effects of prior high-intensity exercise on glucose metabolism in normal and insulin-resistant men. *Diabetes.*; 34: 973-979.

Diez, J.J. and Iglesias, P. (2003):

The role of the novel adipocyte-derived hormone adiponectin in human disease. *Eur. J. Endocrinol.*; 148: 293–300.

Dunstan D, Puddey IB, Beilin LJ, Burke V, Morton AR and Stanton KG. (1998):

Effects of short-term circuit weight training program on glycaemic control in NIDDM. *Diabetes Res Clin Pract* ;40:53–61.

Ebeling, P., R. Bourey, L. Koranyi, J. A. Tuominen, L. C. Groop, J. Henriksson, M. Mueckler, A. Sovijarvi, and V. A. Koivisto (1993):

Mechanism of enhanced insulin sensitivity in athletes. *J. Clin. Invest.*; 92: 1623-1631

Ebihara, K., Ogawa, Y., Masuzaki, H., Shintani, M., Miyanaga, F., Aizawa-Abe, M., Hayashi, T., Hosoda, K., Inoue, G., and Yoshimasa, Y.(2001):

Transgenic overexpression of leptin rescues insulin resistance and diabetes in a mouse model of lipotrophic diabetes.;Diabetes 50: 1440–1448.

Emanuelli, B., Peraldi, P., Filloux, C., Chavey, C., Freidinger, K., Hilton, D.J., Hotamisligil, G.S., and Van Obberghen, E. (2001):

SOCS-3 inhibits insulin signaling and is up-regulated in response to tumor necrosis factor- α in the adipose tissue of obese mice. J. Biol.Chem.; 276: 47944–47949.

Evans, J.L., Goldfine, I.D., Maddux, B.A., and Grodsky, G.M. (2002):

Oxidative stress and stress-activated signaling pathways: A unifying hypothesis of type 2 diabetes. Endocr. Rev.; 23: 599–622.

Ezenwaka CE, Kalloo R, Uhlig M and Eckel J. (2004):

Relationship between adiponectin and metabolic variables in caribbean offspring of patients with type 2 diabetes mellitus. Horm Metab Res.; 36:238–242.

Fain, J.N., Madan, A.K., Hiler, M.L., Cheema, P., and Bahouth, S.W. (2004):

Comparison of the release of adipokines by adipose tissue, adipose tissue matrix, and adipocytes from visceral and subcutaneous abdominal adipose tissues of obese humans. Endocrinology.; 145: 2273–2282.

Fenicchia LM, Kanaley JA, Azevedo JL Jr, et al. (2004):

Influence of resistance exercise training on glucose control in women with type 2 diabetes. *Metabolism*.;53:284-9.

Fontana, L., Eagon, J.C., Trujillo, M.E., Scherer, P.E., and Klein, S. (2007):

Visceral fat adipokine secretion is associated with systemic inflammation in obese humans. *Diabetes*.; 56: 1010–1013.

Fruebis J, Tsao TS, Javorschi S, Ebbets-Reed D, Erickson MR, Yen FT, Bihain BE and Lodish HF. (2001):

Proteolytic cleavage product of 30-kDa adipocyte complement-related protein increases fatty acid oxidation in muscle and causes weight loss in mice. *Proc Natl Acad Sci*.; 98:2005–2010.

Garry J. (2002):

"Banting lecture 2001: dysregulation of fatty acid metabolism in the etiology of type 2 diabetes". *Diabetes*.; 51 (1): 7–18.

Gavi, S., Stuart, L.M., Kelly, P., Melendez, M.M., Mynarcik, D.C., Gelato, M.C., and McNurlan, M.A. (2007):

Retinol-binding protein 4 is associated with insulin resistance and body fat distribution in nonobese subjects without type 2 diabetes. *J. Clin. Endocrinol. Metab*.; 92: 1886–1890.

Giorgino F, Laviola L. and Eriksson JW. (2005):

Regional differences of insulin action in adipose tissue: insights from *in vivo* and *in vitro* studies. *Acta Physiol Scand*.;183:13–30.

Goldstein BJ. And Scalia R (2004):

Adiponectin: a novel adipokine linking adipocytes and vascular function. J Clin Endocrinol Metab.; 89:2563–2568.

Goutham RAO. (2001):

Insulin resistance syndrome: Am Fam Physician.;63:1159-63,1165-6.

Haluzik M, Parizkova J. and Haluzik MM (2004):

Adiponectin and its role in the obesity-induced insulin resistance and related complications. Physiol Res.; 53:123–129.

Haque KA, Pfeiffer RM, Beerman MP, Struewing JP, Chanock JP and Bergen AW (2003):

Performance of high throughout DNA quantification method. BMC Biotechnol.;3:20-24.

Hill JO, Melanson EL and Wyatt HT.(2000):

Dietary fat intake and regulation of energy balance: implications for obesity.J Nutr.;130:284s-8s.

Hirosumi, J., Tuncman, G., Chang, L., Gorgun, C.Z., Uysal, K.T., Maeda, K., Karin, M., and Hotamisligil, G.S. (2002):

A central role for JNK in obesity and insulin resistance. Nature.; 420: 333-336.

Holloszy JO. (2005):

Exercise-induced increase in muscle insulin sensitivity. J Appl. Physiol.;99:338-43.

Hotamisligil, G.S. (2003):

Inflammatory pathways and insulin action. *Int.. Obes. Relat. Metab. Disord.*; 27 (3): S53–S55.

Houmard, J. A., M. H. Shinebarger, P. L. Dolan, N. Leggett-Frazier, R. K. Bruner, M. R. McCammon, R. G. Israel, and G. L. Dohm.(2003):

Exercise training increases GLUT-4 protein content in previously sedentary middle-aged men. *Am. J. Physiol.*;264 (27): E896-E901.

Houstis, N., Rosen, E.D., and Lander, E.S. (2006):

Reactive oxygen species have a causal role in multiple forms of insulin resistance. *Nature.*; 440: 944–948.

Hu E, Liang P and Spiegelman BM. (1996):

AdipoQ is a novel adipose-specific gene dysregulated in obesity. *J Biol Chem.*; 271:10697–10703.

.

Hu Huang, Kaoruko Tada Iida, Hirohito Sone, Tomotaka Yokoo, Nobuhiro Yamada¹ and Ryuichi Ajisaka (2006):

The effect of exercise training on adiponectin receptor expression in KKAY obese/diabetic mice. *Journal of Endocrinology.*; 189: 643-653

Huang Z.,Fasco M.J and Kaninskyl. S(1996):

Optimization of Danse I removal of contaminating DNA fromRNA for use in quantitative RNA-PCR. *Biotechninques.*; 20:1012- 1020.

Hunter SJ and Garvey WT. (1998):

Insulin action and insulin resistance: diseases involving defects in insulin receptors, signal transduction, and the glucose transport effector system. *Am J Med.*;105:331–45.

Inukai K, Nakashima Y, Watanabe M, Takata N, Sawa T, Kurihara S, Awata T and Katayama S (2005):

Regulation of adiponectin receptor gene expression in diabetic mice. *American Journal of Physiology, Endocrinology and Metabolism.*; 288 : E876–E882

Ivy JL(1997):

Role of exercise: training in the prevention and treatment of insulin resistance and non-insulin-dependent diabetes mellitus. *Sports Med.*;24:321–336.

Ivy J. L., Zderic T. W. and Fogt D. L. (1999):

Prevention and treatment on non-insulin-dependent diabetes mellitus *Exerc. Sport Sci. Rev.*, 27: 1-35.

Jack Y. (2003):

Exercise and the regulation of blood glucose *Sport Sci. Rev.*;36:251- 6.

Jacobs HS. and Conway GS. (1999):

Leptin, polycystic ovaries and polycystic ovary syndrome. *Hum Reprod Update.*;5:166–71.

Jamine C. ,Kaufmau M.D.,Tamara J., Harris B.A., Joseph Higginis B.A., Alans. and Maisel M.D (1993):

Exercise induced enhancement of glucose uptake in rats *Circ.*;90:525-532.

John P. Kirwan, Luis F. del Aguila, Jazmir M. Hernandez, David L. Williamson, Donal J. O'Gorman, Rebecca Lewis, and Raj K. Krishnan (2000):

Regular exercise enhances insulin activation of IRS-1-associated PI3-kinase in human skeletal muscle. *J. Appl. Physio.*; 88 (2):797-803.

Kabir, M., Catalano, K.J., Ananthnarayan, S., Kim, S.P., Van Citters, G.W., Dea, M.K., and Bergman, R.N. (2005):

Molecular evidence supporting the portal theory: A causative link between visceral adiposity and hepatic insulin resistance. *Am. J. Physiol. Endocrinol. Metab.* Doi.; 10.1152/ajpendo.00203.2004.

Kadi F. and Thornell LE. (2000):

Concomitant increases in myonuclear and satellite cell content in female trapezius muscle following strength training. *Histochem Cell Biol.*;113:99-103.

Kadowaki T. and Yamauchi T. (2005):

Adiponectin and adiponectin receptors. *Endocr Rev.*; 26:439–451

Kahn CR. (1999):

Diabetes: Causes of insulin resistance. *Nature.*; 373:384–385.

Karhapaa P, Malkki M. and Laakso M. (1994):

Isolated low HDL cholesterol. An insulin-resistant state. *Diabetes.*;43:411-7.

Kelesidis I, Kelesidis T. and Mantzoros CS (2006):

Adiponectin and cancer: a systematic review. *Br J Cancer.*; 94:1221–1225.

Kennedy JW, Hirshman MF, Gervino EV, Ocel JV, Forse RA, Hoenig SJ, Aronson D, Goodyear LJ. and Horton ES.(1999):

Acute exercise induces GLUT4 translocation in skeletal muscle of normal human subjects and subjects with type 2 diabetes. Diabetes.;48:1192–1197.

Kharroubi I, Rasschaert J, Eizirik DL. and Cnop M (2003):

Expression of adiponectin receptors in pancreatic β cells. Biochem. Biophys Res Commun.; 312:1118–1122.

Kim YB, Nikoulina SE, Ciaraldi TP, Henry RR. and Kahn BB. (1999):

Normal insulin-dependent activation of Akt/protein kinase B, with diminished activation of phosphoinositide 3-kinase, in muscle in type II diabetes. J Clin Invest.; 104:733–741.

Koivisto VA, Yki-Järvinen H and DeFronzo RA. (1986):

Physical training and insulin sensitivity Diabetes Metab.Rev.;1 (4):445-81.

Kouichi Inukai, Youhei Nakashima, Masaki Watanabe, Nobuki Takata, Takahiro Sawa, Susumu Kurihara, Takuya Awata, and Shigehiro Katayama (2004):

Regulation of adiponectin receptor gene expression in diabetic mice Am J Physiol Endocrinol Metab.; 288: E876-E882 0193-1849/05.

Koval, J. A., R. A. DeFronzo, R. M. O'Doherty, R. Printz, H. Ardehali, D. K. Granner, and L. J. Mandarino (1998):

Regulation of hexokinase II activity and expression in human muscle by moderate exercise. Am. J. Physiol. Endocrinol. Metab.; 274: E304-E308.

Kumada M, Kihara S, Sumitsuji S, Kawamoto T, Matsumoto S, Ouchi N, Arita Y, Okamoto Y, Shimomura I, Hiraoka H, Nakamura T, Funahashi T, Matsuzawa Y and Osaka CAD Study Group.(2003):

Association of hypoadiponectinemia with coronary artery disease in men. *Arterioscler Thromb Vasc Biol.*; 23:85–89.

Lagouge, M., Argmann, C., Gerhart-Hines, Z., Meziane, H., Lerin, C., Daussin, F., Messadeq, N., Milne, J., Lambert, P. and Elliott, P.. (2006):

Resveratrol improves mitochondrial function and protects against metabolic disease by activating SIRT1 and PGC-1 α . *Cell.*; 127: 1109–1122.

Li, Z., Bowerman, S., and Heber, D. (2005):

Health ramifications of the obesity epidemic. *Surg. Clin. North. Am.*; 85: 681–701

Lumeng, C.N., Bodzin, J.L., and Saltiel A.R. (2007):

Obesity induces a phenotypic switch in adipose tissue macrophage polarization. *J. Clin. Invest.* 117: 175–184.

Luo XH, Guo LJ, Yuan LQ, Xie H, Zhou HD, Wu XP and Liao EY (2005):

Adiponectin stimulates human osteoblasts proliferation and differentiation via the MAPK signaling pathway. *Exp Cell Res.*; 309:99–109

Ma, L.J., Mao, S.L., Taylor, K.L., Kanjanabuch, T., Guan, Y., Zhang, Y., Brown, N.J., Swift, L.L., McGuinness, O.P. and Wasserman D.H. (2004):

Prevention of obesity and insulin resistance in mice lacking plasminogen activator inhibitor 1. *Diabetes.*; 53: 336–346.

Maayan Barnea, Avi Shamay, Aliza H. Stark and Zecharia Madar (2006):

A High-Fat Diet Has a Tissue-Specific Effect on Adiponectin and Related Enzyme Expression. *Obesity.*; 14, 2145–2153;

Maeda N, Takahashi M, Funahashi T, Kihara S, Nishizawa H, Kishida K, Nagaretani H, Matsuda M, Komuro R, Ouchi N, Kuriyama H, Hotta K, Nakamura T, Shimomura I and Matsuzawa Y.(2001):

PPARgamma ligands increase expression and plasma concentrations of adiponectin, an adipose-derived protein. *Diabetes.*;(9):2094-9.

Masuzaki, H., Paterson, J., Shinyama, H., Morton, N.M., Mullins, J.J., Seckl, J.R., and Flier J.S. (2001):

A transgenic model of visceral obesity and the metabolic syndrome. *Science.*; 294: 2166–2170.

Matthews DR, Hosker JP and Rudenski AS (1985):

Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia.*; 28:412-419

Matthias Bluher, John W. Bullen, Jr., Jennifer H. Lee, Susan Kralisch, Mathias Fasshauer, Nora Kloeting, Josef Niebauer, Michael R. Schoen, Catherine J. Williams, and Christos S. Mantzoros (2006):

Circulating Adiponectin and Expression of Adiponectin Receptors in Human Skeletal Muscle: Associations with Metabolic Parameters and Insulin Resistance and Regulation by Physical Training. *Journal of Clinical Endocrinology & Metabolism*.; 91(6):2310–2316.

McGarry, J.D. (2002):

Dysregulation of fatty acid metabolism in the etiology of type 2 diabetes. *Diabetes*.; 51: 7–18.

Meigs JB, Mittleman MA, Nathan DM, Tofler GH, Singer DE and Murphy-Sheehy PM. (2000):

Hyperinsulinemia, hyperglycemia, and impaired hemostasis: the Framingham Offspring Study. *JAMA*; 283:221-8.

Mensink, M., Hesselink, M.K., Russell, A.P., Schaart, G., Sels, J.P., and Schrauwen, P. (2007):

Improved skeletal muscle oxidative enzyme activity and restoration of PGC-1 α and PPAR β/δ gene expression upon rosiglitazone treatment in obese patients with type 2 diabetes mellitus. *Int. J. Obes. (Lond.)* doi: 10.1038/sj.ijo.0803567.

Minokoshi Y, Kim YB, Peroni OD, Fryer LG, Muller C, Carling D and Kahn BB (2002):

Leptin stimulates fatty-acid oxidation by activating AMP-activated protein kinase. *Nature*.; 415:339–343.

Mootha, V.K., Lindgren, C.M., Eriksson, K.F., Subramanian, A., Sihag, S., Lehar, J., Puigserver, P., Carlsson, E., Ridderstrale, M., and Laurila, E. (2003):

PGC-1 α -responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat. Genet.*; 34: 267–273.

Naderali, E. K., Estadella, D. and Rocha, M. (2003):

A fat-enriched, glucose-enriched diet markedly attenuates adiponectin receptors mRNA levels in rat epididymal adipose tissue. *Clin Sci (Lond.)*; 105: 403–408.

Nagasaki M, Nakai N, Oshida Y, Li Z, Xu M, Obayashi M, Murakami T, Yoshimura A, Fujitsuka N, Shimomura Y. and Sato Y.(2000):

Exercise training prevents maturation-induced decreases in insulin receptor substrate-1 and phosphatidylinositol 3-kinase in rat skeletal muscle. *Metabolism.*; 49:954–959.

Nakamura T, Tokunaga K, Shimomura I, Nishida M, Yoshida S, Kotani K, Islam AH, Keno Y, Kobatake T. and Nagai Y(1994):

Contribution of visceral fat accumulation to the development of coronary artery disease in non-obese men. *Atherosclerosis*; 107(2):239-46.

NRC (1994):

National Research council, National Academy of science
Washington , D.C.,USA

Obici, S., Feng, Z., Karkanias, G., Baskin, D.G., and Rossetti, L. (2002):

Decreasing hypothalamic insulin receptors causes hyperphagia and insulin resistance in rats. *Nat. Neurosci.*; 5: 566–572.

Okamoto, H., Nakae, J., Kitamura, T., Park, B.C., Dragatsis, I., and Accili, D. (2004):

Transgenic rescue of insulin receptor-deficient mice. *J. Clin. Invest.*;114: 214–223.

Osei K. (1999):

Insulin resistance and systemic hypertension. *Am J Cardiol.*;84:33J- 6J.

Ouchi N, Kihara S, Arita Y, Okamoto Y, Maeda K, Kuriyama H, Hotta K, Nishida M, Takahashi M, Muraguchi M, Ohmoto Y, Nakamura T, Yamashita S, Funahashi T. and Matsuzawa Y (2000): Adiponectin, an adipocyte-derived plasma protein, inhibits endothelial NF- κ B signaling through a cAMP-dependent pathway. *Circulation.*; 102:1296–1301.

Ouchi N, Ohishi M, Kihara S, Funahashi T, Nakamura T, Nagaretani H, Kumada M, Ohashi K, Okamoto Y, Nishizawa H, Kishida K, Maeda N, Nagasawa A, Kobayashi H, Hiraoka H, Komai N, Kaibe M, Rakugi H, Ogihara T. and Matsuzawa Y.(2003): Association of hypoadiponectinemia with impaired vasoreactivity. *Hypertension.*; 42:231–234.

Ouchi N, Shibata R. and Walsh K (2006):

Cardioprotection by adiponectin. *Trends Cardiovasc Med.*; 16:141–146.

Pajvani UB, Du X, Combs TP, Berg AH, Rajala MW, Schulthess T, Engel J, Brownlee M. and Scherer PE. (2003):

Structure-function studies of the adipocyte-secreted hormone Acrp30/adiponectin. Implications for metabolic regulation and bioactivity. J Biol Chem.; 278:9073–9085

Pajvani UB, Hawkins M, Combs TP, Rajala MW, Doebber T, Berger JP, Wagner JA, Wu M, Knopps A, Xiang AH, Utzschneider KM, Kahn SE, Olefsky JM, Buchanan TA. And Scherer PE. (2004):

Complex distribution, not absolute amount of adiponectin, correlates with thiazolidinedione-mediated improvement in insulin sensitivity. J Biol Chem.; 279:12152–12162.

Paolisso, G. and Giugliano, D. (1996):

Oxidative stress and insulin action: Is there a relationship? Diabetologia.; 39: 357–363.

Patti, M.E., Butte, A.J., Crunkhorn, S., Cusi, K., Berria, R., Kashyap, S., Miyazaki, Y., Kohane, I., Costello, M.,and Saccone, R. (2003):

Coordinated reduction of genes of oxidative metabolism in humans with insulin resistance and diabetes: Potential role of PGC1 and NRF1. Proc. Natl. Acad. Sci.; 100: 8466–8471.

Petersen AM. and Pedersen BK. (2005):

The anti-inflammatory effect of exercise. J Appl Physiol. ;98:1154-62.

Petersen, K.F. and Shulman, G.I. (2006):

Etiology of insulin resistance. Am. J. Med.; 119:123-125.

Petersen, K.F., Befroy, D., Dufour, S., Dziura, J., Ariyan, C., Rothman, D.L., DiPietro, L., Cline, G.W., and Shulman, G.I. (2003):

Mitochondrial dysfunction in the elderly: Possible role in insulin resistance. *Science.*; 300: 1140–1142.

Petersen, K.F., Dufour, S., Befroy, D., Garcia, R., and Shulman, G.I. (2004):

Impaired mitochondrial activity in the insulin-resistant offspring of patients with type 2 diabetes. *N. Engl. J. Med.*;350: 664–671.

Pocai, A., Lam, T.K., Obici, S., Gutierrez-Juarez, R., Muse, E.D., Arduini, A., and Rossetti, L. (2006):

Restoration of hypothalamic lipid sensing normalizes energy and glucose homeostasis in overfed rats. *J. Clin. Invest.* 116: 1081–1091.

Punyadeera C, Zorenc AH, Koopman R, McAinch AJ, Smit E, Manders R, Keizer HA, Cameron-Smith D and van Loon LJ (2005):

The effects of exercise and adipose tissue lipolysis on plasma adiponectin concentration and adiponectin receptor expression in human skeletal muscle. *European Journal of Endocrinology.*; 152 : 427–436.

Qi, Y., Nie, Z., Lee, Y.S., Singhal, N.S., Scherer, P.E., Lazar, M.A., and Ahima, R.S. (2006):

Loss of resistin improves glucose homeostasis in leptin deficiency. *Diabetes.*; 55: 3083–3090.

Reaven G. (2004):

The metabolic syndrome or the insulin resistance syndrome?
Different names, different concepts, and different goals.
Endocrinol Metab Clin North Am.;33:283–303.

Reaven GM. (1988):

Role of insulin resistance in human disease. Banting lecture .
Diabetes.;37: 1595-607.

Reaven GM. (1994):

Syndrome X: 6 years later. J Intern Med Suppl. ;736:13-22.

Reaven GM. (2006):

The metabolic syndrome: is this diagnosis necessary?. Am J Clin
Nutr.;83(6):1237-47.

**Rieusset, J., Bouzakri, K., Chevillotte, E., Ricard, N., Jacquet, D.,
Bastard, J.P., Laville, M., and Vidal, H. (2004):**

Suppressor of cytokine signaling 3 expression and insulin
resistance in skeletal muscle of obese and type 2 diabetic patients.
Diabetes.; 53: 2232–2241.

**Ross R., Dagnone D., Jones P. J., Smith H., Paddags A., Hudson R. and
Janssen I.(2000):**

Reduction in obesity and related comorbid conditions after diet-
induced weight loss or exercise–induced weight loss in men. A
randomized controlledtrial. Ann.Intern.Med.; 133: 92-103.

Rui, L., Yuan, M., Frantz, D., Shoelson, S., and White, M.F. (2002):

SOCS-1 and SOCS-3 block insulin signaling by ubiquitin-mediated
degradation of IRS1 and IRS2. J. Biol. Chem.; 277: 42394–42398.

Saengsirisuwan, V, Perez FR, Kinnick TR, and Henriksen EJ. (2002):

Effects of exercise training and antioxidant R-(+)-lipoic acid on glucose transport in insulin-sensitive rat skeletal muscle. J Appl Physiol.; 92: 50-58.

Salminen A and Vihko V. (1983):

Endurance training reduces the susceptibility of mouse skeletal muscle to lipid peroxidation in vitro. Acta Physiol Scand.;117:109-13

Saltiel, A.R. and Kahn, C.R. (2001):

Insulin signalling and the regulation of glucose and lipid metabolism Nature.; 414: 799–806.

Sato Y. (2000):

Diabetes and life-styles: role of physical exercise for primary prevention. Br J Nutr.; 84:S187–S190.

Scherer PE, Williams S, Fogliano M, Baldini G. and Lodish HF (1995):

A novel serum protein similar to C1q, produced exclusively in adipocytes. J Biol Chem.;270:26746–26749.

Seckl, J.R., Morton, N.M., Chapman, K.E., and Walker, B.R. (2004):

Glucocorticoids and 11 β -hydroxysteroid dehydrogenase in adipose tissue. Recent Prog. Horm. Res.; 59: 359–393.

Sheehan MT. and Jensen MD. (2000):

Metabolic complications of obesity. Med Clin North Am.; 84:363–385.

Shoelson, S.E., Lee, J., and Goldfine, A.B. (2006):

Inflammation and insulin resistance. J. Clin. Invest.; 116: 1793–1801.

Shulman GI. (2000):

Cellular mechanisms of insulin resistance. J Clin Invest.; 106:171–176.

Smith U. (2002):

Impaired (‘diabetic’) insulin signaling and action occur in fat cells long before glucose intolerance--is insulin resistance initiated in the adipose tissue? Int J Obes Relat Metab Disord.;26:897–904.

SoJung Lee, Jennifer L. Kuk, Lance E. Davidson, Robert Hudson, Katherine Kilpatrick, Terry E. Graham and Robert Ross (2005):

Exercise without weight loss is an effective strategy for obesity reduction in obese individuals with and without Type 2 diabetes J Appl. Physiol.; 99: 1220-1225.

Staiger H, Kaltenbach S, Staiger K, Stefan N, Fritsche A, Guirguis A, Peterfi C, Weisser M, Machicao F, Stumvoll M, and Häring HU.(2004):

Expression of adiponectin receptor mRNA in human skeletal muscle cells is related to in vivo parameters of glucose and lipid metabolism. Diabetes.; 53: 2195–2201

Stefann and Stumvol M. (2002):

Adiponectin-its role in metabolism and beyond. Horm.Metab.Res.; 34:469-474.

Stephens, J.M., Lee, J., and Pilch, P.F. (1997):

Tumor necrosis factor- α -induced insulin resistance in 3T3-L1 adipocytes is accompanied by a loss of insulin receptor substrate-1 and GLUT4 expression without a loss of insulin receptor-mediated signal transduction. *J. Biol. Chem.*; 272: 971–976.

Steppan CM.; Baiely ST.; Bhat S.; Brown EJ.; Wright CM. and Lazar MA. (2001):

The hormone resistin links obesity to diabetes mellitus. *Nature*. 409: 307-312.

Steppan, C.M., Bailey, S.T., Bhat, S., Brown, E.J., Banerjee, R.R., Wright, C.M., Patel, H.R., Ahima, R.S., and Lazar, M.A. (2001):

The hormone resistin links obesity to diabetes. *Nature.*; 409: 307–312.

St-Pierre, J., Lin, J., Krauss, S., Tarr, P.T., Yang, R., Newgard, C.B., and Spiegelman, B.M. (2003):

Bioenergetic analysis of peroxisome proliferator-activated receptor coactivators 1 α and 1 β (PGC-1 α and PGC-1 β) in muscle cells. *J. Biol. Chem.* ; 278: 26597–26603.

Stumvoll M, Tschritter O, Fritsche A, Staiger H, Renn W, Weisser M, Machicao F. and Haring H (2002):

Association of the T-G polymorphism in adiponectin (exon 2) with obesity and insulin sensitivity: interaction with family history of type 2 diabetes. *Diabetes.*; 51:37-41.

Suwazono, Y., Sakata, K., Okubo, Y., Harada, H., Oishi, M., Kobayashi, E., Uetani, M., Kido, T., and Nogawa, K. (2006):

Long-term longitudinal study on the relationship between alternating shift work and the onset of diabetes mellitus in male Japanese workers. *J. Occup. Environ. Med.*; 48: 455–461.

Timothy M.cunba, Richard G.Peterson and Troy A. Gobbet (2005):

Differing sources of dietary fat alter the character of metabolic syndrome induced in the C57BL/6 mouse. *Am J Clin Nutr.*; 70:1157.

Trujillo ME and Scherer PE. (2005):

Adiponectin: journey from an adipocyte secretory protein to biomarker of the metabolic syndrome. *J Intern Med.*; 257:167–175

Tsao, T.S.; Lodish, H.F. and Fruebis, J.(2002):

Acrp 30, a new hormone controlling fat & glucose metabolism. *Eur. J. pharmacol.*; 440:213-221.

Tsuchida A, Yamauchi T, Ito Y, Hada Y, Maki T, Takekawa S, Kamon J, Kobayashi M, Suzuki R, Hara K, Kubota N. and Terauchi Y. (2004):

Insulin/Foxo1 pathway regulates expression levels of adiponectin receptors and adiponectin sensitivity. *J Biol Chem.*; 279:30817 - 30822.

Turek, F.W., Joshu, C., Kohsaka, A., Lin, E., Ivanova, G., McDearmon, E., Laposky, A., Losee-Olson, S., Easton, A., and Jensen, D.R. (2005):

Obesity and metabolic syndrome in circadian Clock mutant mice. *Science.*; 308: 1043–1045.

Vivian, V.U.; Michchael, C. and Ridell, G. (2007):

Circulating adiponectin and adiponectin receptor expression in skeletal muscle: effects of exercise. *diabetes/Metabolism research and review.*; 23: 60-611.

Vojarova, B., Weyer, C., Hanson, K., Tataranni, P.A., Bogardus, C., and Pratley, R.E. (2001):

Circulating interleukin-6 in relation to adiposity, insulin action, and insulin secretion. *Obes. Res.*; 9: 414–417.

Wajchenberg BL. (2000):

Subcutaneous and visceral adipose tissue: their relation to the metabolic syndrome. *Endocr Rev.*; 21:697–738.

Wallace MB, Mills BD and Browning CL. (1997):

Effects of cross-training on markers of insulin resistance /hyperinsulinemia. *Med Sci Sports Exerc.*; 29:1170–1175.

Wang CC, Goalstone ML. and Draznin B. (2004):

Molecular mechanisms of insulin resistance that impact cardiovascular biology. *Diabetes.*; 53:2735–40.

Wang H, Zhang H, Jia Y, Zhang Z, Craig R, Wang X, and Elbein SC. (2004):

Adiponectin receptor 1 gene as a candidate for type 2 diabetes an insulin resistance. *Diabetes.*; 53: 2132–2136, 2004

Wellen, K.E. and Hotamisligil, G.S. (2005):

Inflammation, stress, and diabetes. *J. Clin. Invest.*;. 115: 1111–1119.

Wheatcroft SB, Williams IL, Shah AM. and Kearney M.(2003):

Pathophysiological implications of insulin resistance on vascular endothelial function. *Diabet Med.*; 20:255–68.

White MF. (1997):

The insulin signalling system and the IRS proteins. *Diabetologia.*;40 (Supl 2):S2–S17.

Wilfiger, W.W., MACKEY. M. and Chomczynskip (1997):

Effect of PH and ionic strength on the spectrophotometric assessment of nucleic acid purity. *Biotechniques.*; 22:474.

Withers DJ and White M. (2000):

Perspective: The insulin signaling system-a common link in the pathogenesis of type 2 diabetes. *Endocrinology.*;141:1917–21

Wu X, Motoshima H, Mahadev K, Stalker TJ, Scalia R. and Goldstein BJ (2003):

Involvement of AMP-activated protein kinase in glucose uptake stimulated by the globular domain of adiponectin in primary rat adipocytes. *Diabetes.*; 52:1355–1363.

Xu, H., Barnes, G.T., Yang, Q., Tan, G., Yang, D., Chou, C.J., Sole, J., Nichols, A., Ross, J.S., and Tartaglia, L.A. (2003):

Chronic inflammation in fat plays a crucial role in the development of obesity-related insulin resistance. *J. Clin. Invest.*;. 112: 1821–830.

Yamagishi, S.I., Edelstein, D., Du, X.L., Kaneda, Y., Guzman, M. and Brownlee, M. (2001):

Leptin induces mitochondrial superoxide production and monocyte chemoattractant protein-1 expression in aortic endothelial cell increasing fatty acid oxidation via protein kinase A. *J. Biol. Chem.* ; 276: 25096–25100.

Yamauchi T, Kamon J, Minokoshi Y, Ito Y, Waki H, Uchida S, Yamashita S, Noda M, Kita S, Ueki K, Eto K. and Akanuma Y. (2003):

Adiponectin stimulates glucose utilization and fatty-acid oxidation by activating AMP-activated protein kinase. *Nat Med.*; 8:1288–1295.

Yamauchi T, Kamon J, Waki H, Terauchi Y, Kubota N, Hara K, Mori Y, Ide T, Murakami K, Tsuboyama-Kasaoka N, Ezaki O, Yoda M. and Kadowaki T. (2001):

The fat-derived hormone adiponectin reverses insulin resistance associated with both lipodystrophy and obesity. *Nat Med.*; 7:941–946.

Yamauchi T, Nio Y, Maki T, Kobayashi M, Takazawa T, Iwabu M, Okada-Iwabu M, Kawamoto S, Kubota N, Kubota T, Ito Y, Kamon J, Tsuchida A, Kumagai K, Kozono H, Hada Y and Ogata H. (2007):

Targeted disruption of AdipoR1 and AdipoR2 causes abrogation of adiponectin binding and metabolic actions. *Nat Med.*; 13:332–339

Yamauchi T, Waki H, Kamon J, Murakami K, Motojima K, Komeda K, Miki H, Kubota N, Terauchi Y, Tsuchida A, Tsuboyama-Kasaoka N, Yamauchi N, Ide T, Hori W, Kato S, Fukayama M, Akanuma Y, Ezaki O, Itai A, Nagai R, Kimura S, Tobe K, Kagechika H, Shudo K and Kadowaki T (2002):

Inhibition of RXR and PPAR γ ameliorates diet-induced obesity type 2 diabetes. *J Clin Invest.*; 108:1001–1013.

Yang WS, Lee WJ, Funahashi T, Tanaka S, Matsuzawa Y, Chao CL, Chen CL, Tai TY and Chuang LM.(2001):

Weight reduction increases plasma levels of an adipose-derived anti- inflammatory protein, adiponectin. *J Clin Endocrinol Metab.* 86:3815–3819.

Yang, Q., Graham, T.E., Mody, N., Preitner, F., Peroni, O.D., Zabolotny, J.M., Kotani, K., Quadro, L., and Kahn, B.B. (2005):

Serum retinol binding protein 4 contributes to insulin resistance in obesity and type 2 diabetes. *Nature.*; 436: 356–362.

Zeng Q, Fu L, Takekoshi K, Kawakami Y and Isobe K (2007):

Effects of short-term exercise on adiponectin and adiponectin receptor levels in rats. *J Atheroscler Thromb.*;14(5):261-5.

Zimmet, P., Alberti, K.G., and Shaw, J. (2001):

Global and societal implications of the diabetes epidemic. *Nature .*; 414: 782–787.