

REFERENCES

Abbasi, F.; Brown, B.W. Jr.; Lamendola, C.; McLaughlin, T. and Reaven, G.M. (2002):

Relationship between obesity, insulin resistance, and coronary heart disease risk. J. Am. Coll. Cardiol.; 40: 937 – 943.

Adam, M. (2005):

Integrating research and development: the emergence of rational drug design in the pharmaceutical industry. Studies in history and philosophy of biological and biomedical sciences; P. 36.

Aguilera, G.; Kiss, A. and Sunar-Akbasak, B. (1995):

Hyperreninemic hyperaldosteronism after chronic stress in the rat. J. Clin. Invest.; 96(3): 1512 – 19.

Ahijevych, K. and Wewers, M.E. (2003):

Passive smoking and vascular disease. J. Cardiovasc. Nurs.; 18: 69 – 74.

Akhtar, J.; Bejum, S.; Islam, N.; Bejum, N. and Ara, S. (2008):

Study of restrain stress induced changes in serum lipid profile and haemoconcentration and effect sympathetic blocker on it. Mymensingh Med. J.; 17 (2 Suppl.): S 16 – 21.

Al-Ayadhi, L.; Korish, A. and Al-Tuwaijri, A. (2006):

The effect of vitamin E , L-arginine and N-nitro L-arginine methyl ester on endocrine and metabolic changes of rats exposed to acute cold stress. Saudi Med. J.; 27 (1): 17 – 22.

Alexander, K.; Dana, J.; Danila, J. and Olga, K. (2001):

Changes in angiotensin AT1 receptor mRNA levels in the rat brain after immobilization stress and inhibition of central nitric oxide synthase. *Endocrine Regulations*; 35: 65 – 70.

Alexander, R.W. and Dzau, V.J. (2000):

Vascular biology: the past 50 years. *Circulation*; 102(suppl IV): 112 – 116.

Anderson, M.F.; Aberg, M.A.; Nilsson, M. and Erriksson, P.S. (2002):

Insulin- like growth factor-1 and neurogenesis in the adult mammalian brain. *Brain Res. Devel.*; 134(1-2): 115 – 22.

Ando, K. and Fujita, T. (2009):

Metabolic Syndrome and Oxidative Stress. *Free Radic. Biol. Med.*; 47(3): 213 – 8.

Antony, F.S. (2008):

Harrison's principles of internal medicine. McGraw-Hill Medical.

Armando, I.; Carranza, A.; Nishimura, Y.; Hoe, K.; Barontini, M.; Terrón J.A.; Neri, A.F.; Ito, T.; Juorio, A.V. and Saavedra. J.M. (2001):

Peripheral administration of an angiotensin II AT₁ receptor antagonist decreases the hypothalamic-pituitary-adrenal response to isolation stress. *Endocrinology*; 142(9): 3880 – 9.

Armario, A.; Castellanos, J.M. and Balasch. J. (1985):

Chronic noise stress and insulin secretion in male rats. *Physiology & Behavior*; 34(3): 359 – 361.

Armario, A.; Montero, J.L. and Balasch. J. (1986):

Sensitivity of corticosterone and some metabolic variables to graded levels of low intensity stresses in adult male rats. *Physiology & Behavior*; 37(4): 559 – 561.

Armistag, A. (1983):

Textbook of statics.

Arnaud, C.; Dematteis, M.; Pepin, J.L.; Baquet, J.P. and Lévy, P. (2009):

Obstructive sleep apnea, immuno-inflammation and atherosclerosis. *Semin. Immunopathol.*; 31(1): 113 – 25.

Atlas, S.A. (2007):

The renin angiotensin aldosterone system: Pathophysiological role and pharmacologic inhibition. *JMCP*; 13(8): S9 – S20.

Avogadro, A.; Crepaldi, G.; Enzi, G. and Tiengo, A. (1967):

Association of hyperlipidemia, diabetes mellitus with moderate obesity. *Acta Diabetol.*; 4:572 – 590.

Balletshofer, B.M.; Rittig, K.; Enderle, M.D.; Volk, A.; Maerker, E.; Jacob, S.; Matthaei, S.; Rett, K. and Häring, H.U. (2000):

Endothelial dysfunction is detectable in young normotensive first-degree relatives of subjects with type 2 diabetes in association with insulin resistance. *Circulation*; 101: 1780 – 84.

Barnes, V.A.; Davis, H.C.; Murzynowski, J.B. and Treiber, F.A. (2004):

Impact of medication on resting and ambulatory blood pressure and heart rate in youth. *Psychosom. Med.*; 66: 909 – 14.

Barret-Connor, E. and Bush, T.L. (1991):

Estrogen and coronary heart disease in women. *JAMA*; 256: 1861 – 7.

Barton, M.; Carmona, B.; Morawietz, H.; Goettsch, W. and Hillen, C.C. (2000):

Obesity is associated with tissue specific activation of renal angiotensin converting enzyme, in vivo evidence for regulatory role of endothelium. *Hypertension*; 35: 329 – 336.

Barua, R.S.; Ambrose, J.A.; Eales- Reynolds, L.J.; DeVoe, H.C.; Zervas, J.G. and Saha, D.C. (2002):

Heavy and Light cigarette smokers have similar dysfunction of endothelial vasoregulatory activity: an in vivo and in vitro correlation. *J. Am. Coll. Cardiol.*; 39: 1758 – 63.

Basso, N. and Terragno, N.A. (2001):

History about the discovery of the renin-angiotensin system. *Hypertension*; 38(6): 1246 – 9.

Beckman, J.A.; Creager, M.A. and Libby, P. (2002):

Diabetes and Atherosclerosis: epidemiology, pathophysiology and management. *JAMA*; 287(19): 2570 – 2581.

Beck-Nielsen, H. (1999):

General characteristics of the insulin resistance syndrome: prevalence and heritability. European group for the study of insulin resistance (EGIR). *Drugs*; 58(Suppl. 1): 7 – 10 and 75 – 82.

Belchetz, P.E. (1994):

Hormonal treatment of postmenopausal women. *N. Engl. J. Med.*; 330: 1062.

Bhatia, V. and Tandon, R.K. (2005):

Stress and the gastrointestinal tract. *J. Gastroenterol. & Hepatol.*; 20: 332 – 9.

Black, P.H. (2002):

Stress and the inflammatory response. A review of neurogenic inflammation. *Brain Behav. Immun.*; 16: 622 – 53.

Black, P.H. (2003):

The inflammatory response is an integral part of the stress response: implication for atherosclerosis, insulin resistance, type II diabetes and metabolic syndrome X. *Brain Behav. Immun.*; 17: 350 – 64.

Black, P.H. and Garbutt, L.D. (2002):

Stress, inflammation and cardiovascular disease. *J. Psychosom. Res.*; 52(1): 1 – 23.

Blankenhorn, D.H. and Hodis, N.H. (1993):

Atherosclerosis-- reversal with therapy. *West. J. Med.*; 159(2): 172 – 9.

Boden, W.E.; O'Rourke, R.A.; Teo, K.K.; Hartigan, P.M.; Maron, D.J.; Kostuk, W.J.; Knudtson, M.; Dada, M.; Harris, C.L.; Shaw, L.; Gosselin, G.; Nawaz, S.; Title, L.M.; Gau, G.; Booth, D.C.; Bates, E.R.; Spertus, J.A.; Berman, D.S. and Weintraub, W.S. (2007):

Optimal medical therapy with or without PCI for stable coronary disease. *N. Engl. J. Med.*; 356: 1503 – 1516.

Bonnet, F.; Irving, K.; Terra, J.L. ; Nony, P. ; Berthezène, F. and Moulin, P. (2005):

Anxiety and depression are associated with unhealthy lifestyle in patients at risk of cardiovascular disease. *Atherosclerosis*; 178(2): 339 – 44.

Borchard, U. (2001):

The role of the sympathetic nervous system in cardiovascular diseases. *J. Clin. Basic Cardiol.*; 4(3): 175 – 7.

Bouskela, E.; Kraemer de Aguiar, L.G.; Nivoit, P.; Bahia, L.R.; Villela, N.R. and Bottino, D.A. (2007):

Vascular dysfunction in metabolic disorders: evaluation of some therapeutic interventions. Bull. Acad. Natl. Med.; 191(3): 492 – 3.

Bowling, A. (1999):

Ageism in cardiology. BMJ; 319: 1353 – 1355.

Brewster, U.C. and Perazella, M.A. (2004):

Cardiorenal effects of the renin- angiotensin- aldosterone system. Textbook, P. 11 – 20.

Brown, M.J. (2006):

Direct renin inhibition: A new way of targeting the renin system. J. Renin Angiotensin Aldosterone Syst.; 7(2): S7 – S11.

Brunner, E.J.; Hemingway, H.; Walker, B.R.; Page, M.; Clarke, P.; Juneja, M.; Shipley, M.J.; Kumari, M.; Andrew, R.; Seckl, J.R.; Checkley, S.; Rumley, A. and Marmot, M.G. (2007):

Adrenocortical, autonomic, and inflammatory causes of the metabolic syndrome: nested case-control study. Circulation; 106: 2659 – 65.

Brydon, L.; Magid, K. and Steptoe, A. (2005):

Platelets, coronary heart disease and stress. Brain Behav. Immun.; 20 (2): 113 – 9.

Bucolo, G. and David, H. (1973):

Quantitative determination of serum triglycerides by the use of the enzymes. Clin. Chem.; 19: 475.

Burnier, M. and Brunner, H.R. (2000):

Angiotensin II receptor antagonists. Lancet; 355: 637 – 645.

Burtis, C.A. and Ashweed, E.R. (1994):

Tietz textbook of clinical chemistry 2nd Ed. W.B. Saunders Company, Philadelphia. P. 1825 – 27.

Campmany, L.; Pol, O. and Armario, A. (1996):

The effect of two chronic intermittent stressors on brain monoamines. Pharmacology, Biochemistry & Behavior; 53(3): 517 – 523.

Cannon, W.B. (1939):

The wisdom of the body. Norton, New York.

Capuron, L. and Dantzer, R. (2003):

Cytokines and depression: the need for a new paradigm. Brain Behav. Immun.; 17(Suppl. 1): S119 – S124.

Caraway, W.T. and Watts, N.B. (1987):

Carbohydrates in ***Tietz, N.W.*** ed. Fundamentals of clinical chemistry 3rd ed. Philadelphia WB Saunders: P. 422 – 447.

Carey, R.M. and Siragy, H.M. (2003):

Newly recognized component of the renin- angiotensin system: A hypercoagulable state. Endocr. Rev.; 24: 261 – 71.

Carnthon, M.R.; Loria, C.M.; Hill, J.O.; Sidney, S.; Savage, P.J. and Liu, K. (2004):

Risk factors for the metabolic syndrome: the coronary artery risk development in young adults (CARDIA) study, 1985 – 2001. Diabetes Care; 27: 2707 – 15.

Carol Y. (1999):

Stress and the autonomic nervous system "Fight or Flight"

Carr, M.E. (2001):

Diabetes mellitus: a hypercoagulable state. *J. Diabetes Complications*; 15: 44 – 54.

Chen, X.L.; Tummala, P.E.; Olbrych, M.T.; Alexander, R.W. and Medford, R.M. (1998):

Angiotensin II induces monocytes chemoattractant protein-1 gene expression in rat vascular smooth muscle cells. *Circ. Res.*; 83(9): 952 – 9.

Chrousos, G.P. (1995):

The hypothalamic pituitary adrenal axis and immune mediated inflammation. *N. Engl. J. Med.*; 332: 1351 – 62.

Chrousos, G.P. and Gold, P.W. (1992):

The concepts of stress and stress system disorders. Overview of physical and behavioral homeostasis. *JAMA*; 267: 1244 – 52.

Chuang, G.C.; Yang, Z.; Westbrook, D.G.; Pompilius, M.; Ballinger, C.A.; White, C.R.; Krzywanski, D.M.; Postlethwait, E.M. and Ballinger, S.W. (2009):

Pulmonary ozone exposure induces vascular dysfunction, mitochondrial damage and atherogenesis. *Am. J. Physiol. Lung Cell Mol. Physiol.*; 297(2): L209 – 16.

Cottart, C.H.; Laguillier, C.; Nivet-Antoine, V.; Klimczak, C.; Sebban, C. and Beaudeau, J.L. (2009):

Biology of arterial aging and arteriosclerosis. *C. R. Biol.*; 332(5): 433 – 47.

Crepaldi, G.; Tiengo, A. and Manzato, E. (1993):

The plurimetabolic syndrome in textbook of diabetes, obesity and hyperlipidemias. Amsterdam, Netherlands Excerpta Medica, International congress series. Chapter V, P. 1039.

Critchley, J.A. and Capewell, S. (2003):

Mortality risk reduction with smoking cessation in patient with coronary heart disease: Systematic review. JAMA; 290: 86 – 97.

Dallman, M.F.; Pecoraro, N.C. and La Fleur, S.E. (2005):

Chronic stress and comfort food: Self medication and abdominal obesity. Brain Behav. Immun.; 19: 275 – 80.

Dang, A.; Zhang, Y.; Liu, G.; Chen, G.; Song, W. and Wang, B. (2006):

Effects of losartan and irbesartan on serum uric acid in hypertensive patients with hyperuricaemia in Chinese population. J. Hum. Hypertens.; 20(1): 45 – 50.

Daugherty, A.; Manning, M.W. and Gassis, L.A. (2000):

Angiotensin II promotes atherosclerotic lesions and aneurysms in apolipoprotein E-deficient mice. J. Clin. Invest.; 105: 1605 – 12.

Deedwania, P.C. (2003):

Diabetes and vascular disease: common link in the emerging epidemic of coronary artery disease. Am. J. Cardiol.; 91(1): 68 – 77.

De Fronzo, R.A. (1992):

Insulin resistance, hyperinsulinemia, and coronary artery disease: A complex metabolic web. J. Cardiovasc. Pharmacol.; 20 (Suppl. 11): S1 – 16.

De Fronzo, R.A. and Ferrannini, E. (1991):

Insulin resistance. A multifaceted syndrome responsible for NIDDM, obesity, hypertension, dyslipidemia and atherosclerotic cardiovascular disease. *Diabetes Care*; 14: 173 – 94.

De Gasparo, M.; Catt, K.J.; Inagami, T.; Wright, J.W. and Unger, T. (2000):

International union of pharmacology. XXIII. The angiotensin II receptors. *Pharmacol. Rev.*; 52(3): 415 – 72.

De Kloet, R.E. (1991):

Brain corticosteroid receptor balance and homeostatic control. *Front Neuroendocrinol.*; 12: 95 – 164.

De Paulo Brotto, M.A. (2003):

Temporal effects of stress by immobilization and sensitivity of isolated rat pacemaker to isoproterenol: role of corticosterone, neuronal uptake and β -adrenergic homogeneity. *JPET*; 306: 1152 – 58.

De Vriese A.S.; Verbeuren, T.J.; Van de Voorde J.; Lameire N.H. and Vanhoutte, P.M. (2000):

Endothelial dysfunction in diabetes. *Br. J. Pharmacol.*; 130: 963 – 74.

Dhabhar, F.S. and McEwen, B.S. (1999):

Enhancing versus suppressive effects of stress hormones on skin immune function. *PNAS*; 96: 1059 – 64.

Dimitriadis, G.; Leighton, B.; Parry-Billings, M.; Sasson, S.; Young, M.; Krause, U.; Bevan, S.; Piva, T.; Wegener, G. and Newsholme, E.A. (1997):

Effects of glucocorticoid excess on the sensitivity of glucose transport and metabolism to insulin in skeletal muscle. *Biochem. J.*; 321: 707 – 12.

Dimsdale, J.E. (2008):

Psychological stress and cardiovascular disease. *J. Am. Coll. Cardiol.*; 51(13): 1237 – 46.

Di Rossa, M.; Rodomski, M.; Carnuccio, R. and Moncada, S. (1990):

Glucocorticoid inhibits the induction of nitric oxide synthase in macrophages. *Biochem. Biophys. Res. Commun.*; 172: 1246 – 52.

D'mello, A.P.; and Liu, Y. (2006):

Effects of maternal immobilization stress on birth weight and glucose homeostasis in the offspring. *Psychoneuroendocrinology*; 31(3): 395 – 406.

Dronjak, S.; Gavrilović, L.; Filipović, D. and Radojčić, M.B. (2004):

Immobilization and cold stress affect sympatho-adrenomedullary system and pituitary-adrenocortical axis of rats exposed to long term isolation and crowding. *Physiol. Behav.*; 81(3): 409 – 15.

Dronjak, S.; Jezova, D. and Kvetnansky, R. (2006):

Different effects of novel stressors on sympathoadrenal system activation in rats exposed to long term immobilization. *Annals of the New York academy of sciences. Current neuroendocrine and genetic approaches*; 1018: 113 – 123.

Dumont, E.C.; Rafrafi, S.; Laforest, S. and Drolet, G. (1999):

Involvement of central angiotensin receptors in stress adaptation. Neuroscience; 93(3): 877 – 884.

Eckel, R.H. and Krauss, R.M. (1998):

American Heart Association call to action: obesity as a major risk factor for coronary heart disease: AHA Nutrition Committee. Circulation; 97, 2099 – 2100.

Einhorn, D.; Reaven, G.M.; Cobin, R.H.; Ford, E.; Ganda, O.P.; Handelsman, Y.; Hellman, R.; Jellinger, P.S.; Kendall, D.; Krauss, R.M.; Neufeld, N.D.; Petak, S.M.; Rodbard, H.W.; Seibel, J.A.; Smith, D.A. and Wilson, P.W. (2003):

American College of Endocrinology position statement on the insulin resistance syndrome. Endocr. Pract.; 9(3): 237 – 52.

Ellefson, R.D. and Caraway, W.T. (1976):

Fundamentals of clinical chemistry. Ed. **Tietz N.W.**: P. 506.

Esch, T.; Stefano, G.B.; Fricchione, G.L. and Benson, H. (2002):

Stress in cardiovascular diseases. Med. Sci. Monit.; 8(5): RA93 – R101.

Farsang, C. and Fisher, J. (2006):

Optimizing antihypertensive therapy by angiotensin receptor blockers in textbook; analogue-based drug discovery. Wiley- VCH, P. 157 – 167.

Feldeisen, S.E. and Tucker, K.L. (2007):

Nutritional strategies in the prevention and treatment of metabolic syndrome. Appl. Physiol. Nutr. Metab.; 32(1): 46 – 60.

Fenga, C.; Micali, E.; Cacciola, A.; Trimarchi, C. and Germano, D. (2004):

Stressful life events and fibrinogen level in middle aged teachers. Psychopathology; 73(2): 64 – 8.

Ferrario, C.M. (2006):

Role of angiotensin II in cardiovascular disease: Therapeutic implications of more than a century of research. J. Renin Angiotensin Syst.; 7: 3 – 14.

Festa, A.; D'Agostino, R. Jr; Tracy, R.P. and Haffner, S.M. (2002):

Elevated levels of acute-phase proteins and plasminogen activator inhibitor-1 predict the development of type 2 diabetes: the insulin resistance atherosclerosis study. Diabetes; 51: 1131 – 1137.

Fleschner, M.; Watkins, L.R.; Belligrau, D.; Laudenslager, M.L. and Maier, S.F. (1992):

Specific changes in lymphocyte subpopulation: a mechanism for stress induced immunomodulation. Neuroimmunol.; 41: 131 – 42.

Fletcher, G.F.; Balady, C.G.; Blair, S.N.; Blumenthal, J.; Caspersen, C.; Chaitman, B.; Epstein, S.; Froelicher, S.S.; Froelicher, V.F.; Pina, I.L. and Pollock, M.L. (1996):

Statement on exercise: benefits and recommendation for physical activity programs for all Americans: a statement for health professionals by the committee on exercise and cardiac rehabilitation of the council of clinical cardiology, American Heart Association. Circulation; 94: 857.

Flores-Riveros, J.R. (1993):

Insulin downregulate expression of the insulin-responsive glucose transporter (GLUT4): effects on transcription and mRNA turnover. PNAS; 90(2): 512 – 6.

Folkow, B. (1993):

Physiological organization of neurohormonal responses to psychological stimuli: implication for health and disease. Ann. Behav. Med.; 15: 236 – 44.

Folkow, B. (1997):

Physiological aspects of the defense and defeat reaction. Acta. Physiol. Scand.; 161(Suppl 640): 34 – 47.

Ford, E.S.; Giles, W.H. and Dietz, W.H. (2002):

Prevalence of metabolic syndrome among US adults: findings from the third National Health and Nutrition Examination Survey. JAMA; 287(3): 356 – 9.

Fowler, B. (2005):

Homocysteine--an independent risk factor for cardiovascular and thrombotic diseases. Ther. Umsch.; 62(99): 641 – 6.

Fukuchi, S.; Hamaguchi, K.; Seike, M.; Himeno, K.; Sakata, T., Yoshimatsu, H. (2004):

Role of fatty acid composition in the development of metabolic disorders in sucrose-induced obese rats. Exp. Biol. Med.; 229(6): 486 – 93.

Fuster, V.; Aleander, R.W.; Rouke, R.A.; Roberts, R.; King III, S.B. and Ellens, H.J. (2001):

Hurst's The Heart. Textbook, 10th ed., Vol. 1, P. 1131 & 1237.

Garcia, R. (2002):

Stress, synaptic plasticity and psychopathology. *Rev. Neurosci.*; 13(3): 195 – 208.

Gaspaini, L. and Xu, H. (2003):

Potential role of insulin and IGF-1 in Alzheimer's disease. *Trends Neurosci.*; 26(8): 404 – 406.

Gavras, I. and Gavras, H. (2002):

Angiotensin II as a cardiovascular risk factor. *J. Hum. Hypertens.*; 16(Suppl. 2): S2 – S6.

Ginsberg, H.N. (2000):

Insulin resistance and cardiovascular disease. *J. Clin. Invest.*; 106: 453 – 8.

Gold, P.W. and Chrousos, G.P. (2002):

Organization of the stress system and its dysregulation in melancholic and atypical depression: high vs low CRH/NE state. *Mol. Psychiatry*; 7(3): 254 – 75.

Golin, R.M.A.; Gotoh, E.; Said, S.I. and Ganong, W.F. (1988):

Pharmacological evidence that the sympathetic nervous system mediates the increase in renin secretion produced by immobilization and head tilt in rats. *Neuropharmacology*; 27: 1209 – 13.

Goodfriend, T.L.; Elliot, M.E. and Catt, K.J. (1996):

Angiotensin receptors and their antagonists. *N. Engl. J. Med.*; 334: 1649 – 55.

Gosse, P. and Dallochio, M. (1993):

Left ventricular hypertrophy: epidemiological prognosis and associated critical factors. *Eur. Heart J.*; 14: 16 – 21.

Gould, A.L.; Rossouw, J.E.; Santanello, N.C.; Heyse, J.F. and Furberg, C.D. (1998):

Cholesterol reduction yields clinical benefits impact of statin trials. Circulation; 97: 946 – 952.

Green, P.G.; Janig, W. and Levine, J.D. (1997):

Negative feedback neuroendocrine control of inflammatory response in rats is dependent on the sympathetic postganglionic neuron. J. Neurosci.; 17: 3234 – 8.

Grundy, S.M. (2002):

Obesity, metabolic syndrome, and coronary atherosclerosis. Circulation; 105: 2696 – 98.

Grundy, S.M. (2006):

Metabolic syndrome: connecting and reconciling cardiovascular and diabetes worlds. J. Am. Col.; 47: 1093 – 1100.

Grundy, S.M.; Brewer, H.B. Jr; Cleeman, J.I.; Smith, S.C. Jr and Lenfant, C. (2004):

Definition of the metabolic syndrome: report of the National heart, lung and blood institute/ American heart association conference on scientific issue related to definition. Circulation; 163: 427 – 436.

Grundy, S.M.; Hansan, B.; Smith, S.C. Jr; Cleeman, J.I. and Khan R.A. (2004*):

Clinical management of the metabolic syndrome: report of American heart association/ National heart, lung and blood institute/ American diabetes association conference on scientific issue related to management. Circulation; 109: 551 – 556.

Grundy, S.M.; Pasternak, R.; Greenland, P.; Smith, S. and Fuster, V. (1999):

Assessment of cardiovascular risk by use of multiple risk factor assessment equation: A statement for health care professionals from American Heart Association and American College of Cardiology. *Circulation*; 100: 1481 – 92.

Guerci, B.; Bohm, P.; Kearney –Schawartz, A.; Zannad, F. and Drouin, P. (2001):

Endothelial dysfunction and type 2 diabetes. *Diabetes Metab.*; 27(4): 436 – 47.

Haffner, S.M.; Lehto, S.; Ronnemaa, T.; Pyroala, K. and Laakso, M. (1998):

Mortality from coronary heart disease in subjects with type 2 diabetes and in non diabetic subjects with and without prior myocardial infarction. *N. Engl. J. Med.*; 339(4): 229 – 234.

Hakim, A.A.; Curb, J.D.; Petrovitch, H.; Rodriguez, B.L.; Yano, K.; Ross, G.W.; White, L.R. and Abbot, R.D. (1999):

Effect of walking on coronary heart disease in elderly men. *Circulation*; 100: 9 – 13.

Haller, H. (1977):

Epidemiology and associated risk factors of hyperlipoproteinemia. *Z. Gesamte Inn. Med.*; 32(8): 124 – 8.

Hawley, J.A. and Lessard, S.J. (2008):

Exercise training- induced improvements in insulin action. *Acta. Physiol.*; 192(1): 127 – 35.

Hem, L.; Ahuja, G.K. and Narang, A.P. (2002):

Effect of starvation stress on lipid peroxidation and lipid profile in rabbits. Indian J. Physiol. Pharmacol.; 46, 371 – 374.

Higuchi, S.; Ohtsu, H.; Suzuki, H.; Shirai, H.; Frank, G.D. and Eguchi, S. (2007):

Angiotensin II signal transduction through the AT1 receptor: novel insights into mechanisms and pathophysiology. Clin. Sci.; 112(8): 417 – 28.

Hjemdahl, P. (2002):

Stress and the metabolic syndrome. Circulation; 106: 2634.

Hopfner, R.L. and Gopalakrishnan, V. (1999):

Endothelin: emerging role in diabetic vascular complications. Diabetologia; 42: 1383 – 94.

Hu, F.B.; Stampfer, M.J.; Solomon, C.G.; Liu, S.; Willett, W.C.; Speizer, F.E.; Nathan, D.M. and Manson, J.E. (2001):

The impact of diabetes mellitus on mortality from all causes and coronary heart disease in women: 20 years of follow-up. Arch. Intern. Med.; 161(14): 1717 – 23.

Imai, T.; Miyazaki, H.; Hirose, S.; Hori, H.; Hayashi, T.; Kageyama, R.; Ohkubo, H.; Nakanishi, S. and Murakami, K. (1983):

Cloning and sequence analysis of cDNA for human renin precursor. Proc. Natl. Acad. Sci. U.S.A.; 80(24): 7405 – 9.

Innes, K.E.; Vincent, H.K. and Taylor, A.G. (2007):

Chronic stress and insulin resistance- related indices of cardiovascular diseases risk, part 1: Neurophysiological responses and pathological sequelae. Altern. Ther. Health Med.; 13(4): 46 – 52.

Inogushi, T.; Li, P.; Umeda, F.; Yu, H.Y.; Kakimoto, M.; Imamura, M.; Aoki, T.; Etoh, T.; Naruse, M.; Sano, H.; Utsumi, H. and Nawata, H. (2000):

High glucose level and free fatty acid stimulate reactive oxygen species production through protein kinase C– dependent activation of NAD(P)H oxidase in cultured vascular cells. *Diabetes*; 49: 1939 – 45.

Jackson, E.K. (2006):

Renin and Angiotensin in the textbook; the Pharmacological Basis of Therapeutics. New York McGraw Hill, 11th ed. P. 789 – 821.

Jain, S.K.; Pandey, S.N.; Srivastava, R.K. and Ghosh, S.K. (2000):

Stress and serum cholesterol levels. An experimental study. *J. Anat. Soc. India*; 49(2): 165 – 167.

Jesova, D.; Ochedalski, T.; Kiss, A. and Aguilera, G. (1998):

Brain angiotensin II modulates sympathoadrenal and hypothalamic pituitary adrenocortical activation during stress. *J. Neuroendocrinology*, 10: 67 – 72.

John, C.W. and Alexander, S.K. (2002):

Endocrine Responses to Unpredictable Environmental Events: Stress or Anti-Stress Hormones? *Integrative and Comparative Biology*; 42(3): 600 – 609.

Jousilahti, P.; Vartiainen, E.; Tuomilehto, J. and Puska, P. (1999):

Sex, age, cardiovascular risk factors and coronary heart disease: prospective follow up study of 14,786 middle aged men and women in Finland. *Circulation*; 99: 165 – 72.

Joy, T.; Lahiry, P.; Pollex, R.L. and Hegele, R.A. (2008):

Genetics of metabolic syndrome. *Curr. Diab. Rep.*; 8(2): 141 – 8.

Juhan-Vaugh, I.; Morange, P.E. and Alessi, M.C. (2002):

The insulin resistance syndrome: implication for thrombosis and cardiovascular disease. *Pathophysiol. Haemost. Thromb.*; 32(5-6): 269 – 273.

Kamari, Y.; Hrari, A.; Shaish, A.; Peleg, E.; Sharabi, Y.; Harats, D. and Grossman, E. (2008):

Effect of telmisartan, angiotensin II receptor antagonist, on metabolic profile in fructose-induced hypertensive, hyperinsulinemic hyperlipidemic rats. *Hypertens. Res.*; 31(1): 135 – 40.

Karatas, O.F.; Bayrak, O.; Cimentepe, E. and Unal, D. (2009):

An insidious risk factor for cardiovascular disease: Benign prostatic hyperplasia. *Int. J. Cardiol.*; 144(3): 452.

Katzmarzyk, P.T.; Leon, A.S.; Wilmore, J.H.; Skinner, J.S.; Rao, D.C. Rankinen, T. and Bouchard, C. (2003):

Targeting the Metabolic Syndrome with Exercise: Evidence from the HERITAGE Family Study. *Med. Sci. Sports Exerc.* 35(10): 1703 – 9.

Kawano, H.; Do, Y.S.; Kawano, Y.; Starnes, V.; Barr, M.; Law, R.E. and Hsueh, W.A. (2000):

Angiotensin II has multiple profibrotic effects in human cardiac fibroblasts. *Circulation*; 101(10): 1130 – 7.

Kazemi, M.B. Eshraghian, K.; Omrani, G.R.; Lankarani, K.B. and Hosseini, E. (2006):

Homocysteine level and coronary artery disease. *Angiology*; 57(1): 9 – 14.

Kim, K.S.; Own, W.L.; Williams, D. and Adams-Campbell, L.L. (2000):

A comparison between BMI and conicity index on predicting coronary heart disease: The Framingham Heart study. *Ann. Epidemiol.*; 10(7): 424 – 31.

Koko, V. Djordjević, J.; Cvijić, G. and Davidović, V. (2004):

Effect of acute heat stress on rat adrenal glands: a morphological and stereological study. *J. Experimental Biology*; 207: 4225 – 4230.

Kranzhofer, R.; Browatzki, M.; Schmidt, J. and Kubler, W. (1999):

Angiotensin II activates the proinflammatory transcription factor nuclear factor- kappa B in human monocytes. *Biochem. Biophys. Res. Commun.*; 257: 826 – 8.

Krentz, A.J. (2006):

Prevention of cardiovascular complications of the metabolic syndrome: Focus on pharmacotherapy *in metabolic syndrome and related disorders*, textbook. Mary Ann Liebert Inc., Vol. 4, P. 328 – 341.

Kubo, T.; Numakura, H.; Endo, S.; Hagiwara, Y. and Fukumori, R. (2001):

Angiotensin receptor blockade in the anterior hypothalamic area inhibits stress induced pressor response in rat. *Brain Research Bulletin*; 56(6): 569 – 574.

Kyvelou, S.G.; Vyssoulis, G.P.; Karpanou, E.A.; Adamopoulos, D.N.; Zervoudaki, A.I.; Pietri, P.G. and Stefandis, C.I (2006):

Effect of antihypertensive treatment with angiotensin II receptor blockers on lipid profile: An open multi-drug comparison trial. *Hellenic J. Cardiol*; 47: 21– 28.

Laakso, M.; Karjalainen, L. and Lempiäinen-Kuosa, P. (1996):

Effect of losartan on insulin sensitivity in hypertensive subjects. Hypertension; 28(3): 392 – 6.

Lacy, F.; O’connor, D.T. and Schmid- Schonbein, G.W. (1998):

Plasma hydrogen peroxide production in hypertensive and normotensive subjects at genetic risk of hypertension. J. Hypertens.; 16: 291 – 303.

Lakka, T.A. and Laaksonen, D.E. (2007):

Physical activity in prevention and treatment of the metabolic syndrome. Applied physiology, nutrition, and metabolism. 32(1): 76 – 88.

Lander, K.; Hallqwst, J.; Reuterwall, C.; Ahlbam, A. and DeFire, V. (2001):

Family history of coronary heart disease, strong risk factor for myocardial infarction interacting with other risk factors: results from Stockholm Heart Epidemiology Program (SHEEP). Epidemiology; 12(2): 215 – 21.

Landsberg, L. (2001):

Insulin-mediated sympathetic stimulation: role in the pathogenesis of obesity-related hypertension (or, how insulin affects blood pressure, and why). J. Hypertens.; 19: 523 – 8.

Laragh, J.H.; Angers, M.; Kelly, W.J. and Lieberman, S. (1960):

Hypotensive agents and pressor substances: the effect of epinephrine, angiotensin II, and others on the secretory rate of aldosterone in man. JAMA; 174: 234 – 40.

Lazarus, R.S. (1993):

Coping theory and research: past, present and future. Psychosom. Med.; 55: 234 – 47.

Lerch, M.; Teuscher, A.U.; Beissner, P.; Schneider, M.; Shaw, S.G. and Weidmann, P. (1998):

Effect of angiotensin II-receptor blockade with losartan on insulin sensitivity, lipid profile, and endothelin in normotensive offspring of hypertensive parents. J. Cardiovasc. Pharmacol.; 31(4): 576 – 80.

Li, Y.; Woo, V. and Bose R. (2001):

Platelet hyperactivity and abnormal $\text{Ca}^{(2+)}$ homeostasis in diabetes mellitus. Am. J. Physiol. Heart Circ. Physiol.; 280: H1480 – H1489.

Linden, W.; Stossel, C. and Maurice, J. (1996):

Psychological interventions for patients with coronary artery disease: a meta- analysis. Arch. Intern. Med.; 156: 745 – 52.

Loka, T.; Tasaki, H.; Yashiro, A.; Yamashita, K.; Zozumi, K.; Tsotsui, M.; Kouzuma, R.; Okazaki, M. and Nakashima, Y. (2002):

Association between plasma lipoprotein(a) and endothelial dysfunction in normocholesterolemic and non-diabetic subjects with angiographically normal coronary arteries. Circ. J.; 66(3): 267 – 71.

Lonnroth, P. and Smith, U. (1983):

Beta adrenergic dependent down-regulation of insulin binding in rat adipocytes. Biochem. Biophys. Res. Commun.; 112: 972 – 9.

Lorell, B.H. (1999):

Role of angiotensin AT_1 and AT_2 receptors in cardiac hypertrophy and disease. Am. J. Cardiol.; 83: 48H – 52H.

Ludwig, D.S.; Pereira, M.A.; Kroenke, C.H.; Hilner, J.E.; Van Horn, L.; Slattery, M.L. and Jacobs, D.R. Jr. (1999):

Dietary fiber, weight gain, and cardiovascular disease risk factors in young adults. *JAMA*; 282(16): 1539 – 46

Luke, R.H. and Michael, L.R. (2001):

Corticosterone and insulin interact to regulate glucose and triglyceride levels during stress in a bird .*Am. J. Physiol. Regul. Integr. Comp. Physiol.*; 281: R994 – R1003.

Malik, S.; Wong, N.D. and Franklin, S.S. (2004):

Impact of the metabolic syndrome on mortality from coronary heart disease, cardiovascular disease, and all causes in United States adults. *Circulation*; 110: 1245 – 50.

Mallika, V.; Goswami, B. and Rajappa, M. (2007):

Atherosclerosis pathophysiology and the role of novel risk factors: a clinicobiochemical perspective. *Angiology*; 58(5): 513 – 22.

Mandinov, L.; Eberli, F.R.; Seiler, C. and Hess, O.M. (2000):

Diastolic heart failure. *Cardiovasc. Res.*; 45: 813 – 25.

Mar, S., Aguado, F., Sánchez, T. and Saphier, D. (1998):

Neuroendocrine and immunocytochemical demonstrations of decreased hypothalamo-pituitary-adrenal Axis responsiveness to restraint stress after long-term social isolation. *Endocrinology*; 139: 579 – 587.

Maseri, A. and Fuster, V. (2003):

Is there a vulnerable plaque?. *Circulation*; 107(16): 2068 – 71.

Maurer, M.S. (2003):

Age: a non modifiable risk factor?. *JAAC*; 42(8): 1427 – 1428.

McBride, P.E. (1992):

The health and consequence of smoking and cardiovascular diseases. *Med. Clin. North Am.*; 76: 333 – 53.

McEwen, B.S. (1998):

Protective and damaging effects of stress mediators. *N. Engl. Med. J.*; 338: 171 – 9.

McEwen, B.S. (2002):

The neuroendocrinology and neurobiology of stress. Implication for post-traumatic disorder from a basic science prospective. *Psychiatr. Clin. North Am.*; 1018: 424 – 9.

McEwen, B.S. (2002*):

Protective and damaging effects of stress mediators: the good and bad sides of the response to stress. *Metabolism*; 51(6 suppl. 1): 2 – 4.

McEwen, B.S. (2003):

Mood disorders and allostatic load. *Biol. Psychiatry*; 54(3): 200 – 7.

McEwen, B.S.; Magarinos, A. and Reagan, L. (2002):

Studies of hormone action in the hippocampal formation: possible relevance to depression and diabetes. *J. Psychosom. Res.*; 53(4): 883 – 90.

McEwen, B.S. and Stellar, E. (1993):

Stress and the individual. Mechanisms leading to disease. *Arch. Intern. Med.*; 153: 2093 – 2101.

McGarry, J. (2002):

Dysregulation of fatty acid metabolism in the aetiology of type 2 diabetes. *Diabetes*; 51(1): 7 – 18.

Meggs, L.G.; Goupet, J. Huang, H.; Cheng, W.; Li, P.; Capasso, J.M.; Homcy, C.J. and Anversa, P. (1993):

Regulation of angiotensin II receptors on ventricular myocytes after myocardial infarction in rats. *Circ. Res.*; 72(6): 1149 – 62.

Merali, Z.; Hayley, S.; Kent, P.; McIntosh, J.; Bédard, T. and Anisman, H. (2009):

Impact of repeated stressor exposure on the release of corticotrophin releasing hormone, arginine- vasopressin and bombesin like peptides at the anterior pituitary. *Behav. Brain Res.*; 198(1): 105 – 12.

Messerli, F.H.; Weber, M.A. and Brunner, H.R. (1996):

Angiotensin II receptor inhibition. *Arch. Intern. Med.*; 156: 1957 – 65.

MicultKova, L.; krepsova, K.; Sabban E.; Krizanova, O. and Kvetnansky, R. (2004):

Modulation of catecholamine-synthesizing enzymes in rat heart by repeated immobilization stress. *Ann. N.Y. Acad. Sci.*; 1018: 424 – 9.

Misra, A. and Vikram, N.K. (2002):

Insulin resistance syndrome (metabolic syndrome) and Asian Indians. *Curr. Sci.*; 83(12): 1483 – 96.

Mizokami, T.; Wu, L.A.; El-Kaissi, S. and Wall, J.R. (2004):

Stress and thyroid autoimmunity. *Thyroid*; 14: 1047 – 55.

Morgan, I.; Broughton, P.F. and Kalsheker, N. (1996):

Angiotensinogen: molecular biology, biochemistry and physiology. *Int. J. Biochem. Cell Biol.*; 28: 1211 – 22.

Morse, S.A; Zhang, R.; Thakur, V. and Reisin, E. (2005):

Hypertension and the metabolic syndrome. Am. J. Med. Sci.; 330: 303 – 310.

Muhtz, C.; Zyriax, B.C.; Klähn, T.; Windler, E. and Otte, C. (2009):

Depressive symptoms and metabolic risk: Effects of cortisol and gender. Psychoneuroendocrinology; 34(7): 1004 – 1011.

Mulcahy, M.P.; Ahmed, K.; Luzio, S.; Owens, D.R. and Lazarus, J.H. (2002):

Effect of losartan on insulin sensitivity and serum lipids in hypertensive type 2 diabetic patients. Br. J. Diabetes Vasc. Dis.; 2: 67 – 8.

Murali, B. and Goyal, R.K. (2001):

Improvement in insulin sensitivity by losartan in non insulin-dependent diabetic (NIDDM) rats. Pharmacol. Res.; 44(5): 385 – 9.

Nagaraja, H.S.; Anupama, B.K. and Jeganathan, P.S. (2006):

Stress responses in albino rats. J. Physiol. Sci.; 19(2): 8 – 15.

Nagaraja, H.S. and Jeganathan, P.S. (1999):

Influence of different types of stress on selected cardiovascular parameters in rats. Indian J. Physiol. Pharmacol.; 43(3): 296 – 304.

Naito HK (2003):

Coronary artery disease and disorders of lipid metabolism, clinical chemistry: theory. analysis. correlation. 4th ed Kaplan LA and pesce AJ; P.603.

Nakamura, S.; Nakamura, I.; Ma, L.; Vaughan, D.E. and Fogo, A.B. (2000):

Plasminogen activator inhibitor-1 expression is regulated by angiotensin type 1 receptor in vivo. *Kidney Int.*; 58(1): 251 – 9.

Natali, A. and Ferrannini, E. (2004):

Hypertension, insulin resistance and the metabolic syndrome. *Endocrinol. Metab. Clin. North Am.*; 33(2): 417 – 29.

Neaton, J.D. and Wentworth, D. (1992):

Serum cholesterol, blood pressure, cigarette-smoking and death from coronary heart disease: overall findings and differences by age for 316099 white men. *Arch. Intern. Med.*; 152: 56 – 64.

Nickenig, G. and Harrison, D.G. (2002):

The AT(1)-type angiotensin receptor in oxidative stress and atherogenesis: part II: AT(1) receptor regulation. *Circulation*; 105: 530 – 6.

Nickenig, G.; Jung, O.; Strehlow, K.; Zolk, O.; Linz, W.; Scholkens, B.A. and Bohm, M. (1997):

Hypercholesterolemia is associated with enhanced angiotensin AT1-receptor expression. *Am. J. Physiol.*; 272: H2701 – H2707.

Nicole A.J.; Eric J.G.; Boomsma, F.; Anton H. (2006):

Effect of low-density lipoprotein cholesterol on angiotensin II sensitivity: A randomized trial with fluvastatin. *Hypertension*; 47: 1125.

Nygaard, O.; Nordrehaug, J.E.; Refsum, H.; Ueland, P.N.; Farstad, M. and Vollset, S.E. (1997):

Plasma homocysteine levels and mortality in patients with coronary artery disease. *N. Engl. J. Med.*; 337: 230 – 6.

Pacak, K. and Palkovits, M. (2001):

Stressor specificity of central neuroendocrine responses: Implication for stress related disorders. *End. Rev.*; 22 (4): 502 – 48.

Pacak, K.; Palkovits, M.; Yadid, G.; Kvetnasky, R.; Kopin, I.J. and Goldstein, D.S. (1998):

Heterogenous neurochemical response to different stressors: a test of Selye's doctrine of nonspecificity. *Am. J. Physiol.*; 275: R1247 – 55.

Pacifico, L.; Cantisani, V.; Anania, C.; Bonaiuto, E.; Martino, F.; Pascone, R. and Chiesa, C. (2009):

Serum uric acid and its association with metabolic syndrome and carotid atherosclerosis in obese children. *Eur. J. Endocrinol.*; 160(1): 45 – 52.

Park, Y.W.; Zhu, S.; Palaniappan, L.; Heshka, S.; Carnethon, M.R. and Heymsfield, S.B. (2003):

The metabolic syndrome: prevalence and associated risk factor findings in the US population from the Third National Health and Nutrition Examination Survey, 1988–1994. *Arch. Intern. Med.*; 163: 427 – 436.

Paul, M.; Mehr, A.P. and Kreutz, R. (2006):

Physiology of local renin- angiotensin system. *Physiol. Rev.*; 86: 747 – 803.

Pederson, O.D.; Gram, G. and Jespersen, J. (1995):

Plasminogen activator inhibitor type determines plasmin formation in patients with ischemic heart disease. *Thromb. Haemost.*; 73: 835 – 40.

Phillips, G.B. (1978):

Sex hormones, risk factors and cardiovascular disease. *Am. J. Med.*; 65: 7 – 11.

Pilkis, S.J. and Granner, D.K. (1992):

Molecular physiology of the regulation of hepatic gluconeogenesis and glycolysis. *Annu. Rev. Physiol.*; 54: 885 – 909.

Ploghaus, A.; Narain, C. and Beckmann, C.F. (2001):

Exacerbation of pain by anxiety is associated with activity in a hippocampal network. *J. Neurosci.*; 21(24): 9896 – 9903.

Pollex, R.L. and Hegele, R.A. (2006):

Genetic determinants of the metabolic syndrome. *Nat. Clin. Pract. Cardiovasc. Med.*; 3(9): 482 – 9.

Polonsky, K.S. and O'Meara, N.M. (1995):

Secretion and metabolism of insulin, proinsulin, and C-peptide in ***DeGroot, L.J. (ed); Endocrinology. W.B. Saunders Co., Philadelphia, P. 1354 – 1376.***

Porter, J.P. (2000):

Contribution of central ANG. II to acute stress- induced changes in baroreflex function in young rats. *Am. J. Physiol.*; 279(4): R1386 – R1391.

Prasad, A. and Quyyumi, A.A. (2004):

Renin angiotensin system and angiotensin receptor blockers in the metabolic syndrome. *Circulation*; 110: 1507 – 12.

Raasch, W.; Wittmershaus, C.; Dendorfer, A.; Voges, I.; Pahlke, F.; Dodt, C.; Dominiak, P. and Jöhren, O. (2006):

Angiotensin II inhibition reduces stress sensitivity of hypothalamo-pituitary- adrenal axis in spontaneously hypertensive rats. *Endocrinology*; 147(7): 3539 – 46.

Raikkonen, K.; Keltikangas-Järvinen, L.; Adlercreutz, H. and Hänttinen, A. (1996):

Psychological stress and the insulin resistance syndrome. *Metabolism*; 45: 1533 – 8.

Ran, J.; Hirano, T. and Adachi, M. (2004):

Chronic ANG II infusion increases plasma triglyceride level by stimulating hepatic triglyceride production in rats. *Am. J. Physiol. Endocrinol. Metab.*; 287(5): E955 – 61.

Raygada, M.; Shaham, Y.; Nespor, S.M.; Kant, G.J. and Grunberg, N.E. (1992):

Effect of stress on hypothalamic insulin in rats. *Brain Research Bulletin*; 29(2): 129 – 134.

Reaven, G.M. (1988):

Role of insulin resistance in human disease. *Diabetes*; 37: 1595 – 1607.

Reaven, G.M. (1994):

Syndrome X: 6 years later. *J. Intern. Med.*; 736: 13 – 22.

Reaven, G.M. (1995):

Pathophysiology of insulin resistance in human disease. *Physiol. Rev.*; 75(3): 473 – 86.

Reaven, G.M. (2005):

Insulin resistance, the insulin resistance syndrome, and cardiovascular disease. *Panminerva Med.*; 47(4): 201 – 10.

Reaven, G.M.; Abbasi, F. and McLaughlin, T. (2004):

Obesity, insulin resistance and cardiovascular disease. *Recent Prog. Horm. Res.*; 59: 207 – 23.

Reaven, G.M.; Lithell, H. and Landesberg, L. (1996):

Hypertension and associated metabolic abnormalities – the role of insulin resistance and sympathoadrenal system. *N. Engl. J. Med.*; 334: 374 – 381.

Reaven, P. (2004):

Metabolic syndrome. *J. Insur. Med.*; 36(2): 132 – 142.

Redondo, M.; Watzke, H.H.; Stucki, B.; Sulzer, I.; Baisiutti, F.D.; Binder, B.R.; Furlan, M.; Lämmle, B. and Wuillemin, W.A. (1999):

Coagulation factors II, V, VII, and X, prothrombin gene 20210G -->A transition, and factor V Leiden in coronary artery disease: high factor V clotting activity is an independent risk factor for myocardial infarction. *Arterioscler. Thromb. Vasc. Biol.*; 19(4): 1020 – 5.

Renaldi, O.; Pramono, B.; Sinorita, H.; Purnomo, L.B.; Asdie, R.H. and Asdie, A.H. (2009):

Hypoadiponectinemia: a risk factor for metabolic syndrome. *Acta. Med. Indones*; 41(1): 20 – 4.

Reudelhuber, T.L. (2005):

The renin- angiotensin System: Peptides and enzymes beyond angiotensin II. *Curr. Opin. Nephrol. Hypertens.*; 14: 155 – 159.

Reudelhuber, T.L. (2005*):

The continuing saga of the AT₂ receptor: A case of the good, the bad, and the innocuous. *Hypertension* 46(6): 1261 – 2.

Ricart-Jané, D.; Rodríguez-Sureda, V.; Benavides, A.; Peinado-Onsurbe, J.; López-Tejero, M.D. and Llobera M. (2002):

Immobilization stress alters intermediate metabolism and circulating lipoproteins in the rat. *Metabolism*; 51(7): 925 – 31.

Rich E.L. and Romero M. (2005):

Exposure to chronic stress downregulates corticosterone responses to acute stressors. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* 288: R1628 – R1636.

Rizzo, M.; Kotur-Stevuljevic, J.; Berneis, K.; Spinass, G.; Rini, G.B.; Jelic-Ivanovic, Z.; Spasojevic-Kalimanovska, V. and Vekic, J. (2009):

Atherogenic dyslipidemia and oxidative stress: A new look. *Transl. Res.*; 153(5): 217 – 23.

Robert, M.S.; Michael, L.R. and Allan, U.M. (2000):

How do glucocorticoids influence stress responses? integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine Reviews*; 21(1): 55 – 89.

Roberts, C.K. and Seidell, K.K. (2009):

Oxidative stress and metabolic syndrome. *Life Sci.*; 84(21-22): 705 – 12.

Rodnick, K.J.; Mondon, C.E.; Haskell, W.L.; Azhar, S. and Reaven, G.M. (1990):

Differences in insulin induced glucose uptake and enzyme activity in running rats. *J. Appl. Physiol.*; 68: 513 – 19.

Rodriguez-Sureda, V.; López-Tejero, M.D.; Llobera, M. and Peinado-Onsurbe, G.M. (2007):

Social stress profoundly affect lipid metabolism: over-expression of SR-BI in liver and changes in lipids and lipases in plasma and tissues of stressed mice. *Atherosclerosis*; 195(1): 57 – 65.

Ronald, D.K. (2003):

Hormones, brain and stress. *Endocrine Regulations*; 37: 51 – 68.

Rosmond, R. (2003):

Association studies of genetic pleomorphisms in central obesity: a critical review. *Int. J. Obes. Relat. Metab. Disord.*; 27: 1141 – 51.

Rosmond, R. (2005):

Role of stress in the pathogenesis of the metabolic syndrome. *Psychoneuroendocrinology*; 30: 1 – 10.

Rozanski, A.; Blumenthal, J.A. and Kaplan, J. (2000):

Impact of psychological factors on the pathogenesis of cardiovascular disease and implication for therapy. *Circulation*; 101(16): E177 – 8.

Saavedra, J.M.; Ando, H.; Armando, I.; Baiardi, G.; Bregonzio, C.; Jezova, M. and, Zhou, J. (2004):

Brain angiotensin II, an important stress hormone: Regulatory sites and therapeutic opportunities. *Ann. N. Y. Acad. Sci.*; 1018: 76 – 84.

Saavedra, J.M. and Benicky, J. (2007):

Brain and peripheral angiotensin II play a major role in stress. *Stress*; 10(2): 185 – 93.

Sacks, B.D. (1994):

Carbohydrates in **Burtis, C.A. and Ashweed, E.R. (Eds)** Tietz textbook of clinical chemistry 2nd Ed. W.B. Saunders Company, Philadelphia. P. 1815.

Saiki, Y.; Watanabe, T.; Tan, N.; Matsuzaki, M. and Nakamura, S. (1997):

Role of central ANG II receptors in stress- induced cardiovascular and hyperthermic responses in rats. *Am. J. Physiol.*; 272(1 Pt 2): R26 – R33.

Samuel, K.C. (2005):

Atherosclerosis and occlusion of the internal carotid artery. *J. Pathol. & Bacteriol.*; 71: 391 – 401.

Sánchez, O.; Arnau, A.; Pareja, M.; Poch, E.; Ramírez, I. and Soley, M. (2002):

Acute stress-induced tissue injury in mice: differences between emotional and social stress .*Cell Stress Chaperones*; 7(1): 36 – 46.

Schnee, J.M. and Hsueh, W.A. (2000):

Angiotensin II, adhesion and cardiac fibrosis. *Cardiovasc. Res.*; 46: 264 – 8.

Schneider, D.J. (2005):

Abnormalities of coagulation, platelet function and fibrinolysis associated with syndrome of insulin resistance. *Coron. Artery Dis.*; 16: 473 – 6.

Schneider, R.H.; Staggers, F.; Alexander, C.N.; Sheppard, W.; Rainforth, M.; Kondwani, K.; Smith, S. and King, C.G. (1995):

A randomised controlled trial of stress reduction for hypertension in older African Americans. *Hypertension*; 26(5): 820 – 7.

Sebstjen, M.; Zegura, B.; Guize, Z. and Keber I. (2001):

Fibrinolytic parameters and insulin resistance in young survivors of myocardial infarction with heterozygous families hypercholesterolemia. *Hypertension*; 15: 11B(3-4): 13 – 18.

Seidell, J.C. (2000):

Obesity, insulin resistance and diabetes: a worldwide epidemic. *Br. J. Nutr.*; 83(Suppl. 1): S5 – S8.

Seltzer, A.; Bregonzio, C.; Armando, I.; Baiardi, G. and Saavedra, J.M. (2004):

Oral administration of an AT₁ receptor antagonist prevents the central effects of angiotensin II in spontaneously hypertensive rats. *Brain Res.*; 1028: 9 – 18.

Selye, H. (1936):

A syndrome produced by diverse noxious agents. *Nature*; 138: 32.

Selye, H. (1950):

The physiology and pathology of exposure to stress. A treatise based on the concepts of the general adaptation syndrome and the disease of adaptation. Montreal: Acta Inc. *Wisconsin Med. J.*; 49: 6.

Selye, H. (1956):

The stress of life. New York. McGraw-Hill.

Sesso, H.D.; Lee, I.M.; Gazian, J.M.; Rexrode, K.M.; Glynn, R.J. and Buring, J.E. (2001):

Maternal and paternal history of myocardial infarction and cardiovascular risk of cardiovascular disease in men and women. *Circulation*; 104: 393 – 8.

Sheppard, M.R.; Vinay, K.; Abul, K.A. and Nelson, F. (2007):

Robbins Basic Pathology: With student consult online access.
Philadelphia: Saunders. Textbook, 8th edition, P. 345.

Silbernagl, I. (2000):

Color atlas of pathophysiology; P. 268.

Singer, H. (1977):

Diagnosis of primary hyperlipoproteinemia. Z. Gesamte Inn. Med.;
32(9): 129 – 33.

Singhal, A.; Agarwal, D.K and Srivastava, V. (1997):

Effect of immobilization stress on lipid profile. Indian J. of Physiol.
and Allied Sci.; 51(3): 138 – 43.

***Singhal, A.; Farooqi, S.; Cole, T.J.; O’Rahilly, S.; Fewtrell, M.;
Kattenhorn, M.; Lucas, A. and Deanfield, J. (2002):***

Influence of leptin on arterial dispensability: a novel link between
obesity and cardiovascular disease. Circulation; 106(15): 1919 – 24.

Siragy, H.M. and Carey, R.M. (2001):

Angiotensin type 2 receptors: potential importance in the regulation
of blood pressure. Curr. Opin. Nephrol. Hypertens.; 10: 99 –103.

Smith, D.F. and Toft, D.O. (1993):

Steroid receptors and their associated proteins. Mol. Endocrinol.;
7(1): 4 – 11.

***Smith, D.G.; Ben-Sholomo, Y.; Beswick, A.; Yarnell, J.; Lightman, S. and
Elwood, P. (2005):***

Cortisol, testosterone and coronary heart disease. Circulation; 112:
332 – 340.

Snijder, M.B.; van der Heijden, A.A.; van Dam, R.M.; Stehouwer, C.D.; Hiddink, G.J.; Nijpels, G.; Heine, R.J.; Bouter, L.M. and Dekker, J.M. (2007):

Is higher dairy consumption associated with lower body weight and fewer metabolic disturbances?. The hoorn study. *Am. J. Clin. Nutr.*; 85(4): 989 – 95.

Stanton, A. (2003):

Therapeutic potential of renin inhibitors in the management of cardiovascular disorders. *Am. J. Cardiovasc. Drugs*; 3: 389 – 94.

Step toe, A. and Brydon, L. (2005):

Associations between acute lipid stress responses and fasting lipid levels 3 years later. *Health Psychology*; 24(6): 601– 607.

Sterling, P. and Eyer, J. (1988):

Allostasis: a new paradigm to explain arousal pathology, *In Fisher, S. and Reason, J. (Eds.), Handbook of life stress, cognition and health.* Wiley, New York, P. 629 – 649.

Sun, P.; Dwyer, K.M.; Merz, N.B.; Sun, W.; Johnson, C.A.; Schircore, A.M. and Dwyer, J.H. (2000):

Blood pressure, LDL cholesterol and intima-media thickness. *Arterioscler. Thromb. Vasc. Biol.*; 20: 2005.

Third report of the National Cholesterol Education Program (NCEP) expert panel (2002):

On detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III), final report. *Circulation*; 106: 3143 – 3221.

Thogerson, A.M.; Jansson, J.H.; Boman, K.; Nilsson, T.K.; Weinehall, L.; Hutsaari, F. and Hallmans, G. (1998):

High plasminogen activator inhibitor and tissue plasminogen activator levels in plasma precedes a first acute myocardial infarction in both men and women: evidence for the fibrinolytic system as an independent primary risk factor. *Circulation*; 98: 2241.

Timmermans, P.B.; Wong, P.C.; Chiu, A.T; Herblin, W.F.; Benfield, P.; Carini, D.J.; Lee, R.J.; Wexler, R.R.; Saya, J.A. and Smith, R.D. (1993):

Angiotensin II receptors and angiotensin II receptor antagonists. *Pharmacol. Rev.*; 45(2): 205 – 251.

Torpy, J.M. (2007):

Chronic stress and the heart. *JAMA*; 298(14): 1722.

Tracey, K.J. (2002):

The inflammatory reflex. *Nature*; 420(6917): 853 – 859.

Tronvik, E.; Stovner, L.J.; Helde, G.; Sand, T. and Bovim, G. (2003):

Prophylactic treatment of migraine with an angiotensin II receptor blocker: a randomized controlled trial. *JAMA*; 1(289 Pt 1): 65 – 9.

Tschoepe, D.; Roesen, P. and Schwippert, B. (1993):

Platelets in diabetes: the role in the hemostatic regulation in atherosclerosis. *Semin. Thromb. Hemost.*; 19: 122 – 128.

Tsigos, C. and Chrousos, G.P. (2002):

Hypothalamic- pituitary- adrenal axis, neuroendocrine factors and stress. *J. Psychosom. Res.*; 53(4): 865 – 71.

Tuli, J.S.; Smith, J.A. and Morton, D.B. (1995):

Effects of acute and chronic restraint on the adrenal gland weight and serum corticosterone concentration of mice and their fecal output of oocytes after infection with *Eimeria apionodes*. *Res. Vet. Sci.*; 59: 82 – 86.

Tuzcu, E.M.; Kapadia, S.R. and Tutar, E. (2001):

High prevalence of coronary atherosclerosis in asymptomatic teenagers and young adults: evidence from intravascular ultrasound. *Circulation*; 103(22): 2705 – 10.

Uresin, A.Y.; Tonyali, H. and Karamursel, S. (2004):

The effects of losartan and immobilization stress on heart rate variability and plasma corticosterone levels in rats. *Int. J. Neurosci.* 114 (3): 365 – 79.

Uresin, Y.; Erbas, B. and Ozek, M. (2004):

Losartan may prevent the elevation of plasma glucose levels induced by chronic stress. *Pol. J. Pharmacol.*; 56: 271 – 3.

Uresin, Y.; Erbas, B.; Ozek, M.; Ozkok, E. and Gurol, A.O. (2004*):

Losartan may prevent the elevation of plasma glucose, corticosterone and catecholamine levels induced by chronic stress. *J. of RAAS*; 5(2): 93 – 6.

Vaccarino, V. and Bremner, J.D. (2006):

Stress response and the metabolic syndrome. *Cardiology*; Vol. 11, Part 2, P. 1 – 12.

Vanitallie, T. (2002):

Stress: a risk factor for serious illness. *Metabolism*; 51(6 Suppl. 1): 40 – 45.

Vasan, R.S.; Larson, M.G.; Kannd, W.B. and Levy, D. (2000):

Evolution of hypertension from non hypertensive blood pressure levels: rates of progression in Framingham Heart Study. *J. Am. Coll. Cardiol.*; 35: 292A.

Vague, J. (1947):

Factors determinant for obesity. *Presse Med*; 30: 339 – 40.

Vickers, M.; Ikensaio, B. and Breier, B. (2001):

IGF-1 treatment reduces hyperphagia, obesity and hypertension in metabolic disorders induced fetal programming. *Endocrinology*; 142(9): 3964 – 73.

Vinik, A.I.; Erbas, T.; Park, T.S.; Nolan, R. and Pittenger, G.L. (2001):

Platelet dysfunction in type 2 diabetes. *Diabetes Care*; 24: 1476 – 1485.

Visser, M.; Bouter, L.M.; McQuillan, G.M.; Wener, M.H. and Harris, T. (1999):

Elevated C-reactive protein levels in over weight and obese adult. *JAMA*; 282(22): 2131 – 35.

Vitale, C.; Murcuro, G.; Castiglioni, C.; Tulli, M.; Fini, M.; Volterrani, M. and Rosano, G.M.. (2005):

Metabolic effect of telmisartan and losartan in hypertensive patiants with metabolic syndrome. *Cardiovasc. Diabetol.*; 15: 4 – 6.

Vonhoute, P.M. and Boulanger, C.M. (1995):

Endothelium dependent response in hypertension. *Hypertens. Res.*; 18: 87 – 98.

Walsh, C.M.; Hilar, O. and Ezzo, D.C. (2009):

Metabolic syndrome and cardiovascular risk factors. *US Pharm.*; 34(2): HS12 – HS15.

Wang, H.L.; Fan, D.S.; Shen, Y.; Sun, A.P.; Zhang, J. and Yang, (2005):

The relationship between carotid artery atherosclerosis and hyperhomocysteinemia. *Zhonyhua Nei Ke Za Zhi.*; 44 (4): 258 – 61.

Watanabe, T.; Barker, T.A. and Berk, B.C. (2005):

Angiotensin II and the endothelium. Diverse signals and effects. *Hypertension*; 45: 163 – 9.

Watanabe, T.; Fujioka, T.; Hashimoto, M. and Nakamura, S. (1998):

Stress and brain angiotensin II receptors. *Critical Reviews in Neurobiology*; 12(4): 305 – 317.

Watkins, L.O. (2004):

Epidemiology and burden of cardiovascular disease. *Clin. Cardiol.*; 27(6 Suppl. 3): 1112 – 1116.

Weinstock, R.S.; Dai, H. and Wadden, T.A. (1998):

Diet and exercise in the treatment of obesity: effect of 3 interventions on insulin resistance. *Arch. Intern. Med.*; 185: 2477 – 83.

Weir, M.R. and Dzau, V.J. (1999):

The renin- angiotensin aldosterone system: a specific target for hypertension management. *Am. J. Hypertens.*; 12: 205S – 213S.

Weyer, C.; Funahashi, T.; Tanaka, S.; Hotta, K.; Matsuzawa, Y.; Pratley, R.E. and Tataranni, P.A. (2001):

Hypoadiponectinemia in obesity and type 2 diabetes: close association with insulin resistance and hyperinsulinemia. *J. Clin. Endocrinol. Metab.*; 86: 1930 – 35.

Wheatcroft, S.B.; Williams, I.L.; Shah, A.M. and Kearney, M.T. (2003):

Pathophysiological implications of insulin resistance on vascular endothelial function. *Diabet. Med.*; 20(4): 255 – 268.

White, M.; Racine, N.; Ducharme, A. and de Champlain, J. (2001):

Therapeutic potential of angiotensin II receptor antagonists. *Expert Opin. Investing. Drugs*; 10: 1687 – 1701.

Wold, L.E.; Muralikrishnan, D.; Albano, C.B.; Norby, F.L.; Ebadi, M. and Ren, J. (2003):

Insulin like growth factor 1 (IGF-1) supplementation prevents diabetes- induced alterations in coenzymes Q₉ and Q₁₀. *Acta diabetol.*; 40(2): 85 – 90.

Wood, D.; Backer, G.D. and Faergeman, O. (1998):

Prevention of coronary heart disease in clinical practice: recommendations of the second joint task force of European and other societies on coronary prevention. *Eur. Heart J.*; 19: 1434 – 1503.

Woodward, M.; Lowe, G.D.O.; Rumley, A. and Tunstall- peobe, E. (1998):

Fibrinogen as a risk factor for coronary heart disease and mortality in middle aged men and women: The Scottish Heart Health Study. *Eur. Heart J.*; 19: 55 – 62.

World Health Organization (1999):

Definition, diagnosis and classification of diabetes mellitus and its complications: report of a WHO Consultation. Part 1: diagnosis and classification of diabetes mellitus. Geneva, Switzerland.

Yang, G.; Wan, Y. and Zhu, Y. (1996):

Angiotensin II- an important stress hormone. Biol. Signals; 5: 1 – 8.

Yang, G.; Xi, Z.; Wan, Y.; Wang, H. and Bi, G. (1993):

Changes in circulating and tissue angiotensin II during acute and chronic stress. Biol. Signals; 2: 166 – 72.

Young, C.R. and Welsh, C.J. (2005):

Stress, health and disease. Introductory Review. Cell Sci. Rev.; 2(2): 1742.

Zardooz, H.; Zahedi, A.S. and Naseri, M. G. (2006):

Effect of chronic psychological stress on insulin release from rat isolated pancreatic islets. Life Sci.; 79(1): 57 – 62.

Zavaroni, I.; Mazza, S.; Dall'Aglio, E.; Gasparini, P.; Passeri, M. and Reaven, G.M. (1992):

Prevalence of hyperinsulinemia in patients with high blood pressure. J. Intern. Med.; 231: 1235 – 40.