

References

- Ahorlu C.K., Dunyo S.K., Asamoah G., Simonsen P.E; (2001): Consequences of hydrocele and the benefits of hydrocelectomy: A qualitative study in lymphatic filariasis endemic communities on the coast of Ghana, *Acta Tropica* 80(3):215-21, 2001.
- Ali M., Goovaerts P., Nazia N., Haq M. Z., Yunus M. and Emch M; (2006): Application of Poisson kriging to the mapping of cholera and dysentery incidence in an endemic area of Bangladesh, *International Journal of Health Geographics* 2006, 5:45
- Anderson, R.M; (1990): Populations and infectious diseases: ecology or epidemiology? ,*Journal of Animal Ecology* 60: 1-50
- Anderson, R.M. and May, R.M; (1991): *Infectious Diseases of Humans: Dynamics and Control*. Oxford: Oxford University Press, pp.757.
- Anselin, L; (2005): Exploring Spatial Data with GeoDATM, A Workbook, Spatial Analysis Laboratory, p. 138
- Asadi-Lari M, Packham C, Gray D;(2003): Need for redefining needs, *Health Qual Life Outcomes* 2003, 1:34.
- Babu B.V, Nayak A.N, Acharya A.S, Jangid P.K, and Mallick G; (2002): The economic loss due to treatment costs and work loss to individuals with chronic lymphatic filariasis in rural communities of Orissa, India, *Acta Tropica*, 82(1):31-38.
- Babu S. , Porte P., Klei T. R., Shultz L. D. and Rajan T. V;(1998): Host NK Cells Are Required for the Growth of the Human Filarial Parasite *Brugia malayi* in Mice, *The Journal of Immunology*, 1998, 161: 1428-1432.

- Bailey T;(1994): A review of statistical spatial analysis in geographical information systems. In: Fotheringham S, Rogerson P,: Spatial analysis and GIS, London: Taylor and Francis. 1994.
- Balarajan, K. and Bundy, D. A; (2001): Placebo-controlled community trial of four cycles of single-dose diethylcarbamazine or ivermectin against *Wuchereria bancrofti* infection and transmission in India, *Trans R Soc Trop Med Hyg* 95: 336, 2001.
- Barnes S. and Peck A;(1994): Mapping the future of health care: GIS applications in Health care analysis, *Geographic Information systems*,1994;4:31-3.
- Boulos, M.K. and Roudsari, A; (2000): WHO HealthMap,
<http://www.soi.city.ac.uk/~dk708/healthmap.htm>
- Brooker, S. and Michael, E; (2000): The potential of geographical information systems and remote sensing in the epidemiology and control of human Helminth infections, In Hay, S. I., Randolph, S. E. and Rogers, D. J. (eds.), *Remote sensing and geographical information systems in epidemiology*, *Advances in Parasitology*, Vol. 47. New York, NY: Academic Press.
- Brooker, S., Hay, S.I., Tchuem Tchuente, L.A. and Ratard, R; (2002): Using NOAA-AVHRR data to model human helminth distributions in planning disease control in Cameroon, West Africa, *Photogrammetric Engineering & Remote Sensing* 68(2):175-179.
- Brooker S., Rowlands M., Haller L., Savioli L., and Bundy D.A;(2000): Towards an atlas of human helminth infection in sub-Saharan Africa: The use of geographical information systems (GIS), *Parasitology Today* (Personal Ed.) 2000 Jul, 16(7):303-307.
- Burkot T. and Bockarie M; (2004): Vectors, *Am J Trop Med Hyg* 71 (Suppl.): 24–26.

- Carrat F. and Valleron A.J; (1992): Epidemiologic Mapping using the "Kriging" Method: Application to an Influenza-like Illness Epidemic in France, American Journal of Epidemiology Vol. 135, No 11, Copyright 1992 by The Johns Hopkins University School of Hygiene and Public Health Printed in U.S.A

- CDC; (1993): Recommendations of the International Task Force for Disease Eradication, MMWR Recomm Rep. (1993) , 42(RR-16):1-38

- CDC, (2006): Database and statistics software for public health professionals, Components of Epi Info for Windows, <http://www.cdc.gov/epiinfo/>

- CDC/DPD; (2006): Lymphatic Filariasis, http://www.cdc.gov/ncidod/dpd/parasites/lymphaticfilariasis/factsht_lymphatic_filar.htm

- Chung-Yuan, H .; Chuen-Tsai S.; Ji-Lung H. and Holin L;(2004) : Simulating SARS: Small-World Epidemiological Modeling and Public Health Policy Assessments , *Journal of Artificial Societies and Social Simulation* vol. 7, no. 4 <http://jasss.soc.surrey.ac.uk/7/4/2.html>

- Clarke K.C; (1995): Analytical and computer cartography, 2nd ed. Englewood Cliffs, NJ: Prentice-Hall. 1995.

- Clarke K. C., McLafferty S. L., and Tempalsk B. J;(1996): On Epidemiology and Geographic Information Systems: A Review and Discussion of Future Directions, *Emerging Infectious Diseases*, Vol.2, NO.2, April-June 1996.

- Cliff A, Haggett P, and Smallman-Raynor, M;(1993): Measles: an historical geography of major human viral disease from global expansion to local retreat, Oxford, UK: Blackwell Reference, 1993.

- Colorado University;(2006):
http://www.colorado.edu/geography/gcraft/notes/gps/gps_f.html
- Connor, S., Thomson, M., Clarke GP, Craig M, et al; (1999): Forecasting and prevention of epidemic malaria: New perspectives on an old problem, *Parassitologia* 41:439-448.
- Cressie N;(1992): Smoothing regional maps using empirical Bayes predictors, *Geographical Analysis* 1992;24:75-95.
- Crossen E. and Sandoval N;(2004):Filariasis, Stanford University,
<http://www.stanford.edu/class/humbio103/ParaSites2004/Filariasis/>
- Cuenco K.T, Halloran M.E, Charles J.L, and Lammie P.J; (2004): A family study of lymphedema of the leg in a lymphatic filariasis–endemic area, *Am. J. Trop. Med. Hyg.*, 70(2), 2004, pp. 185–190.
- Curran, P. J., Atkison, P. M., Foody, G. M., and Milton, E. J; (2000): Linking remote sensing, land cover and disease, In Hay, S. I., Randolph, S. E. and Rogers, D. J. (eds.). *Remote sensing and geographical information systems in epidemiology*, *Advances in Parasitology*, Vol. 47. New York, NY: Academic Press.
- CSR/ PHE/ RBM; (2006): Using climate to predict disease outbreaks: a review,
<http://www.who.int/globalchange/publications/oeh0401/en/index.html>
- Dadzie K. Y., Basáñez M., AND Richards F; (2004): Epidemiology, parasite biology and modeling, *Am. J. Trop. Med. Hyg.*, 71(5 suppl), 2004, pp. 22-23.
- Das P.K and Subramanian S; (2002): Modelling the epidemiology, transmission and control of lymphatic filariasis, *Annals of Tropical Medicine & Parasitology*, Vol. 96, Supplement No. 2, S153–S164 ,2002.

- Das P.K and Pani SP;(2000): Towards elimination of lymphatic filariasis in India, problems, challenges, opportunities and new initiatives, *J Int Med Sci Acad* 2000, 13:18-26.

- Das, P. K., Subramanian, S., Manoharan, A., Ramaiah, K.D., Vanamail, P., Grenfell, B. T., Bundy, D. A. P. and Michael, E; (1995): Frequency distribution of *Wuchereria bancrofti* infection in the vector host in relation to human host: Evidence for density dependence, *Acta Tropica* 60: 159, 1995.

- Dean, A.G;(1999): EpiInfo and EpiMap: Current Status and Plans for EpiInfo 2000, *Journal of Public Health Management Practice*, 5(4), 54-57, 1999.

- Ding Y. and Fotheringham A.S;(1992): The integration of spatial analysis and GIS, *Computers in Environmental and Urban Systems* 1992:3-19.

- Dister, S.W., Fish, D., Bros, S.M., Frank, D.H., and Wood, B.L; (1997): Landscape characterization of peridomestic risk for Lyme disease using satellite imagery, *American Journal of Tropical Medicine & Hygiene* 57(6):687-692.

- DPDx; (2006): Filariasis
<http://www.dpd.cdc.gov/dpdx/HTML/Filariasis.htm>

- Dreyer G, Noroes J, Figueredo-Silva J, and Piessens WF;(2000): Pathogenesis of Lymphatic Disease in Bancroftian Filariasis: A Clinical Perspective, *Parasitology Today* 2000, 16:544-548.

- Durrheim D.N, Wynd S, Liese B, and Gyapong J.O, (2004): Lymphatic filariasis endemicity—an indicator of poverty? *Trop Med Int Health* 9: 843–845.

- El-Setouhy M, Elaziz K.M, Helmy H, Farid H.A, Kamal H.A, Ramzy R.M, Shannon W.D, and Weil G.J;(2007): The Effect of Compliance on the Impact

of Mass Drug Administration for Elimination of Lymphatic Filariasis in Egypt, *Am J Trop Med Hyg*, 2007; 77: 1069-1073.

- EMRO;(2000): Elimination of Lymphatic Filariasis in the EMR, The Forty-seventh Session of the Regional Committee for the Eastern Mediterranean ,Cairo, Egypt, 2000.

- Escobar F.and Hunter G; (2001): Introduction to GIS, Department of Geomatics, The University of Melbourne, USA,
<http://www.sli.unimelb.edu.au/gisweb>

- Eyles J. and Furgal C;(2002): Indicators in environmental health: identifying and selecting common sets, *Can J Public Health* 2002;93:S62–7.

- Ezzati M., Lopez A.D, and Rodgers A;(2002):Selected major risk factors and global and regional burden of disease, *Lancet* 2002;360:1347–60.

- Ezzati M , Saleh H and Kammen DM;(2000):. The contributions of emissions and spatial microenvironments to exposure to indoor air pollution from biomass combustion in Kenya, *Environ Health Perspect* 2000;108:833–9.

- Ezzati M., Utzinger J., Cairncross S., Cohen A.J. and Singer B.H;(2005): Environmental risks in the developing world: exposure indicators for evaluating interventions, programmes, and policies, *Journal of Epidemiology and Community Health* 2005;59:15-22

- Farid H.A., Kamal S.A., Weil G.J., Adham F.K. and Ramzy R.M.R; (2003): Filariasis elimination in Egypt: impact of low microfilaraemics as sources of infection for mosquitoes, *Eastern Mediterranean Health Journal*, Vol. 9, No. 4, 2003

- Farid H, Morsy Z, Gad AM, Ramzy RMR, Faris R, andWeil G.J; (1997): Filariasis transmission potential of mosquitoes to humans of different age groups, *J Egypt Soc Parasitol* 27: 355– 364.

- Faris R; (1996): Parasite antigenemia without microfilaremia in bancroftian filariasis, *Am J Trop Med Hyg* 55: 333–337.

- Faris R, Hussain O, El Setouhy M, Ramzy RM, Weil G.J;(1998): Bancroftian filariasis in Egypt: visualization of adult worms and subclinical lymphatic pathology by scrotal ultrasound, *Am J Trop Med Hyg*, 1998, Dec;59(6):864-867.

- Faris R, Ramzy RMR, Gad AM, Weil GJ, Buck A.A; (1993): Community diagnosis of bancroftian filariasis, *Trans R Soc Trop Med Hyg* 87: 659–661.

- Fontes G, Rocha E.M, Brito A.C, and Antunes C.M; (1998): Lymphatic filariasis in Brazilian urban area (Maceio, Alagoas), *Mem Inst Oswaldo Cruz* 93: 705-710.

- Gasarasi, D. B., Premji, Z. G., Mujinja, P. G. M. and Mpembeni, R; (2000): Acute adenolymphangitis due to Bancroftian filariasis in Rufiji district, south east Tanzania, *Acta Tropica*, 75,19-28.

- Gatrell A. and Bailey T;(1995): Can GIS be made to sing and dance to an epidemiological tune? Presented at the International Symposium on Computer Mapping and Environmental Health, Tampa, FL, February 1995.

- Gettinby G, Revie C.W. and Forsyth A.J;(2006): Modelling: A review of systems and approaches for vector-transmitted and other parasitic diseases in developing countries, <http://www.fao.org/wairdocs/ilri/x5441e/x5441e0a.htm>

- Glass GE, Schwartz BS, Morgan JM III, Johnson DT, Noy PM, and Israel E;(1995): Environmental risk factors for Lyme disease identified with geographic information systems. *Am J Public Health* 1995;85:944-8.

- Goodchild, M.F., Parks, B.O. and Steyaert, L.T;(1993): Environmental Modeling with GIS. pp. 488, Oxford University Press, New York.

- Gotway, C. A. and Wolfinger, R. D; (2003): Spatial prediction of counts and rates, *Statistics in Medicine*, 22: 1415-1432.

- Gould P;(1993): *The slow plague: a geography of the AIDS epidemic*, Cambridge, UK: Blackwell, 1993.

- Grady C.A, Beau de Rochars M, Direny A.N, Orelus J.N, Wendt J, Radday J, Mathieu E, et al; (2007): Endpoints for Lymphatic Filariasis Programs, *Emerging Infectious Diseases* , www.cdc.gov/eid , Vol. 13, No. 4, April, 2007

- Guthmann J.P, Llanos-Cuentas A., Palacios A. and Hall A.J; (2002): Environmental factors as determinants of malaria risk, A descriptive study on the northern coast of Peru, *Tropical Medicine and International Health*, 7:518–525.

- Gyapong, J.O, Gyapong M, Evans D.B, Aikins M.K, and Adjei S; (1996): The economic burden of lymphatic filariasis in northern Ghana, *Ann Trop Med Parasitol* 90(1):39-48.

- Gyapong J. O. and Remme J. H. F; (2001): The use of grid sampling methodology for rapid assessment of the distribution of bancroftian filariasis, *Transactions of the ROYAL SOCIETY OF TROPICAL MEDICINE AND HYGIENE* (2001) 95,681-686.

- Gyapong J. O. Kumaraswami V., Biswas G. and Ottesen E.A; (2005): Treatment strategies underpinning the global programme to eliminate lymphatic filariasis, *Expert Opin. Pharmacother.* 2005, 6 (2).

- Haddix A.C, Carter S, Michael E, and Benton B; (1999): Cost effectiveness of integrating lymphatic filariasis elimination into APOC onchocerciasis control programs, Report prepared for the Joint Action Forum/Joint Program Committee, December 1999, Den Haag.

- Haddix A.C. and Kestler A; (2000): Lymphatic filariasis: Economic aspects of the disease and programmes for its elimination, *Trans R Soc Trop Med Hyg* 94(6):592-3.
- Hales S. and Hearnden M; (1999): Modelling the potential distribution of disease vectors in New Zealand, Presented at the SIRC 99 – The 11th Annual Colloquium of the Spatial Information Research Centre. University of Otago, Dunedin, New Zealand 13 - 15th December, 1999.
- Harb M, Faris R, Gad A.M, Hafez O.N, Ramzy RMR, and Buck A.A; (1993): The resurgence of lymphatic filariasis in the Nile delta, *Bull World Health Organ* 71: 49–54.
- Hay S. I, and Lennon J.J; (1999): Deriving meteorological variables across Africa for the study and control of vector-borne disease: a comparison of remote sensing and spatial interpolation of climate, *Tropical Medicine and International Health*, 4(1): 58-71.
- Hay, S.I., Randolph, S. E. and Rogers, D. J; (2000): Remote sensing and geographical information systems in epidemiology, *Advances in Parasitology* 47:357. London, UK: Academic Press.
- Helmy H., Fischer P., Farid H. A., Bradley M. H. and Ramzy R. M. R; (2004): Test strip detection of *Wuchereria bancrofti* amplified DNA in wild-caught *Culex pipiens* and estimation of infection rate by a PoolScreen algorithm, *Tropical Medicine and International Health*, volume 9 no 1 pp 158–163 January 2004
- Hengl. T; (2007): A Practical Guide to Geostatistical Mapping of Environmental Variables, Luxembourg: Office for Official Publications of the European Communities, European Communities, 2007
- Henriques W. D; (1998): Introduction to Geographic Information Systems for Environmental Health Protection,
www.atsdr.cdc.gov/GIS/conference98/henriques1/sld001.htm

- Hoskins R. E; (1998): Monte Carlo and Spatial Bayesian Methods applied to Environmental Equity Assessment in Western Washington,
www.atsdr.cdc.gov/GIS/conference98/hoskins-35 mm/ sld001.htm

- IPCC; (2001): Climate change 2001: impacts, adaptation and vulnerability. Contribution of working group II to the third assessment report of the Intergovernmental Panel on Climate Change (IPCC), McCarthy JJ, et al., eds. Cambridge: Cambridge University Press.

- Iqbal J. and Sher A. (2006): Determination of the prevalence of lymphatic filariasis among migrant workers in Kuwait by detecting circulating filarial antigen, *J Med Microbiol* 55 (2006), 401-405

- Janssen B.J.M, and Pruppers, M.J.M;(1998): Integrated Risk Assessment and Management: Overview and State-of-the-Art, *Journal of Hazardous Materials*, 61 (1998) 5-22.

- Jeevan B.S, Valérie O, Garib D. T. and Marcel H; (2003): Mapping of lymphatic filariasis in Nepal, *Filaria Journal* 2003, 2:7.

- Johnson, W.O., Gastwirth, J.L., and Pearson, L.M. (2001): Screening without a gold standard: The Hui-Walter paradigm revisited. *American Journal of Epidemiology* 153, 921-924.18.

- Kahlmeier S. and Braun-Fahrlander C;(2004): Environmental health indicators in policy evaluation. *Eur J Public Health* 2004;14:101–4.

- Kalyobiya: <http://www.kalyobiya.gov.eg/index1>

- Kellyn, S.B;(1997): Mapping the environment, Innovations; environmental health perspectives, volume 105, number 6, June 1997.

- Kirby R.S; (1996): Toward congruence between theory and practice in small area analysis and local public health data. *Statistics in Medicine* 15(17–18):1859–66.
- Kleinschmidt I, Bagayoko M, Clarke GP, Craig M, and Le Sueur D. A;(2000): spatial statistical approach to malaria mapping. *Int J Epidemiol.* 2000; 29:355–61.
- Kreiser P.B. and Nutman T.B; (2002): Update on lymphatic filariasis, *Current Opinion in Infectious Diseases* 4, 65–69.
- Kristensen, T. K, Malone, J. B, and McCarroll, J. C; (2001): Use of satellite remote sensing and geographic information systems to model the distribution and abundance of snail intermediate hosts in Africa: A preliminary model for *Biomphalaria pfeifferi* in Ethiopia. *Acta Tropica* 2001 Apr 27 79(1):73-78.
- Kuhn K., Campbell-Lendrum D, Haines A.,and Cox J; (2005): Using climate to predict infectious disease epidemics, WHO,2005, ISBN 92 4 159386 5 (NLM classification: WB 700)
- Kulldorff M and Nagarwalla N; (1995): Spatial disease clusters: detection and inference, *Stat Med.* 1995;**14**:799–810.
- Kumar D.V.R, Sriram k, Rao K.M, and Murty U.S; (2005): Management of filariasis using prediction rules derived from data mining, *Bioinformation.* 2005; 1(1): 8–11.
- Levy JI, Houseman EA, Ryan L, et al;(2000): Particle concentrations in urban microenvironments. *Environ Health Perspect* 2000;108:1051–1057.
- Li J., Zhou Y. C., Ma Z. and Hyman J. M;(2004): "Epidemiological models for mutating pathogens", *SIAM Journal on Applied Mathematics*, vol. 65, no. 1, pp. 1--23, 2004.

- Liang, S.Y., Linthicum, K.J. and Gaydos, J. C; (2002): Climate change and the monitoring of vector-borne disease, *Medical Student JAMA* 287:2286.

- Lindsay, S. W. and Martens, W. J. M; (1998): Malaria in the African highlands: Past, present and future, *Bulletin of the World Health Organization* 76(1):33-45.

- Lindsay S.W, and Thomas C.J; (2000): Mapping and estimating the population at risk from lymphatic filariasis in Africa, *Trans R Soc Trop Med Hyg* 94: 37–45.

- Loslier L; (1994): Geographical information systems (GIS) from a health perspective, In: De Savigny D, Wijeyaratne P. , editor, *GIS for health and the environment*. Ottawa: IDRC; 1994. pp. 13–20.

- Maizels, R.M. and Lawrence, I.A; (1991) Immunological tolerance: the key feature in human filariasis. *Parasitol. Today* 7, 271–276.

- Martin C, Curtis B, Fraser C, Sharp B;(2002): The use of a GIS-based malaria information system for malaria research and control in South Africa. *Health Place*. 2002;8:227–36.

- Martínez-Piedra R., Loyola-Elizondo E., Vidaurre-Arenas M. and Aguilar P. N;(2004): Software Programs for Mapping and Spatial Analysis in Epidemiology and Public Health, *Epidemiological Bulletin / PAHO*, Vol. 25, No. 4 ,2004.

- Medley, G.F., Guyatt, H.L. and Bundy, D.A.P; (1993): A quantitative framework for evaluating the effect of community treatment on the morbidity due to ascariasis. *Parasitology* 106: 201-221.

- Mellor P.S. and Leake C.J; (2000): Climatic and geographic influences on arboviral infections and vectors. *Revue Scientifique et Technique de l'Office International des Epizooties*, 19:41–54.

- Meyers M.F, Rogers D.J, Cox J., Flanhault A, and Hay S.I; (2000): Forecasting disease risk for increased epidemic preparedness in public health. In Hay S. I., Randolph S. E. and Rogers D. J. (eds.). Remote sensing and geographical information systems in epidemiology, *Advances in Parasitology*, Vol. 47. New York, NY: Academic Press.
- Michael E, Bundy D.A, and Grenfell B.T; (1996): Re-assessing the global prevalence and distribution of lymphatic filariasis, *Parasitology*, 1996, Apr;112 (Pt 4):409-28.
- Moffett A., Shackelford N., and Sarkar S; (2007): Malaria in Africa: Vector Species' Niche Models and Relative Risk Maps, *PLoS ONE*. 2007; 2(9): e824
- Molyneux D.H; (2003): Common themes in changing vector-borne disease scenarios, *Trans R Soc Trop Med Hyg*, 97: 129–132.
- Molyneux D, (2003): Lymphatic filariasis (elephantiasis) elimination: a public health success and development opportunity. *Filaria J* 2: 13.
- Muhondwa E.P.Y, (1983): Community participation in filariasis control: The Tanzania experiment, TDR/SER/SWG(4)/WP/83.13.
- Murray S.A and Graham L.J.C; (1996): Practice based health needs assessment: Use of four methods in a small neighborhood. *British Medical Journal*, 310:1443–8.
- NASA;(2006): Landscape Epidemiology and RS/GIS, <http://geo.arc.nasa.gov/sge/health/landepi.html>
- Nissen M. D. and Walker J. C; (2006): **Filariasis**
<http://www.emedicine.com/ped/topic782.htm>
- Odoi A, Martin S.W, Michel P, Middleton D, Holt J, and Wilson J; (2004): Investigation of clusters of giardiasis using GIS and a spatial scan statistic. *Int J of Health Geog*. 2004;**3**:11–21.

- Olszewski, W. L., Jamal, S., Dworzynski, A., Swoboda, E., Pani, S. P., et al; (1994): Bacteriological studies of skin, tissue fluid and lymph in filarial lymphoedema, *Lymphology*, 27 (Suppl.), 345± 348.
- O'Neill M.S, Jerrett M, Kawachi I, *et al*; (2003): Health, wealth and air pollution: advancing theory and methods, *Environ Health Perspect* ,2003;111:1861–70.
- Ord K. and Getis A;(1995): Local spatial autocorrelation statistics: distributional issues and an application, *Geographical Analysis*, 1995;24:286-306.
- Ottesen E.A; (2000): The global programme to eliminate lymphatic filariasis, *Trop. Med.Int. Health* (2000) 5(9):591-594.
- Palmer T. and Anderson D; (1994): The prospects for seasonal forecasting – a review paper. *Quarterly Journal of the Royal Meteorological Society*, 120: 755-796.
- Pani, S. P., Balakrishnan, N., Srividya, A., Bundy, D. A. P. and Grenfell, B. T; (1991): Clinical epidemiology of Bancroftian filariasis: effect of age and gender, *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 85, 260-264.
- Perry B. and Gesler W;(2000): Physical access to primary health care in Andean Bolivia, *Soc Sci Med*. 2000;50:1177–88.
- Plaisier A.P, Stolk W.A, van Oortmarssen G.J, Habbema J.D.F; (2000): Effectiveness of annual ivermectin treatment for *Wuchereria bancrofti* infection, *Parasitol Today* 16: 298–302
- Plaisier, A. P., Subramanian, S., Das, P. K., Souza, W., Lapa, T., Furtado, A. F., Van der Ploeg C. P., et al; (1998): The LYMFASIM simulation programme for modeling lymphatic filariasis and its control, *Methods Inf Med* 37: 97, 1998.

- Pondicherry,(2006): Mathematical Models,

<http://pondicherry.nic.in/fil-free/vcrc/tools.html>

- Purdy M., (2006): Global program to eliminate elephantiasis has early success in Egypt, Washington University School of Medicine Public release date: 23-Mar-200 No.6314-286-0122

http://www.eurekalert.org/pub_releases/2006-03/wuso-gpt032106.php

- Public Health Agency of Canada; (2006): GIS for public health practice, http://www.phac-aspc.gc.ca/php-ppsp/gis_e.html

- Purdy M, (2006): Global program to eliminate elephantiasis has early success in Egypt, Record, March 31, 2006 ,Vol. 30, No. 27,Washington University in St. Louis

- Pruss A , Corvalan C, Pastides H, *et al*; (2001):Methodologic considerations in estimating burden of disease from environmental risk factors at national and global levels. *Int J Occup Environ Health* 2001;7:58–67.

- Rahmah N, Lim BH, Anuar AK et al; (2001): A recombinant antigen-based IgG4 ELISA for the specific and sensitive detection of *Brugia malayi* infection. *Transactions of The Royal Society of Tropical Medicine and Hygiene* 95, 280–284.

- Rahmah N, Shenoy R. K , Nutman T. B , Weiss N, Gilmour K , Maizels R. M , Yazdanbakhsh M. and Sartono E; (2003): Multicentre laboratory evaluation of *Brugia* Rapid dipstick test for detection of brugian filariasis, *Tropical Medicine and International Health* , volume 8 no 10 pp 895–900 October 2003.

- Rahme E., Joseph, L. and Gyorkos, T; (2000): Bayesian sample size determination for estimating binomial parameters from data subject to misclassification, *Applied Statistics* 49, 119-228.

- Ramaiah K.D, Das P.K, Michael E, and Guyatt H; (2000): The economic burden of lymphatic filariasis in India, *Parasitol Today* 16(6):251-3.

- Ramaiah K.D, Guyatt H, Ramu K, Vanamail P, Pani S.P, and Das P.K;(1999): Treatment costs and loss of work time to individuals with chronic lymphatic filariasis in rural communities in south India, *Trop Med Int Health*, 1999 Jan;4(1):19-25

- Ramaiah K.D, Vijay Kumar K.N, Ramu K, Pani S.P, and Das P.K;(1997): Functional impairment caused by lymphatic filariasis in rural areas of south India, *Trop Med Int Health* 1997, 2:832-838.

- Ramzy R.M.R , El Setouhy M , Helmy H , Ahmed E.S , Abd Elaziz K.M , Farid H.A , Shannon W.D , and Weil G.J ; (2006): Effect of yearly mass drug administration with diethylcarbamazine and albendazole on bancroftian filariasis in Egypt: a comprehensive assessment, *The Lancet* 2006; 367:992-999

- Ramzy R.M.R, Hafez O.N, Gad A.M, Faris R, Harb M, Buck A.A, and Weil G.J, (1994): Efficient assessment of filariasis endemicity by screening for filarial antigenemia in a sentinel population, *Trans R Soc Trop Med Hyg* 88: 41–44.

- Ramzy R.M.R, Helmy H., el-Lethy A. S., Kandil A. M., Ahmed E. S., Weil G. J. and Faris R; (2000): Field evaluation of a rapid-format kit for the diagnosis of bancroftian filariasis in Egypt, *East Mediterranean Health Journal*, 5, 880-887.

- Robertson G.P; (2008): *Geostatistics for the Environmental Sciences*, Copyright 1989-2008 Gamma Design Software, LLC. Plainwell, Michigan USA ISBN number 0-9707410-0-6

- Rogers D.J, Randolph S.E, Snow R.W. and Hay S.I; (2002): Satellite imagery in the study and forecast of malaria, *Nature*, 7 Feb 2002, 415:710-715.

- Samuel P.P, Arunachalam N, Hiriyan J, Thenmozhi V, Gajanana A, and Satyanarayana K; (2004): Host-feeding pattern of *Culex quinquefasciatus* Say and *Mansonia annulifera*, the major vectors of filariasis in a rural area of south India, *J Med Entomol*, 41: 442-446.
- SaTScan,(2006): Spatial and Temporal Scan Tests Program <http://www.satscan.org>
- Schirnding V.Y;(1997): Health in sustainable development: can we rise to the challenge, *Lancet* 2002;360:632–7.
- Schirnding V.Y;(2002): *Health in sustainable development planning: the role of indicators*. Geneva: World Health Organisation, 2002.
- Scholten H.J. and de Lepper M.J;(1991): The benefits of the application of geographical information systems in public and environmental health, *World Health Stat Q*. 1991;44:160–70.
- Schweitzer, A.N. and Anderson, R.M; (1992): Dynamic interactions between CD4+ T cell subsets and parasitic helminths: mathematical models of heterogeneity in outcome. *Parasitology* 105: 513-522.
- Seto E, Liang S, Gong P, Wu W, Davis G, Qui D, Gu X, and Spear R;(2002): The use of remote sensing for predictive modeling of shistosomiasis in China, *Photogrammetric Engineering & Remote Sensing*, Vol. 68, No. 2, pp. 175-179.
- Shanks G.D, Biomndo K, Hay S.I. and Snow R.W; (2000): Changing patterns of clinical malaria since 1965 among a tea estate population located in the Kenyan highlands, *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 94:253–255.
- Shenoy R. K., Kumaraswami V, Suma K, Rajan K. and Radhakuttyamma G;(1999): A double-blind, placebo-controlled study of the efficacy of oral penicillin, diethylcarbamazine or local treatment of the affected limb in preventing acute adenolymphangitis in lymphoedema caused by brugian

filariasis, *Annals of Tropical Medicine & Parasitology*, Vol. 93, No. 4, 367- 377 (1999).

- Simonsen P.E, Meyrowitsch DW, Mukoko D.A, Pedersen E.M, Malecela-Lazaro M.N, Rwegoshora R.T; (2004): The effect of repeated half-yearly diethylcarbamazine mass treatment on *wuchereria bancrofti* infection and transmission in two east African communities with different levels of endemicity, *Am J Trop Med Hyg* 2004;70:63-71.

- Soumyadip, G; (2004): Dependence in Stochastic Simulation Models, Cornell University 2004, www.orie.cornell.edu/~sdghosh/thesis.pdf

- Spinello E.F; (2006): A Public Health Information System for Conducting Community Health Needs Assessment, California State University, Northridge
- Dept. of Health Sciences, Northridge, CA ,
<http://www.rpmconsulting.com/phs.html>

- Sridhar L; (2002): Fighting Filariasis <http://www.boloji.com/wfs/wfs014.htm>

- Srividya A, Pani, S. P, Rajagopalan P. K, Bundy D. A. and Grenfell B. T; (1991): The dynamics of infection and disease in bancroftian filariasis, *Trans R Soc Trop Med Hyg* 85: 255, 1991.

- Stevens A and Gillam S; (1998): Health needs assessment - Needs assessment: from theory to practice, *British Medical Journal* 1998, 316:1448-1452.

- Stolk W.A, de Vlas S.J, and Habbema J.D.F; (2006): Advances and challenges in predicting the impact of lymphatic filariasis elimination programmes by mathematical modeling, *Filaria J*. 2006; 5: 5.

- Subramanian, S., Manoharan, A., Ramaiah, K. D. and Das P. K; (1994): Rates of acquisition and loss of *Wuchereria bancrofti* infection in *Culex quinquefasciatus*, *Am J Trop Med Hyg* 51: 244, 1994.

- Subramanian S, Krishnamoorthy K, Ramaiah K. D, Habbema J. D. F, Das P. K and Plaisier A. P; (1998): The relationship between microfilarial load in the human host and uptake and development of *Wuchereria bancrofti* microfilariae by *Culex quinquefasciatus*. A study under natural conditions. *Parasitology* 116: 243, 1998.

- Subramanian S., Vanamail P., and Das P.K ;(2002): Mathematical Models for optimizing and predicting outcomes of intervention measures for the control of Lymphatic Filariasis, ICMR Bull 32 (2002) 1.

- Subramanian S, Stolk W.A, Ramaiah K.D, Plaisier A.P, Krishnamoorthy K, Van Oortmarssen G.J, Dominic Amalraj D, Habbema J.D, Das P.K ; (2004): The dynamics of *Wuchereria bancrofti* infection: a model-based analysis of longitudinal data from Pondicherry, India, *Parasitology*. 2004; 128(Pt 5):467-82

- Subramanian S, Pani P.S, Ravi R, Krishnamoorthy K, and Das P.K ; (2008): Mathematical models for lymphatic filariasis transmission and control: Challenges and prospects, *Parasit Vectors*, 2008; 1: 2.

- Sujatha V. and Nagendra Rao C.R; (2003): ENVIRONMENTAL CARE IN THE CONTROL OF FILARIASIS – A CASE STUDY, in Martin J. Bunch, V. Madha Suresh and T. Vasantha Kumaran, eds., *Proceedings of the Third International Conference on Environment and Health, Chennai, India, 15-17 December, 2003*. Chennai: Department of Geography, University of Madras and Faculty of Environmental Studies, York University, Pages 529 – 536.

- Tanser F.C. and le Sueur D; (2002): The application of geographical information systems to important public health problems in Africa, *Int J Health Geogr*. 2002 ; 1: 4.

- Taylor M.J; (2003): *Wolbachia* in the inflammatory pathogenesis of human filariasis ,*Ann N Y Acad Sci*. 2003 Jun;990:444-9.

- TDR/Gen; (1996): Investing in health research and development. Report of the AdHoc Committee on health research relating to future intervention options, 1996. Geneva, World Health Organization, TDR/Gen/96.1.

- Tempalski B.J;(1994): The case of Guinea worm: GIS as a tool for the analysis of disease control policy. *Geographic Information Systems* 1994;4:32-8.

- Texas State Data Center and Office of the State Demographer;(2006): Thematic Maps, Institute for Demographic and Socioeconomic Research (IDSER), College of Public Policy, The University of Texas at San Antonio, <http://txsdc.utsa.edu/maps/thematic>

- The Ohio State University ;(2006): Parasites and parasitological resources, <http://www.biosci.ohio-state.edu>

- The University of Iowa, Iowa City, USA;(2005):Improving Public Health Through Geographical Information Systems, An Instructional Guide to Major Concepts and Their Implementation, <http://www.uiowa.edu/~geog/health/GIS2.html>

- Thompson D. F, Malone J. B, Harb M, Faris R, Huh O. K, Buck A.A, and Cline B. L; (1996): Bancroftian Filariasis Distribution and Diurnal Temperature Differences in the Southern Nile Delta, *Emerging Infectious Diseases, Vol. 2, No. 3—July-September 1996*

- Thomson M. C. and Connor S. J; (2001): The development of malaria early warning systems for Africa. *Trends in Parasitology* 17(9):438-445.

- Thrall S.E;(1999): Geographic Information System (GIS). Hardware and Software. *J Public Health Management Practice*, 5(2), 82-90, 1999.

- Tomlin W.R;(1990): Geographic information systems and cartographic modeling, Englewood Cliffs, NJ: Prentice-Hall, 1990.

- Tucker C.J, Wilson J.M, Mahoney R, Anyamba A, Linthicum K. and Myers M.F; (2002): Climatic and ecological context of the 1994-1996 Ebola outbreaks. *Photogrammetric Engineering & Remote Sensing* 68(2):147-152.
- van den Bos G.A. and Triemstra A.H; (1999): Quality of life as an instrument for need assessment and outcome assessment of health care in chronic patients, *Qual Health Care* 1999, 8:247-252.
- VCRC; (1998): The report of a training workshop on the "Collection, Processing and Analysis of data relating to operational aspects of filariasis control with special reference to Mass Drug Delivery (MDD)" organized by Vector Control Research Centre and National Filaria Control program at VCRC, Pondicherry, pp.1-17.
- Vlassoff C. and Bonilla E; (1994): Gender-related differences in the impact of tropical diseases in women: what do we know? *Journal of Biosocial Science*, 26:37-53.
- Waller L. A. and Gotway C. A; (2004): *Applied Spatial Statistics for Public Health Data*. John Wiley & Sons, New York.
- Wang H.J, Zhang R, Cole J, and Chavez F; (1999): El Nino and the related phenomenon Southern Oscillation (ENSO): the largest signal in interannual climate variation, *Advances in Parasitology*, Vol. 23. New York
- WHO; (1998): Research on rapid geographical assessment of bancroftian filariasis. Geneva, Switzerland: World Health organization. TDR/TDF/COMDT/98.2.
- WHO;(2000): The World Health Report 2000. Health Systems: Improving performance. Geneva: World Health Organisation. 2000.
- WHO; (2001): Malaria early warning systems, concepts, indicators and partners: a framework for field research in Africa. Roll Back Malaria/Technical Support Network for Prevention and Control of Malaria, Document WHO/CDS/R.

- WHO; (2004):Eliminating lymphatic filariasis: Egypt begins the final chapter in stopping a dreaded disease, Report; 29 October 2004,
http://www.who.int/features/2004/lymphatic_filariasis/en

- WHO; (2005a): Weekly epidemiological record, No. 23, 2005, 80, 201–212,
<http://www.who.int/wer>

- WHO; (2005b): HealthMapper 4.1, A WHO information and mapping application for public health, Learner's Guide
<http://www.who.int/csr/mapping/tools/healthmapper/healthmapper>

- WHO; (2006a): The disease and its epidemiology,
http://www.who.int/lymphatic_filariasis/epidemiology/en

- WHO; (2006b): Fact sheet N°102, Lymphatic filariasis,
<http://www.who.int/mediacentre/factsheets/fs102/en>

- WHO/CDS/CPE/CEE; (2000): Preparing and Implementing a National Plan to Eliminate Lymphatic Filariasis, WHO/CDS/CPE/CEE/2000, Report 15.

- WHO/CDS/CPE/CEE; (2002): global program to eliminate lymphatic filariasis, Annual Report on Lymphatic Filariasis, 2 0 0 1.

- WHO/TDR; (2004): New simulation model predicts the long-term impact of different treatment strategies ,Report No.43 Lymphatic filariasis, 2004.

- Zaffar, S. M. G. S. M, Ramalingam, M. and Venugopal, L;(2005): GIS for public health management, Anna University, Chennai 2005,
<http://www.gisdevelopment.net/application/health/overview/healtho0006.htm>

- Zagaria N. and Savioli L; (2002): Elimination of lymphatic filariasis: a public-health challenge. *Ann Trop Med Parasitol* 96 (Suppl. 2): S3–S13.