REFERENCES

- Ader R, (2000): Psychosomatic and psychoimmunologic research. psychosom Med 42: 307-321.
- Alfredsson L, Spetz C, Theorell T. (1995): Type of occupation and near future hospitalization for myocardial infarction and some other diagnoses. Int J Epidemiol 34: 378-88.
- Alterman T, Shekelle RB, Vernon SW, Burau KD, (1994): Decision Latitude, Psychologic demand, job strain and coronary heart disease in the western Electric study. Am J Epidemiol; 120; 620-7.
- American Heart Association (2000): 2001 Heart and stroke statistical update. Dallas, Texas, American Heart Association. www.Americanheart.org.
- Anderson KM, Castelli W, Levy D, (1998): Cholesterol and mortality. JAMA 257: 2176-80.
- Anderson KM, Wilson PWF, Odell PM and Kannel WB, (1991): An updated coronary risk profile: A statement for health professionals. Circulation 83: 356-362.
- Andrew MT, Paul MC, and Jean AC (1996): Disorders of lipoproteins metabolism and transport. In: Nilson Text Book of medicine.
- Appel LJ, Moore TJ, Obarzanek E et al., (1997): A clinical trial of the effects of dietary patterns on blood pressure. N Engl J Med 336: 1117-1124.

- Aranjo TM, Aquino E, Menezes G, Santos Co, Aquiar L. (2003): Work psychosocial aspects and psychological distress among nurses. Rev. Saude Publica. 37 (4): 424-33.
- Arthur HM, Garfinkel PE, Irvine J, (1999): Development and testing of a new hostility scale. Canadian journal of Cardiology, Vol. 15, No. 5, pp. 539-544.
- Aslan S, (2001): Stress and age effects on catecholamines in normal subjects. J Psychosom Res. 45: 33-41.
- Aspinwall LG, Taylor SE (1997): A stitch in time; self regulation and proactive copping. Psychological Bulletin, 121, 417-436.
- Assis MA, Kupek E, Nahas MV, Bellisle F, (2003): Food intake and circadian rhythm in shift workers a high work load. Appetite; 40 (2): 175-83.
- Baker DB, (2000): The study of stress at work. Annal Review of Public Health, 6, 367-381.
- Baker GHB, (1997): Psychological factors and immunity'. J Psychosom Res. 41: 1-10.
- Bakker AB, Killmer CH, Siegrist J, Schaufeli WB, (2000): Effort-reward imbalance and burn out among nurses. J adv Nurs; 31: 884-91.
- Bandura A, (1999): Self-efficacy: towards a unifying theory of behvaioural change. Psychological Review, 84: 191-215.

- Barasch E, Benderly M, Graff E, et al., (1995): Plasma fibrinogen levels and their correlates in 6457 coronary heart disease patients. the Bezafibrate infarction Prevention (BIP), Study J Clinic Epidemiol 48: 757-765.
- Barefoot J, Helms M, Mark D (1996): Depression and long term mortality risk in patients with coronary artery disease. Am J Cardiol 78: 613-617.
- Barry H, (1998): Relation of HDL to coronary heart diseases. Am. J. Cardiol 2: 5.
- Beale D, Clarke D, Cox T, Leather P, and Lawrence C, (1999): System memory in violent incident: Evidence from patterns of reoccurrence. Journal of occupational Health psychology, 4 (3): 233-244.
- Beale D, Cox T, Clarke D, Lawrence C, and Leather P, (1998):
 Temporal architecture of violent incidents. Journal of occupational Health psychology, 3: 65-82.
- Behar S, (1999): Lowering fibrinogen levels clinical update. BIP Study Group. Bezafibrate infarction Prevention. Blood Coagul Fibrinolysis. 10: 41-43.
- Bettenhausen KL, (2001): 5 years of groups research- what we have learned and what needs to be addressed. Journal of Management, 17 (12): pp. 345-381.
- Bhalla S, Jones B, and Flynn DM, (2001): Role stress among canadian white -collar workers. Work & Stress, 15, 289-299.
- Bjorntorp P, (1993): Visceral obesity a civillization syndrome'. Obesity Res 1: 206-22.

- Bjorntorp, Rosmond R, (2000): Obesity and cotisol. Nutrition. 16 (10): 924-36.
- Blankenhorn DH, Nessin SA and Johnson RL (1997): Beneficial effects of combined colestipoliniacin therapy on coronary atherosclerosis and coronary venous by-pass grafts JAMA. 257: 3233-40.
- Blumenthol (1995): Contribution of job strains, job status and marital status to laboratory blood pressure in patients with mild hypertension. Psychosomatic medicine, 60, 697-706.
- Blumenthol, JA, Thyrum, ET, Siegel WC, (1995): Contribution of job strain, job status and marita status to laboratory blood pressure in patients with mild hypertension. Psychosomatic medicine, 60, 697-706.
- Boggild H, & Knutsson A, (1999): Shift work, risk factors and cardiovascular disease. Scandinavaian Journal of work Environment of health, Vol. 25, No. 2, pp. 85-99.
- Bojanowska E, (2005): Physiology and pathophysiology of glucagonlike peptide-1 (GLP-1): the role of GLP-1 in the pathogenesis of diabetes mellitus, obesity, and stress. Med Sci Monit Aug; 11 (8): RA 271-278. Epub. 2005.
- Bosma H, Marmot M, Hemingway H, Nicholson A, et al., (1997): Low job control and risk of coronary heart disease in white hall II (prospective cohort) study. BMJ, 314: 558-65.
- Bosma H, Peter R, Siergrist J, and Marmot M. (1998): Two alternative job stress models and the risk of coronary heart disease.

 American Journal of public Health, 88, 68-74.
- Bounhoure GH, Rubins HB, Collins D, et al., (1997): Correlates and consequence of diffuse atherosclerosis in men associated with helicobacter pylori. Br Gen Pract 41: p. 286-288.

- Byle JA, Greig, WR, Jasani MK, Duncan A, Diver M, & Buchanan WW, (1997): Relative role of genetic and environmental factors in the control of serum uric acid levels in normouricaemic subjects. Annals of Rheumatic Diseases, 26, 234-238.
- Bradley G, (1999): Computers and the psychological work Environment.

 Taylor and Francis, London.
- Brand FN, McGee DL, Kannel WB, (1995): Hyperuricemia as a risk factor of coronary heart disease: The Framingham Study. American Journal of Epidemiology, 121, 11-18.
- Bray G, (2000): Epidemiology of obesity. In: progress in obesity research Libbey; 639-43.
- Brenner P, Brim OG, and Rottier, JB (2000): Stress and coping under extreme conditions.' In Dimsdale, JE, (Ed.), Survivors, victims, and perpetrators: essays on the Nazi holocaust. Washington, DC: Hemisphere, PP, 219-258.
- Breslow L and Buell P, (2000): Mortality from coronary heart disease and physical activity of work in california. Journal of chronic diseases 51, 87-91.
- Brindley DN, McCann BS, Niaura, R, Stoney CM, and Suarez EC, (2003): Stress and lipoprotein metabolism: Modulators and mechanism. Metabolism, 52, 3-15.
- Brook A, (2003): Mental stress at work. Practitioner, 210, 500-506.
- Brown DE, James GD, Nordloh L, Jones AA. (2003): Job strain and physiological stress responses in nurses and nurs's aids; Predictors of daily blood pressure viability. Blood press Monit. 8 (6): 237-42.

- Brownlee M, Vlassara H, Cerami A, (1995): Non enzymatic glycosylation products in collagen covalently trap low density lipoportein. Diabetes; 34: P. 938-941.
- Brunner E. (1997): Stress and biology of inedquality. Brit Med J 314: 1472-6.
- Buck V, (2002): Working under pressure. Staples press, London.
- Buel P, and Breslow, L (2000): Mortality from CHD and Physical activity of work in California 'J Chronic Dis 61: 615-626.
- Bushwald H, Varco RL and Natts JP, (2002): Effect of partial ilia by pass surgery on mortality and morbidity from coronary heart disease in patients with hypercholesteroloemia. N Engl J Med. 323: 946-55.
- Cambien F, Chretien JM, (1995): Is the relationship between blood pressure and cardiovascular risk dependent on body mass index? Am J Epidemol 122 (3): 434-42.
- Cappuccio FP, Strazzullo P, Farinaro E and Trevisan M, (2003): Uric acid metabolism and tubular sodium handling: Results from population-based study. Journal of the American Medical Association, 270, 354-359.
- Caraway W, (1996): Determination of serum uric acid with deproteinization Clin. Chem. 12: 18.
- Carney RM, Blumenthal JA, Stein PK et al., (2001): Depression, heart rate variability, and acute myocardial infarction. Circulation. 104: 2024-2028.
- Castelli WP, (1998): Cardiovascular diseases in women. Am J Obstet Gynecol 158 (6 Pt. 2): 1553-1560, 1566-1567.

- Chappel D and Di Martino V, (1998): Violence at work. Geneva: international Labour Office.
- Cheng Y, Kawachi I, Coakley E, et al., (2000): Association between psychosocial work characteristics and health functioning in American women: prospective study. BMJ; 320: 1432-36.
- Chislm GM, Irwin KC, Penn Ms. (1998): Lipoprotein oxidation and lipoprotein induced cell injury in diabetes. Diabetes. 1998; 41 (suppl 2): 61-66.
- Chu NF, Wang DJ, Liou SH, Shieh SM. (2000): Relationship between hyperuricemia and other cardiovascular disease risk factors among adult males in Taiwan. Eur J Epidemiol Jan; 16 (1): 13-7.
- Clarke R, Frost C, Collins R, et al., (1997): Dietary lipids and blood cholesterol: quantitative meta-analysis of metabolic ward studies. Br Med J 314: 112-117.
- Clarkson P, Montgomery HE, Mullen MJ, et al., (1999): Exercise training enhances endothelial function in young men. J Am Coll cardiol 33: 1379-1385.
- Cohen H, Nield GH, Patel R, Machie IJ, Machin, SJ, (2003): Evidence for chronic platelet hyperaggregability and in vivo activation in cyclosporine -treated renal allograft recipients. Thrombos. Res. 49: p. 91-101.
- Cohen SG, and Ledford, GE, (2003): The effectiveness of self-managing teams a quasi- experiment. Human relations, 47 (1), pp. 13-43.
- Conen D, Wietlisbach, V, Bovet P, Shamlaye C, et al., (2004):

 Prevalence of hyperuricemia and relationship of serum uric acid with cardiovascular risk factors in a developing country.

 BMC Public Health, 25; 4 (1): 9.

- Conn H, Kostis J and Saviano G, (1995): Diseases of the coronary arteries CH8. Diseases of the blood vessels. Eds. Horwitz O.; Mc Combs, P.; Roberts, B, Lea and Febiger Co. Philadelphia. P. 130-162.
- Consensus -Building Meeting (2000): Cultural competence Curriculum Models for Family Physicians. Extracts from Health People 2010 for Top Seven Areas of Health disparities. US Department of Health and Human Services. Healthy people 2010 (Conference Edition). Washington, DC.
- Cooper CL, (2002): Work stress. In: P.B, Warr (eds.) Psychology at Work. Penguin, Harmondsworth.
- Cooper CL, and Marshall, (1996): Occupational sources of stress: a review of the literature relating to coronary heart disease and Mental ill health. J Occup Health Pychol 49: 11-28.
- Cooper CL, Sloans J, William S, (1998): Occupational stress indicator management guide. Windsor: NFER- Nelson.
- Cox T, (1999): The recognition and measurement of stress: Conceptual and methodological issues. In: E.N., Corlett & J.
- Cox T, and Griffiths AJ, (1995): The assessment of psychosocial hazards at work. In MJ, Shabracq, JAM, Winnubst and CL Cooper (Eds.) Handbook of work and Health psychology Chichester: Wiley and Sons.
- Cox T, and Thomson L, (2000): Organization healthiness: work -related stress and employee health- in p. DeWe, M. Leiter and T.
- Crawford DW, and Plankenorn DH, (1996): Arterial wall oxygenation, oxyradicals, and atherosclerosis. Atherosclerosis; 89: p. 97-108.

- Csef H, Hefnerg (2005): Stress and myocardial infarction. MMW Fortschr Med. 31; 147 (13): 33-5.
- Danelia M, Trapaidze D, (2005): Psychosocial work environment and coronary heart disease. Georgian Med News. (4): 56-8.
- Danesh J, Wincup P, Walker M et al., (2000): Chalmydia pneumoniae IgG titres and coronary heart disease: Prospective study and Meta analysis. BMJ 321: 208-13.
- Danna K and Griffin RW, (1999): Health and well-being in the workplace: A review and synthesis of the literature. Journal of Management, 25, 357-384.
- Davidson DM, (1994): Cardiovascular disease in occupation. Cardiovascular Rev Rep 5: 503.
- Davidson MJ, and Cooper CL (2001): A model of occupational stress. J Occup. Med. 43: 564-574.
- De Bacquer D, Pelfrene E, Clays E, et al., (2005): Perceived job stress and incidence of coronary events: 3 years follow-up of the Belgian Job Stress project Cohort. Am J Epidemiol. 1; 161 (15): 434-41.
- Dewe, P, (2000): Measure of coping with stress at work: a review and critique. Inp. Dewe, U, Leiter and T. Cox (Eds) Coping, Health and organizations. London: Taylor and Francis
- Di Lorenzol, Depergola G, Zocchetti C, L'Abbate N, et al., (2003): Effect of shift work on body mass index: results of a study performed in 319 glucose- tolerant men working in a southern Italian industry. Int J Obes, Relate Metab Disord, 27 (1): 353-8.

- Diament, J and Byers, SO, (2002): Aprecise catecholamine assay for small samples, Journal of laboratory and clinical Medicine, 679-693.
- Dimsdale JE, and Herd JA, (2002): Variability of plasma lipids in response to emotional arousal. Psychosomatic Medicine, 84, 413-427.
- Dorian B, Garfinkel P, Brown G, (2002): Aberrations in lymphocyte subpopulations and functions during Psychological stress". Clin Experimen Immunol. 70: 132-138.
- Dorn JM, Schisterman EF Winkelstein WJ, Trevisan M (1997): Body Mass index and mortality in a general population sample of men and women: The buffalo health study. Am J of Epidemiol 146 (11): 919-931.
- Drexler H, Zeiher AM, Meinzer K and Just H, (1996): Correction of endothelial dysfunction in coronary microcirculation of hypercholesterolemic patients by L-arginine. Lancet, 338: 1546-1550.
- Dublin LI, (2003): Relation of obesity to longevity. N. Engl. J. Med., 348: 971-4.
- Dubrow R, Burnett AA, Gute DM, and Brockert JE, (1998): Ischaemic heart disease and acute myocardial infarction among police officers. J occup, Med, 50: 650.
- Eden D, (2002): Critical job events, acute stress, and strain: A multiple interrupted time series. Organization Behavior and Human Performance, 30, 312-329.
- Ekehammer B, (2000): Interactionims in personality from a historical perspective. Psychological Bulletin, 81, 1026.

- Elisaf M, (2001): The treatment of coronary heart diseases: an update: part 1: an overview of the risk Factors for cardiovascular Disease. Curr Med Res Opin 17 (1): 18-26
- Elmer PJ, (1994): Obesity and cardiovascular diseases: practical approaches for weight loss in clinical practice. In pearson TA, Crique MH, Luepker RV, et al: Primer in preventive cardiology. Dallas: American Heart Association; 189-204.
- Emmerson BT, (2001): Identification of the causes of persistent hyperuricemia. Lancet; 347: 1461-6.
- Endresen IM, Vaernes RJ, Ursin H, and Tonder O, (1995):

 Psychological stress factors and concentration of immunoglobulins and complement components in Norwegian nurses. Work & Stress, 1, 365-375.
- Epel E, McEwen B, Seeman T, Mahhews K, et al., (2000): Stress and body shape. Psychosom Med. 62 (5): 623-32.
- Fahey TD (2002): Coronary Heart Disease and Exercise. California state University, Chico Exercise Physiology Laboratory, Chico CA, USA. WWW.sportsci. org/encye/drafts.
- Fang J, Alderman MH, (2000): Serum uric acid and cardiovascular mortality the NHANESI epidemiological follow up study, 1971-1992. National Health and nutrition Examination survey. JAMA; 10; 283 (18): 2404-10.
- Ferguson D, (2001): A study of occupational stress and health. Ergonomics, 16, 649-633.
- Ferketich AK, Schwartzbaum JA, Frid DJ, Moeschberger ML (2000):

 Depression as an Antecedent to Heart Disease Among
 Women and Men in the NHANEs 1 study. Arch intern 160:
 1261-1268.

- Fielden SL and Peckar C.J, (1999): Work stress and hospital doctors: a comparative study. Stress Medicine, Vol., 15, no. 3, pp. 137-141.
- Fishel R, Shai S, Narc K, (1996): Endothelial cell angiotensin coverting enzyme expression is unregulated by fluid shear stress (Abstract) FASEB J, 5: A: P. 1247.
- Folkard S and Monk TH, (2001): Hours of work-temporal factors in work scheduling. Wiley and Sons, Chichester.
- Fossat P, (1992): Determination of triglycerides by enzymatic calorimetric method. Clin Chemi 28: 2077
- Frankenhaeuser M, (2000): Comparison between type A and type B males and females. Biologic psychol. 30-79-91.
- Franklin SS, Khan SA, Wong ND, Larson MG, Levy D, (1999): Is pulse pressure useful in predicting risk for coronary heart disease? The Framingham heart study. Circulation 100: 354-60.
- Fredrikson, M, and Matthews KA, (2000): Cardiovascular responses to behavioral stress and hypertension: A meta-analytic review. Annals of Behavioral Medicine, 22, 30-39.
- French JRP, and Caplan RD, (2002): Organizational stress and individual strain. In: A Marrow (ed) the Failure of Success. AMACOM, New York.
- French JRP, Caplan RD, and Van Harrison R, (1992): The mechanisms of job stress and strain. New York: wiley.
- Friedman MD, Rosenmann RD, and Carroll V, (1998): Changes in serum cholesterol and blood clotting time in men subjects to cyclic variation of occupational stress. Circulation 67: 852-861.

- Frohlich ED, (2003): Uric Acid: A risk factor for coronary heart diseases. Journal of the American Medical Association, 270 (3), 378-379.
- Froom L, Kristal Boneh E, Melamed S, Harari G, et al., (2000): Serum total choelsterol and cardiovascular mortality in Israel: Males: the CORIS study. Cardiovascular occupational risk factor, Determination in Israeli Industry. Isr Med Assoc J, 2 (9): 668-71.
- Garcia MI, MicNamara PM & Gordon T. (1996): Morbidity and mortality in diabetics in the Framingham population. Diabetes 23: 105.
- Gardell B, (2000): Alienation and mental heatlh in the modern industrial environment. In: L. Levi (ed) Society, Stress and Disease Vol. 6, oxford university press, oxford.
- Gardell B, Gustavsen B, (2000): Work environment research and social change: current developments in scandinavia. J Occup Behav 16: 13-17.
- Genest JJ, Marcil M, Denis M et al., (1999): High—density lipoprotein in health and disease. J. Invest. Med. 47: 31-42.
- Georges E, Wear ML, Muller WH. (2002): Body fat distribution and Job stress in Mexican American men of the hispanic Health and Nutrition Examination survey. Am J Hum Biol; 14: 657-67.
- Giamurce SH, (1998): Apolipoprotein E mediates uptake of S.F. 100-400 Hypertriglyceridemic VLDL by LDL receptors pathway in normal human fibroblast. J Biochem Chemi, 250-452.
- Gordon DJ (2000): Role of circulating high-density lipoprotein and triglycerides in coronary artery diseases: risk and prevention. Endocrinol Metab Clin North Am Jun 19 (22): 299-309.

- Gordon DJ and Rifkind BM (1999): High density lipoprotein. The clinical implication of recent studies. N Engl J Med 321: 1311.
- Gotto AM (1998): Triglycerides the forgettin risk factor. Circulation 97: 1027.
- Gotto AM, and Farmer JA, (1998): Risk factor for CAD. In Braunwald E.W.B. Heart disease. Vol. 2. Text Book of Cadiovascular Medicine, 3rd (Eds). Saunders Company. Philadelphia, London, 1153.
- Gray -Toft P, and Anderson JG, (2001): Stress among hospital nursing staff: its causes and effects. Social sciences and Medicine, 45: 639-647.
- Greiner BA, Krause N, Ragland D, et al., (2004): Occupational stressors and hypertension: amulti-method study using observer based job analysis and self-reports in urban transit operators Soc Sci Med. 59 (5): 1081-94.
- Grundy SM, Balady GJ, Criqui MH and et al., (1998): Primary prevention of coronary Heart Disease: Guidance From Framingham. Circulation 97: 1876-1887.
- Grundy SM, Goodman DW, Rifkind BM, et al., (1999): The place of HDL in cholesterol management. A perspective from the National Cholesterol Education Program. Arch Intern Med. 149 (3): 505-10.
- Guppy A, Weatherstone L, (1997): Coping strategies, dysfunctional attitudes and pscychological well-being in white collar public sector employees. Work and stress, 11, 58-67.

- Hallquist J, Diderichsen F, Theorell T, et al., (1998): The SHEEP study group: Is the effect of job strain on myocardial infarction due to international between high psychological demands and low decision latitude? Results from stockholm Heart epidemiology program (SHEEP). Soc Sci Med 46: 1405-15.
- Hamilton L- and Gregor F, (1996): Self directed learning in a critical care: nursing program. J. contin. Educ. Nurs. 17: 94-99.
- Harrell J, (2000): Psychological factors and hypertension: A status report. Psychol. Bull. 87: 482-501.
- Harrington JM, (1998): Shift work and Health: A critical review of the Literature. HMSO, London.
- Harris B, Zinman B, Hanley A et al., (2003): The impact of diabetes on cardiovascular risk factors and outcomes in a native canadian population. Daibetes Res. Clin pract, 55 (2): 1657-3.
- Haudeschild CC, Prescott MF, chobanian AV, (1998): Aortic endothelial and subendothelial cells in experimental hypertension and aging. Hypertension 3 (suppl. 1): 148-153.
- Hauschildt E, (2002): Fish and Omega -3 Fatty Acid intake and risk of coronary heart disease in Women. JAMA 287: 1815-1821.
- Haynes SG, and Feinleib M, (2000): Women, Work, and CHD. Prospective findings from the Framingham Heart study. Am. J. Public Health. 90: 133-141.
- Health and Safety executive (2000): Organizational inter ventions for work stress: a risk management approach. Norwich: Her Majesty's stationery office, 2000.

- Heany CA, Price RH, Reffeoty J. (1995): Increasing coping resources at work: a field experiment to increases social support, improve work team. Am J Of Epidemiol (1): 37-58.
- Hemingway H, Marmot M. (1999): Psychosocial factors in the etiology and prognosis of coronary heart disease: systematic review of prospective cohort studies. Brit Med J 318: 1460-7.
- Herlitz J, Karlson WB, Lindqvist J, et al., (2001): Important factors for the 10-Year Mortality Rate in patients with Acute Chest pain or other symptoms consistent with acute myocardial infarction with Particular Emphasis on the influence of Age. Am Heart J 142 (4): 624-632.
- Hertz RP, Unger AN, Mc Donald M, et al., (2004): The impact of obesity on work limitation and cardio vascular risk factors in the U.S. work force. J Occup Environ Med. 46 (12) 1196-203.
- Hoieggen A, Alderman MH, Kjeldsen SE, et al., (2004): The impact of serum uric acid on CV outcomes in the LIFE study. Kidney Int. 66 (4): 1714-5.
- Holland PJ, (1998): Psychiatric aspects of occupational medicine. 'In Mccunney, R.J. Ed. Hand Book of occupational Medicine, Little, Brown and Company, Boston/Toronto, p. 237.
- Holvoet P, Johan V, Jannse S and Collen D (1998). Oxidised LDL in patients with acute coronary syndromes. Circulation 98: 1487-1494.
- House J, (1994): Occupational stress and coronary heart disease: a review and theoretical integration. J Hlth Soc Behav 35: 12-27.
- House JS, (2001): Work stress and social support. Reading, MA: Addison Wesley.

- Hulst V, (2003): Long work hours and health. Scand J Work Environ Health. Jun; 29 (3): 167-9.
- Hunt D, (1997): Etiology and pathogenesis of CAD, smoking hyperlipidemia and primary prevention of CAD. In Carlisle, C. Hurd, D. and Slombi G, Coronary care work book, 3rd (Eds). Boehringer Ingelheim. P. 49.
- Neidhammer I, Flert and MJ Marne, (1996): Prevalence of over weight and weight gain in relationship work in nurse's cohort. Int J Obes Relat Metab Disord 20, 625-633.
- Ibrahim MM, (2001): Epidemiology of hypertension in Egyptians. In: Manual of hypertension, Chapter: 15, pp. 182.
- Ingersoll GL, Cook JA, Fogel S, Applegate M, Frank B (1999): The effect of patient -focused redesign on mid level nurse managers' role responsibilities and work environment. Journal of nursing Administration, Vol. 29: No. 5. pp. 21-27.
- International Labour Organization (1996): Psychological factors at work: Recognition and Control. Occupational safety and Health series No: 56: International labour office, Geneva.
- Ishizaki M, Morikawa Y, Nakagawa H, Honda R, et al., (2004): The influence of work characteristics on body mass index and waist hip ratio in Japanese employees. Ind Health. 42 (1): 41-9.
- James GD, and Brown DE, (1997): The biological stress response and lifestyle: Catecholamines and blood pressure. Annual Review of Anthropology, 26: 313-353.
- James GD, Broege, PA and Schlussel YR, (1996): Assessing cardiovascular risk and stress-related blood pressure variability in young women employed in wage jobs. (American Journal of human biology), 8, 743-749.

- Jenkins CD, (1995): Psychologic and Social precursors of coronary disease: N Engl J Med 284: 255; 307-317.
- Jenkins CD, (2003): Psychosocial risk factors of CHD. Act Med Scand, Suppl 660: 123-136.
- Jensen J, Blankenhorn, DH, Chin HP, Strugeon P and Ware AG, (1995): Serum lipids and serum uric acid in human twins. Journal of Lipids Research, 6, 193-205.
- Jeppsen J, Hein HO, Suadiciani P and Gyantelberg F, (1998): Triglycerides concentrations and ischemic heart diseases. Circulation. 97: 1029-1036.
- John A, Farmer Antonio M, Gotto et al., (2002): Risk factors for coronary artery diseases. Braunwald heart disease 4th edition. Volume, 2: P1151.
- Joint National Committee on Detection, Evaluation and Treatment of High Blood Pressure (2003): The seventh report of the Joint National Committee on Detection, Evaluation and Treatment (JNC 7). JAMA, 289 (19): 2560-2572.
- Jones JR, Hodogson JT, Clegg TA, and Elliot RC, (1998): Self-reported work related illness in 1995: results from a household survey. Subdury: HSE Books.
- Jousilahti P, Tuomilehto J, Vartiainen E, Pekkanen J, Puska P, (1996):

 Body weight, cardiovascular risk factors, and coronary mortality, 15-year follow-up of middle-aged men and women in eastern Finland. Circulation 93: 1379-1379.
- Kagan NI, Kagan H, Watson MG. (2000): Stress reduction in the work place: the effectiveness psychoeducational programs. J counselling psychol; 52: 71-8.

- Kahn RL, Wolfe DM, Quinn RP, Snoek JD, and Rosenthal RA, (2002):
 Organizational stress: studies in Role conflict and Ambiguity.
 Wiley and Sons, New York.
- Kang MG, Koh SB, ChaBS et al., (2005): Job stress and cardiovascular risk factors in male workers. Prev Med. 2005; 40 (5): 583-8.
- Karasek RA, (1990): Job characteristics, occupation, and coronary heart disease. 'cincinnati, OH: NIOSH.
- Karasek RA, and Theorell T, (1990): Healthy work: stress, productivity and the reconstruction of working life. New York, NY, Basic Books.
- Karbapas P, Hilkki M, Laakso M, (1994): Isolated low HDL cholesterol an insulin-resistant state. Diabetes 3: 411-417.
- Karlsson BH, Knutsson AK, Lindahl BO, Alfredsson LS. (2003): Metabolic disturbances in male workers with rotating three-shift work. Results of the WOLF study. Int Arch occup Environ Health. 76 (6): 424-30.
- Kasl SV, (1996): The influence of the work environment on cardiovascular health a historical, conceptual and methodological perspective. J Occup Health Psychol; 1; 42-56.
- Kasl SV, (1997): Surveillance of psychological disorders in the workplace. In GP Keita and SL Sauter (eds) Work and Well. Being: An Agenda for the 1990s. American psychological association, Washing ton DC.
- Kawachi I (1998): Passive smoking and coronary heart disease. Cardiologia 43 (7): 667-675.

- Kawachi I, Colditz GA, Stampfer MJ, et al., (1997): Prospective study of shift work and risk of coronary heart disease in women. Circulation 92, 3178-3182.
- Kelley WN, and palella TD (1997): Gout and other disorders of purine metabolism. In E.Braunwald, K.J. Isselbacher, R.G. Petersdorf et al., (eds) Harrison's principles of internal medicine. (pp. 1623-1632). New York: McGraw-Hill.
- Kissebath AH, Peiris AN, (1999): Biology of regional body fat distribution; relationship to non insulin- dependent diabetes. Diabetes Metabol., Rev., 15: 527-35.
- Kivimaki M, (2002): Work stress doubles heart risk Occup Environ med 59: 75-77.
- Knutsson A and Boggild H, (2000): Shift work, risk factors and cardiovascular disease: Review of disease mechanisms. Rev Environ Health 15, 359-372.
- Knutsson A, Karlsson B, Ornkloo K et al., (2002): Post pradnial response of glucose, insulin and triglycerides influence of the timing of meal intake during night work. Nutr Health.; 16 (2): 133-41.
- Kobasa SC, Maddi SR, Puccetti MC, and Zola MC, (1995): Effectiveness of hardiness, exercise and social support as resources against illness. J Psychosom Res. 39: 525-533.
- Kobayashi F, Furui H, Akamatsu Y, Watanabe T, Horibe H, (1999):
 Changes in psychophysiological functions during night shift in nurses-influence of changing from a full-day to a half-day work shift before night duty. International Archives of occupational and Environmental Health, Vol., 69, No. 2, pp. 83-90.

- Kornhauser M, Kittel F, and Debacker G, (2001): The Belgian Heart Diseases prevention project: type "A" behavior pattern and the prevalence of CHD. Psychosom Med 63: 133-145.
- Krantz DS, and Manuck SB, (1994): Acute psychophysiologic reactivity and risk of cardiovascular diseases a review and methodologic critique "Psychol Bull 96: 435-464.
- Krauss RM, (1997): Relationship of intermediate and low density lipoproteins to risk of coronary artery heart diseases Am Heart J 113: 578.
- Kristensen TS, (1996): Job stress and cardiovascular disease; A theoretical and critical review. Journal of Occupational Health psychology, 1, 246-260.
- Kujala UM, Kaprio J, Sarna S, Koskenvuo M, (1998): Relationship of leisure-tem physical activity and mortality; the Finnish Twin Cohort JAMA 279: 440-444.
- Kushi LH, Meyer KA, Jacobs DR (1999): Cereals, legumes, and chronic disease risk reduction: evidence from epidemilogical studies. Am J Clin Nutr 70 (Suppl): 451S-458S.
- Kyriacou C, (2000): Coping actions and occupational stress among teachers. Res. In Educ. 44: 57-61.
- La Rocco JM, House JS, and French JRP, (2000): Social support, occupational stress, and health. J Health and social behavir 41: 202-208.
- Laasko M, (1996): Insulin resistance and coronary heart disease. Curr Opin Lipidol 7, 217-226.

- LaCroix A, Haynes S. (1994): Occupational exposure to high demand/low control work and coronary heart disease incidence in the Framing ham cohort. Am J Epidemiol 130; 481.
- Lakka TA, Venalaienen JM, Rauramaa R, et al., (1994): Relation of leisure- time physical activity and cardiorespiratory fitness to the risk of acute myocardial infarction in men. N. Engl. J. Med. 330: 1549-1554.
- Landbergis P, and Theorell T, (2000): Measurements of psychosocial work place exposure variables. Occup Med. 15: 163-88.
- Landy FJ, (1999): The psychology of work Behaviour. Brook, Cole, Monterey, California.
- Lapidus L, Bengtsson C, Larsson B, et al., (1999): Distribution of adipose tissue and risk of cardiovascular diseases and death: 12th year follow-up of participants in the population study of women in Gotebiog, Sweden Br Med J 289: 1257-61.
- Larsson B, Svardsudp K, Welin L, et al., (1994): Abdominal adipose tissue distribution, obesity and risk of cardiovascular disease and death: 13 year follow up of participants in the study of men Br Med J 288: 1401-4.
- Lazarus RS, (1999): Stress and emotion: A new synthesis. New York New York; Springer.
- Leather P, Lawrence C, Beale D, Cox T, and Dickson R, (1998): Exposure to occupational violence and the buffering effects of intra- organizational support. Work and Stress, 12 (2): 161-178.

- Lee J, Sparrow D, Vokonas PS, Landsberg L and Weiss ST, (1995):
 Uric acid and coronary heart disease risk: Evidence for a role
 of uric acid in the obesity- insulin resistance syndrome.
 American Journal of Epidemiology, 142 (3), 288-294.
- Lee S, Graham Coldditz, Lisa Berkman and Ichiro Kowachi. (2002): A prospective study of job strain and coronary heart disease in US women. International Journal of Epidemiology, 31: 1147-53.
- Lefkowitz RJ, Willerson JT, and Willerson MD (2001): Prospects for cardiovascular Research. JAMA Vol 285. 5, February 7.
- Legrand D, Legrand V, (2005): Coronary artery disease associated to diabetes Mellitus Rev Med. Liege.; 60 (5-6): 526-30.
- Lennernas M, Akerstedt T, Hambraeus L. (1994): Nocturnal eating and serum cholesterol of three-shift works. Scand J Work Environ Health; 20 (6): 401-6.
- Levenstein S, Margot W Swith, George Kaplan (2001): Psychosocial predictors of hypertension in men and women. Arch Intern Med, 161: 1341-1346.
- Levey R, and Feinleib M, (1998): Risk factors CAD and their management. CH. 35. Heart disease, Vol. 2. A text book of cardiovascular medicine. 5th ed. Ed By: E. Braunwald. W.B. Saunders Company. Philadelphia, Lodon. P. 1205-1234.
- Levi L, (2002): Stress and distress in response to psychosocial stimuli.

 Acta Medica Scandinavica, 191, Supplement: 528.
- Leviene GN, Keany JF, (1995): Cholesterol reduction in cardiovascular disease. N Engl J Med. 332: P. 512-521.

- Lim JK, (1998): Hyperlipidemia, classification, Pathology and pathophysiology in: Hyperlipidemia. Supernaw, RB, (ed). Kansas Pharmat Inc. p. 1.
- Lissner L, Odell PM, D'Agostino RB, et al., (2001): Variability of body weight and health outcomes in the framing ham population. N Engl J Med 324: 1839-44.
- Liu Y, and Tanaka H, (2002): Overtime work, in sufficient sleep, and risk of non fatal acute myocardial infarction in Japanese men.

 Occup Environ Med 60: 67-75.
- Liuzzo G, Biasucci LM Galimore R, et al., (1994): The prognostic value of C-reactive protein and serum amyloid a protein in severe unstable angina. N Engl J Med 331-417-424.
- Liveshit G, Weistart J, Meshulam N, and Bruner D, (1999): Multivariate analysis of the twenty year follow up of Donolo-TaelA in prospective coronary artery disease study and the usefulness of high density lipoprotein cholesterol percentage. Am. J. Cardiol. 63: p 676-681.
- Lokk J, Arnetz B, (1997): Psychophysiological concomitants of organizational change in health care personnel: effects of a controlled intervention study. J Psychosom; 66: 74-7.
- Luepker R, (1998): Heart Disease: From Text book maxcy Rosenau-Last, Public Health and preventive Medicine York Production services USA: 927-948.
- Luepker RV, Rosamond WD, Murphy R, et al., (2003): Socioeconomic status and coronary heart disease risk factor trends. The Minnesota Heart Survey. Circulation 88: 2172-2179.
- Lynch J, (2002): Psychosocial factors strengthening the evidence base. BMJ, 324: 1251.

- Lyness, SA (2003): Predictors of differences between type A and B individuals in heart rate and blood pressure reactivity. Psychological Bulletin, 124, 266-28.
- MacLennan A, (2000): Physical activity and coronary heart disease risk in men, does the duration episodes predict risk. Circulation 102: 981.
- Malcolm RM, Harrison J, Forster H. (2003): Effects of changing the pattern of referrals in a local authority. Occup Med 63: 211-15.
- Malloy MJ, Kane JP (2001): A risk factor for atherosclerosis: triglyceride-rich lipoprteins. Adv. Intern. Med. 47: 111-136. castelli WP (1994): Epidemiology of coronary heart disease: the Framingham study. Am J of Med. 27: 4-12.
- Mangin EL, Kugiyama JR, and Nguy JH, (1998): Effects of isolipids and oxidatively modified low density lipoprotein on endothelium- dependent relaxation of rabbit aorta. Circ Res, 72: 161-166.
- Manson JE, Colditz GA, Stampfer MJ, et al., (1997): A prospective study of maturity onset diabetes mellitus and a risk of coronary heart disease and a stroke on women. Arch Intern Med 151: 1141-1147.
- Manson JE, Hu FB, Rich-Edwards JW, et al, (1999): A prospective study of walking as compared with vigorous exercise in the prevention of coronary heart disease in women. N Engl J Med 341: 650-658.
- Margolis B, L, & Kroes, WH, (1994): Work and the Health of man. In: J. O'Toole (ed) Work and the Quality of Life. MIT Press, Cambridge, Mass.

- Margolis BL, Kroes WH, ad Quinn RP, (2003): Job stress: an unlisted occupational hazard. Journal of occupational Medicine, 31, 652-661.
- Markovitz JH, Matthews KA, Whooley M, Lewis CE, Greenlund kJ, (2004): Increases in job strain are associated with incident hypertension in the CARDIA study. Ann Behav Med. 28 (1):1-3.
- Martin JR, Tato F, Mostaza JM, Vega GL, Grandy SM (1996): Effectiveness of low -dose crystalline nicotinic acid in men with low high -density lipoprotein cholesterol levels. Arch Intern Med 156: 1081-1088.
- Matsubara K, MatsuZawa Y, Jia S, et al., (1999): Relationship between hypertriglyceridemia and uric acid production in primary gout. Metabolism. 48: 688-701.
- Mattiasson I, Lindgarde F, Lilsson JA, and Theorell T, (2000): Threat of unemployment and cardiovascular risk factors: longitudinal study of quality of sleep and serum cholesterol concentrations in men threatened with redundancy. British Medical Journal, 31, 461-466.
- Mayes PA, (1998): Lipids transport and storage in: Harpar, s Biochemistery Text book. Murray PR, Granner, DK, Mayes PA, (eds). East Norwalk, Appleton and Lange Co. p. 251.
- Meade TW, (2001): Cardiovascular disease-linking pathology and epidemiology. Int J epidemiol 30: 1179-83.
- Melamed S, Ben- Avil I, Luz J, Green MS, (1995): Repetitive work, work under load and coronary heart disease risk factors among blue-collar workers- the CORDIS study. Cardiovascular Occupational risk factors Determination in Israel. J Psychosom Res. 39 (1): 19-29.

- Melamed S, Kushnir T, Strauss E, et al., (1997): Negative association between reported life events and cardiovascular disease risk factors in employed men: The CORDIS Study. Cardiovascular occupational risk factors: Determination in Israel. J Psychosom Res. 43 (3): 247-58.
- Mendelson ME, Craeger MA, Cooke JP, (2000): C-reactive protein and its relation to cardiovascular risk factors. B.M.J 312: P. 1061-1065.
- Mesler R, Capobianco M, (2001): Psychosocial factors associated with job stress. Stress news, 13 (4), 543-458.
- Messerli F.H. (2002): Cardiovascular effects of obesity and hypertension. Lancet, 1165-8.
- Messerli FH, Frohlich ED, Dreslinski GR, et al. (2009): Serum uric acid in essential hypertension: an indication of renal vascular involvement. Ann Intern Med. 113: 817-21.
- Michael J, (2001): Primary and secondary prevention of coronary heart disease In: Heart disease. A textbook of cardiovascular medicine, 6th edition W.B. Sounders CO.
- Michie S, (2002): Causes and management of stress at work. Occup Environ Med 59: 67-72.
- Mikhailidis DP, Ganotakis ES, Spyropoulos KA, et al., (1998):

 Prothrombotic and lipoprotein variables in patients attending a cardiovascular risk management clinic: response to ciprofibrate or lifestyle advice Int angiol 17: 225-233.
- Mikkelsen WM, Dodge HJ, Valkenburg H, et al., (1995): The distribution of serum uric acid values in a population unselected as to gout or hyperuricemia. Am J Med 69: 242-51.

- Miller NE (1997): Association of high density lipoprotein subclasses and apoliporprteins with ischemic heart disease and coronary atherosclerosis. Am Heart J 113: 589-97.
- Mina H and Park J (2005): Shift work and Metabolic risk factors of cardiovascular disease. J Occup Health; 47: 89-95.
- Mitchell GF, Moye LA, Braunwald E, et al., (1997): Sphygmomanometrically determined pulse pressure is a powerful independent predictor of recurrent events after myocardial infarction in patients with impaired left ventricular function. SAVE investigators, Survival and Ventricular Enlargment. Circulation 96: 4254-4260.
- Mohsen MI, (2002): Lecture presented at the clinical society classroom, Heart J Vol., 5.
- Mueller EF, Kasl SV, Brooks GW, and Cobb S, (2000): Psychosocial correlates of serum urate levels. Psychological Bulletin, 73, 238-257.
- Najem GR, Seebode JJ, Samedy AJ, Feuerman M, and Freidman L, (1997): Stressful events and the risk of symptomatic kidney stones. International Journal of Epidemiology, 26, 1017-1023.
- Nakanishi N, Okamota M, Makino K, Suzuki K, Tatarak (2002):
 Distribution and cardiovascular risk correlates of serum triglycerides in young Japanese adults. Ind Health. 40 (1): 28-35.
- Neary J, Elliott KV, and Toohey J, (2002): The causes of workplace stress and strategies for management. Paper presented at the international Symposium on Work-related Diseases: prevention and health Promotion, Linz Austria.

- Neaton JD, Blackburn H, Jacobs D et al., (1992): Serum cholesterol level and mortality findings for men secreened in the Multiple Risk factor Intervention trial. Arch Intern Med. 152: 1490-1500.
- Niaura R, Stoney CM, and Herbert PN, (2003): Lipids in psychological research: The last decade. Biological psychology, 34, 1-34.
- Niedhammer I, Goldberg M, Lecler CA, Davids, Bugel I, Landre MF (1998): Psychosocial work environment and cardiovascular risk factors in an occupational cohort in France. J Epidemiol Community Health. 52 (2): 93-100.
- NIH Consensus Conference (1999): Triglycerides, high density lipoprotein, and coronary disease. Journal of the American Medical Association, 269, 505-510.
- Onat A, Sansoy V, Uysal O. (1999): Waist circumference and waist-to-hip ratio in Turkish adults: Interrelation with other risk factors and association with cardiovascular disease Int J Cardiol; 1; 70 (1): 43-50.
- Orth -Gomer K, Chesney MA, Wenger NK, (1998): Vomen, Stress and Heart Disease. Lawrence Erlbaum Associates Inc., New Jersey USA.
- Palmar RM, Ferrige AG, Moncada S, (2000): Nitric oxide release accounts for the biological activity of endothelium derived relaxing factor. Nature (London), 327: p 524-6.
- Panagiotakos DB, Chrysohoou CH, pitsavosch et al., (2003): The association between occupational stress and the risk of developing acute coronary syndrome: The CARDIO 2000 study. Cent Eur J public Health. 11 (1): 25-30.
- Panza JA, Meredith IT, Anderson TJ, et al., (1998): Role of impaired endothelium dependent vasodilitation. Hypert. 14: p 673-5.

- Parkes CM, Bejamin B and Fitzgerold RG (1999): Broken Heart a statistical study of increased mortality among widowers. Br. Med J 31: 740-743.
- Parkes KR, (2002): Shift work and age as interactive predictors of body mass index among off shore workers. Scand J Work Environ Health. 28 (1): 64-71.
- Parkinson R, (2002): Managing health promotion in the work place; Guidelines for implementations and evaluation. Palo Alto, CA: Moyfield.
- Patton JF, Vogel JA, Damokosh AI, and Mello KP (1999): Effects continuous military operations on physical fitness capacity and physical performance. Work and Stress 3, 69-77.
- Pearlin LI, and Schooler C. (1998): The structure of coping. J Health Soc. Behav, 39: 2-21.
- Physician's weekly (2003): Cholesterol Ratio more indicative Than LDL. Physician's weekly Vol. XX, No. 27.
- Pieper C, Warren K, Pickering TG. (2003): A comparison of ambulatory blood pressure and heart rate at home and work on work and non-work days. J Hypertens 11: 177-183.
- Pihl E and Jurimae T (2001): Cardiovascular diseases risk factors in males with normal body weight and high v/aist-to-hip ration. J Cardiovasc risk 8 (5): 299-305.
- Pikering TG, (1995): Hypertension. In A.J. Goreczny (Ed.), Handbook of health rehabilitation. Psychology (pp. 219-243). New York, Plenum.
- Platt R, (2002): Coronary Heart and Modern stress. Int J Epidemiol 31: 1108-1109.

- Puddu PE, Lanti M, Menotti A, et al., (2001): Serum uric acid for short term prediction of cardiovascular disease incidence in the Gubbio Population study. Acta Cadiol., 56 (4): 243-51.
- Pudel V, (1999): Weight loss by changing eating behaviour: the target goal in therapy of obesity. Int J Obes 20: 297-99.
- Puffer JC, (2001): Exercise and Heart Disease Clin Cornerstone 3 (5): 1-7.
- Pyorala K, Laakso M and Usitupa M, (1997): Diabetes and atherosclerosis: an epidemiologic view. D.abetes Metab Rev. 3: 463-524.
- Ravussin E, Swin Burn BA, (2002): Pathophysiology of obesity. Lancet. 360: 404-8.
- Reed DM, Lacroix A, Karasek RA, Miller D, Maclean CA. (1999):
 Occupational strain and the incidence of coronary heart disease. Am J Epidemiol; 129: 495-502.
- Rexrode KM, Carey VJ, Hennekens CH, et al., (1998): Abdominal adiposity and coronary heart disease in women. JAMA 280: 1843-1848.
- Riddell LJ, Chisholm A, Williams et al., (2000): Dietary strategies for lowering homocytsteine concentrations. Am J Clin Nurt 71: 1448-54.
- Ridker PM, Genest J, and Libby P, (1999): Risk factor for Atherosclerotic disease. Heart disease. In Text Book of Cardiovascular Medicine. Braundwald Zipes libby. 6th Edition. Vol. 2: 1010-1039.
- Rifai N, (1999): Hypercholesterolaemia in children. J of Int fedr Clin Chem 1: 142.

- Rimm B, Stampfer MJ, Giovannucci E, et al., (1995): Body size and fat distribution as predictors of coronary heart disease among middle-aged and older US men. Am J Ep demiol 141, 1117-1127.
- Rodriguez F, Bautista Castano I, Bello LuJan L, Hernandez Bethencourt L, et al., (2004): Nutritional evaluation of health shift workers from the canary Islands. Nutr Hosp; 19 (5): 286-91.
- Romon M, Nuttens MC, Fievet C, et al., (1992): Increased triglyceride levels in shift workers. Am J Med, 93 (3): 259-62.
- Rose GA, Blackburn H, Gillum RF, and Prineas RJ (1997): Cardiovascular survey methods in: WHO Monograph Series No. 56. Geneva, Switzerland: World Health Organization.
- Rosenman RH, (1998): Psychological influences on the variability of plasma cholesterol. Homeostasis in Health and Disease. 34: 129-136.
- Rosenman RH, Friedman M, Straus, R, Wurm M, Kositchek R, Hahn W, and Werthessen N.T. (1994): A predictive study of coronary heart disease: appendix. Journal of the American Medical Association, 189, 1-4.
- Rosenson RS, tangney CC, Casey LC, (1999): Inhibition of proinflammatory cytokine production by pravastatin. Lancet 353: 983-984.
- Rosenstock L, (1997): Work organization research at the National Institute for Occupational Safety and Health. Journal of Occupational Health Psychology, 2 (1): 7-100.
- Rosmond R, Bjorntorp P. (2000): The hypothalamic-pituitary adrenal axis activity as a predictor of cardiovascular disease, type 2 diabetes and stroke. J Intern Med; 247 (2): 88-97.

- Ross R (2000): Hypothesis of atherogenesis. In: Hurst text book of Heart disease. Willis J Hurst 7th (ed), McGrow Hill information services complex. new York 880-8.
- Ross R, (1999): Mechanism of diseases- atherosclerosis- an inflammatory disease. N Engl J Med 340: 115-126.
- Rossen E, Buschmann M (1995): Mental illness in late life: the neurobiology of depression. Arch Psychiatr Nurs 9: 130-136.
- Rosvall M, Ostergran PO, Hedblad B, Isacsson SO, Janzon L, Berglund G, (2002): Occupational status, educational level and the prevalence of carotid atherosclersis in a general population sample of middle-aged swedish men and women: results from the malmo diet and cancer study, Arn J epidemiol 152: 334-46.
- Rudiger HW, (2004): Health problems due to night shift work and jet lag. Internist (Berl). 45 (9): 1021-5.
- Russek HI, and Zohman BL, (1998): Relative significance of heredity, diet and occupational stress in CHD of young adults. American journal of Medical Sciences, 235, 266-275.
- Ryman DH, Naitoh P, and Englund CE, (1999): Perceived exertion under conditions of sustained work and seep loss. Work & stress. 3, 5-68.
- Sales SM and House J (2001): Job dissatisfaction as a possible risk factor in coronary heart diseases. J Chronic Dis. 53: 861-873.
- Samuelson O, Wilhelmsen L, Anderson OK, et al., (1997): Cardiovascular mortality in relation to changes in blood pressure and serum cholesterol levels in treated hypertension. JAMA, 255: p. 1768.

- Sauter SL, Hurrel JJ and Cooper CL (1999): Job Control and worker Health. Wiely & Sons, chichester.
- Saw S, and Rajan U, (1997): The epidemiology of obesity: are view Ann. Acad Med Singapare; 26 (4): 489-92.
- Schalling DS (1995): Types of anxiety and types of stressors as related to personality. In spielberger, CD, and sarason IG (Eds), stress and anxiety. Washington, DC: Hemispheric publishing corporation, pp, 279-283.
- Schaubroeck J and Merritt DE (1997): Divergent effects of job control on coping with work stressors: The key role of self-efficacy. Academy of Management journal, 40: 738-754.
- Scheuch K, (1998): Psycyosoziale Faktoren im arbietsproze B and Gesundheit: Einfuhrung. Z. Ges. Hyg. 36: 403-407.
- Schnall PL, Schwartz JE, Landsbergis, PA, Warren K, Pickering TG, (1998): A Longitudinal study of job strain and ambulatory blood pressure: Results from a three- year follow-up. Psychosomatic Medicine, 60, 697-709.
- Schwartz JE, Pickering TG, and Landsbergis, PA, (1996): Work-related stress and blood pressure: Current theoretical models and considerations from a behavioral medicine perspective. Journal of occupational Health psychology, 1, 287-310.
- Scott MG (1999): Primary prevention of coronary heart disease. The Am J of Med 107: 1-5 supp.
- Selye H, (1996): Stress of life. McGraw-Hill, New York.

- Shaper AG, Wannamethee SG, Walker M, (1997): Body weight: implications for the prevention of coronary heart diseases storke and diabetes mellitus in a cohort study of middle aged men. BMJ 314:1311-1317.
- Shirom A, Melamed S, & Nir-Dotan M, (2000): The relationships among objective and subjective environmental stress levels and serum uric acid: the moderating effects of perceived control Journal of Occupational Health Psychology, 5, 374-386.
- Shirom A, Westman M, Shamai O and Carel RS, (1997): Effects of work overload and burnout on cholesterol and triglycerides levels: the moderating effects of emotional reactivity among male and female employees. Journal of Occupational Health psychology, 2 (4): 275-288.
- Siegrist J, (2002): Work stress and coronary heart disease agender (Role) specific association; Int J Epidemiol, 31: 154.
- Siegrist J, Matschinger H, Cremer P, and Seidel D, (1998): Atherogenic risk in men suffering. From occupational stress. Atherosclerosis, 89, 211-218.
- Siest G, Henny J & Schiele F (1995): Interpretation des examens de laboratoire. Karger ed P. 206-233.
- Silverstone JT, (2000): Obesity. In Bodlgy Scott, R, and Milnes Walker, R (Eds), the Medical Annual. Bristol. John Wright, pp. 72-81.
- Simon RI, (1999): Chronic posttraumatic stress disorder: A review and checklist of factors influencing prognosis. Harvard Review of psychiatry, Vol. 6, No. 6, pp. 304-312.

- Smoot SL, Gonzales JL, (1995): Cost-effective communication skills training for state hospital employees. Psychiatr serv. 46: 819-22.
- Stammer J, Wentworth D and Neartor JD (1996): For the MRFIT group: Is relation between serum cholesterol and risk to premature death from coronary artery disease continous and graded JAMA. 256: 2823-8.
- Stampi C, (1999): Polyphasic sleep strategies imporve prolonged sustained performance: a field study on 99 sailors. Work & Stress, 3, 41-55.
- Standing H, & Nicolini D, (1997): Review of Work-Related Violence. Health & Safety Executive contract Research Report 143/1997. Sudbury, Suffolk, U.K., HSE Books.
- Steenland K, Fine L, Belkic K, et al., (2000): Research findings linking work place factors to CVD outcomes. Occup Med. 15: 7-68.
- Stein EA (1997): Lipids lipoproteins and apoliproteins in: Fundimentals of clinical chemistery. Teitz NW, (ed). Philadilphia, WB. Saunders CO P, 448.
- Steinberg D, (1999): The cholesterol controversy is over. Why did it take so long. Circulation 97: 1070-8.
- Steinberg D, Lewis A and Aston (1997): LDL and atherogenesis. Circulation 95: 1062-71.
- Steptoe J, Siegrist C, Kirschbaum and Marmot M, (2004): Effort-Reward imbalance, overcommitment, and Measures of cortisol and Blood pressure over the Working Day. Psychosom Med 1, 66 (3): 323-329.

- Stoney CM, and West S (1997): Lipids personality, and stress: Mechanisms and modulators. In M. Hillbrand and R.T. Spitz (Eds.), Lipids and human behaviour (pp47-66). Washington, DC: American Psychological association.
- Stoney CM, Niaura R, Bausserman L, and Matacin M, (1999): Lipid reactivity to stress: I Comparison of chronic and acute stress responses in middle- aged airline pilots. Flealth Psychology, 18, 241-250.
- Strickland BR, (1999): Internal external control expectancies from contingency to creativity. Am psychol. 54: 1-12.
- Su CT, (2001): Association between job strain and cardiovascular risk in a population of Taiwanese white-collar workers. Jpn Circ J. 65: 509-513.
- Su CT, Yang HJ, Lin CF, et al., (2001): Arterial blood pressure and blood lipids as cardiovascular risk factors and occupational stress in Taiwan. Int J Cardiol., 81 (2-3): 131-7.
- Sutherland and Cooper (2000): Occupational stress: understanding, recognition and prevention. Series: 5. champman and Hall, London.
- Tenkanen L, Sjoblom T, Harma M. (1998): Joint effect of shift work and adverse life style factors on the risk of coronary heart disease; scand J Work Environ Health; 24 (5): 351-7.
- Tenkanen L, Sjoblom T, Kalimo R, Alikoski T, Harme M (1997): Shift work, occupation and coronary heart disease over 6 years of follow up in the Helsinki heart Study. Scand J Work Environ Health 23 (4): 257-65.

- Terry DJ, and Jimmieson NL, (1999): Work control and employee wellbeing: A decade review. In cooper & Robertson (Eds) international review of industrial and organizational psychology, 1999, Vol. 14. (pp. 95-England UK; American Ethnological press.
- Theorell T, and Emulund N, (2003): On physiological effects of positive and negative life changes- a longitudinal study. J. of Psychosomatic Research 37: 653-659.
- Theorell T, and Rahe RH, (1994): Psychological characteristics of subjects with myocardial infarction in stockholm. In Gunderson E, and Rahe RH (Eds), life stress and illness. Springfield, Illinois charles C, Thomas, PP. 90-104.
- Theorell T, and Rahe RH, (2002): A longitudinal study of 21 subjects with coronary heart diseases: life changes, catecholamine excretion and related bio-chemical reactions. Psycho Som Med 54: 505-516.
- Trevisan M, O'Leary E, Farinaro E, Jossa F, Galasso R, Clentano E, Scottoni A, Fusco G, and Panico S, (1997): Short- and long term association between uric acid and a natural disaster. Psychosomatic Medicine, 59, 109-113.
- Troxiler RG, Park WM, (1995): Hyperlipidemia in childhood in: Pediatric cardiology. Park MK (ed) Louris, Mobsy year book. P. 430.
- Tsutsumi A, Kayaka A, Ishikawa S, Gotoh T, Nago N (2003): The jichi medical school cohort study group. Job characteristics and serum lipid profile in Japanese rural workers: the Jichi Medical school cohort study. J Epidemiol; 13: 63-71.
- Tsutsumi A, (2005): Psychosocial factors and health: community and work place study. J Epidemiol. 15 (3): 65-9

- of the Jichi medical school cohort study group. Association between job strain and prevalence of hypertension: a cross sectional analysis in a Japanese working population with a wide range of occupations: the Jichi medical School cohort study. Occup Environ Med. 58: 367-373.
- Turner RC, Millns H, Neil HA, et al., (1998): For the United kingdom Prospective Diabetes study Group. Risk factors for coronary artery diseases in non-insulin dependent diabetes mellitus: UK prospective diabetes study (UKPD: 23). Br Med J 316: 823-828.
- Walker BR, Soderberg S, lindahl B et al., (2000): independent effects of obesity and cortisol in predicting cardiovascular risk factors in men and women. J intern. Med 247 (2): 198-204.
- Uris A, (2002): How mangers ease job pressures. International Management, June, 45-46.
- Vaernes RJ, Myhre G, Asa H, Homnes, T, Hansen I, and Tonder O (1999): Relationships between stress, psychological factors, health and immune levels among military aviators. Work and Stress, 5, 5-16.
- Vague P, Juhan-Vague L, Albiaud MF, et al., (1996): Correlation between blood fibrinolytic activity, plasminogen activator inhibitor level, plasma insulin level and relative body weight in normal and obese subjects. Metabolism, 2: 250-3.
- Van Italli E, (1995): Health implications of overweight united states. Ann Intern Med 103: 983-8.
- Van Raaij MTM, Ortgiesen M, Timmerman HH, Dobbe CJG, & Vanloveren, H, (1996): Time-dependent differential changes of immune function in women exposed to chronic intermittent noise. Physiology & Behavior, Vol. 60, No. 6, pp. 1527-1533.

- Vane JR Anggard EE Botting RM, (2000): Regulatory function of the vascular endothelium. N Engl J Med 323: P. 27-36.
- Verdecchia P, Schillaci G, Reboldi GP, et al., (2000): Relation between serum uric acid and risk of cardiovascular disease in essential hypertension. Hypertension, 36: 1072-1078.
- Vrijkotte T, Lorenz J, Pvan Doornen; Ecoj, C. de Geus. (2000): Effects of work stress on Ambulatory blood pressure; heart rate and heart rate variability. Hypertension; 35: 880-890.
- Wadden TA, Stunkard AJ, (2003): Social and psychological consequences of obesity Ann Intern Med 123: 1062-7.
- Wang TD, Chen WJ, Chien KL, et al., (2001): Efficacy of cholesterol levels and ratios in predicting future coronary heart disease in a Chinese population. Am J Cardiol 88 (7): 737-43.
- Wannamethee SG, Shaper HG, Walkers M (1998): Changes in physical activity, mortality and incidence of coronary heart disease in oldermen. Lancet 351: 1603- 1608.
- Wardell W.I, Hyman M, and Bahnson CB, (2003): Stress and coronary heart disease in three field studies. Journal of chronic diseases, 57, 73-84.
- Warr PB, (1997): Job features and excessive stress. In: R. Jenkins and N. Coney (eds) Prevention of Mental III Health at work. HMSO, London.
- Warshaw LJ, (2000): Managing Stress. Addison- Wesley, Reading, Mass.
- Wasserman, J Whitmer RW, Bazzarre TL, et al., (2000): Gender-specific effects of modifiable health risk factors on CHD and related expenditures. J Occup Environ Med 42 (11): 1060-90.

- Weiner H, (1997): The psychobiology of human illness. "New York, Elsevier, pp. 79-95.
- Wenger NB, Miller NE, (1994): Hypertension a major risk factor for coronary atherosclerosis. Clin Endocrinal Metabol 1: P. 603-622.
- Westin L, Carlsson R, Israelsson B, Willenheimer R, Cline C McNeil T. (1997): Quality of life in patients with ischemic heart disease: a prospective controlled study. J Intern Med. 242: 239-247.
- WHO (1995): Prevention and control of cardiovascular disease: World Health organization, Regional office for Eastern Mediterranean, Alexandria, Egypt. EMRO technical publication, Eastern Mediterranean series 22.
- WHO (2002): The work of WHO in Eastern Mediterranean Region, Annual report of the regional director 1 January 31 December 2001. World Health Organization, Regional Office for the Eastern Mediterranean, El-Zahara, Arab Mass Media, Cairo, Egypt. 700.
- Willett WC, Manson JE, stampfer MJ, et al., (1995): Weight change and coronary heart disease in women. Risk within the 'normal' weight range JAMA 273-461-465.
- William S, Michie S, Patani S, (1998): Improving the health of the NHS work force. London: the Nuffield trust.
- Williams RB, Barefoot JC, Blumenthal JM, Helms Michael J, (1997):
 Psychosocial correlates of job strain in a sample of working women. Archives of General psychiatry, 1997 June, 54 (6): 543-548.
- Windel A, Zimolong B, (1999): Group work and performance in business. Gruppendynamik- Zeitschrift Fur Angewandte Sozialpsychologie. 28 (4): pp. 333-35.

- Wing RR, Jeffery RJ, Burton LR, et al., (2002): Change in waist-to-hip ratio with weight loss and its association with change in cardiovascular risk factors. Am J Clin Nut 55: 545-51.
- Witzum JL, Mahoney EM, Branks MJ, et al., (2002): Non enzymatic glycosylation of human low density lipoprotein is more sensitive to oxidation: implications for diabetic patient. Atherosclerosis. 102: p. 63-67.
- World Health Organization (2000): The World health report 2000, health system: improving performance. Geneva.
- World Health Organization (2002): The world health report: 2002: reducing risk, promoting healthy life. Publication of the World Health organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland.
- World Health Organization (WHO) (2000): Constitution of the world health organization. In: Basic Documents (36thed). World Health organization, Geneva.
- World Health organization (WHO) (2000): Obesity: preventing and managing the global epidemic. WHO technical report series Geneva.
- Yoshimaus k, Liu Y, Kodama H, e al., (2000): Job strain, Type A behavior pattern, and the prevalence of coronary atherosclerosis in Japanese working men. J Psychoms Res 49 (1): 77-83.
- Yu LC, Packard JS, Vicary J and Mc Cool W, (1999): "Occupational stress among nurses in hospital setting". AOHN journal 37 (4): 121-128.

- Yu SF, Li KR, Yang Y, et al., (2003): The relationship between occupational stress and cardiovascular disease risk factors. Zhonghua Lao Dong Wei Sheng Zhi Ye Bing Zazhi. 21 (1): 12-5.
- Zhijie Yu, Aulikki Nissinen, Erkki Vartiainen, et al., (2000):
 Associations between socioeconomic status and cardiovascular risk factor in an urban population in China.
 Bulletin of the World Health organization; 78: 1296-1305.
- Zorbas YG, Yaroshenko YN, Federenko YF, (1996): Serum urate and cholesterol levels in endurance trained. Volunteers during acute and rigorous bed test conditions. Panminerva Medicine, 38, 223-228.
- Fahmy, S.I. and El Sherbini, A.F. (1983): Determining simple parameters for social classification for health research bulletin of H.I.P.H.