

REFERENCES

- Adams-Chapman I. and Stoll B. (2002):** Prevention of nosocomial infections in the neonatal intensive care unit. *Curr Opin Pediatr*, 14(2):157-164.
- Ahmed A. S., Chowdhury M. A., Hoque M. and Darmstadt G. (2002):** Clinical and Bacteriological Profile of Neonatal Septicemia in a Tertiary level Pediatric Hospital in Bangladesh. *Indian Pediatr*; 39:1034-1039.
- Ananthakrishnan S. and Gunasekaran D. (2009):** Etiology and risk factors for early onset neonatal sepsis. *Indian Journal of Medical Microbiology*. 27 : 279
- Andreas H. and Karin K. (2007):** Time between Membrane Rupture and Delivery and Septicemia in Term Neonates. *Obstetrics & gynecology*, 110, 612-618
- Annette H., Denise G., Ronda. L., Lisa A., (2001):** Prevalence of nosocomial infections in neonatal intensive care unit patients: Results from the first national Point-Prevalence survey. *Jour. Of Ped.*, 139, 6.
- Arafa K. (2003):** Study of nosocomial infection in neonatal intensive care unit at Benha university hospital . M. Sc. Thesis in pediatrics, Benha faculty of medicine, Benha university.
- Auriti C., Maccallini A., and Dilsio D., (2003):** risk factors for nosocomial infection in a neonatal intensive care unit . *hosp. Infct. J.* 53: 25-30

- Badrawi N., Bashir M., Iskander I., Saied D., (2005):** Neonatal infections in NICU: magnitude of the problem in NICU of Kasr El-Aini Maternity HospitaL.J Kasr El-Aini Medical; 11: 180-195.
- Berner R., Tüxen B., Clad A., (2002):** Elevated gene expression of interlukin-8 in cord blood is. a sensitive marker of neonatal infection. J Eur Pediatr, 159: 71-79.
- Bhaumik S., Ghos, S., Haldar K. , Mitra P. , and Manna B. (2000):** Risk of Early Onset Neonatal Septicemia in Babies Born to Mothers with Pre-Eclampsia. Indian Pediatr; 37: 775- 779.
- Bischoff W. , Reynolds T. , Sessler C. , Edmond M. and Wenzel R. (2000):** Handwashing compliance by health care workers: the impact of introducing an accessible, alcohol-based hand antiseptic. Arch Intern Med; 160: 1017 1021.
- Boal WL., Leiss JK., Sousa S., Lyden JT., Li J., and Jagger J. (2008):** The National Studyto prevent blood exposure in paramedics: exposure reporting. Am JInd Med;51:213-22
- Borghesi A. and Stronati M. (2008):** Strategies for the prevention of hospital-acquired infections in the neonatal intensive care unit. Journal of Hospital Infection, 68, 293-300
- Boyce J. and Pittet D. (2002):** Guideline for Hand Hygiene in Health-Care Settings. Centers for Disease Control and Prevention. Morbidity and Mortality Weekly Report (MMWR); 51:1-45.

Brodie S. B.; Sands K., Gray J. , Parker R. , Goldmann D. , Davis R. and Richardson D. (2000): Occurrence of nosocomial bloodstream infections in six neonatal intensive care units. *Pediatr Infect Dis J*; 19(1):56-65.

Buttery, J. P. (2002): Blood cultures in newborns and children: optimising an everyday test. *Arch Dis Child Fetal Neonatal Ed*; 87:25-28.

Calfee D. and Farr B. (2002): Comparison of four antiseptic preparations for skin in the prevention of contamination of percutaneously drawn blood cultures: a randomized trial. *J Clin Microbiol*; 40:1660–5.

Capretti M., Sandri F., Tridapalli E., Galletti S., Petracci E., and Faldella G.,(2008): Impact of a standardized hand hygiene program on the incidence of nosocomial infection in very low birth weight infants. *AJIC*, 36:430-435

Catalin S., Vineet B., Yiping W. , Antonette T., Margaret A., Joseph A. and Irina A. (2009): Using proteomics in perinatal and neonatal sepsis: hopes and challenges for the future. *Current Opinion in Infect Dis* 22:235–243

CDC (2003): NNIS System. National nosocomial infections surveillance (NNIS) system report data summary from January 1992 through June 2003, issued August 2003. *Am J Infect Control*;31:481-98

CDC-NNIS, (2001): Centers for Disease Control and Prevention (CDC). National Nosocomial Infections Surveillance (NNIS) System report, data summary from January 1992--June 2001, issued August 2001. *Am J Infect Control*; 6:404-21.

- Centers for Disease Control and Prevention (CDC) (2002):*** Guidelines for the Prevention of Intravascular Catheter-Related Infections. MMWR; 51(No. RR-10).
- Chen K., Tuomata R. and Cohen A. (2002):*** No increase in rates of early onset neonatal sepsis by non- group B strepto coccus or ampicillin-resistant organisms. J Obstet Gynecol (185): 854-58.
- Chiesa C., Panero A., Osborn J. (2004):*** Reliability of procalcitonin concentrations for the diagnosis of sepsis in critically ill neonates. J Clin Infect Dis (26) 664-72.
- Christopher D. (2006):*** Catheter-Related Bloodstream Infections in the Pediatric Intensive Care Unit. Pediatr Infect Dis 17:20-24
- Clark R., Powers R., White R., Bloom B., Sanchez P., and Benjamin D. (2004):*** Prevention and Treatment of Nosocomial Sepsis in the NICU. Journal of Perinatology; 24:446-453.
- Corbin D., Brian P., Daniel K. and Benjamin Jr. (2010):*** Risk factors and prevention of late-onset sepsis in premature infants. EHD, article in press
- Cordero L., Rau R., Taylor D., and Ayers L. (2004):*** Enteric gram-negative bacilli bloodstream infections: 17 years' experience in a neonatal intensive care unit. Am J Infect Control; 32(4):189-195.
- Corless, C. , Guiver, M., Borrow, R., Edwards-Jones V., Fox A., and Kaczmarek E., (2001):*** Simultaneous detection of Neisseria meningitidis,

Haemophilus influenzae, and Streptococcus pneumoniae in suspected cases of meningitis and septicemia using real-time PCR. J Clin Microbiol; 39:1553–8.

Daniel K., Kelly R., Richard A. and randall G. (2000): When to suspect fungal infection in neonates; A clinical comparison of candida albicans and candida parapsilosis fungaemia with coagulase negative staphylococcal bacteraemia. pediatrics 106: 712-18.

Deep A., Ghildiyal R., Kandian S., Shinkre N. (2004): Clinical and microbiological profile of nosocomial infections in the pediatric intensive care unit (PICU). Indian Pediatr; 41: 1238–46

Dement J., Epling C., Østbye T., Pompeii L., Hunt D. (2004): Blood and body fluid exposure risks among health care workers: results from the Duke health and safety surveillance system. Am J Ind Med;46:637-48

Doebbeling B., Vaughn T., McCoy K., Beekmann S., Woolson R., Ferguson K. (2003): Percutaneous injury, blood exposure, and adherence to standard precautions: are hospital-based health care providers still at risk? Clin Infect Dis;37:1006-13

Duerink D., Roeshadi D., Wahjono H. (2006). Surveillance of healthcare-associated infections in Indonesian hospitals. J Hosp Infect; 62: 219–29

Dutta S. and Narang A. (2003): Drug resistant neonatal candida tropicalis septicemia. Does it cause diaphragmatic hernia?. J Indian Pediatrics; 40: 261-264.

Eggimann P., and Pittet D.(2001): Infection Control in the ICU. Chest; 120:2059-2093.

El-Fiky, O., El-Mahdy, M., Shebl A., Abdel-Maksoud Y. and Abdel-Rhahman S. (2003): Epidemiological study of nosocomial infection of neonates admitted to Benha university neonatal intensive care unite. Benha M J; 20:539-556.

El-Sadik A. (2005): Comparative study between conventional blood culture method versus polymerase chain reaction assay for diagnosis of neonatal septicemia. M. Sc. Thesis in microbiology & immunology, Zagazig university.

El-Sayed, H. (2002): Study of some diagnostic methods for early detection of neonatal sepsis. MD Thesis (Clinical and Chemical Pathology).Zagazig Faculty of Medicine, Zagazig University.

Flidel-Rimon O., Friedman S., Lev E., Juster-Reicher A., Amitay M., Shinwell E. (2004): Early enteral feeding and nosocomial sepsis in very low birthweight infants. Arch Dis Child Fetal Neonatal Ed 2004;89:F289-F292.

Foca, M., Jakob, K., Whittier S., Della, L. , Factor S., Rubenstein D., and Saiman L. (2000): Endemic Pseudomonas aeruginosa infection in a neonatal intensive care unit. N Engl J Med; 343(10):695-700.

Friedman N. , and Sexton D. (2001): General principles of infection control. In: Rose, B. D. UpToDate. Wellesley, MA: UpToDate.

- Garland, J. , Henrickson K., and Maki D. , (2002):** The 2002 Hospital Infection Control Practices Advisory Committee Centers for Disease Control and Prevention guideline for prevention of intravascular device related infection. *Pediatrics*; 110(5):1009-1013.
- Gerdes J. (2004):** Diagnosis and management of infections in the neonate. *Pediatr Clin North Am* 2004;51:939-59
- Giannacopoulou C., Hatzidaki E., Kokori H., Giannacopoulos K., Margari K. , Manoura A., and Sbirakis S. (2000):** Evaluation of the micro-erythrocyte sedimentation rate in relation to hemoglobin and hematocrit in newborn infections. *Haematologia (Budap)*; 30(3): 215- 220.
- Giles M., Harwood H., Gosling D., (2006):** What is the best screening method to detect *Serratia marcescens* colonization during an outbreak in a neonatal intensive care nursery? *J Hosp Infect*; 62: 349–52
- Girou, E., Loyeau S., Legrand P., Oppein F., and Brun-Buisson C., (2002):** Efficacy of handrubbing with alcohol based solution versus standard handwashing with antiseptic soap: randomised clinical trial. *BMJ*; 325: 362.
- Godambe S., Shah P., and Shah V. (2005):** Breast milk as a cause of late onset neonatal sepsis. *Pediatr Infect Dis J*; 24 (4): 38 1-2.
- Gordts B. (2005):** Models for the organization of hospital infection control and prevention programmes. *Clin Microbiol Infect*;11(1 1):19-23

- Gotoff, S. (2000):** The fetus and the newborn. In: Behrman, R. E., Kliegman, R. M. and Jenson, H. B. (eds.). Nelson Text Book of Pediatrics. 16th ed. Philadelphia: W.B. Saunders. Pp: 425-428.
- Gray J. (2004):** A 7-year study of bloodstream infections in an English children's hospital. Eur J Pediatr; 163: 530–35
- Guerina J., (2004):**bacterial and fungal infections In: Cloherty J, Stark A (Eds). Manual of Neonatal Care .5 th d.271-300
- Guida J. , Kunig A. , Leef K., McKenzie S. , and Paul D. (2003):** Platelet Count and Sepsis in Very Low Birth Weight Neonates: Is There an Organism-Specific Response. Pediatrics; 111(6):1411-1415.
- Harbath S., Pittet D., Grady L., and Goldmann D. (2001):** Compliance with hand hygiene practice in pediatric intensive care. Pediatr Crit Care Med; 2 :311 –314.
- Hayes R. , Trick W. , Vernon M. ,; Nathan C., Peterson B. , Segreti J., Pur S. , Schmitt B. , Rice T. , Welbel S. and Weinstein R. (2001):** Ring use as a risk factor (RF) for hand colonization in a surgical intensive care unit (SICU) [Abstract K-1333]. In: Program and abstracts of the 41st Interscience Conference on Antimicrobial Agents and Chemotherapy. Washington, DC: American Society for Microbiology, 2001.
- Harris J. and Goldmann D. (2001):** Infections acquired in the nursery: Epidemiology and control. in Infectious Diseases of the fetus and Newborn Infant 5th ed: 1371-418. Edited by Remington J.S. and Klein J.O. By W.B. Saunders.

Hobson R. (2003): The global epidemiology of invasive Candida infectious—is the tide turning? J Hosp Infect;55:159-68.

Horbar J., Rogowski J., Plsek P. , Delmore P., Edwards W. , Hocker J., Kantak A. , Lewallen P., Lewis W., Lewit E., McCarroll C. , Mujsce D., Payne N. , Shiono P., Soll R. , Leahy K. and Carpenter J. (2001): Collaborative Quality Improvement for Neonatal Intensive Care. Pediatrics; 107: 14 - 22.

Hudson H. (2003): Stethoscopes and infection control: a study into the use of stethoscopes in a paediatric ward and their possible contamination. Child Health Care; 7: 142–143

Ihn S., and Sunmi P.(2009): Use of hands-free technique among operating room nurses in the Republic of Korea Association for Professionals in Infection Control and Epidemiology, Inc. Am J Infect Control ;37:131-5.

Inweregbu K. (2005): Nosocomial infections. Continuing Education in Anaesthesia, Critical Care & Pain; 5(1):14-17.

Jain A., Roy I., Gupta M. , Kumar M., and Agarwal S. (2003): Prevalence of extended-spectrum beta-lactamase-producing Gram-negative bacteria in septicemic neonates in a tertiary care hospital. J Med Microbiol; 52: 421-425.

Jarvis W. (2004): Controlling healthcare-associated infections: the role of infection control and antimicrobial use practices. Semin Pediatr Infect Dis; 15:30–40

- Jeannie P., Janet H., Lisa S., Elaine L., (2008):*** Impact of Staffing on Bloodstream Infections in the Neonatal Intensive Care Unit. *arch pediatr adolesc med* 160, 832-836
- Jones, R. D.; Jampani H. B.; Newman J. L. and Lee A. S. (2000):*** Triclosan: a review of effectiveness and safety in health care settings. *Am J Infect Control*; 28:184-96.
- Joshi S. G., Ghole V. , and Niphadkar K.. (2000):*** Neonatal gram-negative bacteremia. *Indian Pediatr*; 67:27-32.
- Kampf G. and Kramer A. (2004):*** Epidemiologic background of hand hygiene and evaluation of the most important agents for scrubs and rubs. *Clin Microbiol Rev*; 17:863-893.
- Karam G. and Heffner J. (2000):*** Emerging Issues in Antibiotic Resistance in Blood-borne Infections. *Am J Respir Crit Care Med*; 162:1610-1616.
- Karam N. (2002):*** Study of intercellular adhesion molecule-1 (ICAM-1) and E-selectin in neonatal infection. M.D. thesis in pediatrics. Zagazig University.
- Karen M. (2004):*** Bacterial and fungal infections. In: Cioherty J Stark A Eichenwald (Eds) *manual of neonatal care*. 4th ed. Phyladeiphia, 287-313.

Kaufman D. and Fairchild K. (2004): Clinical Microbiology of Bacterial and Fungal Sepsis in Very-Low-Birth-Weight Infants. Clin Microbiol Rev; 17; 638–680.

Kawagoe J. , Segre C. , Pereira C. , Cardoso M. , Silva C. , and Fukushima J. (2001): Risk factors for nosocomial infections in critically ill newborns: A 5-year prospective cohort study. Am J Infect Control; 29(2): 109-114.

Kennedy K. , Tyson J. and Chamnanvanikij S. (2000): Early versus delayed initiation of progressive enteral feedings for parenterally fed low birth weight or preterm infants. Cochrane Database Syst Rev; (2):CD001970.

Kenneth E. , Gray J. , Parker R. , Goldmann D. , Davis R. and Richardson D. (2000) : Occurrence of nosocomial blood stream infections in six neonatal intensive care units. Pediatrics. 19: 59-62.

Kilbride H., Wirtschafter D., Powers R., Sheehan M. (2003): Implementation of evidence-based potentially better practices to decrease nosocomial infections. Pediatrics; 111: e519–e533

Kilbride H., Powers R., Wirtschafter D., Sheehan M., Charsha D., LaCorte M., Finer N. and Goldmann D. (2003): Evaluation and Development of Potentially Better Practices to Prevent Neonatal Nosocomial Bacteremia. Pediatrics; 111;504-518.

Kilgore M., Ghosh K., Beavers C., Wong D., Hymel P., Brossette S. (2008): The costs of nosocomial infections. Med Care ;46:101-4

- Klara M., Zerr D., Pittet D., (2008):** Infection control in paediatrics, *Lancet Infect Dis* ; 8: 19–31
- Klein J., and Remington J., (2003):** Bacterial sepsis and meningitis. In: Klein and Remington (Eds) . Infectious diseases of the fetus and newborn infants. WB Saunders 7th ed (4): 538-890
- Kollef M., Morrow L., Baughman R., Craven D., McGowen J., Micek S., (2008):** Health care-associated pneumonia (HCAP): a critical appraisal to improve identification, management, and outcomes. Proceedings of the HCAP Summit. *Clin Infect Dis*;46(Suppl):S296-334
- Kollef M., Shorr A., Tabak Y., Gupta V., Lui L., Johannes R. (2005):** Epidemiology and outcomes of health care-associated pneumonia: results from a large US database of culture-positive pneumonia. *Chest*;128: 3854-62.
- Koti M., Zllang G., Daggard G., (2003):** Late-onset and recurrent neonatal Group B streptococcal disease associated with breast-milk transmission. *Pediatr Dev Pathol*;6(3):25 1-6.
- Kwak Y., Lee O., Kim Y., Kim K., Park S., (2010):** Risk factors for device-associated infection related to organizational characteristics of intensive care units: findings from the Korean Nosocomial Infections Surveillance System. *Journal of Hospital Infection. J Hosp Infect.*;75(3):195-9.

- Lam B. , Lee J. and Lau Y. , (2004):** Hand Hygiene Practices in a Neonatal Intensive Care Unit: a Multimodal Intervention and Impact on Nosocomial Infection. *Pediatrics*; 114:e565-e571.
- Larson E. (2001):** Hygiene of the skin: when is clean too clean? *Emerg Infect Dis*;7:225-30
- Larson E., Lin S., Gomez-Pichardo C., Della-Latta P. (2004):** Effect of antibacterial home cleaning and handwashing products on infectious disease symptoms. A randomized, double-blind trial. *Ann Intern Med*;140:321-9
- Li'lian A., Lizandra F. and Paulo P. (2009):** Changes in hands microbiota associated with skin damage because of hand hygiene procedures on the health care workers. Association for Professionals in Infection Control and Epidemiology, Inc. *Am J Infect Control*; 37:155-9
- Lin F. , Weisman L. , Troendle J., and Adams K. (2003):** Prematurity is the major risk factor for late-onset group B streptococcus disease. *J Infect Dis*; 188:267-271.
- Linhart Y., Bashiri, A., Maymon E., Shoham-Vardi I., Furman B., Vardi H. and Mazor M. (2000):** Congenital anomalies are an independent risk factor for neonatal morbidity and perinatal mortality in preterm birth. *Eur J Obstet Gynecol Reprod Biol*; 90(1): 43- 49.
- Long S., and Stevenson D. (2005):** Reducing Candida infections during neonatal intensive care: management choices, infection control, and fluconazole prophylaxis. *J Pediatr*; 147: 135–141

Lopez S. , Coto C. , and Fernandez C. , (2002): Neonatal sepsis of nosocomial origin: an epidemiological study from the "Grupo de Hospitales Castrillo". J Perinat Med; 30(2):149-157.

Lossius I., Inger E., Lorentz M. Arild R., Tore G., Sverre M., Hallvard R., (2005): Septicemia in the First Week of Life in a Norwegian National Cohort of Extremely Premature Infants. Pediatrics;115;262-268

Lott J. (2003): Neonatal bacterial sepsis. Crit Care Nurs Clin North Am; 15(1): 35-46.

Ludo M., Nienke K., Jozef J., Hilde J., Margaretha M. (2004): Decreasing incidence of neonatal nosocomial bloodstream infections in a neonatal intensive care unit: antenatal corticosteroid treatment an innocent bystander?. Eur J Pediatr 163: 151–157

Lumbiganon P., Thinkhamrop J., Thinkhamrop B. and Tolosai E. (2005): Vaginal chlorhexidine during labour for preventing maternal and neonatal infections (excluding Group B Streptococcal and HIV) (Cochrane Review). The Cochrane Library; Issue 2. Chichester, UK: John Wiley & Sons, Ltd.

MacCannell D., Louie T., Gregson D., Laverdiere M., Labbe A., Henwick S, (2006): Molecular analysis of *Clostridium difficile* PCR ribotype 027 isolates from eastern and western Canada. J Clin Microbiol;44:2147-52

Macharashvili N., Kourbatova E., Butsashvili M., Tsertsvadze T., McNutt L., Leonard M. (2009): Etiology of neonatal blood stream infections in Tbilisi, Republic of Georgia. *Int J Infect Dis.* official publication of the International Society for Infectious Diseases ; 13(4):499-505.

Macrae B., Shannon K., Rayner D., Kaiser A., Hoffman P., and French G. (2001): A simultaneous outbreak on a neonatal unit of two strains of multiply antibiotic resistant *Klebsiella pneumoniae* controllable only by ward closure. *J Hosp Infect*; 49(3):183-92.

Maki, D. G. (1994): Infections caused by intravascular devices used for infusion therapy: pathogenesis, prevention, and management. In: Bisno, A. L. and Waldvogel, F.A. (eds.). *Infections associated with indwelling medical devices.* 2nd ed. Washington, DC: American Society for Microbiology Press. Pp: 155-212.

Maury E., Alzieu M., Baudel J., Haram N., Barbut F., Guidet B. (2000): Availability of an alcohol solution can improve hand disinfection compliance in an intensive care unit. *Am J Respir Crit Care Med*; 162:324-7.

McNeil S. , Foster C. , Hedderwick S., and Kauffman C. (2001): Effect of hand cleansing with antimicrobial soap or alcohol-based gel on microbial colonization of artificial fingernails worn by health care workers. *Clin Infect Dis*; 32(3):367-372.

Mehall J. , Kite C. , Saltzman D. , Wallett T., Jackson R. and Smith S. (2002): Prospective study of the incidence and complications of bacterial

contamination of enteral feeding in neonates. J Pediatr Surg; 37:1177-1182.

Miho S., Yuichi I., Hiroyoshi K., Takashi J., Hiroe K. (2009): Factors affecting performance of hospital infection control in Japan. Association for Professionals in Infection Control and Epidemiology, Inc., Am J Infect Control;37:136-42.

Moolenaar R., Crutcher J., San-Joaquin V., Sewell L., Hutwagner L., Carson L., Robison D., Smithee L. and Jarvis W. (2000): A prolonged outbreak of *Pseudomonas aeruginosa* in a neonatal intensive care unit: did staff fingernails play a role in disease transmission? Infect Control Hosp Epidemiol; 21(2):80-85.

Moreira M.(2005):Controversies about the mangment of invasive fungal infection in very low birth weight infants.J Pediatr (Rio); 81 (1 Suppl): S 52- 8.

Morrison J. (2004): for Health Canada, Nosocomial and Occupational Infections Section. Development of a resource model for infection prevention and control programs in acute, long term, and home care settings: conference proceedings of the Infection Prevention and Control. Alliance. Am J Infect Control;32:2-6

Muhlemann K., Franzini C., and Aebi C. (2004) : Prevalence of nosocomial infections in Swiss children's hospitals. Infect Control Hosp Epidemiol ; 25: 765–71.

- Musoke R., and Revathi G. (2000):** Emergence of multidrug-resistant gram-negative organisms in a neonatal unit and the therapeutic implications. *J Trop Pediatr*; 46:86-91.
- Muto C., Giannetta E., Durbin L., Simonton B., Farr B. (2002):** Cost-effectiveness of perirectal surveillance cultures for controlling vancomycin-resistant *Enterococcus*. *Infect Control Hosp Epidemiol*;23:429-35
- Nagata E., Brito A. and Matsuo T. (2002):** Nosocomial infections in a neonatal intensive care unit: incidence and risk factors. *Am J Infect Control*; 30(1):26-31.
- Nazemi K. J.; Buescher E.S.; Kelly Jr. R. E. and Karlowicz M. G. (2003):** Central venous catheter removal versus in situ treatment in neonates with enterobacteriaceae bacteremia. *Pediatrics*; 111(3):e269-e274.
- Ng P. (2004):** Diagnostic markers of infection in neonates. *Arch Dis Child Fetal Neonatal Ed*; 89:F229.
- Nicolle L. (2001):** Infection control programmes to contain antimicrobial resistance. WHO; WHO/CDS/CSR/DRS/2001/7.
http://www.who.int/entity/csr/resources/publications/drugresist/en/infection_control.pdf
- Nizet V., Ferrieri P., and Rubens C. , (2000):** Molecular pathogenesis of group B streptococcal disease in newborns, p. 180-221. In: D. Stevens (ed.), *Streptococcal infections: clinical aspects, microbiology, and molecular pathogenesis*. Oxford University Press, New York, N.Y.

Nosocomial Infection National Surveillance Service (NINSS) (2002):

Surveillance of Hospital-Acquired Bacteraemia in English Hospitals
1997-2002. [www.hpa.org.uk/infections/
publications/ninns/hosacq_HAB_2002.pdf](http://www.hpa.org.uk/infections/publications/ninns/hosacq_HAB_2002.pdf)

O'Boyle C., Jackson M., Henly S. (2002): Staffing requirements for
infection control programs in US health care facilities: Delphi project.
Am J Infect Control; 30:321-33.

O'Boyle C., Henly S., Larson E. (2001): Understanding adherence to hand
hygiene recommendations: the theory of planned behavior. *Am J Infect
Control*;29:352-60.

***O'Grady N. , Alexander M., Dellinger E. , Gerberding J. , Heard S. ,
Maki D. , Masur H., McCormick R. , Mermel L. , Pearson M. , Raad
I. , Randolph A. and Weinstein R. (2002):*** Guidelines for the prevention
of intravascular catheter-related infections. Centers for Disease Control
and Prevention. *MMWR Recomm Rep*; 51(RR-10):1-29.

Oh H., Chung H., Kim J., and Cho S. (2006): National survey of the status
of infection surveillance and control programs in acute care hospitals with
more than 300 beds in the Republic of Korea. *Am J Infect Control*;34:223-
33

Omar N. , El Sallab S., El Shahat T. and Shokeir M.(2006): Intravascular
Catheter Related Infections in Neonatal Care Unit at Mansoura University
Children's Hospital . 6th Int. Conf. of the Hospital Infection Society

- Opilla M., Kirby D., Edmond M. (2007):** Use of ethanol lock therapy to reduce the incidence of catheter-related bloodstream infections in home parenteral nutrition patients. *J Parent Ent Nutr*;31: 302-5
- Osrin D., Vergnano S. and Costello A. (2004):** Serious bacterial infections in newborn infants in developing countries. *Curr Opin Infect Dis*; 17(3): 217-224.
- Pai M., Pendland S. and Danziger L. H. (2001):** Antimicrobial-coated/bonded and -impregnated intravascular catheters. *Ann Pharmacother*; 35(10):1255-1263.
- Parry G., Tucker J., and Tarnow-Mordi W. (2005):** Relationship between probable nosocomial bacteraemia and organisational and structural factors in UK neonatal intensive care units. *Qual Saf Health Care*; 14: 264–69
- Patel A., White C., Mary R., Woolard C. and Perz F. (2009):** Infection Control Practices in Assisted Living Facilities: A Response to Hepatitis B Virus Infection Outbreaks. *Infection Control & Hospital Epidemiology*. 30(3):209-214.
- Payne N. , Carpenter J. , Badger G. , Horbar J. and Rogowski J. (2004):** Marginal Increase in Cost And Excess Length of Stay Associated With Nosocomial Bloodstream Infections in Surviving Very Low Birth Weight Infants. *Pediatrics*; 114(2):348-355.

- Pellowe C. , Macqueen S., and Coe L. (2004):** Preventing healthcare associated infections in the neonatal unit: The use of evidence-based infection control guidelines. *Journal of neonatal nursing*; 10(5): 171-175.
- Pierce C. , Wade A. and Mok Q. (2000):** Heparin-bonded central venous lines reduce thrombotic and infective complications in critically ill children. *Intensive Care Med*; 26(7):967-972.
- Pittet D. (2000):** Improving compliance with hand hygiene in hospitals. *Infect Control Hosp Epidemiol*; 21:381-6.
- Pittet D. (2001):** Compliance with hand disinfection and its impact on hospital acquired infections. *J Hosp Infect*;48:40-6
- Polak J. , Ringler N. and Daugherty B. (2004):** Unit Based Procedures: Impact on the Incidence of Nosocomial Infections in the Newborn Intensive Care Unit. *Newborn and Infant Nursing Reviews*; 4(1): 38-45.
- Pronovost P., Needham D., Berenholtz S., Sinopoli D., Fraser V., and Warren D. (2006):** An intervention to decrease catheter-related bloodstream infections in the ICU. *N Engl J Med*;355:2725-32
- Raimundo O., Heussler H., Bruhn J. , Suntrarachun S., Kelly N., Deighton M. and Garland S. (2002):** Molecular epidemiology of coagulase-negative staphylococcal bacteraemia in a newborn intensive care unit. *J Hosp Infect*; 51:33-42.

- Rocha V., Benjamin A., Procianoy R. (2001):** The effect of recombinant human erythropoietin on the treatment of anemia of prematurity J Pediatr (Rio J); 77(2):75-83.
- Rønnestad A., Abrahamsen T. , Medbø S., Reigstad H., Lossius K., Kaaresen P. , Egeland T., Engelund I. , Irgens L. and Markestad T. (2005):** Late-Onset Septicemia in a Norwegian National Cohort of Extremely Premature Infants Receiving Very Early Full Human Milk Feeding. Pediatrics; 115;269-276.
- Rotter M. (1999):** Hand washing and hand disinfection [Chapter 87]. In: Mayhall C. G. (ed.) Hospital epidemiology and infection control. 2nd ed. Philadelphia PA: Lippincott Williams & Wilkins.
- Rubin L., Sánchez P., Siegel J., Levine G., Saiman L., (2002):** Evaluation and Treatment of Neonates With Suspected Late-Onset Sepsis: A Survey of Neonatologists' Practices. Pediatrics; 110; 42
- Saiman L. (2002):** Risk factors for hospital-acquired infections in the neonatal intensive care unit. Semin Perinatol; 26(5):315-21.
- Saiman L., Ludington E., Pfaller M., Rangel-Frausto S., Wiblin R. , Dawson J., Blumberg H. , Patterson J. , Rinaldi M., Edwards J., Wenzel R. and Jarvis W. (2000):** Risk factors for candidemia in neonatal intensive care unit patients. Pediatr Infect Dis J; 19:319-324.

- Sakr M., El-Kholy A., El-Rholy O., (1998):** Comparison between conventional blood culture and serum tumor necrosis factor-alpha (TNF-cc) for early diagnosis and follow up to newborns with clinical diagnosis of septicemia. J Gaz Ped Ass. 44(1): 33-9.
- Saqui O., Raman M., Chang A., and Allard J. (2007):** Catheter-related infections in a Canadian home total parenteral nutrition program: a prospective study using US Centers for Disease Control and Prevention criteria. J Assoc Vascular Access; 12:85-8
- Sartor C., Jacomo V., Duvivier C., Tissot-Dupont H., Sambuc R. and Drancourt M. (2000):** Nosocomial *Serratia marcescens* infections associated with extrinsic contamination of a liquid nonmedicated soap. Infect Control Hosp Epidemiol; 21:196-9.
- Sattar S., Tetro J., Springthorpe V. and Giulivi A. (2001):** Preventing the spread of hepatitis B and C viruses: where are germicides relevant? Am J Infect Control; 29:187-97.
- Schelonka R., Scruggs S., Nichols K., Dimmitt R. and Carlo W. (2006):** Sustained reductions in neonatal nosocomial infection rates following a comprehensive infection control intervention Journal of Perinatology 26 176–179
- Schrag S., Zywicki S., Farley M., Reingold A., Harrison L., Lefkowitz L., Hadler J., Danila R., Cieslak P., and Schuchat A. (2000):** Group B streptococcal disease in the era of intrapartum antibiotic prophylaxis. N Engl J Med; 342: 15-20.

Schrag S., Gorwitz R., Fultz-Butts K. and Schuchat A. (2002): Prevention of perinatal group B streptococcal disease. Revised guidelines from CDC. *Morb Mortal Wkly Rep Recomm Rep*; 51:1-22.

Schuchat A., Schrag S., Gorwitz R., Fultz-Butts K. (2002): Prevention of perinatal group B streptococcal disease. Revised guidelines from CDC. *Morb Mortal Wkly Rap Recomm Rep*; 51:1-22.

Scott H. (2002): Nurses are not washing their hands well enough. *Br J Nurs*; 9 :2264.

Seale C., Michael M., James A. (2009): Maternal and early onset neonatal bacterial sepsis: burden and strategies for prevention in sub-Saharan Africa. *Lancet Infect Dis*9: 428–38

Seaward P. (2004): Evaluation of infection born to patients with premature rupture of membranes at term. *J Obstet Gyne Col*;(179):35-9

Sekimoto M., Imanaka Y., Kobayashi H., Okubo T., Kizu J., Kobuse H., (2008): Impact of hospital accreditation on infection control programs in teaching hospitals in Japan. *Am J Infect Control*;36:212-9

Shang S., Chen Z., and Yu X. (2001): Detection of bacterial DNA by PCR and reverse hybridization in the 16S rRNA gene with particular reference to neonatal septicemia. *Acta Paediatr*; 90(2): 179- 183.

Shapey I., Foster M., Whitehouse T., Jumaa P., and Bion J. (2009): Central venous catheter-related bloodstream infections: improving post-insertion catheter care. *Journal of Hospital Infection*. 71 117-122

- Shorr A., Tabak Y., Killian A., Gupta V., Lui L., and Kollef M. (2006):** Healthcare associated blood stream infection: a distinct entity? Insights from a large US database. Crit Care Med;34:2588-95
- Siegel J., Rhinehart E., Jackson M., and Chiarello L. (2007):** for the Healthcare Infection Control Practices Advisory Committee. 2007 guidelines for isolation precautions: preventing transmission of infection agents in healthcare settings June 2007. Available from: <http://www.cdc.gov/ncidod/dhqp/pdf/isoation2007.pdf>. Accessed June 2008
- Singh N., Patel K., Leger M., Short B., Sprague B. M., Kalu N. and Campos J. (2002):** Risk of resistant infections with Enterobacteriaceae in hospitalized neonates. Pediatr Infect Dis J; 21:1029-1033
- Snydman D. (2003):** Shifting patterns in the epidemiology of nosocomial Candida infections. Chest;123:500-3
- Sohn A. , Garrett D. , Sinkowitz-Cochran R., Grohskopf L. , Levine G., Stover B., Siegel J. and Jarvis W. (2001):** Prevalence of nosocomial infections in neonatal intensive care units: results from the first national point-prevalence survey. J Pediatr; 139(6):821-827.
- Srivastava S., and Shetty N. (2007):** Healthcare-associated infections in neonatal units: lessons from contrasting worlds J Hosp Infect 65 292-306

- Steiner L., Bizzarro M., Ehrenkranz R., and Gallagher PG. (2007):** decline in the frequency of neonatal exchange transfusions and its effect on exchange-related morbidity and mortality. *Pediatrics* ;120:27-32
- Steinmann J. (2001):** Some principles of virucidal testing. *J Hosp Infect*; 48(supplA):S15-S17.
- Stevenson K., MurphyC., SamoreM., Hannah E., MooreJ., Barbera J. (2004):** Assessing the status of infection control programs in small rural hospitals in the western United States. *Am J Infect Control*;32:255-61
- Stoll B. (2004):** Infections of the neonatal infant. In: Behrman R. E.; Kleigman R. M. and Jenson H. B. (eds.). *Nelson Text Book of Pediatrics*. 17th ed. Philadelphia: W.B. Saunders.Pp: 623-640.
- Stoll B. , Hansen N., Fanaroff A. , Wright L. , Carlo W. , Ehrenkranz R. , Lemons J. , Donovan E., Stark A., Tyson J. , Oh W., Bauer C. , Korones S. , Shankaran S., Laptook A. , Stevenson D. , Papile L. and Poole W. (2002):** Changes in pathogens causing early-onset sepsis in very-low- birth-weight infants. *N Engl J Med*; 347:240-247.
- Stoll B. , Hansen N., Fanaroff A., Wright L., Carlo W. , Ehrenkranz R. , Lemons J., Donovan E., Stark A., Tyson J., Oh W., Bauer C., Korones S., Shankaran S., Laptook A., Stevenson D. , Papile L. and Poole W. (2002b):**Late-onset sepsis in very low birth weight neonates: the experience of the NICHD Neonatal Research Network.*Pediatrics*; 110:285-291.

- Stringer B., Haines T., Goldsmith C., Blythe J., and Harris K. (2006):** Perioperative use of the hands-free technique: a semistructured interview study. *AORN J*;84:233-48
- Stringer B., Infante-Rivard C., and Hanley J. (2002):** Effectiveness of the hands-free technique in reducing operating theatre injuries. *Occup Environ Med*;59:703-7
- Trick W., Vernon M., Hayes R., Nathan C., Rice T. and Peterson B. (2003):** .Impact of ring wearing on hand contamination and comparison of handhygiene agents in a hospital. *Clin Infect Dis*; 36: 1383–1390
- Turnberg W. , Daniell W., Simpson T. , Van Buren J., Seixas N., Lipkin E. and Duchin J. (2009):** Personal Healthcare Worker (HCW) and Work-Site Characteristics That Affect HCWs' Use of Respiratory-Infection Control Measures in Ambulatory Healthcare Settings. *Infection Control & Hospital Epidemiology*. 30(1):47-52.
- Urrea M., Iriondo M., and Thio M. (2003):** A prospective incidence study of nosocomial infections in a neonatal care unit. *Am J Infect Control*; 31: 505–07
- Urrea A., Pons O., Krauel V., Latorre O., Martí'n M., and Campins M. (2007):** Nosocomial infections in paediatric and neonatal intensive care units. *Journal of Infection*. 54 212-220
- Venberghé A., Laterre P., Goenen M., Reynaert M., Wittebole X. and Simon A. (2002) :** Surveillance of hospital-acquired infections in an

intensive care department: the benefit of the full-time presence of an infection control nurse. J Hosp Infect;52:56-9

Vergnano S., Sharland M., Kazembe P., Mwansambo C. and Heath P. (2005): Neonatal sepsis: an international perspective. Arch Dis Child Fetal Neonatal Ed; 90:F220-FF224.

Watt J., Schuchat A., and Erickson K. (2001): Group B streptococcal disease prevention practices of obstetriciangynecologists. J Obstet Gynecol (98): 7-13.

Weinstein J. (2001): Isolation guidelines for hospitals. In: Rose B. D. (Ed.). UpToDate. Wellesley MA: UpToDate.

White R. (2002): Recommended standards for newborn ICU design - Report of the Fifth Consensus Conference on Newborn ICU Design. January 2002 Clearwater Beach Florida.
<http://www.nd.edu/~kkolberg/DesignStandards.htm>

WHO (2004): Practical Guidelines for Infection Control in Health Care Facilities WPRO Regional Publication; Publication No. 41: 10-18

WHO (2008): INFECTION control
[www.wpro.who.int/health_topics/infection control/ overview.htm](http://www.wpro.who.int/health_topics/infection_control/overview.htm)

Winnefeld M., Richard M. A., Drancourt M. and Grobb J. (2000): Skin tolerance and effectiveness of two hand decontamination procedures in every day hospital use. Br J Dermatol; 143:546-50.

- Wolff M. , Schmitt J., Rahaus M. and König A. (2001):** Hepatitis A virus: a test method for virucidal activity. *J Hosp Infect*; 48(suppl A):S18—S22.
- Wunderink R. (2006):** Healthcare-associated bacteremia: stirring the mud. *Crit Care Med*; 34:2685-6.
- Yadav A. , Wilson C., Prasad P. and Menon P. (2005):** Polymerase Chain Reaction in Rapid Diagnosis of Neonatal Sepsis. *Indian Pediatr*; 42; 681-685.
- Yuh L. (2009):** The Changing Characteristics of Neonatal Sepsis in the Neonatal Intensive Care Unit: A Never-ending Challenge. *Pediatr Neonatol* 2009;50(3):83–84
- Zafar A., Sylvester L., Beidas S. (2002):** *Pseudomonas aeruginosa* infections in a neonatal intensive care unit. *Am J Infect Control*; 30:425-9
- Zaidi A., Huskins W., Thaver D., Bhutta Z., Abbas Z. and Goldmann D. (2005):** Hospital-acquired neonatal infections in developing countries. *Lancet*; 365:1175—1188
- Zaki M. and Hesham S. (2009):** Evaluation of Microbiologic and Hematologic Parameters and E-Selectin as Early Predictors for Outcome of Neonatal Sepsis. *Arch Pathol Lab Med*—133 1291-1296
- Zoutman D., Ford B., Bryce E., Gourdeau M., He´bert G., Henderson E., (2003):** for the Canadian Hospital Epidemiology Committee Canadian Nosocomial Infection Surveillance Program Health Canada. The state of infection surveillance and control in Canadian acute care hospitals. *Am J Infect Control*; 31:266-72