

References

A

Abraham J. and Zujewski A.J., (2001): Breast cancer. In: Abraham J, Allegra CJ. The Bethesda Handbook of Clinical Oncology. New York, NY: Lippincott, Williams & Wilkins; 127.

Acs G., (2007): Cancer Facts and Figures. Breast cancer: Statistics on Incidence, Survival, and Screening. Am. J. Surg. Pathol.; 34(6):132-40.

Acs G., Simpson J.F., Bleiweiss I.J. et al, (2003): Microglandular adenosis with transition into adenoid cystic carcinoma of the breast. Am. J. Surg.Pathol.; 27(8):1052-60.

Adsay N.V., Merati K., Nassar H. et al, (2003): Pathogenesis of colloid (pure mucinous) carcinoma of exocrine organs: coupling of gel-forming mucin (muc2) production with altered cell polarity and abnormal cell-stroma interaction may be the key factor in the morphogenesis and indolent behavior of colloid carcinoma in the breast and pancreas. Am. J. Surg. Pathol; 27(5):571-8.

Agoff S.N. and Lawton T.J., (2004): Papillary Lesions of the Breast with and Without Atypical Ductal Hyperplasia. Am J Clin Pathol; 122: 440-443

Amit N., Dinker R.P., Jagdish S. et al, (2004): Alveolar soft part sarcoma: Aunique tumor with disputed histogenesis. Indian Journal of Cancer; 40 (2): 82-83.

American Cancer Society, (2003): ACS News Center web site <http://www.cancer.org/downloads/STT/CAFF2003PWSecured.pdf>.

Andrson B.O., Daling J.R. and Moe R., (2002): Changing incidence of lobular carcinoma in situ of the breast. Breast Cancer Res Treat; 75(3):259-268.

Antoniou A.C., Pharoah P.D., Narod S. et al, (2005): Breast and ovarian cancer risks to carriers of the BRCA1 5382insC and 185delAG and BRCA2 6174delT mutations: a combined analysis of 22 population based studies. *Journal of Medical Genetics*; 42 (7): 602-603.

Ariga R., Bloom K., Reddy V.B. et al, (2002): Fine needle aspiration of clinically suspicious palpable breast masses with histopathologic correlation. *Am. J. Surg*; 184: 410-413.

Arriola E., Marchio C., Tan D. et al, (2008): Genomic analysis of the HER2/TOP2A amplicon in breast cancer and breast cancer cell lines. *Laboratory Investigation*; 88: 491-503.

B

Baldus S.E., Wienand J.R., Werner J.P., (2005): Expression of MUC1, MUC2 and oligosaccharide epitopes in breast cancer: prognostic significance of a sialylated MUC1 epitope. *Int J Oncol.*; 27(5):1289-97.

Barbareschi M., Pecciarini L., Cangi M.G. et al, (2001): p63, a p53 homologue, is a selective nuclear marker of myoepithelial cells of the human breast. *Am. J. Surg. Pathol.*; 25(8):1054-60

Bardou V.J., Arpino G., Elledge R.M. et al, (2003): Progesterone receptor status significantly improves outcome prediction over estrogen receptor status alone for adjuvant endocrine therapy in two large breast cancer databases. *J. Clin. Oncol.*; 21: 1973-1979.

Baruch A., Hartmann M., Yoeli M. et al., (1999): The Breast Cancer-associated MUC1 Gene Generates Both a Receptor and Its Cognate Binding Protein. *Cancer Research*; 59: 1552-1561.

Bayer-Garner I.B. and Reed J.A., (2004): Immunolabeling pattern of syndecan-1 expression may distinguish pagetoid Bowen's disease, extramammary Paget's disease, and pagetoid malignant melanoma in situ. *J Cutan Pathol.*; 31(2):169-73.

Berg W.A., Gutierrez L., NessAiver M.S. et al, (2004): Diagnostic accuracy of mammography, clinical examination, US, and MR imaging in preoperative assessment of breast cancer. *Radiology*; 233(3): 830-49.

Bishop J., Cooke B., Coleman M. et al, (2004): Breast fine needle aspiration cytology and needle core biopsy. A guide for practice, first edition. National breast cancer center, Australia-Sidney.

Bhargava R., Gerald W.L., Li A.L. et al, (2005): [*EGFR*](#) gene amplification in breast cancer: correlation with epidermal growth factor receptor mRNA and protein expression and HER-2 status and absence of *EGFR*-activating mutations. *Modern Pathology*; 18:1027-1033.

Bharti A., Chen D., Gong J. et al., (1998): Interaction of glycogen synthase kinase 3 β with the DF3/MUC1 carcinoma-associated antigen and β -catenin. *Mol. Cell. Biol.*; 18: 7216–7224.

Bleiweiss I., Nagi C. and Jaffer S., (2005): Epithelial Displacement in Breast Lesions: A Papillary Phenomenon. *Archives of Pathology and Laboratory Medicine*: 129 (11): 1465–1469.

[Bobo J.K., Lee N.C. and Thames S.F., \(2000\): Findings from 752,081 clinical breast examinations reported to a national screening program from 1995 through 1998. J Natl Cancer Inst 92 \(12\): 971-6.](#)

Bracke M.E., van Roy F.M., and Mareel M. M., (1996): *Curr. Top. Microbiol. Immunol.*; 213: 123-161.

[Bratthauer G.L., Moinfar F., Stamatakis M.D., et al, \(2002\):](#) Combined E-cadherin and high molecular weight cytokeratin immunoprofile differentiates lobular, ductal, and hybrid mammary intraepithelial neoplasias. [*Hum Pathol*](#); 33(6):620-7.

Britton P. (1999): Fine needle aspiration or core biopsy. *The breast J.*; 8: 1-4.

[Bulgaresi P., Cariaggi M.P., Bonardi L. et al, \(2005\):](#) Analysis of morphologic patterns of fine-needle aspiration of the breast to reduce false-negative results in breast cytology. *Cancer*; 105 (3):152-7.

Cai G., Simsir A. and Cangiarella J., (2004): "Invasive mammary carcinoma with osteoclast-like giant cells diagnosed by fine-needle aspiration biopsy: review of the cytologic literature and distinction from other mammary lesions containing giant cells. [Diagn. Cytopathology](#); 30 (6):396-400.

Cangiarella J., Waisman J. and Shapiro R.L., (2001): Cytologic features of tubular adenocarcinoma of the breast by aspiration biopsy. [Diagn. Cytopathology](#); 25: 311-315.

Capó M., Reboredo J.M., and Tomas D., (2002): Fine needle aspiration cytology of mammary adenomyoepithelioma, a case report. Acta clin. Croat; 41:135-186.

Castro M.R. and Gharib H., (2005): Continuing controversies in management of thyroid nodules. Ann Intern Med; 142: 926-931.

Carter M.R., Hornick J.L. and Lester S., (2006): Spindle Cell (Sarcomatoid) Carcinoma of the Breast: A Clinicopathologic and Immunohistochemical Analysis of 29 Cases. Am J Surg Pathol.; 30 (3):300-309.

Chaiwun B., Sukhamwang N., Lekawanfiji S. et al, (2005): Atypical and suspicious categories in fine needle aspiration cytology of the breast: histological and mammographical correlation and clinical significance. Singapore Med. J.; 64 (12): 706-709.

Chang G.J., Rodriguez-Bigas M.A., Skibber J.M. et al, (2002): Lymph Node Evaluation and Survival after Curative Resection of Colon Cancer: Systematic Review. J Natl Cancer Inst; 99(6): 433 - 441.

Chang S., Brenner R.J. and Giuliano A.E., (2004): Atypical lobular hyperplasia: is excision necessary? Poster presentation abstracts. Las Vegas; 10: 18-20.

Chuo C. and Corder A., (2003): Core biopsy vs fine needle aspiration cytology in a symptomatic breast clinic¹. European Journal of Surgical Oncology; 29 (4): 374 – 378.

Ciatto S., Del Turco M.R., Risso G. et al, (2003): Comparison of standard reading and computer aided detection (CAD) on a national proficiency test of screening mammography. Eur J Radiol 45 (2): 135-8.

Ciborowski P. and Finn O.J., (2002): Non-glycosylated tandem repeats of MUC1 facilitate attachment of breast tumor cells to normal human lung tissue and immobilized extracellular matrix proteins (ECM) in vitro: potential role in metastasis. Clin Exp Metastasis; 19(4):339-45.

Cibull M., Lele S. and Kesler M., (2004): Distinguishing well-differentiated hepatocellular carcinoma from benign liver by fine-needle aspirates. Mod Pathol.; 17: 798-802.

Cohen E.G., Patel S.G. and Lin O., (2004): Fine-needle aspiration biopsy of salivary gland lesions in a selected patient population. Arch Otolaryngol Head Neck Surg.; 130(6): 773-8

Collins L.C. and Schnitt S.T., (2005): HER2 protein overexpression in estrogen receptor-positive ductal carcinoma in situ of the breast: frequency and implications for tamoxifen therapy. Modern Pathology; 18: 615-620.

Cokkinda V, Bandi P, Siegel R et al, 2007: Cancer prevention and early detection facts and figures 2007. American Cancer Society-Texas.

Copson E.R., White H.E., Blaydes J.P. et al, (2006): Influence of the MDM2 single nucleotide polymorphism SNP309 on tumour development in BRCA1 mutation carriers. BMC Cancer; 6: 80-89.

Corfield A.P., Myerscough N., Longman R. et al., (2000): Mucins and mucosal protection in the gastrointestinal tract: new prospects for mucins in the pathology of gastro- intestinal disease. Gut; 47: 589-594.

Costas A., Castro P. and Martin-Granizo R., (2000): Fine needle aspiration biopsy (FNAB) for lesions of the salivary glands. Br J Oral Maxillofac Surg; 38 (5): 539-42.

Croce M.V., Isla-Larrain M.T., Demichelis S.O. et al, (2003): Tissue and serum MUC1 mucin detection in breast cancer patients. Breast cancer research and treatment; 81: 195-207.

Cummings M.C., Walsh M.D., Hohn B.G. et al, (2002): Occult axillary lymph node metastases in breast cancer do matter: results of 10-year survival analysis. Am J Surg Pathol.; 26 (10):1286-95.

D

Daniel B.L., Birdwell R.L., Butts K. et al, (2001): Freehand iMRI-guided large-gauge core needle biopsy: a new minimally invasive technique for diagnosis of enhancing breast lesions. J Magn Reson Imaging.; 13: 6: 896-902.

Das D.K. and Sheikh Z.A., (2003): Breast Carcinoma with Neuroendocrine Differentiation: Diagnosis of a Case by Fine Needle Aspiration Cytology and Immunocytochemistry. Acta Cytologica J.; 23 (16):186-195.

Dawson A.E. and Mulford D.K., (1998): Fine needle aspiration of mucinous (colloid) breast carcinoma—nuclear grading and mammographic and cytologic findings. Acta Cytol; 42: 668–72.

Deb S.J., Zehr K., Shields R.C., et al, (2001): Effectiveness of breast cancer screening by mammography in younger womenBreast. Cancer.org.; 80: 1500-1502.

de Cremoux P., Extra J. M., Denis M. G. et al, (2000): Detection of MUC1-expressing Mammary Carcinoma Cells in the Peripheral Blood of Breast Cancer Patients by Real-Time Polymerase Chain Reaction. Clinical Cancer Research; 6: 3117-3122.

Delva D., (2002): Practice Tips: Fine needle aspiration of breast lumps. Canadian family physician, Ann R Coll Surg Engl; 83(2):110-2.

Denley H., Pinder S.E., Elston C.W. et al, (2001): Preoperative assessment of prognostic factors in breast cancer. J Clin Pathol; 54:20-24.

Diallo R., Schaefer K.L., Bankfalvi A. et al, (2003): Secretory carcinoma of the breast: a distinct variant of invasive ductal carcinoma assessed by comparative genomic hybridization and immunohistochemistry. *Hum Pathol.*; 34(12):1299-305.

Diaz N.M., Cox C.E., Ebert M. et al, (2004): Benign mechanical transport of breast epithelial cells to sentinel lymph nodes. *Am J Surg Pathol.*; 28(12):1641-5.

Di Tommaso L., Arizzi C., and Rahal D., (2001): Anatomic Location of Breast Cancer Micrometastasis in Sentinel Lymph Node Predicts Axillary Status *Ann Surg.*; 243 (5): 706–707.

Downs-Kelly E., Yoder B.J., Stoler M. et al, (2005): The influence of polysomy 17 on HER2 gene and protein expression in adenocarcinoma of the breast: a fluorescent in situ hybridization, immunohistochemical, and isotopic mRNA in situ hybridization study. *Am J Surg Pathol.*; 29(9):1221-7.

Dunne B., Lee A.H. and Pinder S.E., (2003): An immunohistochemical study of metaplastic spindle cell carcinoma, phyllodes tumor and fibromatosis of the breast. *Hum.Pathol.*; 34(10):1009-15.

Dyomin V.G., Palanisamy N., Lloyd K.O. et al., (2000): MUC1 is activated in a B-cell lymphoma by the t (1; 14) (q21; q32) translocation and is rearranged and amplified in B-cell lymphoma subsets. *Blood*, 95(8): 2666-2671.

£

Eccles D.M. and Pichert G., (2005): Familial non-BRCA1/BRCA2-associated breast cancer. *The lancet oncology*; 6 (9): 705-711.

Eckert R. and Howell L.P., (1999): Number, size, and composition of cell clusters as related to breast FNA adequacy. *Diagn Cytopathol*; 21(2): 105-11.

Elattar I., (2005): Breast cancer: magnitude of the problem. Egyptian Society of Surgical Oncology Conference.

El-Bolkainy, (1998): Topographic pathology of cancer. NCI, Cairo Univerity.

Elledge R.M., Green S., Pugh R. et al, (2000): Estrogen receptor (ER) and progesterone receptor (PgR), by ligand-binding assay compared with ER, PgR and pS2, by immuno-histochemistry in predicting response to tamoxifen in metastatic breast cancer: a Southwest Oncology Group Study. *Int J Cancer*; 89: 111–117.

Ellis I.O., Humphreys S., Michell M. et al, (2001): Guidelines for non-operative diagnostic procedures and reporting in breast cancer screening. NHS Cancer screeing programme, NHSBSP Publication (5).

Ellison J.H., Wu X.C., Howe H.L. et al, (2005): Cancer in North America, 1998-2002, Volum 3: NAACCR Combined Incidence Rates. Springfield IL: North America Association of Central Cancer Registries.

Elmore J.G. and Carney P.A., (2004): Computer-aided detection of breast cancer: has promise outstripped performance? *J. Natl. Cancer Inst.*; 96 (3): 162-3.

Elmore J.G., Armstrong K., Lehman C.D. et al, (2005): Screening for breast cancer. *JAMA*; 293(10): 1245-56.

F

Fabian C.J., Kimler B.F., Brady D.A. et al, (2002): Phase II Breast Cancer Chemoprevention Trial of Oral α -Difluoromethylornithine. *Clinical Cancer Research*; 8: 3105-3117.

Favret A.M., Carlson R.W., Goffinet D.R. et al, (2001): Locally advanced breast cancer: is surgery necessary? *Breast J.*; 7 (2): 131-7.

Feng H., Ghazizadeh M., Konishi H. et al, (2002): Expression of MUC1 and MUC2 Mucin Gene Products in Human Ovarian Carcinomas. *Japanese Journal of Clinical Oncology*; 32: 525-529.

Fentiman I.S., (2001): Fixed and modified risk factors for breast cancer. *Int. J. Clin. Pract.*; 55 (8): 527-530.

Fernandez Y., Martinez A., Torregrosa A., et al, (2002): "Inhibition of Apoptosis in Human Breast Cancer Cells: Role in Tumor Progression to the Metastatic State." *Int. J. Cancer*; 101: 317-326.

Fiegelson H.S., Jonas C.R., Teras L.R. et al, (2004): Wight gain, body mass index, hormonal replacement therapy, and postmenopausal breast cancer in a large prospective study. *Cancer Epidemiol Biomarker Prev.*; 13 (2): 220-224.

Fitzgibbons P.L., Page D.L., Weaver D. et al, (2002): Prognostic factors in breast cancer: College of American Pathologists consensus statement 1999. *Arch Pathol Lab Med*; 124: 966-978.

Frost F.A., Sterrett G.F., Whitaker D et al, (2001): Core imprint cytology: technique used in a breast assessment center. Data presented at the 27th Annual Scientific Meeting of Australian Division of The International Academy of Pathology Limited. Sidney 2001.

Fujita K., Denda K., Yamamoto M. et al, (1999): Expression of MUC1 mucins inversely correlated with post-surgical survival of renal cell carcinoma patients. *British Journal of Cancer*; 80: 301-308.

G

Gaemers I.C., Vos H.I., Volders H.H., et al, (2001): A STAT-responsive Element in the Promoter of the Episialin/MUC1 Gene Is Involved in Its Overexpression in Carcinoma Cells. *J. Biol. Chem.*, 276 (9): 6191-6199

Geradts J. and Ingram C.D., (2000): Abnormal expression of cell cycle regulatory proteins in ductal and lobular carcinomas of the breast. *Modern Pathology*. 13(9): 945-953.

Gilles F., Goy A., Remache Y. et al, (2000): MUC1 dysregulation as the consequence of a t(1;14)(q21;q32) translocation in an extranodal lymphoma *Blood*; 95: 2930-2936

Gong Y., Gilcrease M., and Sneige N., (2005): Reliability of chromogenic in situ hybridization for detecting HER-2 gene status in breast cancer: comparison with fluorescence in situ hybridization and assessment of interobserver reproducibility. Mod Pathol.; 18(8):1015-21.

Gram I.T., Fankhouser E. and Nordgard L., (2002): Oral contraceptive use and mammographic patterns. Eur J Cancer Prev.; 11(3): 265-270.

Gray W. and McKee G.T., (2003): Diagnostic Cytopathology, 2nd edition Elsevier Science. J.Clin. Path.; 56: 556-560.

Groisman G.M., Amar M. and Livne E., (2002): CD10: a valuable tool for the light microscopic diagnosis of microvillous inclusion disease (familial microvillous atrophy). Am J Surg Pathol.; 26 (7): 902-7.

Gui G.P., Hogben R.K., Walsh G. et al, (2001): The incidence of breast cancer from screening women according to predicted family history risk: Does annual clinical examination add to mammography? Eur J Cancer 37 (13): 1668-73.

Gunther K., Merkelbach-Bruse S. and Amo-Takyi B.K., (2001): Differences in genetic alterations between primary lobular and ductal breast cancers detected by comparative genomic hybridization. J Pathol; 193: 40-47.

Gupta D., Croitoru C.M., Ayala A.G. et al, (2002): E-cadherin immunohistochemical analysis of histiocytoid carcinoma of the breast. Ann Diagn Pathol; 6(3):141-7

Gupta R.K., (2007): Needle aspiration cytodiagnosis of the breast with abundant acellular calcific material mimicking soft tissue calcinosis. Acta Cytol; 51(1):33-6.

Gürbüz Y. and Özkara S.K., (2003): Clear cell carcinoma of the breast with solid papillary pattern: a case report with immunohistochemical profile. Journal of Clinical Pathology; 56: 552-554.

H

Hamajima M., Hirose K., Tajima K. et al, (2002): Alcohol and breast cancer-collaborative reanalysis of individual data of 53 epidemiological studies. Br J Cancer; 87(11): 1234-1245.

Hansen N.M., Ye X., Grube B.J. et al. (2004): Manipulation of the primary breast tumor and the incidence of sentinel node metastases from invasive breast cancer. Arch Surg.; 139:634-640

Hasebe T., Sasaki S., Imoto S. et al, (2002): Prognostic significance of fibrotic focus in invasive ductal carcinoma of the breast: a prospective observational study. Mod Pathol.; 15(5):502-16.

Hasebe T., Sasaki S., Imoto S. et al (2002): Characteristics of tumors in lymph vessels play an important role in the tumor progression of invasive ductal carcinoma of the breast: a prospective study. Mod Pathol.; 15(9):904-13.

Hattrup C.L., and Gendler S. J., (2006): MUC1 alters oncogenic events and transcription in human breast cancer cells. Breast Cancer Res.; 8 (4): R37.

Hernandez O., Oweity T. and Ibrahim S. (2005): Is an increase in CD4/CD8 T-cell ratio in lymph node fine needle aspiration helpful for diagnosing Hodgkin lymphoma? A study of 85 lymph node FNAs with increased CD4/CD8 ratio. CytoJournal, 2:14 doi: 10.1186/1742-6413-2-14

Higashi M., Yonezawa S., Ho J.L. et al., (1999): Expression of MUC1 and MUC2 mucin antigens in intrahepatic bile duct tumors: its relationship with a new morphological classification of cholangiocarcinoma. Hepatology, 30: 1347-1355.

Hill C.B., and Tien Yeh I., (2005): Myoepithelial Cell Staining Patterns of Papillary Breast Lesions from Intraductal Papillomas to Invasive Papillary Carcinomas. Am. J. Clin. Pathol.; 123:36-44.

Hirasawa Y., Kohno N., Yokoyama A. et al, (2000): Natural autoantibody to MUC1 is a prognostic indicator for non-small cell lung cancer. Am. J. Respir. Crit. Care Med.; 161: 589-594.

Honma N., Sakamoto G., Ikenaga M. et al, (2003): Mucinous Cystadenocarcinoma of the Breast: A Case Report and Review of the Literature. Archives of Pathology and Laboratory Medicine; 127(8): 1031–1033.

Hughes O.D., Bishop M.C., Perkins A.C. et al., (2000): Targeting superficial bladder cancer by the intravesical administration of copper-67-labeled anti-MUC1 mucin monoclonal antibody C595. J. Clin. Oncol.; 18: 363-70.

Hulka B.S. and Moorman P.G., (2001): Breast cancer: Hormones and other risk factors. Maturita.; 38(1): 103-113.

Hummel L.P., Waisman J., Yang G.C.H. et al, (2003): Aspiration cytology of cystic carcinoma of the breast. Diagn. Cytopathol.; 28: 39-44.

Humphreys L.L., Helfand M. and Chan B.S., (2002): Breast cancer screening summary of the evidence for the U.S. preventing services task force. Ann Intern Med; 137:347-360.

J

Irwig L., Houssami N. and van Vliet C., (2004): New technologies in screening for breast cancer: a systematic review of their accuracy. Br J Cancer; 90(11): 2118-22

L

Jackson V.P. et al., (1999): Stereotactic fine-needle aspiration biopsy for non palpable breast lesions. AJR Am J Roentgenol; 174: 1196-1197.

Jayaram G., Elsayed E.M. and Yacob R.B., (2007): Papillary breast lesions diagnosed on cytology. Profile of 65 cases. Acta Cytol; 51(1): 3-8.

Jeffrey S.S. and Pollack J.R., (2003): The diagnosis and management of pre-invasive breast disease: promise of new technologies in understanding pre-invasive breast lesions. *Breast Cancer Res.*; 5: 6: 320-8.

Jemal A., Murray T., Ward E. et al, (2005): Cancer statics, 2005. *Cancer J. Clin.*; 55(1): 10-30.

Jemal A., Siegel R, Ward E. et al, (2007): Cancer facts and figures 2007. American cancer society-Atlanta.

Johnson S.J. and Wadehra V., (2001): The importance of intracytoplasmic DPAS positivity in fine-needle aspirates of breast lesions. *J. Clin. Pathol*; 54:146-151.

K

Kamiya M., Mizuguchi K., Yoshimoto M. et al; (1998): Cytologic diagnosis of signet-ring cell carcinoma of the breast. *Acta Cytol*; 42: 650–6.

Kanhoush R., Jorda M., Gomez-Fernandez C. et al, (2004): Atypical and suspicious diagnosis in breast aspiration cytology. *Cancer*; 102: 164-7.

Kawamoto T., Shoda S., Irimura T. et al., (2001): Expression of MUC1 Mucins in the Subserosal Layer Correlates with Postsurgical Prognosis of Pathological Tumor Stage 2 Carcinoma of the Gallbladder. *Clinical Cancer Research*; 7, 1333-1342.

Khan S.A., Wiley E.L., Rodriquez N., et al, (2004): Ductal lavage findings in women with known breast cancer undergoing mastectomy. *J Natl Cancer Inst*; 96:1510-7.

Kim M.J., Gong G., Joo H.J. et al, (2005): Immunohistochemical and clinicopathologic characteristics of invasive ductal carcinoma of breast with micropapillary carcinoma component. *Arch Pathol Lab Med.*; 129 (10):1277-82.

Kleer C.G., van Golen K.L. and Braun T., (2001):Persistent E-cadherin expression in inflammatory breast cancer. Mod Pathol.; 14 (5):458-64.

Klein S., 2005: Evaluation of Palpable Breast Masses. American Academy of Family Physicians afpserv@aafp.org.

Klemi P.J., Joensuu H. and Nylamo E., (1991): Fine needle aspiration biopsy in the diagnosis of thyroid nodules. Acta Cytol.; 35 (4): 434-8.

Klijanienko J. Katsahian S. and Vielh P., (2004): [Stromal infiltration as a predictor of tumor invasion in breast fine needle aspiration biopsy. Diagn. Cytopathol.; 30: 182-186.](#)

Klimstra D.S., Adsay V.N., Andea A. et al., (2004): MUC1 and MUC2 in pancreatic neoplasia. Journal of Clinical Pathology; 57:456-462.

[Knoop A.S., Bentzen S.M., Nielsen M.M. et al, \(2001\): Value of epidermal growth factor receptor, HER2, p53, and steroid receptors in predicting the efficacy of tamoxifen in high-risk postmenopausal breast cancer patients. J. Clin. Oncol.; 19: 3376–3384.](#)

Kohlgraf G.K., Gawron A.J., Higashi M. et al, (2003): Contribution of MUC1 tandem repeat and cytoplasmic tail to invasive and metastatic properties of pancreatic cancer cell line. Cancer Research; 63:5011-5020.

Koker M.M. and Kleer C.G. (2004): p63 expression in breast cancer: a highly sensitive and specific marker of metaplastic carcinoma. Am. J. Surg. Pathol.; 28(11):1506-12

Kolb T.M., Lichy J. and Newhouse J.H., (2002): Comparison of the performance of screening mammography, physical examination, and breast US and evaluation of factors that influences them: an analysis of 27,825 patient evaluations. Radiology; 225(1): 165-75.

Kollur S.M. and El Hag I.A., (2006): FNA of breast fibroadenoma: observer variability and review of cytomorphology with cytohistological correlation. Cytopathology; 17(5): 239-44.

Kontzoglou K., Moulakakis K.G. and Konofaos P., (2005): the role of liquid-based cytology in the investigation of breast lesions using fine-needle aspiration: a cytohistopathological evaluation. J. Surg. Oncol.; 89(2):75-80.

Korkola J.E., Devries S. and Fridlyand J. et al, (2003): Differentiation of lobular versus ductal breast carcinomas by expression microarray analysis. *Cancer Res*; 63: 7167–7175.

Kroman N., Jensen M.B., Wohlfahrt J. et al, (2000): Factors influencing the effect of age on prognosis in breast cancer: population based study. *BMJ*; 320: 474-478.

Kurihara T., Brough E.D., and Kovesdi I., (2000): Selectivity of a replication-competent adenovirus for human breast carcinoma cells expressing the MUC1 antigen. *J. Clin. Invest.*; 106 (6): 763-771.

Kuwahara H., Carraway K. L., Kufe D. et al., (2001): The EGF receptor regulates interaction of the human DF3/MUC1 carcinoma antigen with c-Src and β -catenin. *J. Biol. Chem.*, 276: 35239–35242.

L

Landis M.D., Seachrist D.D., and Abdul-Karim F.W., (2006): Sustained trophism of the mammary gland is sufficient to accelerate and synchronize development of ErbB2/Neu-induced tumors. *Oncogene*; 25: 3325–3334.

Lee M.J., Lee H.S., Kim W.H. et al., (2003): Expression of mucins and cytokeratins in primary carcinomas of the digestive system. *Mod Pathol*; 16: 403–10.

Lee S., Berg K.D., Sherman M.E. et al, (2001): Microsatellite Instability Is Infrequent in Medullary Breast Cancer. *Am. J. Clin. Pathol.*; 115: 823-827.

Leibl S., Denk H., Moinfar F. et al, (2005): Metaplastic breast carcinomas: are they of myoepithelial differentiation? *Am J Surg Pathol.*; 29(3):347-53.

Leroy X., Devisme L., and Buisine M.P., (2003): Expression of human mucin genes during normal and abnormal renal development. Am. J. Clin. Pathol.; 120(4):544-50).

Lichtenstein P., Holm N.V., Verkasalo P.K. et al, (2000): Hereditary and environmental factors in causation of cancer-Analysis of cohorts of twins from Sweden, Denmark and Finland. N. Engl. J. Med.; 343 (2): 78-85.

Lieske B., Ravichandran D. and Wright D., (2006): Role of fine-needle aspiration cytology and core biopsy in the preoperative diagnosis of screen-detected breast carcinoma. British Journal of Cancer; 95:62-66.

Li Y., Yamamoto M., Bharti A., et al., (1997): Interaction of the DF3/MUC1 breast carcinoma-associated antigen and β -catenin in cell adhesion. J. Biol. Chem., 272: 12492–12494.

Li Y., Kuwahara H., Ren J. et al., (2001): The c-Src tyrosine kinase regulates signaling of the human DF3/MUC1 carcinoma-associated antigen with GSK3 β and β -catenin. J. Biol. Chem.; 276: 6061–6064.

Li Y., Ren J., Chen W. et al, (2003): Heregulin Targets γ -Catenin to the Nucleolus by a Mechanism Dependent on the DF3/MUC1 Oncoprotein. Molecular Cancer Research; 1:765-775.

Lim J.C., Al-Masri H., Salhadar A. et al, (2004): The significance of the diagnosis of atypia in breast fine-needle aspiration. Diagn. Cytopathol; 31(5):285-8.

Lloyd J. and Flanagan A.M., (2000): Mammary and extramammary Paget's disease. Journal of Clinical Pathology; 53(10):742-9.

Loman N., Johansson O., and Kristofferson U., (2001): Family history of breast and ovarian cancer and BRCA1&BRCA2 mutation in a population based series of early-onset breast cancer. J. Natl. Cancer Inst.; 93 (16):1215-1223.

Lopes C.V., Pereira-Lima J. and Hartmann A., (2006): Expression of mucin-2 and correlation with PAS-alcian blue stain in Barrett's esophagus. *Hepatogastroenterology*.; 53 (70): 511-5.

Lyda M.H., Tetef M., Carter N.H. et al, (2000): Keratin immunohistochemistry detects clinically significant metastases in bone marrow biopsy specimens in women with lobular breast carcinoma. *Am. J. Surg. Pathol.*; 24(12):1593-9.

M

Madhavan M., Srinivas P., Abraham E. et al, (2001): Cadherins as predictive markers of nodal metastasis in breast cancer. *Mod Pathol.*; 14: 423-427.

Makretsov N., Gilks C.B., Coldman A.J. et al, (2003): Tissue microarray analysis of neuroendocrine differentiation and its prognostic significance in breast cancer. *Hum Pathol.*; 34(10):1001-8.

Makretsov N., He M., Hayes M. et al, (2004): A fluorescence in situ hybridization study of ETV6-NTRK3 fusion gene in secretory breast carcinoma. *Genes Chromosomes Cancer*; 40(2):152-7.

Matsukita S., Nomoto M., Kitajima S. et al, (2003): Expression of mucins (MUC1, MUC2, MUC5AC and MUC6) in mucinous carcinoma of the breast: comparison with invasive ductal carcinoma. *Histopathology*.; 42(1):26-36.

McTiernan A., Kooperberg C. and White E., (2003): Recreational of physical activity and the risk of breast cancer in postmenopausal women: the Women Health Initiative Cohort Study. *JAMA.*; 290(10): 1331-1336.

Mehrotra R., Singh M., Singh P.A. et al, (2007): Should fine needle aspiration biopsy be the first pathological investigation in the diagnosis of a bone lesion? An algorithmic approach with review of literature. *CytoJournal*: <http://www.cytojournal.com/content/4/1/9>.

Meyer J.S., Alvarez C., Milikowski C. et al, (2005): Breast carcinoma malignancy grading by Bloom-Richardson system vs proliferation index: reproducibility of grade and advantages of proliferation index. Mod Pathol.; 18(8):1067-78.

Miksicek R.J., Myal Y., Watson P.H. et al, (2002): Identification of a novel breast and salivary gland-specific, mucin-like gene strongly expressed in normal and tumor human mammary epithelium. *Cancer Res.*; 62: 2736–2740.

Miller A.B., To T., Baines C.J. et al, (2000): Canadian National Breast Screening Study-2: 13-year results of a randomized trial in women aged 50-59 years. *J. Natl. Cancer Inst.*; 92 (18): 1490-9.

Miller C., Elmetts C.A., Vinson R.P. et al, (2005): Cowden Disease (Multiple Hamartoma Syndrome). eMedicine from Web site MD.

Mitchell G., Antoniou A.C., Warren R. et al, (2006): Mammographic density and breast cancer risk in BRCA1 and BRCA2 mutation carriers. *Cancer Research*; 66 (3): 1866-1872.

Mohsin S.K., Weiss H., Havighurst T. et al, (2004): Progesterone receptor by immunohistochemistry and clinical outcome in breast cancer: a validation study. *Modern Pathology*; 17: 1545-1554

Mokhter N., Gouda I. and Adel I., (2007): Cancer pathology registry 2003-2004.

Mora J. Puig P., Boadas J. et al, (1998): K-ras gene mutations in the diagnosis of fine-needle aspirates of pancreatic masses: prospective study using two techniques with different detection limits. *Clinical Chemistry*; 44:2243-2248.

Morandi L., Pession A., Marucci G.L. et al, (2003): Intraepidermal cells of Paget's carcinoma of the breast can be genetically different from those of the underlying carcinoma. *Hum Pathol.*; 34(12):1321-30.

Mouriquand J., Gozlan-Fior M., Villemain D. et al, (1986): Value of cytoprognostic classification in breast carcinomas. J Clin Pathol.; **39**(5):489–496.

Moriya T., Sakamoto K., Sasano H. et al, (2000): Immunohistochemical analysis of Ki-67, p53, p21, and p27 in benign and malignant apocrine lesions of the breast: its correlation to histologic findings in 43 cases. Mod Pathol.; 13(1):13-8

Mridha A.R., Iyer V.K. and Kapila K., (2006): Value of scoring system in classification of proliferative breast disease on fine needle aspiration cytology. Indian J. Pathol Microbiol; 49 (3):334-40.

Muller S. and Hanisch F.G., (2002): Recombinant MUC1 probe authentically reflects cell-specific O-glycosylation profiles of endogenous breast cancer mucin. J. Biol. Chem., 277(29): 26103-26112.

N

Nakagawa K., Akagi J., Takai E. et al. (1999): Prognostic values of MUC-1 molecule expressing cytokine receptor-like epitope and DF3 in patients with gastric carcinoma. Int J Oncol.; 14:425-435

Newcomb P.A., Titus-Ernstoff L., Egan K.M. et al, (2002): Postmenopausal estrogen and progesterone use in relation to breast cancer risk. Cancer Epidemiol biomarker Prev.; 11(7): 539-600.

Newman L.A., Carolin K., Simon M. et al, (2001): Impact of breast carcinoma on American-African women: The Detroit experience. Cancer J.; 91 (9): 1834-1843.

Ng W.K., (2002): fine needle aspiration cytology findings of an uncommon micropapillary variant of pure mucinous carcinoma of the breast. Cancer; 27: 214-217.

Nonni A., Lazaris A.C. and Patsouris E., (2005): Invasive Papillary Breast Carcinoma of Solid Variant. Archives of Pathology and Laboratory Medicine; 129(5):e128–e129.

Nyante S.J., Devries S., Chen Y.Y., (2004): Array-based comparative genomic hybridization of ductal carcinoma in situ and synchronous invasive lobular cancer. Hum Pathol.; 35(6):759-63.



O'Connor J., Julian J. and Lim S.D., (2005): MUC1 expression in human prostate cancer cell lines and primary tumors. Prostate Cancer and Prostatic Diseases; 8: 36–44.

Oh D.S., Troester M.A., Usary J. et al (2005): Estrogen-regulated genes predict survival in hormone receptor-positive breast cancers. J Clin Oncol 24:10.1200/JCO. 03.2755.

O'mally C.D., Le G.M., Glaser S.L. et al, (2003): Socioeconomic status and breast carcinoma survival in four racial/ethnic groups. Cancer; 97(5): 1303-1311.

Omar S., Khaled H. Gaafar R., (2003): Breast cancer in Egypt: a review of disease presentation and detection strategies. La Revue de Santé de la Méditerranée orientale, 9(3): 137-140

Osslon H.L., Ingvar C. and Bladstorm A., (2003): Hormon replacement therapy containing progestins and given continuously increases breast carcinoma risk in Sweden. Cancer; 97(6): 1387-1392.

Ostean R.T., Lenhard R.E. and Gansler T., (2001): Osteon R Breast Cancer. Clinical Oncology. Atlanta, GA: American Cancer Society: 251-268



Page D.L., (2003): Special types of invasive breast cancer, with clinical implications. Am J Surg Pathol; 27: 832–835.

Palacios J., Sarrio D., Garcia-Macias M.C. et al, (2003): Frequent E-cadherin gene inactivation by loss of heterozygosity in pleomorphic lobular carcinoma of the breast. Mod Pathol; 16: 674–678.

Palmieri C., Cheng G. J., Saji S. et al, (2002): Estrogen receptor β in breast cancer. Endocr. Relat. Cancer, 9: 1-13.

Papadimitriou T.J., Burchell J.M., Plunkett T. et al., (2002): MuC1 and the immunobiology of cancer. J Mammary Gland Biol Neoplasia; 7:209–21.

Parkin D.M., Bray F. and Ferlay J., (2001): Estimating the world cancer burden. Int J Cancer; 94:153-156.

Patel K., Sakamoto K., Jones G.R. et al, (2006): Li-Fraumeni Syndrome. eMedicin from WebMD.

Patel S.G., Lin O. and Cohen E.G. (2004): Fine-needle aspiration biopsy of salivary gland lesions in a selected patient population. Arch Otolaryngol Head Neck Surg; 130(6): 773-8

Pettinato G., Manivel C.J., Panico L. et al, (2004): Invasive micropapillary carcinoma of the breast, clinicopathologic study of 62 cases of a poorly recognized variant with highly aggressive behavior. Am J Clin Pathol 121(6):857-866.

Pinto-de-Sousa et al, 2002: Mucins MUC1, MUC2, MUC5AC and MUC6 expression in the evaluation of differentiation and clinico-biological behaviour of gastric carcinoma. Virchows Arch; 440:304–10.

Prolla and Diehl, (2002): Prolla List of Cases Cytopathology Atlas; <http://www.geocities.com/jcprolla/listofcases>.

Poremba C, Shroyer RK, Frost M et al, (1999): Telomerase is a highly sensitive and specific molecular marker in fine-needle aspirates of breast lesions. J. Clin. Oncol.; 17:2020-2026.

Putti T.C., Abd El-Rehim D., Rakha E. et al, (2005): Estrogen receptor-negative breast carcinomas: a review of morphology and immunophenotypical analysis. Modern Pathology; 18: 26-35.

Q

Qayyum A., Birdwell R.L., Daniel B.L. et al, (2002): "MR imaging features of infiltrating lobular carcinoma of the breast: histopathologic correlation." *AJR Am J Roentgenol.*; 178 (5): 1227-32.

R

Raaschou-Nielsen O., Pavuk M., LeBlanc A. et al, (2005): Adipose organochlorine concentration and risk factor of breast cancer among postmenopausal Danish women. *Cancer Epidemiol Biomarkers*; 14(1): 67-74.

Rahn J.J., Dabbagh L., Pasdar M., et al, (2001): The importance of MUC1 cellular localization in patients with breast carcinoma: an immunohistologic study of 71 patients and review of the literature. *Cancer*; 91:1973–1982.

Rakha E., Boyce R.W., Abd El-Rehim D. et al, (2005): Expression of mucins (MUC1, MUC2, MUC3, MUC4, MUC5AC and MUC6) and their prognostic significance in human breast cancer. *Modern Pathology*; 18: 1295–1304.

Raveendran Pillai K., Jayasree, K., Jayalal K.S. et al, (2007): Mucinous carcinoma of breast with abundant psammoma bodies in fine-needle aspiration cytology: A case report. *Diagnostic Cytology*; 35(4): 230-233.

Raynor M., Stephenson S., Walsh D. et al, (2002): Optimisation of the RT-PCR detection of immunomagnetically enriched carcinoma cells. *BMC Cancer*; 2:14-17.

Reboredo J.M., Barreiro A.V. and Kraveva S., (2002): Fine needle aspiration cytology of pleomorphic adenoma of the breast, Case report. *Acta clin. Croat*; 41: 135-186.

Reis-Filho J.S., Fulford L.D., Lakhani S.R. et al, (2003): Lipid-Rich Breast Carcinoma. Archives of Pathology and Laboratory Medicine; 127(9): e396–e398.

Reis-Filho J.S., Milanezi F., Amendoeira I. et al, (2002): P63 staining of myoepithelial cells in breast fine needle aspirates: a study of its role in differentiating in situ from invasive ductal carcinomas of the breast J. Clin. Pathol.; 55(12): 936 – 939

Reis-Filho J.S., Savage K., Lambros M.B. et al, (2006): Cyclin D1 protein overexpression and CCND1 amplification in breast carcinomas: an immunohistochemical and chromogenic in situ hybridisation analysis. Modern Pathology; 19: 999-1009

Renshaw A.A., (2004): Minimal (< or =0.1 cm) invasive carcinoma in breast core needle biopsies. Incidence, sampling, associated findings, and follow-up. Arch Pathol Lab Med.; 128 (9):996-9.

Requena L., Sanguenza M. and Sanguenza O.P., (2002): Pigmented Mammary Paget Disease and Pigmented Epidermotropic Metastases From Breast Carcinoma. Am J Dermatopathol; 24 (3):189-198.

Resetskova E., Padula A., Albarraccin C.T. et al, (2004): Pathologic Quiz Case: A Large, Ill-Defined Cystic Breast Mass; Pathologic Diagnosis: Invasive Cystic Hypersecretory Duct Carcinoma. Archives of Pathology and Laboratory Medicine; 129 (3): e79–e80.

Rhan J.J., Dabbagh L., Pasdar M. et al, (2001): The importance of MUC1 cellular localization in patients with breast carcinoma: An immunohistochemical study of 71 patients and review of literature. Cancer; 91: 1973-1982.

Ries L.G., Eisner M.P., Kosary C.L. et al, (2005): *SEER Cancer Statistics Review, 1975-2002*, National Cancer Institute. Bethesda, MD, http://seer.cancer.gov/csr/1975_2002/, based on November 2004 SEER data submission, posted to the SEER web site.

Ringel J. and Lohr M., (2003): The MUC gene family: their role in diagnosis and early detection of pancreatic cancer. Mol Cancer; 2: 9-11.

Robinson I.A., McKee G., Nicholson A., et al, (1994): Prognostic value of cytological grading of fine-needle aspirates from breast carcinomas. Lancet; 343: 947–9.

Robinson I.A., McKee G., Kissin M.W., et al, (1995): Typing and grading breast carcinoma on fine needle aspiration: Is it clinically useful information? Diagn. Cytopathol.; 13: 260-265.

Rutledge H., Davis J., Chiu R. et al, (2005): Sentinel node micrometastasis in breast carcinoma may not be an indication for complete axillary dissection. Modern Pathology; 18: 762-768.



Saad R.S., Kanbour-Shakir A. and Syed A., (2006): Sclerosing papillary lesion of the breast: a diagnostic pitfall for malignancy in fine needle aspiration biopsy. Diagn Cytopathol; 34 (2):114-8.

Sahoo S. and Recant W.M., (2003): Matrix-Producing Metaplastic Carcinoma of the Breast. Archives of Pathology and Laboratory Medicine; 127 (10): 1385–1386.

Saji S., Omoto Y., Shimizu C. et al, (2002): Expression of Estrogen Receptor (ER) β cx Protein in ER α -positive Breast Cancer. Am. J. Surg.; 82: 129-31.

Sarli G., Brunetti B. and Benazzi C., (2006): Mammary mucinous carcinoma in the cat. Vet. Pathol.; 43 (5): 667-73.

Saxe A., Phillips E. and Husain M., (2001): Role of sample adequacy in fine needle aspiration biopsy of palpable breast lesions. Am. J. Surg.; 182: 369-71.

Schiller A.B., Tadros T.S., Birdsong G. et al, (2001): Cellular Dyscohesion in Fine-Needle Aspiration of Breast Carcinoma. Am. J. Clin. Pathol.; 115(2):219-223.

Schmitt S., (2002): Advances in breast FNA. Acta CLIN Croat; 41(2):150-153

Schnitt S.J. and Guidi A.J., (2000): Pathology of invasive breast cancer. In: Harris JR, Lippman ME, Morrow M, Osborne CK, eds. Diseases of the Breast, 2nd edition. Philadelphia: Lippincott, Williams and Wilkins; 425-470.

Schroeder J., Thompson M., Gardner M. et al., (2001): Transgenic MUC1 interacts with epidermal growth factor receptor and correlates with mitogen-activated protein kinase activation in the mouse mammary gland. J. Biol. Chem.; 276: 13057–13064.

Sethi S., Cajulis R.S., Gokaslan S.T. et al, (1997): Diagnostic significance of signet ring cells in fine-needle aspirates of the breast. Diagn Cytopathol; 16 (2):117-21.

Shousha S. and Baqai T., (2003): Oestrogen receptor negativity as a marker for high grade ductal carcinoma in situ of the breast. Histopathology; 42: 440-447.

Shousha S. and Scawn R., (2002): Morphologic spectrum of estrogen receptor-negative breast carcinoma. Arch Pathol Lab Med.; 126 (3):325-30.

Sidaway M.K., Tabbara S.O., Bryan J.A. et al, (2001): The spectrum of cytologic features in nonproliferative breast lesions. Cancer Cytopathol.; 93: 140-145.

Simon B., Syed A.H., Kim R.G. et al, (2004): E-Cadherin–Negative Primary Small Cell Carcinoma of the Breast Report of a Case and Review of the Literature. Am J Clin Pathol; 121:117-121.

Singh N. and Wells C.A., (2003): Inadequate rates are lower when FNAC are taken by cytopathologist. Cytopathology; 14: 327-31.

Singletary S.E., Allred C., Ashley P., et al, (2002): Revision of the American Joint Committee on Cancer Staging System for Breast Cancer. J. Clin. Oncol.; 20: 3628–3636.

Singletary S.E. and Connolly J.L., (2006): Breast Cancer Staging: Working With the Sixth Edition of the AJCC Cancer Staging Manual. CA Cancer J. Clin.; 56:37-47.

Singhal H., Kaur K. and Thomson S., (2006): Breast Cancer Evaluation. eMedicen from Web site MD.

Smigal C., Siegel R. and Jemal A., (2005): Breast cancer facts & figures 2005-2006. National home office: American cancer society, Clifton, Atlanta.

Smith R., Saslow D. and Sawyer K., (2003): American cancer society guidline for breast cancer screening. CA Cancer J. Clin.; 53(3): 141-169.

Snead D.J., Vryenhoef P., Pinder S.E. et al., (1997): Routine audit of breast fine needle aspiration (FNA) cytology specimens and aspirator inadequate rates. Cytopathology; 8:236-47.

Spicer A. P., Rowse G. J., Lidner T.K. et al, (1995): Delayed mammary tumor progression in MUC-1 null mice. J. Biol. Chem.; 270: 30093-30101.

Sohn J.H., Kim L.S., Chae S.W. et al, (2001): Fine needle aspiration findings of breast mucinous neoplasm: differential diagnosis between mucocele-like tumor and mucinous carcinoma. Acta Cytol.; 45: 723-729.

Song Z.B., Gao S., Yi Z.N. et al., (2003): Expression of MUC1 in esophageal squamous-cell carcinoma and its relationship with prognosis of patients from Linzhou city, a high incidence area of northern China. World J Gastroenterol; 9(3):404-407

Stalsberg H. and Hartmann W.H., (2000): The delimitation of tubular carcinoma of the breast. Hum. Pathol.; 31(5):601-7.

Stelow E.B., Bardales R.H., Crary G.S. et al., (2005): interobserver variability in thyroid fine needle aspiration interpretation of lesions showing predominantly colloid and follicular groups. Am J Clin Pathol; 124(2):239-244.

Tamada S., Goto M., Nomoto M. et al, (2002): Expression of MUC1 and MUC2 mucins in extrahepatic bile duct carcinomas: its relationship with tumor progression and prognosis. *Pathol Int*; 52:713–23

Tan P.H., Yip G., Bay B.H. et al, (2005): Cytokeratins in papillary Lesions of the Breast: Is There a Role in Distinguishing Intraductal papilloma from papillary ductal carcinoma In Situ? *Am J Surg Pathol.*; 29(5):625-632.

Tanaka M., Kitajima Y., Sato S. et al., (2003): Combined evaluation of mucin antigen and E-cadherin expression may help select patients with gastric cancer suitable for minimally invasive therapy. *Br J Surg*; 90:95-101.

Tanimoto T., Tanaka S., Haruma K. et al, (1999): MUC1 expression in intramucosal colorectal neoplasms: possible involvement in histogenesis and progression. *Oncology*; 56:223-231

Tavassoli F.A., et al, (2003): Adapted from WHO, Tumors of the Breast and Female Genital Organs. IARC Press, 2003

Thor A.D., Eng C., Devries S. et al, (2002): Invasive micropapillary carcinoma of the breast is associated with chromosome 8 abnormalities detected by comparative genomic hybridization. *Hum Pathol*; 33(6):628-31

Tomas A.R., Praca M.J., Fonseca R. et al, (2004): Assessing HER-2 status in fresh frozen and archival cytological samples obtained by fine needle aspiration cytology. *Cytopathol.*; 15: 311-314.

U

Ureta, G., Barreiro A.V., Reboredo M. et al, (2002): Fine needle aspiration cytology of pleomorphic adenoma of the breast. Case report. 13th Ljudevit Jurak International Symposium on Comparative Pathology, Zagreb, Croatia June 7-8, juraks@kbsm.hr. *Acta clin Croat* 2002; 41:135-186 Conference papers.

Ureta G. and Barreiro A.V., (2002): Fine needle aspiration cytology of adenoid cystic carcinoma of the breast. Report of three cases with thin-prep and immunocytochemical studies. 13th Ljudevit Jurak International

Symposium on Comparative Pathology, Zagreb, Croatia June 7-8,
juraks@kbsm.hr. Acta clin Croat; 41:135-186 Conference papers.

V^o

van de Vijver M.J., He Y.D., van't Veer L.J. et al (2002): A gene-expression signature as a predictor of survival in breast cancer. N Engl J Med 347:1999-2009.

Varder E., Ozkok G., Cetnile M. et al, (2005): Primary breast lymphoma cytology diagnosis. Arch Pathol Lab; 129: 694-696.

Ventura K., Cangiarella J., Lee I. et al, (2003): Aspiration Biopsy of Mammary Lesions with Abundant Extracellular Mucinous Material. Am J Clin Pathol 120(2): 194-202.

Volante M., Sapino A. and Croce S., (2003): Heterogeneous versus homogeneous genetic nature of multiple foci of in situ carcinoma of the breast. Hum Pathol.; 34(11):1163-9.

von Smitten K., (2000): Surgical management of breast cancer in the future. Acta Oncol; 39(3): 437-9.

W^o

Waldman F.M., Hwang E.S., Etzell J. et al, (2001): Genomic alterations in tubular breast carcinomas. Hum Pathol.; 32(2):222-6.

Walsh M. M. and Bleiweiss I. J., (2001): Invasive micropapillary carcinoma of the breast: Eighty cases of an underrecognized entity. Hum Pathol; 32:583-589.

Wang R.Q. and Fang D.C., (2003): Alterations of MUC1 and MUC3 expression in gastric carcinoma: relevance to patient clinicopathological features. J Clin Pathol.; 56(5): 378–384.

Wang X., Mori I., Tang W. et al, (2001): Metaplastic Carcinoma of the Breast: p53 Analysis Identified the Same Point Mutation in the Three Histologic Components. Mod Pathol; 14:1183-1186

Weigelt B., Verduijn P., Bosma A.J. et al, (2004): Detection of metastases in sentinel lymph nodes of breast cancer patients by multiple mRNA markers. Br. J. Cancer 90, 1531–1537.

Welsh P.L. and King M.C., (2001): BRCA-1 and BRCA-2 and the Genetics of Breast and Ovarian Cancer. Human Molecular Genetics; 10(7): 705-713.

Wesseling J., van der Valk S. W., and Hilken, J. (1996): Mol. Biol. Cell 7, 565-577.

Wheeler D.T., Tai L.H., Brathauer G.L. et al, (2005): Tubulolobular carcinoma of the breast: an analysis of 27 cases of a tumor with a hybrid morphology and immunoprofile. Am J Surg Pathol.; 29(7):980.

Williams S.J., Wreschner D.H., Tran M. et al, (2001): Muc13: a novel human cell surface mucin expressed by epithelial and hemopoietic cells. J Biol Chem; 276: 18327-18336

Wong P.W., Stefanec T. and Brown K., (2002): Role of Fine-Needle Aspirates of Focal Lung Lesions in Patients with Hematologic Malignancies. Chest; 121:527-532.

Woodward W.A., Strom E.A., Tucker S.L. et al, (2003): Changes in the 2003 American Joint Committee on Cancer Staging for Breast Cancer Dramatically Affect Stage-Specific Survival. Journal of Clinical Oncology, 21(17): 3244-3248.

World Health Organization (WHO), (2006): Fact sheet No. 297: Cancer. Retrieved on 2007-04-26 Weidner-breast.

Wreesmann V.B., Sieczka E.M., Socci N.D. et al., (2004): Genome-Wide Profiling of Papillary Thyroid Cancer Identifies MUC1 as an Independent Prognostic Marker. *Cancer Research*; 64: 3780-3789.

Wrench M.R., Petrakis N.L., Miike R. et al, (2001): Breast cancer risk in women with abnormal cytology in nipple aspirates of breast fluid. *J Natl Cancer Inst*; 93(23):1791-8

X

Xu R., Feiner H., Li P. et al, (2003): Differential amplification and overexpression of Her-2/Neu, p53, MIB1, and estrogen receptor/progesterone receptor among medullary carcinoma, atypical medullary carcinoma, and high-grade invasive ductal carcinoma of breast. *Archives of Pathology and Laboratory Medicine*; 127 (11): 1458–1464.

Xu Y., Kimura N., Yoshida R. et al, (2001): Immunohistochemical study of Muc1, Muc2 and human gastric mucin in breast carcinoma: relationship with prognostic factors. *Oncol Rep*; 8: 1177-1182.

Y

Yeoh P.S., Cheung S.Y., and Chan M.B., (1999): Fine-needle Aspiration Cytology of Mucoclelike Tumors of the Breast. *American Journal of Surgical Pathology*. 23(5):552-559

Yang G.C., Yang G.Y. and Tao L.C., (2004): Distinguishing well-differentiated hepatocellular carcinoma from benign liver by the physical features of fine-needle aspirates. *Mod Pathol.*; 17:798-802.

Younberg G.A., Hussain Z. and Martin A., (2001): Blastomyces dermatitis with large yeast forms. Arch Pathol Lab Med; 125:663-664.

Z

Zhang H.Z., Zhang Q.M., Zhao T.H. et al, (2004): Expression of mucins and E-cadherin in gastric carcinoma and their clinical significance. World J Gastroenterol; 10(20):3044-3047