

REFERENCES

Abell, M.R. (1957):

Leiomyosarcomas of inferior vena cava : review of literature and report of two cases.

Am. J. Clin. Pathol. 28 : 272 .

Adam, L.R.; Davison, E.V. and Malcolm, A.J. et al. (1991) :

Cytogenetic analysis of a congenital fibrosarcoma .

Cancer Genet Cytogenet 52:37.

Alguacil-Garcia, A, Unni K.K. and Goellner, J.R. (1977):

Malignant giant cell tumor of soft parts : Ultrastructural study of four cases .

Cancer 40-244 .

Albores-Saavedra, J, Buter, J.J. and Martin, R.G. (1965):

Rhabdomyosarcoma : clinicopathological considerations and report of 85 cases. In Tumors of bone and soft tissue, Chicago, year Book Medical Publishers.

Albus-Lutter C.E., de Stefain, E. And Van Unnik, J.A.M. (1986) :

Clinicopathologic relations in soft tissue sarcomas . In Management of soft tissue and bone sarcomas.

New York, Raven Press.

Angervall, I., Kindblom, L.G. and Rydholm, A. et al. (1986):

The diagnosis and prognosis of soft tissue tumors.

Semin Diagn Pathol 3 : 240.

Arlen , M. Higinbotham, N.L. and Huvos, A.G. et al. (1971):

Radiation – induced sarcoma of bone.

Cancer 28 : 1087.

Aurelio, J., Chenail, B. and Gerstein, H. (1984) :

Foreign-body reaction to bone wax.

Oral Surg , 58:98-100.

Azumi, N., Curtis, J. and Kempson, R.I. Hendirickson, M.R. (1987) :

Atypical and malignant neoplasms showing lipomatous differentiations: a study of III cases.

Am. J. Surg Pathol 11:161.

Bale, P.M. and Reye, R.D. (1975):

Rhabdomyosarcoma in childhood.

Pathology 7:101.

Barr, F.G., Chatten, J. and D'Cruz, C.M. et al. (1995):

Molecular assays for chromosomal translocations in the diagnosis of pediatric soft tissue sarcomas.

JAMA , 273 : 553 .

Battifora, H. and Nunez-Alonso, C. (1980):

Myxoid liposarcoma : study of 10 cases.

Ultrasound Pathol 1:157.

Becker, R.L., Venzon, D. and Lack, E.E. et al. (1991):

Cytometry and morphometry of malignant fibrous histiocytoma of the extremities : prediction of metastasis and mortality.

Am. J. Surg. Pathol. 15:87.

Breslin, P. , Kang, H.K. and Lee, Y. et al. (1988):

Proportionate mortality study of US Army and US Marine Corps veterans of the Vietnam war.

J. Occup Med 30:412.

Budach, W.; Budach, V.; Dinges, S.; Stuschke, M. and Sack; H. (1997):

Correlation between primary chemical and radiation sensitivity in panel of highly malignant human soft tissue sarcoma grafts .

Radiother Oncol. , 42 : 181-187.

Burke , A.P. and Virmani, R. (1993):

Sarcomas of the great vessels : a clinicopathologic study.

Cancer 71:1761.

Burns, W.A., Kanhouwand, S. and Tillman, L. et al. (1972):

Fibrosarcoma occurring at the site of a plastic vascular graft.

Cancer 29:66.

Cardesa, A., Ribalta, T. and Von Schilling, B. et al. (1989):

Experimental model of tumors associated with neurofibromatosis.

Cancer 63:1737.

Carter, R.L., Mccarthy, K.P,m and Machin, L.G. (1989):

Éxpression of desmin and myoglobin in rhabdomyosarcomas and in developing skeletal muscle .

Histopathology. 15 : 585 – 595.

Cavazzana, A.O., Schmidt, D and Ninfo, V. et al. (1992):

Spindle cell rhabdomyosarcoma : a prognostically favorable variant of rhabdomyosarcoma.

Am. J. Surg. Pathol. 16:229.

Carstens, P.H.B. (1986):

Immunohistochemistry and diagnostic electron microscopy (editorial).
Ultrastructural.
Pathol. 10 : 111 .

Chang A.E., Schancer, E.G. and Conkel, D.M. et al. (1979):

Evaluation of computed tomography in the detection of pulmonary metastases.
Cancer 43:913-916.

Chang, Y., Cesarman, E. and Pessin, M. et al. (1994):

Identification of herpesvirus-like DNA-sequences in AIDS-associated Kaposi's sarcoma.
Science 266:1865.

Chase, D.R. and Enzinger, F.M. (1985):

Epithelioid sarcoma : diagnosis. Prognostic indicators. And treatment.
Am. J. Surg. Pathol ; 9:241-63.

Choudhury , L.N.; Swerdlow, M. and Wellington, J. et al. (1980):

Post irradiation malignant fibrous histiocytoma of the lung.
Demonstration of alpha-1- antitrypsin like material in neoplastic cells.
Am. J. Clin Pathol. 74 : 820 – 826 .

Chung, E.B. and Enzinger, F.M. (1976):

Infantile fibrosarcoma .
Cancer 38:729.

Churg, A. and Ringus, J. (1978):

Ultrastructural observations on the histogenesis of alveolar rhabdomyosarcoma.

Cancer 41:1355.

Coindre, J.M., Trojani, M. and Contesso, G. et al. (1986):

Reproducibility of a histopathologic grading system for adult soft tissue sarcoma.

Cancer 58:306.

Collins, J.J. Strauss, M.E. and Levinskas, G.J. et al. (1993):

The mortality experience of workers exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin in a tricholophenol process accident.

Epidemiology 4:7.

Copper, C.S. and Stratton, M.R. (1991):

Soft tissue tumours : The genetic basis of development Carcinogenesis
12 : 155 – 165 .

Copeland, L.J., Sneige, N. and Stringer a, et al. (1985):

Alveolar rhabdomyosarcoma of the female genitalia.

Cancer 56:849.

Costa, J., Wesley, R.A. and Glatstein, E. et al. (1984):

The grading of soft tissue sarcoma : results of a clinicohistopathologic correlation in a series of 163 cases.

Cancer 53:530.

Corson, J.M., Weiss, L.M., Banks-Schlegel, S.P. and Pinkus, G.S. (1984):
Keratin proteins and carcinoembryonic antigen in synovial sarcomas :
an immunohistochemical study of 24 cases.
Hum. pathol. , 15:615-21.

Corzat, A., Aman, P. and Mandahl, N. et al. (1993):
Fusion of CHOP to a novel RNA-binding proteins in human myxoid
liposarcoma.
Nature 363:640.

Craighead, J.E. and Mossman, B.T. (1982):
The pathogenesis of asbestos associated diseases.
N. Engl. J. Med. 306:1446.

Crocker, J. and Egan, M.J. (1988):
Correlation between NOR sizes and numbers in non – Hodgkin's
lymphoma .
J. Pathol. 156 : 233 – 239 .

Crocker, J. and Nar, P. (1987):
Nucleolar organizes regions in lymphomas.
J. Pathol. 151 : 111-118.

Crocker, J., McCarthey, J.C. and Smith, P.G. (1988):
Correlation between D.N.A. Flow cytometric and molecular organizer
region data in non Hodigkin's .
lymphoma Pathol. 154 : 151-160.

Crocker, J., Boldy, D.H.R. and Egan, M.J. (1989):
How should we count AgNORs ? proposals for a standardized
approach.

J. Path. 158:185-188.

Cutler, S.J., Young, I.L. (1975):

Third National Cancer Survey : incidence data.
NCI Monogr 41:1 .

Dahl, I. and Angervall, L. (1974):

Cutaneous and subcutaneous leiomyosarcoma – a clinicopathologic study of 47 patients.
Pathologia Eur. 9:307.

Dal Cin, P., Brock, P. Casteels-Yan and Daele, M. et al. (1991):

Cytogenetic characterization of congenital or infantile fibrosarcoma.
Eur. J. Pediatr. 150 : 579.

Dalcin, P.; Sciote, R.; Fletcher, M.; Samson, I.; De Vos, R.; Mandahl, N.;

Willen, H.; Larsson, O. and Van De Berghe, H. (1998):

Inflammatory leiomyosarcoma may be characterized by specific near – haploid chromosomal changes.
J. Pathol. 185 : 112-115.

Damjanovic, I. And Linder, J. (1996):

Anderson's pathology.
Tenth Ed. Mosby company pp. 2489 – 2505.

Dawson – Saunders, B. and Trapp, P.G. (1990):

Basic and clinical biostatistics, ds.
Appleton and Lange .

Daya , D.A. and Scully, R.E. (1988):

Sarcoma botryoides of the uterine cervix in young women : a clinicopathologic study of 13 cases.
Gynecol oncol 229:290.

Dehner, L.P. (1988):

Malignant fibrous histiocytoma : nonspecific morphologic pattern.
Specific pathologic entity or both ? (editorial).
Arch pathol Lab Med. 112:236.

DeJong, A.S., Van Kessel-Van, M. and Albus-Lutter, C.E. (1987):

Pleomorphic rhabdomyosarcoma in adults : immunohistochemistry as a tool for its diagnosis.
Hum Pathol , 18:298.

Delattre, Zucman, J. and Plougastel, S. et al. (1992):

Gene fusion with an LTR DNA-binding domain caused by chromosome translocation in human tumors.
Nature 359 : 162 .

De Leeuw, B., Suijkerbuijk, R.F. and Balemans, M. et al. (1993):

Sublocalization of synovial sarcoma - associated t (X ; 18) chromosomal breakpoint in X P 11. 2 using cosmid cloning and in situ hybridization.
Oncogene 8 : 1457 .

Derenzini, M., Betts, C.M. Ceccarelli, C. and Eusebi, V. (1986):

Ultrastructural organization of nucleoli in benign naevi and malignant melanomas.
Virchows Arch B Cell Pathol, 5:343-352.

Dervan, P.A.; Gilmartin, L.G.; Loftus, B.M. and Carney, D.N. (1989):
Argyrophilic Nucleoles organizer regions counts correlate with Ki. 67
scors.
Am. J. Clin Pathol. 92 : 401 – 407.

Dias, P.; Kumar, P. and Marsden, H.B (1987):
Evaluation of desmin as a diagnostic and prognostic marker of
childhood rhabdomyosarcoma and embryonal sarcomas .
Br. J. Cancer 56 : 361 – 365 .

**Dias, P.; Parham, D.M.; Shapiro, D.N.; Webber, B.L. and Houghton, P.J.
(1990):**
Myogenicregulatory Protein (myoDI) expression in childhood solid
tumors : utility in rhabdomyosarcoma.
Am. J. Pathol. 137:1283-1291.

Diaz-Cascajo, C., Weyers, W., Rey – Lopez, A. and Bordhis, S. (1998):
Deep dermatofibrosarcoma protubrance : a subcutaneous variant .
Histopathology 32 : 552-555.

Donner, L., De Lanerolle, P. and Costa, J. (1983):
Immunoreactivity of paraffin embedded normal tissues and
mesenchymal tumors of smooth muscle origin.
Am. J. Clin Pathol. 80 : 667-681.

Downing, J.R.; Head, D.R. and Parham, D.M. et al. (1993):
Detection of (11 ; 22) (q 24 ; q 12) translocation of Ewing's sarcoma
and peripheral neuro-ectodermal tumor by reverse transcription
polymerase chain reaction.
Am. J. pathol. 143 : 1294-1300.

Drury, R.A.B. and Wallington, E.A. (1980):

Carletons Histologic techniques 5th ed. Oxford, New York. Oxford University Press 237-247.

Du Boulay, C.E.H. (1982 A) :

Demonstration of alpha-1- antitrypsin and alpa-1-antichymotrypsin in fibrous histocytomas using the immunoperoxidase technique.
Am. J. Surg. Pathol. 6:559-564.

Du Boulay, C.E.H. (1982 B):

Demonstration of fibronectin in soft tissue tumors using the immunoperoxidase .
Diag. Histopathol. 5:283-288.

Du Boulay, C.E.M. (1985):

Immunohistochemistry of soft tissue tumors : A review .
J. Pathol. 146 : 77-94.

Edel, G., Wuisman, P. and Erlemann, R. (1993):

Spindle cell (leiomyomatous) rhabdomyosarcoma, a rare variant of embryonal rhabdomyosarcoma.
Pathol. Res. Pract. 189 : 102 .

Egan, M. and Smith, K. (1987) :

Expression of S 100 protein in breast lesions.
J. Clin. Patho. 40 : 1485 – 1486.

Egan, M., Raafat, F., Crocker, J. and Smith, K. (1988):

Nucleolar organizer regions in fibrous proliferations of childhood and infantile fibrosarcoma .

J. Clin. Pathol. 41 : 31 – 33 .

Enjoji, M., Hashimoto, H. and Iwaski, H., (1980):

Malignant fibrous histocytoma : a clinicopathologic study of 130 cases.

Acta Pathol. Jpn. 30 : 727 .

Enterine, H.T. (1981):

Histopathology of sarcomas.

Seminars in Oncology 8 : 133-155.

Enzinger, F.M. (1977):

Recent developments in the classification of soft tissue sarcoma. In :

Management of primary bone and soft tissue sarcomas.

Chicago. Year book medical publishers.

Enzinger, F.M. (1986):

Malignant fibrous histocytoma 20 years after Stout.

Am. J. Surg. Pathol 10 (suppl 1) : 43 .

Enzinger, F.M. and Shiraki, M. (1969):

Alveolar rhabdomyosarcoma : an analysis of 110 cases.

Cancer 24 : 18.

Enzinger, F.M. and Weiss, S.W. (1983):

Soft tissue tumors .

The C.V. Mosby Company, 166 – 198 .

Enzinger, F.M. and Weiss, S.W. (1996):

Soft tissue tumors 3rd ed. St Louis.

Toronto. London : CV Mosby Company pp. 1 – 15; 269 – 270; 351 – 359; 341; 438 ; 493 ; 545 .

Erlandson, R.A. Woodruff, J.M. (1982):

Peripheral nerve sheath tumors. AN electron microscopic study of 43 cases.

Cancer 49 : 273-87.

Escalona-Zapata J. (1981):

Tumors of the soft tissue : cytology and growth pattern of fibrosarcomas and related tumors.

Pathologica 73 : 119.

Evans, H.L. (1979):

Liposarcoma : a study of 55 cases with a reassessment of its classification.

Am. J. Surg pathol. 3:507.

Evans, D.M. Williams, W.J. and Jung, I.T. (1983):

Angiosarcoma and hepatocellular carcinoma in vinyl chloride workers.

Histopathology 7 : 377.

Evans, H.L. (1985):

Smooth muscle tumors of the gastrointestinal tract : a study of 56 cases followed for a minimum of 10 years.

Cancer 56 : 2242.

Evans, H.L. (1993):

Low grade fibromyxoid sarcoma : a report of 12 cases.

Am. J. Surg pathol. 17 : 595.

Feldman, F. and Lattes, R. (1977):

Primary malignant fibrous histiocytoma (fibrous xanthoma) of bone.
Skeletal Radiol 1 : 145.

Fields, J.P. and Helwig, E.B. (1981):

Leiomyosarcoma of the skin and subcutaneous tissue.
Cancer 47 : 156.

Field, D.H., Filzgerald, P.H. and Sin F.Y.T. (1984):

Nucicolar silver – staining patterns related to cell cycle phase and cell generation of PHA-stimulated human lymphocytes.
Cytobias 41 : 23-33 .

Fisher, C. (1990):

The value of electronmicroscopy and immunohistochemistry in the diagnosis of soft tissue sarcomas : a study of 200 cases.
Histopathology 16:144-55.

Fisher, C. (1996):

Soft tissue sarcoma : diagnosis, classification and prognostic factors.
Br J. plasturg. 49 : 27-33 .

Fletcher C.D. (1987):

Soft tissue sarcomas apparently arising in chronic tropical ulcers,
Histopathology 11 : 501.

Gabbiani, G.; Kaparrci, Y. and Barazzone, P. (1981):

Immunohistochemical identification of intermediate filament sized filaments in human neoplastic cells. A diagnostic aid for the surgical pathologist .

Am. J. Pathol. 104 : 206 – 216.

Gaffney, E.F., Dervan, P.A. and Fletcher, C.D.M. (1993):

Pleomorphic rhabdomyosarcoma in adulthood : analysis of 11 cases with definition of diagnostic criteria.

Am. J. Surg. Pathol. 17 : 601 .

Gerdes, J., Lenke, H. and Baishc, H. et al. (1984):

Cell cycle analysis of a cell proliferation associated human nuclear antigen defined by the monoclonal antibody Ki-67.

J. Immunol. 133 : 1710-1715.

Goodlad, F.R., Mentzel, T. and Fltcher, C.D.M. (1995):

Low grade fibromyxoid sarcoma : clinicopathological analysis of eleven new cases in support of a distinct entity.

Histopathology 26 : 229-237.

Gonzalez-Crussi, F., DeMello, D.E. and Sotelo-Avila, C. (1983):

Omental-mesenteric myxoid hamartomas.

Am. J. Surg. Pathol. 7 : 567.

Guccion, J.G., Enzinger, F.M. (1972) :

Malignant giant cell tumor of soft parts : an analysis of 32 cases.

Cancer 29 : 1518 .

Gusterson, B.A., Cowley, G. and McIlhinney, J. et al. (1985):

Evidence for increases epidermal growth factor receptors in human sarcomas.

International Journal of Cancer 36 : 689-693.

Hachisuga, T., Hashimoto, H. and Enjoji, M. (1984):
Angioleiomyoma : a clinicopathologic reappraisal of 562 cases.
Cancer 54 : 126 .

Hardy, T.J. An T. Brown, P.W. and Terz, J.J. (1978):
Postirradiation sarcoma (malignant histiocytoma) of axilla.
Cancer 42 : 118-124 .

Harms, D. Schmidt, D. and Treuner, J. (1985):
Soft tissue sarcomas in childhood : a study of 262 cases including 169
cases rhabdomyosarcoma.
Z. Kinderchir 40 : 140 .

Harris, M. (1981) :
Differential diagnosis of spindle cell tumors by electron microscopy –
personal experience and a review.
Histopathology 5 : 81 – 105.

Hashimoto, H. and Enjoji, M. (1982):
Liposarcoma : a clinicopathologic subtyping of 52 cases.
Acta Pathol. Jpn. 32 : 933 .

Hashimoto, H., Tsuneyoshi, M. and Enjoji, M. (1985):
Malignant smooth muscle tumors of the retroperitoneum and mesentery
: a clinicopathologic analysis of 44 cases.
J. Surg. Oncol. 28 : 177.

Hashimoto, H., Daimaru, H., Tsuneyoshi, M. and Enjoji, M. (1986):

Leiomyosarcoma of the external soft tissues : a clinicopathologic, immunohistochemical and electron microscopic study.
Cancer 57 : 2077.

Hatfield, P.M. and Schulz, M.D. (1970):
Postirradiation sarcoma. Including 5 cases after therapy of breast carcinoma.
Radiology 96 : 593-602.

Hays, D.M. Shimada, H. and Raney, R.B. Jr. et al. (1988):
Clinical staging treatment results in rhabdomyosarcoma of the female genital tract among children and adolescents.
Cancer 61 : 1893 .

Heerde, P.; van, Felkamp, C.A. and Hart, A.A.M. (1984):
Malignant histiocytosis and related tumors. A clinico - pathological study of 42 cases using cytological, histochemical and ultrastructural parameters.
Hematol Oncology, 2 : 13 - 32 .

Henricks, W.H., Chu- Y.C., Goldbelun, J.R. and Weiss, S.W. (1997):
Dedifferentiated liposarcoma : a clinicopathological analysis of 155 cases with a proposal for an expanded definition of dedifferentiation .
Am. J. Surg. Pathol. 21 : 271 - 281 .

Hernandez - Verdun, D. Hubert J., Bourgeois, C.A. and Bouteille, M. (1980):
Ultrastructural localization of AgNOR stained proteins in the nucleolus during the cell cycle and in other nucleolar structures.
Chromasoma 79 : 349-362 .

Herrera, G.A. and De Morases, H.P. (1983):

Neurogenic sarcomas in patients with neurofibromatosis. Light, electron microscopy and immunohistochemical study.

Virchows Arch. (Path Anat) 401 : 361-376.

Hirose, T. Sano, T. and Hizawa, K. (1988):

Ultrastructural study of the myxoid area of malignant fibrous histiocytomas.

Ultrastruct Pathol. 12 : 621.

Hoffman, M.A. and Dickersin, G.R. (1983):

Malignant fibrous histiocytoma : an ultrastructural study of eleven cases.

Hum. Pathol. 14 : 913 – 922.

Holund, B., Clemmensen, I., Junker, P. and Lyon, H. (1982):

Fibronectin in experimental granulation tissue.

Acta Patho. Microbiol Immunol Scand 90 : 159-165.

Hsu, S.M.; Raine, L. and Fanger, H. (1981):

Use of Avidin – Biotin Peroxidase complex (ABC) in immunoperoxidase techniques : a comparison between ABC and unlabeled antibody (PAP) procedures .

J. Histochem. Cytochem. 29 : 577 – 588.

Hvolris, J.J., Edvadsen, L. and Hansen, L.B. (1997):

Lipomas and liposarcomas 10 years incidence in a local areas.

Ugeskr – Laeger 3 ; 159 : 736 – 739.

Huang, W.L., Ordonez, N.G. and Mackay, B. (1989):

Myxoid malignant fibrous histiocyoma with erythrophagocytosis .
Utrastruct Pathol. 13 : 315 .

Isoda, M. and Yasumoto, S. (1986):

Eosinophil chemotactic factor derived from a malignant fibrous
histiocyoma.
Clin Exp. Dermatol. 11:253.

Iwasaki, H. and Enjoji, M. (1979):

Infantile and adult fibrosarcomas of the soft tissues.
Acta. Pathol. Jpn. 29 : 377 .

**Jenning, T.A.; Peterson, L.; Axiotis, C.A.; Friedlander, G.E.; Cooke, R.A.
and Rosai, J. (1988):**

Angiosarcoma associated with foreign body material .
Cancer 62 : 2436 – 2444 .

Jones, M.W., Norris, H.J. and Wargotz, E.S. et al. (1992):

Fibrosarcoma – malignant fibrous histocyoma of the breast : a
clinicopathological study of 32 cases.
Am. J. Surg. Pathol 16 : 667 .

Kahn, L.B. (1973) :

Retroperitoneal xanthogranuloma and xanthosarcoma (malignant
fibrous xanthoma).
Cancer 31 : 411.

Kahn, H.J., Huang, S.N. and Hanna, W.M. et al. (1982):

Immunohistochemical localization of epidermal and Mallory body
cytokeratin in undifferentiated epithelial tumors.
Am. J. Clin Pathol 81 : 184-191.

Kamat, M.R., Kulkarni, J.N. and Tongaonkar, H.B. et al. (1991):
Rhabdomyosarcoma of the bladder and prostate in children.
J. Surg. Oncol. 48 : 180.

Kasolo, F.C., Mpablwani, E. and Gompels, U.A. (1997):
Infection with AIDS – related hyperviruses – in human immuno
deficiency virus negative infants and evidence childhood kaposi's
sarcoma in Africa .
J. Gen. Virol. 78 : 847-855 .

Kay, S. and McNeill, D.D. (1969):
Leiomyosarcoma of the retroperitoneum.
Surg. Gynecol. Obstet. 129 : 285 .

Kelly, S.J. and Guidotti, T.I. (1989 - 1990):
Phenoxyacetic herbicides and chlorophenols and the etiology of
lymphoma and soft tissue neoplasms.
Publ. Health Rev. 17 : 1.

Kevorkian, J. Cento J.P. (1973):
Leiomyosarcoma of large arteries and veins.
Surgery 73 : 39 .

Kindblom, L.G., Angervall, L. and Svendsen, P. (1975):
Liposarcoma : a clinicopathologic , radiographic, and prognostic study.
Acta Pathol Microbiol Scand 253 : (suppl) 1 .

Kindblom, L.G. Seidal, T. and Karlsson, K. (1982 a) :

Immunohistochemical localization of myoglobin in human muscle tissue and embryonic and alveolar rhabdomyosarcoma.
Acta. Pathol. Microbiol. Immunol. Scand (A) 90 : 167-174 .

Kindblom, L.G., Jacobsen, G.K. and Jacobsen, M. (1982 b) :

Immunohistochemical investigations of tumors of supposed fibroblastic-histiocytic origin.
Hum Pathol 13 : 834-840.

Kraus, M.D., Guillou, L. and Fletcher, C.D. (1997) :

Well differentiated liposarcoma : an uncommon early over looked variant of a common sarcoma.
Am. J. Surg. Pathol. May, 12 : 518-27.

Kuratsu, S. Myoui, A. and Tomita, Y. et al. (1993):

Usefulness of argyrophilic nucleolar organizer staining for histologic grading of soft tissue sarcomas .
J. Surg. Oncol. 54 : 139 .

Kyriakos, M., Kempson, R.L. (1976):

Inflammatory fibrous histiocytoma : an aggressive and lethal lesion,
Cancer 37 : 1584 .

Lack, E.E. Perez-Atayde, A.R. and Schuster, S.R. (1981):

Botryoid rhabdomyosarcoma of the biliary tree.
Am. J. Surg. Pathol. 5 : 143 .

Lack, E.E., Steinbery, S.M. White, D.E., Kinsella, T. and Glaststen, E.

(1989):

Extremity of soft tissue sarcomas : analysis of prognostic variables in 300 cases and evaluation of tumor necrosis as a factor in stratifying higher - Grade sarcomas .

J. Surg. Oncol. 41 : 263 - 273 .

Lagace , R., Delage, C. and Seeemayer, T.A. (1979):

Myxoid variant of malignant fibrous histiocyoma : ultrastuctural observations.

Cancer 43 : 526 .

Lagace, R.; Schurch, W. and Seemayer, T.A. (1980):

Myofibroblasts in soft tissue sarcomas.

Virchows Archiv. (Pathol. Anat. Histol.) 389 : 1-11.

Laskin, W.B., Silverman, T.A. and Enzinger, F.M. (1991):

Postirradiation soft tissue sarcomas : an analysis of 53 cases.

Cancer 68 : 524 .

Lattes, R. (1982):

Malignant fibrous histiocyoma. A review article.

Am. J. Surg. Pathol. 6 : 761-771.

Lawson, C.W.; Fisher, C. and Gatter, K. (1987):

An immunohistochemical study of differentiation in malignant fibrous histocyoma.

Histopathology, 11 : 375 - 383 .

Leong, A.S.-Y. and Gilham, P. (1989):

Sliver staining of nucleolar organizer regions in malignant melanoma and melanotic nevi .
Hum. Pathol. 20 : 257 – 262 .

Leu, H. G. and Makek, M. (1986):
Intramural venous leiomyosarcomas.
Cancer 57 : 1395 .

Leuschner, I., Newton, W.A. Jr. and Schmidt, D. et al. (1993):
Spindle cell variants of embryonal rhabdomyosarcoma in the paratesticular region : a report of the Intergroup Rhabdomyosarcoma Study.
Am. J. Surg. Pathol. 17 : 221 .

Lloyd, R.V., Hajdu, S.I. and Knapper, W.H. (1983):
Embryonal rhabdomyosarcoma in adults.
Cancer 51 : 557 .

Lyng, E. Storm, H.H. and Gensen, O.M. (1987):
The evaluation of trends in soft tissue sarcoma according to diagnostic criteria and consumption of phytochemicals.
Cancer 60 : 1896 – 1901 .

Lundgren, L., Angervall, L. and Stenman, G. et al. (1993):
Infantile rhabdomyofibrosarcoma : a high grade sarcoma distinguishable from infantile fibrosarcoma and rhabdomyosarcoma .
Hum. Pathol. 24 : 785.

Markhede, G., Angervall, L. and Stener, B. (1982):

A multivariate analysis of the prognosis after surgical treatment of malignant soft tissue tumors.
Cancer 49 : 1721 .

Martin, R.G. et al. (1976) :

Soft tissue sarcoma . In Clark, R.L. and Howe, C.D., editors : Cancer patient care at M.D. Anderson .Hospital and Tumor Institute.
The University of Texas. Chicago year book Medical Publishers. Inc.

Mason, D.Y. and Taylor, C.R. (1975):

The distribution of murimadase (lysozyme) in human tissues.
J. Clin. Pathol. 28 : 124 – 132.

MaClay, E.F. (1989):

Epidemiology of bone and soft tissue sarcomas.
Semin Oncol 16 : 264.

Meis J.M. and Enzinger, F.M. (1991):

Inflammatory fibrosarcoma of the mesentery and retroperitoneum : a tumor closely simulating inflammatory pseudotumor.
Am. J. Surg. Pathol 15 : 1146 .

Meister, P. (1988):

Malignant fibrous histiocytoma : history, histology, histogenesis .
Pathol. Res. Pract 183 : 1 .

Meister, P. (1995):

Malignant fibrous histiocytoma : a “ fibrohistiocytic “ or primitive, fibroblastic sarcoma. In : Harms D. Schmidt D. eds Current topics in pathology. Vol. 89. Berlin, Heidelberg.

Springer – Verlag, 193 – 214 .

Mendelsohn, G.; Eggleston, J.C. and Mann, R.B. (1980):

Relation ship of lysozyme to histocytic differentiation in malignant histocytosis .

Cancer , 45 : 237 – 279 .

Mentzel, T., Katenkamp, – D. and Fletcher C.D. (1996):

Low malignancy myxofibrosarcoma versus low malignancy fibromxoid sarcoma.

Patho. 17 : 116 – 121 .

Mierau, G.W. and Favara, B.E. (1980):

Rhabdomyosarcoma in children : ultrastructural study of 31 cases.

Cancer 46 : 2035.

Miettinen, M., Lehto, V.P. Badely, R.A. and Virtanen, I. (1982):

Alveolar rhabdomyosarcoma. Demonstration of the muscle type of intermediate filament protein desmin as a diagnostic aid.

Am. J. Pathol 108 : 246 – 251 .

Miettinen, M., Foidart, J.M. and Ekblom, P. (1983):

Immunohistochemical demonstration of laminin the major glycoprotein of basement membranes as an aid in the diagnosis of soft tissue tumor.

Am. J. Clin Pathol 79 : 306 – 311 .

Miettinen, M. (1988):

Rhabdomyosarcoma in patients older than 40 years of age.

Cancer 62 : 2060 .

Miettinen, M. (1990):

Immunohistochemistry of soft tissue tumors : possibilities and limitations in surgical pathology.

Pathol. Annu 23 (Part 1) : 1 .

Mingoli, A.; Cavallaro, A.; Sapienza, P.; Di – Marzo L.; Feldhaus, R.J. and Cavallar, N. (1995):

International registry of inferior vena cava leiomyosarcoma analysis of a world series on 218 patients.

Anticancer – Res. 16 : 3201 – 3205 .

Mirra, J.M., Gold, R.H. and Marafioti, R. (1977):

Malignant (fibrous) histiocytoma arising in association with a bone infarct in sickle cell disease : coincidence or cause – and – effect ?

Cancer 39 : 186 .

Mokhtar, N. (1991):

Cancer Pathology registry 1985 – 1989 .

National Cancer Institute . Cairo University . Mokhtar N.(ed.) pp.54–61

Molenaar, W.M., Oosterhuis, J.W. and Kamps, W.A. (1984):

Cytological “ differentiation “ in childhood rhabdomyosarcoma following polychemotherapy.

Hum. Pathol. 15 : 973 .

Monckman, G.R., Orwoll, G. and Ivins, J.C. (1974):

Trauma and oncogenesis.

Mayo. Clin. Proc. 157 – 163.

Moreland, R.B. and McNamara, W.L. (1942):

Liposarcoma : report of nine cases.

Arch. Surg. 45 : 164 .

Morimitsu, Y.; Hisaoka, M.; Okamoto, S.; Hashimoto, H. and Ushijima, M. (1998):

Desomatofyrosarcoma protubance and its fibrosarcomatus variant with areas of myxoid differentiation : a report of three cases.

Histopathology 32 : 547 – 551 .

Mukai, K., Rosai J. and Burgdorf W.H.C. (1980):

Localization of factor VIII rag in vascular cells using an immunoperoxidase method.

Am. J. Surg. Pathol. 4 : 273 – 276 .

Myhre Jensen, O., Kaae S. and Madsen, E.H. et al. (1983):

Histopathological grading in soft – tissue tumors : relation to survival in 261 surgically treated patients.

Acta. Pathol. Microbiol. Immuno. Scand 91 A : 145 .

Myhre Jensen, O. Hogh, J. and Ostgaard, S.E. et al. (1991):

Histopathological grading of soft tissue tumors : prognostic significance in a prospective study of 278 consecutive cases.

J. Pathol. 163 : 19 .

Nakamura, S.; Murakami – Mori, K.; Rao N.; Weich, H.A. and Rajeev, B. (1997):

AIDS – associated kaposi's sarcoma – derived spindle cells .

J. immunol 158 : 4992 – 5001 .

- Nakanishi, K.; Hirol, S.; Kawap, T.; Suzuki, M. and Torikata, C. (1998):***
 Argyrophilic nucleolar – organizer region, count and DNA status in
 bronchioloalveolar epithelial hyperplasia and adenocarcinoma of lung.
 Human. Pathol. 29 : 235 – 239 .
- Nakashima, I., Kojiro, M. (1987):***
 Hepatocellular carcinoma : An atlas of its pathology,
 Tokyo : Springer-Verlag.
- Navarro, S., Noguera, R. and D'Orazi, G. et al. (1990):***
 Expression of the Myo D1 gene and protein in various myogenous and
 non-myogenous tumors.
 Laboratory Investigation 62 : 73 A (Abstract).
- Nistal, M., Raniagua, R. and Picazo, M.L. et al. (1980):***
 Granular changes in vascular leiomyosarcoma.
 Virchows Arch (Pathol Anat.) 386 : 239.
- Ochs R.L. and Busch, H. (1984):***
 Further evidence that phosphoprotein C23 clonk D/pt 5,1 is the nucleolar
 silver staining proteins.
 Exp. Cell Res. 152 : 260 – 265 .
- O'Connell, P. Cawthon, R. and Xy, G.H. et al. (1992):***
 The neurofibromatosis type (NF1) gene : identification and partial
 characterization of a putative tumor suppressor gene.
 J. Dermatol 19 : 881 .
- O'Connell, J.X. and Trotter, M.J. (1996):***
 Fibrosarcomatous dermatofibrosarcoma protuberans : a variant.

Mod. Pathol. 9 : 273 – 278 .

Ohno, T. Rao, V.N. Reddy, E.S.P. (1993):
EWS/Fe chimeric proteins is a transcriptional activator.
Cancer Res. 53 : 5859 .

Ooms, E.C.M. and Veldhuizen, R.W. (1989):
Argyrophilic proteins of the nucleolar organizer region in bladder tumors .
Virchows Archiv.(A) Pathol. Anat., 365 .

Orson, G.G., Sim F.H. and Reiman, H.M. et al. (1987):
Liposarcoma of the musculoskeletal system.
Cancer 60 : 1362 .

Osborn, M.; Debused, E. and Weber, K. (1984):
Monoclonal antibodies specific for vimentin .
Eur. J. Cell Biol., 34 : 137 .

Pack, G.T. and Ariel, I.M. (1952):
Fibrosarcoma of the soft somatic tissues : a clinical and pathologic study.
Surgery. 31 : 443.

Patton, R.B. and Horn, R.C. Jr. (1962):
Rhabdomyosarcoma : clinical and pathological features and comparison with human fetal and embryonal skeletal muscle.
Surgery 52 : 572 .

Perosio, P.M. and Brooks, J.J. (1988):

Expression of nerve growth factor receptor in paraffin embedded soft tissue tumors.

Am. J. of Pathol. 132 : 152 – 160 .

Perosio, P.M. and Brooks, J.J. (1989):

Expression of growth factor and growth factor receptor in soft tissue tumors.

Implications for the autocrine hypothesis.

Laboratory Investigation 60 : 245 – 253.

Pezzi, C.M. Rawlings, M.S. Jr. and Esgro, J.J. et al. (1992):

Prognostic factors in 227 patients with malignant fibrous histiocytoma.

Cancer 69 : 2098.

Pierce, E.R., Veisbord, T. and Mckasick, V.A. (1970):

Gardner's syndrome formal genetics and statistical analysis of large canadian .

Kindred. Clin. Genet 1 : 65.

Pilotti, S., Torre, G.D. and Lavarions, C. et al. (1998):

Molecular abnormalities in liposarcoma : rol of MDN2 and CDK4 – containing amplicons at 12413 – 22.

J. Pathol. 185 : 188 – 190 .

Ploton, D.; Menager, M.; Jennesson, P.; Himber, G.; Pigeon, J. and Adnet, J.J. (1986):

Improvement in the staining and in the visualization of the argyrophilic proteins of the nucleolar organizer region at the optical level.

Histochem. J. 18 : 5 – 14 .

Popper, H., Thomas, L.B. and Telles, N.C. et al. (1978):

Development of hepatic angiosarcoma in man induced by vinyl chloride, Thorotrast, and arsenic : comparison with cases of unknown etiology.

Am. J. Pathol. 92 : 349 .

Pratt, C.B., Hustu, H.O. and Kumar, A.P.M. et al. (1981):

Treatment of childhood rhabdomyosarcoma at St. Jude Children's Research Hospital 1962 – 1978.

Natl Cancer Inst. Monogr. 56 : 93 .

Pritchard, D.J., Soule, E.H. and Taylor, W.F. et al. (1974):

Fibrosarcoma : a clinicopathologic and statistical study of 199 tumors of the soft tissues of the extremities and trunk.

Cancer 33 : 888 .

Riccardi, V. (1981):

Von Recklinghausen's neurofibromatosis. N. Engl. J. Med. 305 : 1617 – 26 .

Robinson, E., Neugut, A.I. and Wylie, P. (1988):

Clinical aspects of postirradiation sarcoma.

J. Natl Cancer Inst. 80 : 233 .

Rodary, C., Gelhan, E.A. and Flamant, F. et al. (1991):

Prognostic factors in 951 nonmetastatic rhabdomyosarcomas in children : a report from the international rhabdomyosarcoma workshop.

Med. Pediatr. Oncol. 19 : 89 .

Roholl, P.J.M. Kleyne, J., and Van Basten, C.D.H. et al. (1985):

A study to analyze the tumor cells in malignant fibrous histiocytomas :
a multiparametric characterization.
Cancer 56 : 2809 .

Rooser, B., Willen, H. and Gustafson, P. et al. (1991):

Malignant fibrous histiocytoma of soft tissue : a population – based
epidemiologic and prognostic study of 137 patients.
Cancer 67 : 499 .

Ross, J.A., Severson, R.K., Davis, S. and Brooks, J.J. (1993):

Trends in the incidence of soft tissue sarcomas in the united states from
1973 through 1987.
Cancer 72 : 486 .

Rubin, B.P. and Fletcher, C.D.M. (1997):

The cytogenetics of lipomatous tumors.
Histopathology 30 : 507 – 511 .

Russell, W.O., Cohen, J. and Enzinger, F.M. et al. (1977):

A clinical and pathological staging system for soft tissue sarcomas.
Cancer 40 : 1562 .

Salm, R. and Evans, D.J. (1985):

Myxoid leiomyosarcoma.
Histopathology 9 : 159 .

Schmidt, D., Reimann, O. and Treuner, J. et al. (1986):

Cellular differentiation and prognosis in embryonal rhabdomyosarcoma : a report from the Co-operative soft tissue sarcoma study 1981 (CWS 81). Virchows. Arch (Pathol. Anat) 409 : 183 .

Schmookler, B.M. and Enzinger, F.M. (1981):

Pelomorphie lipoma : A benign tumor simulating liposarcoma.
Cancer 42 : 126 – 133 .

Schmookler, B.M. and Enzinger, F.M. (1983):

Liposarcoma occurring in children : an analysis of 17 cases and review of the literature.
Cancer 52 : 567.

Schmookler, B.M. and Lauer, D.H. (1983):

Retroperitoneal leiomyosarcoma : a clinicopathologic analysis of 36 cases.
Am. J. Surg. Pathol. 7 : 269 .

Sciot, R., Akerman, M. and Dal, C. P. et al. (1997):

Cytogenetic analysis of subcutaneous angiolipoma further evidence supporting its difference from ordinary lipoma. A report of the CHAMP study group.
Am. J. Surg. Pathol. In Press.

Scott, S.M., Reiman, H.M. and Pritchard, D.J. et al. (1989):

Soft tissue fibrosarcoma : a clinicopathologic study of 132 cases.
Cancer 64 : 925 .

Scrabble, H., Witte, D. and Shimada, H. et al. (1989):

Molecular differential pathology of rhabdomyosarcoma.
Genes, Chromosomes and Cancer 1 : 23 – 35 .

Selikoff, I.J., Churg, J. and Hammond, E.C. (1964):

Asbestos exposure and neoplasia.

JAMA 288 : 22 .

Serke, S. Brenner, M. and Zimmerman, R. et al. (1986):

Malignant fibrous histiocytoma associated with peripheral blood eosinophilia : in vitro studies demonstrating tumor – derived eosinophilopoietic activity.

Oncology 43 : 230 .

Shipley, J. and Fisher, C. (1998):

Chromosomal translocation in sarcomas and the analysis of paraffin – embedded material.

J. pathol. 184 : 1 – 3 .

Sorrenti, S.J., Cumming, W.J. and Miller, D. (1984):

Reaction of the human tibia to bone wax.

Clin. Orthop. 182 : 293 – 296 .

Sotello – Avila, C., Conzaes – Crussi, F. and de Mello, D. et al. (1986):

Renal and extra renal rhabdoid tumors in children : a clinicopathologic study of 14 patients .

Semin Diagn. Pathol. 3 : 151 .

Smith, R. and Crocker, J. (1988):

Evaluation of nucleolar organizer region associated proteins in breast malignancy.

Histopathology, 12 : 113 – 125 .

Smith, T.A., Easley K.A. and Gddblam, J.R. (1996):

Myxoid / round cell liposarcoma of the extyermities a clinicopathologic study of 29 cases with particular attention to extent of round cell liposarcoma.

Am. J. Sarg. Pathol. 20 : 171 – 1780 .

Snover, D.C., Sumner, H.W. and Dehner, L.P. (1982):

Variability of histologic pattern in recurrent soft tissue sarcomas originally diagnosed as liposarcoma.

Cancer 49 : 1005.

Sreekantaiah, C., Karakousis, C.P., Leong S.P. and Sandberg, A.A. (1991):

Trisomy 8 as a non-random secondary change in myxoid liposarcoma.

Cancer Genet Cytogenet 51 : 195.

Sreekantaiah, C., Karakousis, C.P., Leong, S.P. and Sandberg, A.A. (1992):

Cytogenetic findings in liposarcoma correlate with histopathologic subtypes.

Cancer 69 : 2484.

Stern, R. (1981):

Current concepts in the diagnosis of human soft tissue sarcoma.

Hum. Patho. 12 : 777 – 631 .

Stringer, B.D. (1977):

Leiomyosarcoma of artery and vein.

Am. J. Surg. 134 : 90 .

Strong, L.C., Williams, W.R. and Tainsky, M.A. (1992):

Li-Fraumeni syndrome : from clinical epidemiology to molecular genetics.

Am. J. Epidemiol. 135 – 190 .

Stuart, F.P. and Baker, W.H. (1973):

Palliative surgery for leiomyosarcoma of the inferior vena cava.

Ann. Surg. 177 : 237 .

Suit, H.D., Russell, W.O. and Martin, R.G. (1973):

Management of patients with sarcoma of soft tissue in an extremity.

Cancer 31 : 1247.

Suster, S., Wong, T.Y. and Moran, C.A. (1993):

Sarcomas with combined features of liposarcoma and leiomyosarcoma : study of two cases of an unusual soft-tissue tumor showing dual lineage differentiation.

Am. J. Surg. Pathol. 17 : 905 .

Sutow, W.W. (1981):

Malignant solid tumors in children : a review.

New York., Raven Press.

Sutow, W.W., Lindberg, R.D. and Gehan, E.A. et al. (1982):

Three – year relapse – free survival rates in childhood rhabdomyosarcoma of the head and neck : report from the intergroup rhabdomyosarcoma study.

Cancer 49 : 2217 .

Swanson, P.E., Stanley, M.W. and Scheithauer, B.W. et al. (1988):

Primary cutaneous leiomyosarcoma : a histologic and immunohistochemical study of 9 cases with ultrastructural correlations.
J. Cutan. Patho. 15 : 129 .

Swanson, P.E.; Wick, M.R. and Dehner, et al. (1991):

Leiomyosarcoma of somatic soft tissue of childhood : an immunohistochemical analysis of six cases with ultrastructural correlation .
Hum. Pathol. 22 : 559 .

Swann, M. (1984):

Malignant soft – tissue tumor at the site of a total hip replacement.
J. Bone. Joint. Surg. (Br) 66 – B : 629 – 631 .

Taheri , S.A. and Conner, G.W. (1983):

Leiomyosarcoma of iliac veins.
Surgery 94 : 516 .

Takahashi, K., Toshiaki, I. And Ohtsuki, Y. et al. (1984):

Immunohistochemical localization and distribution of S-100 proteins in the human lymphoreticular system.
Am. J. patho. 116 : 497 – 503.

Takahashi, K., Kimura, Y. and Naito, M. et al. (1989):

Inflammatory fibrous histiocytoma presenting leukemoid reaction.
Pathol. Res. Pract 184 : 498.

Taxy, J.B. and Battifora, H. (1977):

Van Dam, P.A.; Lowe, D.G. and Mckenzie, B. (1990):

Reteroperitoneal soft tissue sarcomas : a review of literature .
Obstet . & Gynecol. 45 : 670 – 682 .

Wang-Wuu, S.; Jacobs D. and Soukup, S. (1990):

Comparison of chromosome analysis to DNA content by flow
cytometry for pediatric tumors.
Pediatr Pathol. 10 : 671 .

**Watanable K.; Hiraki H.; Ohishi M.; Mushiko K.; Saginoya H. and Suzuki
T. (1996):**

Uterine leiomyosarcoma with osteoclast - like giant cell :
histopathological .
Pathol. Int. 46 : 656 – 660 .

Weber, P.C. (1986):

Epithelioid sarcoma in association with total knee replacement : A
cases report.
J. Bone Joint Surg. (Br.) 68 : 824 – 826 .

Weeks, B.S. and Cooney, R. (1997):

AIDS – associated Kaposi's sarcoma cells secrete a neurotrphic growth
factors .
Biochem Biophys. Res. Commun. 233 : 408 – 412 .

Weidner, N. and Tjoe, J. (1994):

Immunohistochemical profile of monoclonal antibody 013 : antibody
that recognizes glycoprotein p30 / 32 MIC2 and is useful in diagnosiing
Ewing's sarcoma and peripheral neuroepithelioma.

Am. J. Surg. Pathol. 18 : 486 – 494.

Weiss, S.W. and Enzinger, F.M. (1977):

Myxoid variant of malignant fibrous histiocytoma .
Cancer 39 : 1672 .

Weiss, S.W. and Enzinger, F.M. (1978):

Malignant fibrous histiocytoma : an analysis of 200 cases.
Cancer 41 : 2250 .

Weiss, S.W., Brattauer, G.L. and Morris, P.A. (1988):

Postirradiation malignant fibrous histiocytoma expressing cytokeratin :
implications for the immunodiagnosis of sarcomas.
Am. J. Surg. Pathol. 12 : 554.

Weiss, S.W. and Rao, V.K. (1992):

Well – differentiated liposarcoma (atypical lipoma) of deep soft tissue
of the extremities, retroperitoneum, and miscellaneous sites : a follow –
up study of 92 cases with analysis of the incidence of
“dedifferentiation” .
Am. J. Surg. Pathol. 16 : 1051 .

Wiatrak, B.J. and Pensak, M.L. (1989):

Rhabdomyosarcoma of the ear and temporal bone.
Laryngoscope 99 : 1188 .

Wiklund, T.A., Blomqvist, C.P. and Raty, J. et al. (1991):

Post irradiation sarcoma : analysis of a nationwide cancer registry
material.
Cancer 68 : 524 .

Wile, A.G., Evans H.L. and Romsdahl, M.M. (1981):

Leiomyosarcoma of soft tissue : a clinicopathologic study.
Cancer 48 : 1022.

Wingren, G. et al. (1990):

Soft tissue sarcoma and occupational exposures.
Cancer 66 : 806.

Young, J.L. Jr. and Miller, B.W. (1975):

Incidence of malignant tumors in U.S. children.
J. pediatr. 86 : 254 .

Yung, P.; Hirose, T.; Hasegawa, T.; Hizawa, K. and San, O.T. (1998) :

Molecular fluorescence in site hybridization analysis of synovial sarcoma .
J. Patho. 184 : 7-13.