

REFERENCES

1- Ahmed, N. and Mukhtar, H. (1999):

Green tea polyphenols and cancer : biologic mechanisms
and practical implications.

Nutr. Rev., 57 : 78 -83.

2- Akoh, C C. and Min, D B. (2008):

Food Lipids. In : Chemistry, Nutrition and Biotechnology.

3rd edition. PP. 615-617.

Marcel Dekker, New York.

3- Alessio, H.M.; Hagerman, A.E.; Romanello, M.; Carando, S.;

**Threlkeld, M.S.; Rogers, J.; Dimitrova, Y.; Muhammed,
S. and Wiley, R.L. (2003):**

Consumption of green tea protects rats from exercise –
induced oxidative stress in kidney and liver.

Nutr. Res., 22 : 1177 – 1188.

4- Al-Okbi, S.Y.; Metwalli, O. M. and Abbas, A. E. (1989):

Effects of Dietary Fibres on Blood Glucose and Liver
Glycogen in Rats.

Arch. Pharm. Res., 12(2): 125-127.

5- Alschuler, L. (1998):

Green Tea : Healing tonic.

Am. J. Natur. Med., 5 : 28 – 31.

- 6- Aneja, A.; El-Alat, F.; McFarlane, S.I. and Sowers, J.R. (2004):**
Hypertension and Obesity.
Recent progress in Hormone Research, 59 : 169 – 205.
- 7- Arkan, M.C.; Hevener, AL.; Greten, F.R.; Maeda. S.; Li, Z. W.;
Long, J. M.; Wynshaw-Boris, A.; Poli, G.; Olefsky, J. and
Karin, M.(2005):**
IKK-beta links inflammation to obesity-induced insulin
resistance.
Nat. Med., 11(2):191-198.
- 8- Bains, R. K.; Wells, S. E.; Flavell, D. M.; Fairhall, K. M.;
Strom, M.; Tissier, P. L. and Robinson, C. (2003):**
Visceral Obesity without Insulin Resistance in Late-Onset
Obesity Rats.
Endocrinology, 145 (6): 2666-2679.
- 9- Bancroft, J.D. (2008):**
Theory & Practice of histological techniques; 6th ed. Ch. 7
and 12. PP. 99 – 101 and 199.
Charchill livingstone of El Sevier, Philadelphia.
- 10- Baumann, C. A.; Ribon, V.; Kanzaki, M.; Thurmond, D.C.;
Mora, S.; Shigematsu, S.; Bickel, P.E.; Pessin, J.E.; and
Saltiel, A.R. (2000):**
CAP defines a second signalling pathway required for
insulin-stimulated glucose transport.
Nature, 407: 202–207.

11- Brown, C.A.; Bolton – Smith, C.; Woodward, M. and Tunstall-Pedoe, H. (1993):

Coffee and tea consumption and the prevalence of coronary heart disease in men and women: results from the Scottish Heart Health Study.

J. Epidemiol. Community health, 47 : 171 -175.

12- Casteilla, L.; Planat-Benard, V.; Cousin, B.; Silvestre, J.S.; Laharrague, P.; Charriere, G.; Carriere, A. and Pendicaud, L. (2005):

Plasticity of adipose tissue : a promising therapeutic avenue in the treatment of cardiovascular and blood disease.

Arch. Mal. Coeur. Vaiss., 98 (9): 922 – 926.

13- Chan, T.P.; Fong, W.P.; Cheung, Y.L.; Ho, W.K. and Chen, Z.Y. (1999):

Jasmine green tea epichatechins are hypolipidemic in hamsters fed a high fat diet.

J. Nutr., 129 : 1049 – 1101.

14- Cormack, H. (1987):

Ham's Histology. 9th ed. PP. 652-668.

Lippincott Co., Philadelphia and London.

15- David, J.; Guo, P.; Nai, S.; Zong, G.; Yge, H.; Hui, C.H.; Jian, Q.; Xue, J.; Bert, C.; Wouters, M. and Jian, Z. (2003):

Cholesterol lowering effect of a the aflavin – enriched green tea extract.

Arch. Intern. Med., 163 : 1448 – 1453.

16- Diamond, F.B. and Eichler, D.C.(2002):

Leptin and the adipocyte endocrine system.

Crit. Rev. Clin. Lab. Sci., 39(4-5):499-525.

17- Dulloo, A.; Duret, C.; Rohrer, D.; Girardier, L.; Mensi, N.; Fathi, M.; Chantre, P. and Vandermander, J. (1999):

Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24 – h energy expenditure and fat oxidation in humans.

Am. J. Clin. Nutr., 70: 1040 – 1045.

18- Dulloo, A.G. (2002):

Sympathetic Defense against Obesity.

Science, 2 : 780-783

19- Fawcett, D.W. (1994):

Bloom and Fawcett. A Textbook of Histology. 12th ed. PP: 170-181.

Chapman and Hall, New York and London.

20- Fujita, T.; Tanaka, K. and Tokunaga, J. (1981):

Scanning electron microscopy, atlas of cells and tissues.

1st ed. Ch. 9. PP. 318 – 332.

Igaku-Shoin Medical Publishers, Tokyo and New York.

21- Furuyashiki, T; Nagayasu, H; Aoki, Y; Bessho, H; Hashimoto, T; Kanazawa, K. and Ashida, H.(2004):

Tea catechin suppresses adipocyte differentiation accompanied by down-regulation of PPA γ 2 and C/EBP α in 3T3-L1 cells.

Biosci. Biotechnol. Biochem., 68(11):2353-2359.

22- Garaulet, M.; Hernandez – Morante, J.J.; Lujan, J.; Tebar, F.J. and Zamora, S. (2006):

Relationship between fat cell size and number and fatty acid composition in adipose tissue from different fat depots in overweight/obese humans.

Int. J. Obes. (Lond), 30 (6) : 899 – 905.

23- Gartner, L. and Hiatt, J. (2007):

The adipose tissue. In : color textbook of histology.

3rd ed. Ch 6. PP. 115 – 117 and 127 – 129.

Elsevier Saunders, Philadelphia.

24- Goldberg, I. J. (1996):

Lipoprotein lipase and lipolysis: central roles in lipoprotein metabolism and atherogenesis.

Journal of Lipid Research, 37: 693-707.

25- Graham, H.N. (1992):

Green tea composition, consumption, and polyphenol chemistry.

Prev. Med., 21 : 334 – 350.

26- Gregoire, F.M.; Smas, C.M. and Sul, H. S. (1998):

Understanding adipocyte differentiation.

Physiol Rev., 78 (3):783-809.

27- Guebre, F.; Bernhard, J.; Funahashi, T.; Hadj-Aissa, A. and Fouque, D. (2005):

Adiponectin in chronic kidney disease is related more to metabolic disturbances than to decline in renal function.
Nephrol. Dial. Transplant., 20(1):129-134.

28- Gunstone, F.D.; Harwood, J.L. and Dijkstra, A.J. (2007):

The lipid handbook. 3rd ed. PP. 220 – 225.
Chapman and hall, New York.

29- Hall, J.E.; Jones, D.w.; Kuo, J.J.; Da-Silva, A.; Tallam, L.S. and Liu, J. (2003):

Impact of the obesity epidemic on hypertension and renal disease.
Curr. Hypertension. Rep., 5 (5): 386 – 392.

30- Hasegawa, N.; Yamda, N. and Mori, M. (2003):

Powdered green tea has antilipogenic effect on Zucker rats fed a high – fat diet.
Phytother. Res., 17 : 477 – 480.

31- Hausman, D. B.; Fine, J. B.; Tagra, K.; Fleming, S.; Martin, R. J. and DiGirolamo, M. (2003):

Regional Fat Pad Growth and Cellularity in Obese Zucker Rats: Modulation by Caloric Restriction.
Obesity Research, 11: 674–682.

32- Hayat, M.A. (2000):

Principles and Techniques of electron microscopy.

4th ed. PP. 96-123.

Cambridge University Press, UK.

33- Juhel, C.; Armand, M.; Pafumi, Y.; Rosier, C.; Vandermader, J. and Lairon, D. (2000):

Green tea extract inhibits lipolysis of triglycerides in gastric and duodenal medium in vitro.

J. Nutr. Biochem., 11 : 45 – 51.

34- Kao, Y.; Hiipakka, R. A. and Liao, S. (2000):

Modulation of obesity by a green tea catechin.

American Journal of Clinical Nutrition, 72 (5): 1232-1233.

35- Klatsky, A.L.; Petitti, D.B.; Armstrong, M.A. and Friedman, G.D. (1985):

Coffee, tea and cholesterol.

Am. J. Cardiol., 55 : 577 – 578.

36- Koutnikova, H. and Auwerx, J. (2001):

Regulation of adipocyte differentiation.

Ann. Med., 33(8):556-561.

37- Langin, D.; Dicker, A.; Tavernier, G.; Hoffstedt, J.; Mairal, A.; Ryden, M.; Arner, E.; Sciard, A.; Jenkins, C.M.; Viguerie, N.; Van-Harmelen, V.; Gross, R.W.; Holm, C. and Arner, P. (2005):

Adipocyte lipases and defect of lipolysis in human obesity.

Diabetes; 54 (11): 3190 – 7.

38- Lee, M. S.; Kim C T; Kim I H and Kim Y. (2009):

Inhibitory effects of green tea catechin on the lipid accumulation in 3T3-L1 adipocytes.

Phytotherapy Research, 23 (8): 1088 – 1091.

39- Lin, Y.S.; Tsai, Y.J.; Tsay, J.S. and Lin, J.K. (2003):

Factors affecting the levels of tea polyphenols and caffeine in tea leaves.

J. Agric. Food Chem., 51 (7) : 1864 -1873.

40- Luft, J.H. (1961):

Improvements in epoxy resin embedding methods.

J. Biophys. Biochem. Cytol., 9: 407-414.

41- Maassen, J. A.; Romijn, J. A. and Heine, R. J.(2007):

Fatty acid-induced mitochondrial uncoupling in adipocytes as a key protective factor against insulin resistance and beta cell dysfunction: a new concept in the pathogenesis of obesity-associated type 2 diabetes mellitus.

Diabetologia, 50 (10): 2036-2041.

42- MacDougald, O.A. and Lane, M.D. (1995):

Adipocyte differentiation. When precursors are also regulators.

Curr. Biol., 15 (6):618-621.

- 43- Marques, B. G.; Hausman, D. B.; Latimer, A.M.; Kras, K.M.; Grossman, B.M. and Martin, R.J. (2000):**

Insulin-like growth factor I mediates high-fat diet-induced adipogenesis in Osborne-Mendel rats.

Am. J. Physiol., 278: R654–R662.

- 44- Matsubara, Y.; Kano, K.; Kondo, D.; Mugishima, H. and Matsumoto, T. (2009):**

Differences in Adipocytokines and Fatty Acid Composition between Two Adipocyte Fractions of Small and Large Cells in High-Fat Diet-Induced Obese Mice.

Ann. Nutr. Metab., 54:258-267.

- 45- Mescher, A.L. (2010):**

Adipose tissue. In : Junqueira's basic histology, text and atlas. 12th ed. PP. 109 – 113.

McGraw – Hill Companies, United States of America.

- 46- Murase, T.; Nagasawa, A.; Suzuki, J.; Hase, T. and Tokimitsu, I. (2002):**

Beneficial effects of tea catechins on diet – induced obesity : stimulation of lipid catabolism in the liver.

Int. J. Obes. Relat. Metab. Disord., 26 : 1459 – 1464.

- 47- Naim, M.; Brand, A. G.; Morley, A. R.; Kare, A. and Carpenter, G. (1985):**

Energy Intake, Weight Gain and Fat Deposition in Rats Fed Flavored, Nutritionally Controlled Diets in a Multichoice "Cafeteria", Design.

J. Nutr., 115: 1447-1458.

48- Nelson, M. and Poulter, J. (2004):

Impact of tea drinking on iron status in the UK : a review.

J. Hum. Nutr. Diet., 17 : 43 -54.

49- Parton, R.G.; Molero, J. C.; Floetenmeyer, M.; Green, K. M. and James, D. E. (2002):

Characterization of a Distinct Plasma Membrane Macrodome in Differentiated Adipocytes.

The Journal of Biological Chemistry, 277(48):46769–46778.

50- Princen, H.; Dyvenvoorde, W.; Buytenhek, R.; Blonk, C.; Tijburg, L.B.; Langius, J.A.; Meinders, A.E. and Pijl, H. (1998):

No effect of consumption of green and black tea on plasma lipid and antioxidant levels and on LDL oxidation in Smokers.

Arterioscler. Thromb. Vasc. Biol., 18 : 833 – 841.

51- Raederstorff, D.G.; Schlachter, M.F.; Elste, V. and Weber, P. (2003):

Effect of EGCG on lipid absorption and plasma lipid levels in rats.

J. Nutr. Biochem, 14 : 326 – 332.

52- Rah, D.K.; Han, D.W.; Baek, H.S.; Hyon, S.H.; Park, B.W. and Park, J.C. (2007):

Protection of rabbit kidney from ischemia / reperfusion injury by green tea polyphenol pretreatment.

Arch. Pharm. Res., 30 (11): 1447 – 1454.

53- Reiffel, J.A. and McDonald, A. (2006):

Antiarrhythmic effects of omega-3 fatty acids.

Am. J. Cardiol., 40 (12) : 97 - 100.

54- Reynolds, E.S. (1963):

The use of lead citrate as an electron opaque stain in electron microscopy.

J. Cell Biol., 17:208.

55- Ross, M.H. and Pawlina, W. (2006):

Adipose tissue. In : Histology, a text and atlas. 5th ed. Ch. 6.
pp. 238 - 246.

Lippincott Williams and Wikins, Philadelphia.

56- Sabu, M. C.; Smitha, K. and Kuttan, R. (2002):

Anti-diabetic activity of green tea polyphenols and their role in reducing oxidative stress in experimental diabetes.

Journal of ethnopharmacology, 83(1-2):109-116

57- Sakamoto, Y.; Mikuriya, H.; Tayama, K.; Takahashi, H.; Nagassawa, A.; yano, N.; Yuzawa, K.; Ogala, A. and Aoki, N. (2001):

Goitrogenic effects of green tea extract catechins by dietary administration in rats.

Arch. Toxicol., 75:591–596.

- 58- Satoh, K.; Sakamoto, Y.; Ogata, A.; Nagai, F.; Mikuriya, H.; Numazawa, M.; Yamada, K. and Aoki, N. (2002):**
Inhibition of aromatase activity by green tea extract catechins and their endocrinological effects of oral administration in rats.
Food Chem. Toxicol., 40 : 925 – 933.
- 59- Shepherd, J.; Cobbe, S.M.; Ford, I.; Isles, C.G.; Lorimer, A.R. and Packard C.J. (1995):**
Prevention of coronary heart disease with pravastatin in men with hypercholesterolemia.
N. Engl. J. Med., 333 : 1301 – 1307.
- 60- Shimotoyodome, A.; Haramizu, S.; Inaba, M.; Murase, T. and Tokimitsu, I. (2005):**
Exercise and green tea extract stimulate fat oxidation and prevent obesity in mice.
Med. Sci. Sports Exerc., 37(11):1884-1892.
- 61- Tholpady, S.S.; Katz, A.J. and Ogle, R.C. (2003):**
Mesenchymal stem cells from rat visceral fat exhibit multipotential differentiation in vitro.
Anat. Rec. A Discov. Mol.Cell Evol. Biol., 272(1):398-402.
- 62- Tong, Q. and Hotamisligil, G.S. (2001):**
Molecular mechanisms of adipocyte differentiation.
Rev. Endocr. Metab. Disord., 2: 349–355.

- 63- Weisberg, S.P.; McCann, D.; Desai, M.; Rosenbaum, M.; Leibel, R.L. and Ferrante, A.W. (2003):**
Obesity is associated with macrophage accumulation in adipose tissue.
J. Clin. Invest. Dec., 112(12):1785-1788.
- 64- Wilson-Fritch, L. N. ; Chouinard, M.; Lazar, M. A.; Chui, P. C.; Leszyk, J.; Straubhaar, J.; Czech, M. P. and Corvera, S. (2004):**
Mitochondrial remodeling in adipose tissue associated with obesity and treatment with rosiglitazone.
J. Clin. Invest., 114(9): 1281-1289.
- 65- Wolf, G.; Chen, S.; Han, D. and Ziyadeh, F. (2002):**
Leptin and renal disease.
American Journal of Kidney Diseases, 39(1): 1-11
- 66- Wu, L.Y.; Juan, C.c.; Ho, L.T.; Hsu, Y.P. and Hwang, L.S. (2004):**
Effect of green tea supplementation on insulin sensitivity in Sprague – Dawley rats.
J. Agric. Food Chem., 52:643 – 648.
- 67- Yang, J.A.; Choi, J.H. and Rhee, S.J. (1999):**
Effects of green tea catechin on phospholipase A2 activity and antithrombus in streptozotocin diabetic rats.
J. Nutr. Sci. Vitaminol., 45 : 337 – 346.

68- Yokozawa, T.; Nakagawa, T. and Kitani, K. (2002):

Antioxidative activity of green tea polyphenol in cholesterol – fed rats.

J. Agric. Food Chem., 50 : 3549 – 3552.

69- Yokozawa, T.; Nakagawa, T.; Lee, K.I.; Cho, E.J.; Terasawa, K. and Takeuchi, S. (1999):

Effects of green tea tannin on cisplatin – induced nephropathy in LLC- PK1 cells and rats.

J. Pharm. Pharmacol., 51 : 1325 – 1331.

70- Zagorsky, J.L. (2004):

Is obesity as Dangerous to your Wealth as to Your Health ?

Res. Aging, 26 : 130 – 152.

71- Zeyuan, D.; Bingying, T.; Xiaolin, L.; Jinming, H. and Yifeng, C. (1998):

Effect of green tea and black tea on the metabolisms of mineral elements in old rats.

Biol. Trace Elem. Res., 65 : 75 – 86.