

REFERENCE

- Abdel-Moneim, S.M. and Sultan, D.M. (2008):** Genetic characterization of *G.lamblia* isolates from Egyptian patients with relation to clinical giardiasis. J. Egypt. Soc. Parasitol., 38 (2): 547 – 560.
- Abravaya, K. Huff J. and Marshall, R. (2003):** Molecular beacons as diagnostic tools technology and applications. J. Clin. Chem. Lab. Med., 41(4):468-74.
- Adam, R.D. (2001):** Biology of *G.lamblia*. J. Clin. Microbiol. Rev., 14: 447-75.
- Adrabbo, K. and Peura, D. (2002):** Giardiasis: A Review. J. Gastroenterol., 15-29.
- Alles, A.J.; Waldron, M.A.; Sierra, L.S. and Mattia, A.R. (1995):** Prospective comparison of direct immunofluorescence and conventional staining methods for detection of *G.lamblia* and *Cryptosporidium* spp. in human faecal specimens. J. Clin. Microbiol., 33:1632-1634.
- Al-Mekhlafi, M.S., Azlin, M.; Nor Aini, U.; Shaik,A.; Ismail, M. G.; Firdaus, M.S.; Rozlida, A.R. and Norhayati, M. (2005):** Giardiasis as a predictor of childhood malnutrition in Orang Asli children in Malaysia. J. Trop. Med. Hyg., 99: 686-691.
- Al-Shammari, S.; Khoja, T.; El-Khwasky, F. and Gad, A. (2008):**Intestinal parasitic diseases in Saudi Arabia. J. Trop. Med 6 :184 – 189.

- Amar, C.F.; Dear, P.H. ; Pedraza-Diaz, S.; Looker, N.; Linnane, E. and McLauchlin, J. (2002):** Sensitive PCR-restriction fragment length polymorphism assay for detection and genotyping of *G.lamblia* in human faeces. J. Clin. Microbiol., 40 : 446-452.
- Amar, C.F.; East, C.; Maclure, E.; McLauchlin, J.; Jenkins, C.; Duncanson, P. and Wareing, D.R. (2004):** Blinded application of microscopy, bacteriological culture, immunoassays and PCR to detect gastrointestinal pathogens from faecal samples of patients with community- acquired diarrhea. Eur. J. Clin. Microbiol. Infect. Dis., 23:529 – 534.
- Amin, O.M.(2002):** Seasonal prevalence of intestinal parasites in the United States. Am. J. Trop. Med. Hyg., 66:799-803. .
- Ander, P.; Ervin, M.; Talvavera, F.; Halamka, J. and Dronen, S. (2004):** A study on giardiasis. J. Clin. Microbiol. Infect. Dis., 128-135.
- Aucolt, J. (1996):** Giardiasis and other Protozoal Diseases. in Nelson Textbook of Pediatrics, 15th edition, Volume 5., 970 - 972.
- Aydin, A.F.; Besirbelliglu, B.A.; Avci, I.Y.; Tanyuksel, M.; Araz, E. and Phasa, A. (2004):** Classification of *G.lamblia* parasites in Turkey into two groups A and B using restriction fragment length polymorphism. J. Diagn. Microbiol. Infect. Dis., 50: 147 – 151.
- Azazy, A.A.; Al-Mahbashi, T. Y. and Al-Mekhlafi, H.M. (2002):** Prevalence of intestinal and blood parasites among school children in Sana'a and Al-Mahweet provinces, Yemen. J. Med., 4: 50 – 55.

- Baquai, R.; Kazim, S.U. and Qureshi, H. (2002):** Role of cytokines in Giardiasis. J. Pak. Med. Associ., 50:113-115.
- Beaver, P.C. (2004):** Protozoa and protozoan infection. J.Clin. Parasitol., 9: 135-217.
- Berkman, D.S.; Lescone, A.G.; Gilman, R.H.; Lopez, S.L. and Black, M.M. (2006):** Effect of stunting, diarrreal diseases, and protozoal infection during infancy on cognition in late childhood: a follow up study. Lancet., 359 : 564-571.
- Bernal, R.M.; Tovar, R.; Santos, J.I. and Munoz, M.L. (1998):**Possible role of calmodulin in excystation of *G.lamblia*. J. Parasitol. Res., 84(9): 687-93.
- Bertrand, I.; Albertini, L. and Schwartzbrod, J. (2005):** Comparison of two target genes for detection and genotyping of *G. lamblia* in human faeces by PCR and PCR-restriction fragment length polymorphism. J. Clin. Microbiol., 43 (12): 5940 – 5944.
- Bialek, R.N.; Binder, K.; Diettz, A.; Joachin, J.; Nobloch, D. and Zelck, M.E. (2002):** Comparasion of fluorescence, Antigen and PCR assay to detect *C.parvum* in faecal sample. J. Diagn. Microbiol. Infect. Dis., 43: 283-288.
- Boom, R.; Sol, C.J.A. Salimans, M.M.M.; Jansen, C.L.; Wertheim-van Dillin, P.M. and Van der Noordaa, J. (1990):** Rapid and simple method for purification of nucleic acids. J. Clin. Microbiol., 28:495-503.
- Boom, R.; Sol, C. and Weel, J. (2000):** Detection and quantitation of human *Cytomegalo virus* DNA in faeces . J. Virol ., 84:1-14.

- Boyce, E.G.; Cookson, E.T.; and Bond, W.S. (1990):** Persistent metronidazole - induced peripheral neuropathy. J. Clin. Microbiol., 24:19-21.
- Brieva, L.; Ara, J.R.; Bertol, V.; Canellas, A. and Del-Agua, C. (1998):** Polyneuropathy caused by vitamin B 12 deficiencies secondary to chronic atrophic gastritis and giardiasis. J. Rev. Neurol., 26 (541): 1019-1020.
- Buret, A. (1994):** Pathogenesis - How does *G.lamblia* cause disease? In: *G.lamblia*: From molecules to disease. Edited by Thompson R.C.A., Reynoldson J.A. and Lymbery A.J. P., 293-302.
- Buret, A.G. (2007):** Mechanisms of epithelial dysfunction in giardiasis. J. Parasitol., 56: 316 – 317.
- Burke, J.A. (1977):** The clinical and laboratory diagnosis of giardiasis. J. Clin .Lab., 7: 373 – 391.
- Caccio, S.M. (2004):** New methods for the diagnosis of *C.parvum* and *G.lamblia*. J. Parasitol., 46(1-2): 151-5.
- Caccio, S.M.; DeGiacomo, M. and Pozio, E.(2002):** Sequence analysis of the β -giardin gene and development of a polymerase chain reaction restriction fragment length polymorphism assay to genotype *G.lamblia* cysts from human faecal samples. Int. J. Parasitol., 32: 1023-1030.
- Calderaro, A, ;Gorrini, C.; Montecchini, S.; Peruzzi, S.; Piccolo, G.; Rossi,S.; Gargiulo, F.; Manca, N.; Dettori,G. and Chezzi, G. (2010):** Evaluation of a real-time polymerase chain reaction assay

for the laboratory diagnosis of giardiasis. J. Clin. Microbiol., 66:261-267.

Carroccico, A.; Montalto, G.; Iacono, G.; Ippolito, S.; Soresi, M. and Notarbartolo, A. (1997): Secondary impairment of pancreatic function as a cause of severe malabsorption in the intestinal giardiasis: A case report. Am. J. Trop. Med. Hyg., 56 (6): 599-602.

Cartwright, C.P. (1999): Utility of multiple stool specimen ova and parasite examinations in a high prevalence setting with giardiasis. J. Clin Microbiol.; 37 : 2408 – 2411.

Casemore, D.P; Sands, R.T. and Curry, A. (1985): *Cryptosporidium* species a new human pathogen. J.Clin. Pathol., 38:1321-1326.

Chavez-Munguia, B.; Cedillo-Rivera, R. and Martinez-Palomo, A. (2004):The ultrastructure of the cyst wall of *G.lamblia*. J.. Microbiol., 51(2): 220-6.

Cheesbrough, M. (1987): Medical laboratory manual for tropical countries. 2nd ed. Vol. 1, Tropical Health Technology, Butterworth-Heinemann., 88:365-389.

Cheesbrough, M. (1991):Techniques used to identify parasite In: Medical Laboratory Manual for Tropical Countries. C. Halley and H. Jordan (eds.), Britain, Butterworth-Heinemann Ltd., 175-185.

Chien, A.; Edgar, D.B. and Trela, J.M. (1976): Deoxyribonucleic acid polymerase from the extreme thermophile *Thermus aquaticus*. J. Bacteriol. 174: 1550–1557.

- Corsi, A.; Nucci, C.; Knafelz, D.; Bulgarini, D.; Di-Iorio, L.; Polito, A.; De-Risi, K.; Ardenti-Morini, F. and Paone, F.M. (2004):** Ocular changes associated with *G.lamblia* infection in children. Br.J. Ophthalmol., 82(1): 59-62.
- Cox, FEG. (2007):** Classification and introduction to the parasitic protozoa. In: Cox FEG, Wakelin D, Gillespie SH, Despommier DD, eds. Topley, Wilson's Parasitology. 10th edn. London: Edward Arnold; 186-199.
- Craft, J.C. and Nelson, J.D. (1982):** Diagnosis of giardiasis by counterimmunoelectrophoresis of faeces. J. Infect. Dis., 145(4): 499–504.
- Das, S.; Schteingart, C.D.; and Hofmann, A.F., (1997):** *G.lamblia*: evidence for carrier-mediated uptake and release of conjugated bile salts. J. Parasitol; 87: 133 – 141.
- Davies, R.B. and Hibler, C.P. (2005):** Animal reservoirs and cross-species transmission of *G.lamblia*, In W. Jakubowski and J.C. Hoff (ed.), waterborne transmission of giardiasis. Environmental Protection Agency, Cincinnati., 104-126.
- DeLong, E.R.; DeLong, D.M. and Clark Pearson, D.L. (1988):** Comparing the areas under two or more correlated Receiver operating characteristic curves. A non Parametric Approach, Biometric, 44, 837-854.
- Elmendorf, H.G.; Dawson, S.C.; and McCaffery, J.M. (2003):** The cytoskeleton of *G.lamblia*. Int. J. Parasitol., 33 (1): 3 – 28.

- El-Naggar, S.M.; El-Bahy, M.M.; Abd Elasis, J. and El-Dardiry, M.A. (2006):** Detection of protozoal parasites in the stools of diarrhoeic patients using different techniques. J.Egypt. Soc. Parasitol., 36(1):7-22.
- El-Shazly, A.M.; Mowafy, N.; Soliman, M.; El-Bendary, M.; Morsy, ATA.; Ramadan, NII. and Arafa, WAS. (2004):** Egyptian genotyping of *G.lamblia*. J .Egypt. Soc. Parasitol ., 34:265–280.
- Farthing, M.J.G. (1994):** Giardiasis as a Disease. In: *G. lamblia* from Molecule to Disease. Reynoidson, J.A. Thompson, R.C.A. and Lymbery, A. (eds.). London: CAB International., 15-37.
- Farthing, M.J.G. (2001):** Giardiasis. J. Gastreenterol. Clin. North. Am., 25(3): 493-515.
- Faubert, G.M. (2000):** Immune response to *G.lamblia*. J. Clin. Microbial. Rev., 13(1):35-54.
- Favennec, L.; Magne, D. and Gobert, J. G. (1997):** Cytopathogenic effect of *G.lamblia* in vitro. Parasitology. Today., 141-147.
- Fawzi, M.; El-Sahn, A.A.; Ibrahim, H.F. and Shehata, A.I. (2004):** Vegetable-transmitted parasites among inhabitants of El-Prince and Alexandria and its relation to housewives' knowledge and practices. J. Egypt. Public Health Assoc., 79:13-29.
- Filice, F. P. (1952):** Studies on the cytology and life history of *G.lamblia* from the laboratory rat. Univ. Calif. Publ. Zool. 57: 53 - 146.
- Friedman, G.D. (1994):** Primer of Epidemiology, 4th edition, Chapter 3 (observations used in epidemiology)., P:33.

- Furness, B.W.; Beach, M.J. and Roberts, J.M. (2000):** Giardiasis surveillance-United States, 1992-1997. CDC. Surveill. Summ. 2000 Aug., 1149 (7):1-13.
- Galli, S.J. (2005):** Mast cells in the development of adaptive immune responses. J. Nat. Immunol., 6: 135 – 142.
- Garcia, L.S. (1990):** Laboratory methods for diagnosis of parasitic Infections. In: Diagnostic Microbiology. Baron, E.J. and Tenover, S.M. (eds), 8 ed. Bialy and Scott's, The C.V. Mosby Co., St. Louis., 78-83.
- Garcia, L.S. and Bruckner, D. A. (1997):** Intestinal Protozoa Flagellates and Ciliates. J. Diag. Med. Parasitol., 2: 13-39.
- Garcia, L.S. and Garcia, J.P. (2006):** Detection of *G.lamblia* antigens in human faecal specimens by a solid-phase qualitative immunochromatographic assay. J. Clin. Microbiol., 44 (12): 4587 – 4588.
- Garcia, L.S.; Shimizu, R.Y. and Bernard, C.N. (2000):** Detection of *G.lamblia*, *E. histolytica*, *E.dispar* and *C. parvum* antigens in human faecal specimens using the Triage Parasite Panel enzyme immunoassay. J. Clin. Microbiol., 38 (9): 3337 – 3340.
- Gardener, T.B. and Hill, D.R. (2001):** Giardiasis. J. Clin. Microbiol. Rev., 14(1): 114-128.
- Gendrel, D.; Treluyer, J.M. and Richard-Lenoble, D. (2003):** Parasitic diarrhea in normal and malnourished children. J. Fundam. Clin. Pharmacol., 17 : 189 – 198.

- Ghosh, S.; Debanth, A.; Sil, A.; De, S.; Chattopadhyay, D.J. and Das, P. (2005):** PCR detection of *G.lamblia* in stool intergenic space region of multicopy rRNA gene. J. Mol. Biol., . 14(3):181-189.
- Gillin, F.D.; Reiner, D.S. and Boucher, S.E. (2000):** Small intestinal factors promote encystation of *G.lamblia* in vitro. J. Infect. Immunol., 56: 705-707.
- Goka, A.K.J.; Rolston, D.D.K.; Mathan, V.I. and Farthing, M.J.G. (1990):** The relative merits of faecal and duodenal juice microscopy in the diagnosis of giardiasis. Tran. R. Soc. Trop. Med. Hyg., 84: 66-67.
- Goldstein, F.; Thornton, J.J. And Szydlowski, T. (2005):** Biliary tract dysfunction in giardiasis. Am. J. Dig. Dis., 23(6):559- 560.
- Goodgame, R. W. (1996):** Understanding intestinal spore-forming protozoa: *C. parvum*, *Microsporidia*, *Isospora* and *Cyclospora*. J. Ann. Intern. Med., 124: 429-441.
- Gordets, B.; Retore, P.; Cadranet, S.; Heneihofm, W.; Rahman, A. and Butzler, J.P. (1984):** Routine culture of *G.lamblia* trophozoites from human duodenal aspirate. Lancet., 111:137.
- Gottstein, B. and Nash, T.E. (1991):** Antigenic variation of *G.lamblia*: Infection of congenitally a thymic mice. J. Immunol., 13: 649 - 659.
- Grace, M.B.; McLeland, C.B. and Blakely, W.F.(2002):** Real-time quantitative RT-PCR assay of GAD 45 gene expression

changes as a biomarker for radiation biodosimetry. Int. J. Radiat. Biol., 78(11):1011-21.

Green, E.L.; Miles, M.A. and Warhurst, D.C. (1985): Immunodiagnostic detection of *G.lamblia* antigen in faeces by a rapid visual enzyme-linked immunoassay. Lancet., 28: 691-693.

Guimaraes, S. and Sogayar, M.I. (2002): Detection of anti- *G.lamblia* serum antibody among children of day care centers. Rev. Saude Publica., 36 (1): 63 – 68.

Guimaraes, S.; Sogayar, M.I. and Franco, M. (2002): Analysis of proteins from membrane and soluble fractions of *G.lamblia* trophozoites of two Brazilian axenic strains. Rev. Inst. Med. Trop. Sao. Paulo., 44(5): 239-44.

Gurwith, M. Wenman, W. Hinde, D. Feltham, S. and Greenberg, H . (1997): A prospective study of *rotavirus* infection in infants and young children. J . Infect. Dis., 144: 218-224.

Guy, R.A.; Xiao C. and Horgen, P.A. (2004): Real-Time PCR assay for detection and genotype differentiation of *G.lamblia* in stool specimens. J. Clin. Microbiol., 42 (7): 3317 – 3320.

Handoussa, A. E.; Elmy, M.M.; Hussein, E.M. and El-Nimr, H.I. (2005): Degree of symptoms versus copro-antigen levels in *G.lamblia* infection. J. Egypt. Soc. Parasitol., 35 : 69 – 81.

Hanson, K.L. and Cartwright, C.P. (2001): Use of an enzyme immunoassay does not eliminate the need to analyze multiple stool specimens for sensitive detection of *G.lamblia*. J. Clin. Microbiol., 39 (2): 474-7.

- Haque, R.; Roy, S.; Kabir, M.; Stroup, S.E.; Mondel, D. and Houpt, E. R. (2005):** *G.lamblia* assemblage A infection and diarrhea in Bangladesh. J. Infect. Dis., 192: 2171-2173.
- Hardin, J.A.; Buret, A.G.; Olson, M.E.; Kimm, M. and Gall, D. (1997):** Mast cell hyperplasia and increased macromolecular uptake in an animal model of giardiasis. J. Parasitol., 83(5): 908-912.
- Hart, C.A.; Gatei, W.; Wamae, C.N.; Mbae, C.; Waruru, A.; Mulinge, E.; Waithera, T.; Gatika, S.M.; Kamwati, S.K. and Revathi, G. (2006):** Giardiasis prevalence, genotype analysis and symptoms associated with infections in children in Kenya. Am. J. Trop. Med. Hyg., 75(1): 78-82.
- Hashknes, P.J.; Spira, D.T.; Deckelbaum, R.J. and Granot, E. (1994):** Salivary IgA antibodies to *G.lamblia* in day care center children. J. Ped. Inf. Dis., 13(11): 953-958.
- Helmy, M.M; Abdel-Fattah, H.S. and Rashad, L. (2009):** Real time PCR/RFLP assay to detect *G.lamblia* genotypes in human isolates with diarrhea in Egypt. J. Parasitol., 95(4):1000-4.
- Heresi, G.P.; Murphy, J.R. and Cleary, T.G. (2000):** Giardiasts, J. Ped. Inf. Dis., 11: 189 – 195.
- Hiatt, R.A.; Markell, E.K. and Ng, E. (1995):** How many stool examinations are necessary to detect pathogenic intestinal protozoa? Am. J. Trop. Med.; Hyg., 53: 36 – 39.

- Higuchi, R.; Dollinger, G.; Walsh, PS. and Griffith, R. (2004):** Simultaneous amplification and detection of specific DNA sequences. *J. Bio.*, 10 (4):413-7.
- Hill, D.R. (2000): *G.Lamblia*.** In: Mandell, G.L.; Bennett, J.E. and Dolin, R., eds. *Mandell, Douglas and Bennett's: Principles and practice of infectious diseases*, 5th ed. Philadelphia, Churchill Livingstone, pp. 2888 – 2894.
- Homan, W.L. and Mank, T.G. (2001):** Human giardiasis genotype linked differences in clinical symptomatology. *Int. J. Parasitol.*, 31: 822-826.
- Hopkins, R.M.; Meloni, B.P.; Groth, D.M.; Wetherall, J.D.; Reynoldson, J.A. and Thompson, R.C. (1997):** Ribosomal RNA sequencing reveals differences between the genotypes of *G.Lamblia* isolates recovered from humans and dogs living in the same locality. *J. Parasitol.*, 83: 44 – 51.
- Isaac-Renton, J.; Blatherwick, J.; Bowie, W.R.; Fyfe, M.; Khan, M.; Li, A.; King, A.; McLean, M.; Medd, L.; Moorehead, W.; Ong, C.S. and Robertson, W. (1999):** Epidemic and endemic seroprevalence of antibodies to *C.parvum* and *G.lamblia* in residents of three communities with different drinking water supplies. *Am. J. Trop. Med. Hyg.*, 60(4): 578-83.
- Jimeanez, J.C.; Fantanie, J.; Grzych, J.M.; Defcase, E. and Capron, M. (2004):** Systemic and mucosal response to oral administration of excretory and secretory antigens from *G.lamblia*. *Clin. Diagnosis. Lab. Immunol.*, 11(1):157-160.

- Johnston, J.P.; Ballard, M.M.; Beach, M.J.; Cawser, L. and Wilken S.P.P. (2003):** Evaluation of three commercial assays for detection of *G.Lamblia* and *C.parvum* in faecal specimens. J. Clin. Microbiol., 41: 626-632.
- Joseph, S.; and David, W. (2001)::** In vitro Amplification of DNA by the Polymerase Chain Reaction. J. Clin. Chemi., J. 51(12) :2415-2418.
- Kader, S.A.; Mansour, A.M.; Mohran, Z.; El-Taoil, A. and Abdalla, K.F. (1998):** A study on the relation between proton pump inhibitor and gastric giardiasis. J. Egyp. Soc. Parasitol ., 128:149.
- Karabay, O.; Tamer, A.; Gunduz, H.; Kayas, D.; Arinc, H. and Celebi, H. (2004):** Albendazole versus metronidazole treatment of adult giardiasis: An open randomized clinical study. World J. Gastroenterol., 10(8): 1215-7.
- Karanis, P. and Ey, P.L. (2002):** Characterization of axenic isolates of *G.Lamblia* established from humans and animals in Germany. Parasitol. Res., 84:442-449.
- Katanik, M.T.: Schneider, S.K.:Roseablatt, J.E.; Hall, G.S. and Procop, G.W.(2001):** Evaluation of ColorPAC *Giardia* /*Cryptosporidium* rapid assay and ProSpecT *Giardia* /*Cryptosporidium* microplate assay for detection of *Giardia* and *Cryptosporidium* in faecal specimens. J. Microbiol., 39(12): 4523-4525.
- Katellaris, P.H.;and Farthing, MJG (2000):** Diarrhea and malabsorption in giardiasis: J. Infect.Dis., 33: 295 – 297.

- Katellaris, P.H.; Naeem, A. and Farthing, M.J. (1995):** Attachment of *G.lamblia* trophozoite to a cultured human intestinal cell line. *J. Gastroenterol.*, 37(4): 512-518.
- Kaucner, C. and Stinear, T. (2001):** Sensitive and rapid detection of viable *G.lamblia* cysts and *C.parvum* oocysts in large-volume water samples with wound fiberglass cartilage filters and reverse transcriptase-PCR. *Appl. Environ. Microbiol.*, 64 (5): 1743-1749.
- Keister, D.B. (1983):** Axenic cultivation of *G.Lamblia* in TYI-S33 medium supplemented with bile. *Trans. R. Soc. Trop. Med. & Hyg.*, 77: 487 – 488.
- King, M. (1975):** A medical laboratory for developing countries; ELBS, Oxford University. Press. vol. 15., 45-55.
- Kleppe, K.; Ohtsuka, E.; Kleppe, R.; Molineux, I. and Khorana, HG. (1971):** Studies on polynucleotides: Repair replications of short synthetic DNA's as catalyzed by DNA polymerases. *J. Mol. Biol.* 56: 341–361.
- Korman, S.H.; Hais, E. and Spira, D.T. (1990):** Routine in vitro cultivation of *G.Lamblia* by using the string test. *J. Clin. Microbiol.*, 28 (2): 368 – 369.
- Kosek, M.; C.Bern, and R.L. Guerrant (2003):** The global burden of diarrheal 6-disease as estimated from studies published between (1992) and (2000). *Bull.W.H.O.* 81:197-204.
- Lai ,P.S. and Mayer. L. (1997):** Gastrointestinal manifestations of primary immunodeficiency disorders. *Semin. Gastrointest. Dis.*,8:22-23.

- Lane, S. and D. Lloyd (2002):** Current trends in research into the waterborne parasite *G.Lamblia*. Crit.Rev.Microbiol.28:123-124.
- Lanfredi-Rangel, A.; Kattenbach, W.M.; Diniz, J.A .Jr. and de Souza, W. (1999):** Trophozoites of *G.Lamblia* may have a Golgi-like structure. FEMS. Microbiol. Lett., 181(2): 245-51.
- Langfod, T.D. (2002):** Central importance of immunoglobulin A in host defense against *G.Lamblia*. J. Infect. Immun., 70: 11 – 18.
- Lebwohl, B.; Deckelbaum, R.J. and Green, P.H. (2003):** Giardiasis. J. Gastrointest. Endo., 57 L 906 – 913.
- Lengerich, E.J.; Addiss, D.G. and Juranek, D.D. (1994):** Severe giardiasis in the United States. J. Clin. Infect. Dis., 18:760-763.
- Lequin, R. (2005):** Enzyme immunoassay (EIA) and enzyme linked immunosorbent assay (ELISA). J. Clin. Chemi. 51(12) :2415-2418.
- Li, E. (2004):** Mast cell-dependent control of *G.Lamblia* infections in mice. J. Infect. Immun., 72 : 6642 – 6649.
- Livak, K.J. and Schmittgen, T.D. (2001):** Analysis of relative gene expression data using real-time quantitative PCR and the 2(Delta CT) Method. Nucleic Acids Res., 25(4):402-8.
- Ljungstrom, I. and Castor, B. (1992):** Immune response to *G.Lamblia* in a water-borne outbreak of giardiasis in Sweden. J. Med. Microbiol., 36: 247- 352.
- Logan, J.; Edwards, K. and Saunders, N. (2009):**Real-Time PCR: Current Technology and Applications. J. immun., 88-91.

- Mahbubani, M.; Bej, A.K.; Perhn, M.; Schaefer, K.; Jakubwski, W. and Atlas, R.M. (1991):** Detection of *G.Lamblia* cysts by using the polymerase chain reaction and distinguishing live from dead cysts. Appl. Environ. Microbiol., 57(12):3456.
- Mank, T.G. (2001):** Diagnostic advantages and therapeutic options for giardiasis. Expert. Opin. Investig. Drugs., 10(8):1513-1519.
- Marshall, J. and Holberton, D.V. (1993):** Sequence and structure of a new coiled coil protein from a microtubule bundle in *G.Lamblia*. J. Mol. Biol., 23:521-30.
- Masser,C. and Kagnof, M.F.(2002):** Role of the intestinal epithelium in innate and adaptive mucosal immunity. J. Z.Gastroenterol.,40(7):525-529.
- Mayta, H.;Gilman. R.H.; Prendergast, E.;Castillo, J.P.; Tinco, Y.O.; Garcia.H.H.; Gonzalez,A.E. and Stering, C.R. (2008):** Nested PCR for the specific diagnosis of *Taenia solium*. J.Clin.Microbio., 46:386-289.
- McGlade, T.R.; Robertson, I.D.; Elliot, A.D. and Thompson, R.C.A. (2003):** High prevalence of *G.lamblia* detection by PCR. J. Vet. Parasitol., 110:197-205.
- Mhlanga, M.M. and Malmberg, L. (2001):** Using molecular beacons to detect single-nucleotide polymorphisms with real-time PCR. Methods. J. Mol. Biol .,25(4):463-71.
- Misra, P.K.; Kumar, A.; Agarwal, V. and Jagota, S.C. (1995):** A comparative clinical trial of albendazole versus metronidazole in children with giardiasis. J. Indian Pediatr., 32(7): 779-82.

- Miyahara, T.; Kubokawa, M.; Koyanagi, S.; Migita, Y.; Oogoshi, K.; Sakai, H.; Ito, T.; Nakano, I. and Nawata, H. (1997):** A case of successfully treated giardiasis in pancreas. Fukuoka Igaku Zasshi., 88(9):313-318.
- Moncada, D.M.; Kammanadiminti, S.J. and Chadee, K. (2003):** Mucin and T cell like receptors in host defense against intestinal parasites Trends. J. Parasitol., 19: 305-311.
- Monis, P.T.; Mayrhofer, G.; Andrews, R.H.; Homan, W.L.; Limper, L. and Ey, P.L. (2003):** Molecular genetic analysis of *G.ilamblia* isolates at the glutamate dehydrogenase locus. J. Parasitol., 112: 1-12.
- Morimoto, N.; Komatsu, C.; Nishida, M. and Sugiura, T. (2001):** Detection of *G.lamblia* cyst in non fixed long term stored human faeces by direct immunoflorescent stain. J. Infect. Disease., 54(2): 72-74.
- Morrison, T.B.; Weis, J.J. and Wittwer, C.T. (1998):** Quantification of low-copy transcripts by continuous SYBR Green I monitoring during amplification. J. Bio. Tech., 24: 960-964.
- Moucada, D.M.; Kammanadiminti, S.J. and Chadee, K. (2003):** Mucin and T-cell like receptors in host defense against intestinal parasites. Trends. J. Parasitol., 19 : 305 -311.
- Muller, N. and Allmen, N. (2005):** Recent insights into the mucosal reactions associated with *G.Lamblia* infections. Int. J. Parasitol., 35: 1339 – 1347.

- Nacapunchai D, Tepmongkol M, Tharavanij S, Thammapalerd N, and Subchareon A. (1986):** A comparative study of four methods for detecting antibody in asymptomatic giardiasis. South east Asian J. Trop. Med. Public. Health. Mar; 17(1):96–100.
- Nantavisai, K.; Mungthin, M.; Tan-ariya, P.; Rangsin, R.; Naaglor, T. and Leelayoova, S. (2007):** Evaluation of the sensitivities of DNA extraction and PCR methods for detection of *G.Lamblia* in stool specimens. J. Clin. Microbiol., 45 (2): 581 – 583.
- Nash, T.E. (2001):** Treatment of *G.lamblia* infections. Con. Rev. J. Ped. Infect. Dis., 20 (2) : 193 - 5.
- Ng, C.T.; Gilchrist, C.A.; Lane, A.; Roy, S.Haque, R. and Houpt, E.R. (2005):** Multiplex Real-Time PCR assay using scorpion probes and DNA capture for genotype-specific detection of *G.Lamblia* on faecal samples. J. Clin. Microbiol., 43 (3): 1256 – 1260.
- Nolan, T.; Hands, R.E. and Bustin, S.A.(2006):** Quantification of mRNA using real-time RT-PCR. J. Biochem , 1: 1559–1582.
- Norhayati, M. (1998):** Prevalence and risk factors of *G.lamblia* infection in a rural community. J. Trop. Med., 29(4):735–738.
- O'Handley, R.M.; Olson, M.E.; Fraser, D.; Adam, P. and Thompson, R.C. (2000):** Prevalence and genotypic characterization of *G.lamblia* in dairy calves from Western Australia and Western Canada. J. Vet.Parasitol.,90:193-200.
- Orth, T.; Protzer, U.; Mayet, W.J.; Meyer, Z. and Buschenfelde, K.H. (1995):** Long-term damage to duoddenal mucosa in

malabsorption syndrome as a sequel of *G.lamblia* infection. J. Z. Gastroenterol. 33(3):166-169.

Osman, G.A.; Makled, K.M.; El-Shakankiry, H.M.; Metwali, D.M.; Abdel-Aziz, S.S. and Saafan, H.H. (1999): Coccidian parasites as a cause of watery diarrhea among protein energy malnourished and other immunocompromised Egyptian children. J. Egypt. Soc. Parasitol., 29: 653 – 668.

Oster, N.; Gehrig-Feistel, H.; and Jung, H., (2006): Evaluation of the immunochromatographic CORIS strip test for rapid diagnosis of *G.Lamblia*. J. Eur. Parasitol., 25: 112 – 115.

Palm, D.; Weiland, M.; McArthur, A.G., Winiecka-Krusnell, J.; Cipriano, M.J.; Birkeland, S.R.; Pacocha, S.E.; Davids, B.; Gillin, F., Linder, E. and Svard, S. (2005): Developmental change in the adhesive disk during *G.lamblia* differentiation.. J. Mol. Biochem. Parasitol., 141: 199 – 207.

Pavlov, AR.; Pavlova, N.V.; Kozyavkin, SA. and Slesarev, AI. (2004):Recent developments in the optimization of thermostable DNA polymerases for efficient applications. Trends Biotechnol., 22: 253–260.

Payne, P. ; Dryden, M. and Ridley, R. (2002): Evaluation of the efficacy of some parasitological techniques to eliminate cyst shedding in dogs naturally infected with *G.lamblia*. J. Microbiol., 220(3):330–333.

Petri, W.A. (2005): Treatment of Giardiasis. Curr. Treat. Options. J. Gastroenterol., 8(1): 13-17.

- Pettoello Mantovani, M.; Giardino, I.; Magli, A.; di Martino, L. and Guandalini, S. (1990):** Intestinal giardiasis associated with ophthalmologic changes. J. Pediatr. Gastroenterol. Nutr., 11(2): 196-200.
- Pickering, L.K. (2000):** Giardiasis and Balantidiasis. In : Nelson Textbook of Pediatrics, Section 13 Protozoan diseases, 16th edition, Volume 3., 1036-1038.
- Ponce-Macotella, M. M. N. ; Martinez-Gordillo, R. M. Bermudez-Cruz, P. M.; Salazar-Schettino, G.; Ortega-Pierres, and Ey. P. L. (2002):** Unusual prevalence of the *G.lamblia* A-II subtype amongst isolates from humans and domestic animals in Mexico. International J. Parasitol., 32:1201–1202.
- Qin, L.X.; Beyer, R.P.; Hudson, F.N.; Linford, N.J.; Morris, DE. and Kerr, KF. (2006):** Evaluation of methods for oligonucleotide array data via quantitative real-time PCR. BMC bioinformatics., 7:23.
- Rabbani, G.H. and Islam, A. (1994):** Giardiasis in human: population at risk and prospect for control in: *G.lamblia* from molecule to disease. Edited by Thompson R.C.A., Reynoldson J.A., Lymbery A..J., 217-249.
- Rabinow, Paul (1996):** Making PCR. J. Mol Biotechnol., 12(1):96-99..
- Raeymaekers, L. (2000):** Basic principles of quantitative PCR. Mol. Biotechnol., 15(2):115-122.
- Read, C.M.; Monis, P.T. and Thompson, R.C. (2004):** Discrimination of all genotypes of *G. lamblia* at the glutamate dehydrogenase locus using PCR-RFLP. J. Infect. Genet. ., 4 : 125 – 130.

- Reynoldson, J.A.; Behnke, J.M. and Gracey, M. (1998):** Efficacy of albendazole against *G.lamblia* and hookworm in the north of Western Australia. J. Trop., 71: 27 – 44.
- Ridley, D.S. and Hawgood B.C. (1989):** The value of formol-ether concentration of faecal cysts and ova. J. Clin. Pathol., 9: 74 – 76.
- Roberts, L.S. and Janovy, J.J.R. (2002):** Foundations of parasitology .5th ed. W.M.C. Brown . publ., 82-86.
- Roberts-Thomas, J.C.; Anders, R. and Bhathal, P. (1982):** Granulomatous hepatitis and cholangitis associated with giardiasis..J. Gastroenterol., 480-483.
- Rodriguez, O.L.; Hage, I.I.; Gonzalez, Y.; Roque, M.E.; Vasquez, N.; Lopez, E. and Diprisca, M.C. (2004):** Secretory IgA Ab response in venezuelar children infected with *G.lamblia*. J. Trop. Pediatr., 50 (2):68-72.
- Rosenthal, P.I. and Leibman, W.M. (1980):** Comparative study of stool examination, duodenal aspiration and pediatric enterotest for giardiasis in children. J. Pediatr., 96:278-279.
- Rosken, H. and Erlandsen, S. L. (2002):** Inhibition of invitro attachment of *G.lamblia* trophozoite by mucin. J. Parasitol., 88: 869-873.
- Rosoff , J.D.; Sanders, C.A. and Sonnad, S.S (1989):** Stool diagnosis of giardiasis using a commerically available enzme immunoassay to detect *G.lamblia* specific antigen 65 (GSA-65). J. Clin.Microbiol., 26-27.

- Roxstrom-Lindquist, K.; Palm, D.; Reiner, D.; Ringqvist, E. and Svard, S.G. (2006):** *G.lamblia* immunity-an update. Trends J. Parasitol., 22 (1): 26 – 31.
- Rustia, M. and Shubik, P. (1990):** Induction of lung tumors and malignant lymphoma in mice by metronidazole. J. Nat. Cancer Institute 48 : 721 – 1972.
- Rychlik, W.; Spencer, W.J. and Rhoads, R.E. (2003):** Optimization of the annealing temperature for DNA amplification in vitro. Nucl. Acids Res. 18: 6409–6412.
- Saha, B.K.; Tian, B.; and Bucy, R.P. (2001):**Quantitation of HIV-1 by real-time PCR with a unique fluorogenic probe. J. Virol .Methods ;93(1-2):33-42.
- Sails, A.D. (2009):** Applications in Clinical Microbiology. Real-Time PCR: Current Technology and Applications. J.Microbiol., 54-59.
- Schmidt, G.D. and Roberts, L.S. (2000):** Foundations of Parasitology, 6th edition, Mc Graw-Hill, New York, NY.
- Schultz, M.G. (1980):** Giardiasis. JAMA. 233: 1383 – 1384.
- Scott, K.G. (2002):** Intestinal infection with *Giardia* spp. Reduces epithelial barrier function in a myosin light chain kinase-dependent fashion.J. Gastroenterol., 123: 1179 – 1190.
- Scott, K.G. (2004):** Role of CD8⁺ and CD4⁺T-lymphocytes in jejunal mucosal injury during murine giardiasis. J. Infect. Immun., 72: 3536 – 3542.

- Seow, F.; Katelaris, P.H. and Ngu, M. (1993):** The effect of *G.lamblia* trophozoites on trypsin, chymotrypsin and amylase in vitro. J.Parasitol.,106 : 233 – 238.
- Shalta, H.M.; El-Hodhod, M.T.; Mohsen, D.M. and Salah El-din, M.Y. (2004):** Potential diagnosis of *G.lamblia* infection through Ab detection in saliva, J. Egypt. Soc. Parasitol., 34(2):621-630.
- Sharkey, E.R.; Scalice, K.G.; Christy Jr., S.M. Atwood, and Daiss J.L. (2002):** Antibodies as Thermolabile Switches: High Temperature Triggering for the Polymerase Chain Reaction. J. Bio. Technol., 12: 506–509.
- Shetty, N.; Narasimha, M.; Elliott, E.; Raj, L.S. and Macaden, R. (1992):** Age-specific sero-prevalence of amoebiasis and giardiasis in southern Indian infants and children .J. Trop. Pediatr.,38 (2):57-63.
- Singer, S.M. and Nash, T.E. (2000):** The role of normal flora in *G.lamblia* infections in mice. J. Infect. Dis., 181: 1520 – 1522.
- Smith, H.V. and Paget, T. (2007):** *Giardia*. Infectious Disease: Foodborne Diseases. J. Infect. Dis.,55-60.
- Soliman, M.M.; Ti-Kilani, R.; Abou-Shady, A.F.; El-Mageid, S.A.; Handousa, A.A.; Hegazi, M.M. and Belosevic, M. (1998):** Comparison of serum antibody responses to *G.lamblia* of symptomatic and asymptomatic patients. Am. J. Trop. Med. Hyg., 58(2):232-239.

- Solinas, A. ; Brown, L.J, and McKeen, C. (2001).** Duplex Scorpion primers in SNP analysis and FRET applications. *Nucleic Acids Res.*, 29(20): 96-100.
- Sotto, A.; Alvarez, J.L.; Garcia, B.; Pomar, F. and Cendan, A (1990):** Acute hepatic lesion caused by *G.lamblia*. *Rev Esp. Enferm. Dig.*, 77(1):24-28.
- Sousa,M. C.; Goncalves,C.A.; Bairos,V.A. and Poiares-da-silva,J. (2001):** Adherence of *G.lmablia* trophozoite to Int-407 human intestinal cells. *J. Cli. Diagn. Lab.Immunol.*, 8:258-265.
- Souza, D.S.; Barreiros, J.T.; Papp, K.M.; Steinded ,M.; Simoes, C. M. and Barardi, C.R.(2003):**Comparison between immunomagnetic separation coupled with immunofluorescence for the diagnosis of *G. lamblia* cysts in human faeces .*J. Trop, Med.*,45(6):339-342.
- Stuart, J.M.; Orr, H.J.; Warburton, F.G.; Jeyakanth, S.; Pugh, C. and Morris, I. (2003):** Risk factors for sporadic giardiasis: a case-control study in southwestern England. *Emerg. Infect. Dis.*, 9:229-233.
- Sulaiman , I.M. and Cama, V.(2006):** The biology of *G.lamblia* parasites. In: Ortega, Y.R. (ed), *Foodborne parasites*. Springer Press, USA. p.15-32.
- Sullivan, P.B.; Lunn, P.G. and Northrop-Clewes, C.A. (1992):** Parasitic infection of the gut and protein-losing enteropathy. *J. Pediatr .Gastroenterol. Nutr.*,15: 404 -407.

- Sullivan, P.B.; Marsh, M.N.; and Phillips, M.B., (1991):** Prevalence and treatment of giardiasis in chronic diarrhea and malnutrition. Arch. Dis. Child., 66: 304 – 306.
- Sullivan, R.; Linneman, C.C. Jr.; Clark, C.S. and Walzer, P.D. (1987):** Seroepidemiologic study of giardiasis patients and high-risk groups in a midwestern city in the United States. Am. J. Public Health., 77(8): 960-3.
- Sun, T. (1980):** The diagnosis of giardiasis. Am. J. Surg. Pathol., 4(3):265-271.
- Sun, T. (1988):** Diagnostic Parasitology. Igaku-Shoin Medical Publishers, Inc., New York., 32-38.
- Taylan Ozkan, A.; Mungan, M.; Kilic, S.; Babur, C.; Demircen, F. and Esen, B. (2005):** Giardiasis, *Giardia/Cryptosporidium* DFA yönteminin kullanımı. IV Ulusal Sindirim Yolu ile Bulaşan Enfeksiyonlar Simpozyumu, Mersin. Sözlü sunum no:11, Kongre kitabı., p:344 (In Turkish).
- Tellez, A.; Palm, D.; Weiland, M.; Aleman, J.; Winiecka-Krusnell, J.; Linder, E. and Svard, S. (2005):** Secretory antibodies against *G. lamblia* in lactating Norwegian women. J. Immunol., 27 (5): 163 – 169.
- Terry, C.F.; Shanahan, D.J.; Ballam, L.D.; Harris, N. McDowell, D.G. and Parkes, H.C. (2002):** Real-time detection of genetically modified soya using Lightcycler and ABI 7700 platforms with TaqMan, Scorpion, and SYBR Green I chemistries. J. Biochemis., 85(4):938-44.

- Thompson, R.C.A. (2001):** The future impact of social and cultural factors on parasitic disease: some emerging issues. J. parasitol., 31:949–959.
- Thompson, R.C. and Monis, P.T. (2004):** Variation in *G.lamblia*: implications for taxonomy and epidemiology. J. Adv. Parasitol., 58: 69-137.
- Thompson, R.C.; Hopkins, R.A. and Homan, W.L. (2000):** Nomenclature and genetic groupings of *G.lamblia* infection mammals. J. Parasitol., 16 : 210 – 218.
- Thompson, R.C.A. (2005):** The zoonotic significance and molecular epidemiology of *G.lamblia* and Giardiasis. J. Vet. Parasitol., 126(1-2):15-35.
- Thompson, R.C.A.; Reynoldson, J. A. and Mendis, A. H.W. (1993):** *Giardia* and giardiasis. J. Adv. parasitol., 32: 71-160.
- Udvardi, M.K.; Czechowski, T. and Scheible, W.R. (2008):** Eleven Golden Rules of Quantitative RT-PCR. J. Immunol., 20: 1736–1737.
- Van Keulen, H.; Gutell, R.; Campbell, S.; Erlandsen, L. and Meyer, E. (1993):** Unique phylogenetic position of diplomonadida based on complete small subunit ribosomal RNA sequence of *G.lamblia*. J. Immunol., 7:223-231.
- VanGuilder, H.D.; Vrana, K.E. and Freeman, W.M. (2008):** Twenty-five years of quantitative PCR for gene expression analysis. J. Bio. techn. 44: 619–626.

- Verweij, J.J.; Blange, R.A.; Templeton, K.; Schinkel, J.; Brien, E. A.; Van Rooyen, M.A.; Van Lieshout, L. and Polderman, A.M. (2004):** Simultaneous Detection of *E.histolytica*, *G.lamblia* and *C. parvum* in faecal samples by using Multiplex Real-Time PCR. J. Clin. Microbiol., 42 (3): 1220 – 1223.
- Verweij, J.J.; Pit, D.S.S. and Van Lieshout, L., (2001):** Determining the prevalence of *Oesophagostomum bifurcum* and *Necator americanus* infection using specific PCR amplification of DNA from faecal samples. J. Trop. Med. in Health., 6: 726 – 31.
- Verweij, J.J.; Schinkel, J.; Laeijendecker, D.; Van Rooyen, M.A.; Van Lieshout, L. and Polderman, A.M. (2003):** Real-time PCR for the detection of *G.lamblia*. J. Mol. Biol., 17: 223 – 225.
- Vet, J.A. and Marras, S.A. (2005):** Design and optimization of molecular beacon real-time polymerase chain reaction assays. Methods. J. Mol. Biol., 288:273-90.
- Volotao, A.C.; Costa-Macedo, L.M.; Haddad, F.S.; Brandao. A.; Peralta, J.M. and Fernandes, O. (2007):** Genotyping of *G.lamblia* from human and animal samples from Brazil using beta giardin gene: a phylogenetic analysis. J. Mol. Biol., 102(1): 10-9.
- Wallis, P.M. (1994):** Abiotic transmission is water really significant? In: *G.lamblia*: From molecule to disease . edited by Thompson .R.C.A.; Reynoldson, J.A. and Lymbery, A. J., p.99-122.
- Weiland, M.E.; McArthur, A.G.; Morrison, H.G.; Sogin, M.L. and Svard, S.G. (2005):** Annexin-like alpha giardins: a new

cytoskeletal gene family in *G.lamblia*. Int. J. Parasitol., 35 (6): 617 – 626.

Weitzel, T.; Dittrich, S. and Mohl, I. (2006): Evaluation of seven commercial antigen detection tests for *G.lamblia* and *C.parvum* in stool samples. J. Clin. Microbiol Infect; 12: 656 – 659.

Wittwer, C.T.; Ririe, K.M.; Andrew, R.V.; David D.A.; Gundry, R.A.; and Balis, U.J. (2000): The LightCycler: a microvolume multisample fluorimeter with rapid temperature control. J. Biotech., 22(1):176-81.

Wolfe, M.S. (1992): Symptomatology, Diagnosis and Treatment. In: *Giardia* and Giardiasis. Edited by Erlandsen S. and Meyer E. Plenum Press, New York., P:147.

World Health Organization (WHO), (1991): Basic laboratory methods in medical parasitology. Reagents and solutions and their preparation., 100-105.

World Health Organization (WHO), (1996): The world health report, Geneva., 354-378.

Wright, J.M.; Dunn, L.A.; Upcroft P. and Upcroft, J.A. (2003): Efficacy of anti giardial drugs. Expert Opin. Drug Saf., 2(6): 529-41.

Wright, S.G. (2000): Giardiasis In Hunter's Tropical Medicine and Emerging Infectious Diseases. Eighth edition, edited by Strickland G.S., 589-593.

- Yassin, M.M.; Shubair, M.E.; Al-Hindy, A.I. and Jadallah, S. (1999):** Prevalence of intestinal parasites among school children in Gaza City. J. Egypt. Soc. Parasitol., 29(2):365-373.
- Yereli, K.; Balcioglu, I.C.; Ertan, P. ;Limoncu, E. and Onag, A. (2004):** Albendazole as an alternative therapeutic agent for childhood giardiasis in Turkey. J. Clin. Microbiol. Infect. ,10(6): 527- 9.
- Young, K.H.; Bullock, S.L.; Melvin, D.M. and Spruill, C.L. (1979):**Ethyl acetate as a substitute for diethyl ether in the Formalin-ether sedimentation technique. J. Clin. Microbiol., 10:852-853.
- Zaher, S.M. (1993):** Laboratory diagnosis of infection with *C.parvum*. M. Sc. Thesis, Faculty of Medicine, Zagazig University.
- Zajac, A.M., Johnson, J. and King, S.E. (2002):** Evaluation of the importance of centrifugation as a component of zinc sulfate faecal flotation examinations. Journal of the American Animal Hospital Association 38:221–224.
- Zaki, A.M.; Dupont, H.L.; El-Alamy, M.A.; Arafat, R.R.; Amin, K.; Awad, M.M.; Bassiouni, L.; Imam, I.Z.; El-Malih, G.S.; El-Marsafie, A.; Mohieldin, M.S.; Naguib, T.; Rakha, M.A.; Sidaros, M.; Wasef, N.; Wright, C.E. and Wyatt, R.G. (1986):** The detection of enteropathogens in acute diarrhea in a family cohort population in rural Egypt. Am. J. Trop. Med. Hyg., 35:1013-1022.
- Zalipaeva, T.L. (2002):** Clinical symptoms of *G. lamblia* infection in children. Med. Parasitol. (Mosk), 3: 29 - 32.