




References

References

- Abath, F.G. and Werkhauser, R.C. (1996):** The tegument of *Schistosoma mansoni*: functional and immunological features, Parasite Immunol. 18: 15–20.
- Abdel Aziz, S. S. and el-Badawy, N. M. (2000):** Experimental trials of an artemisinin derivative in treatment of *Schistosoma mansoni* infected mice. J. Egypt. Soc. Parasitol. 30:295-303.
- Abdel-Hamid, H.F.(2004):** Investigation of induced biochemical and histopathological parameters of acetonitrile extract of *Jatropha carcus* in albino rats. J. Egypt. Soc. Parasitol., 34(2): 397-406.
- Abdulla, M.H.; Lim, K.C.; Sajid, M.; McKerrow, J.H. and Caffrey, C.R. (2007):** Schistosomiasis mansoni: Schistosomiasis mansoni: novel chemotherapy using a cysteine protease inhibitor, PLoS Med. 4 :14.
- Achan, J.; Tibenderana, J.K.; Kyabayinze, D.; Wabwire, F.; Kamya, M.R.; Dorsey, G.; D'Alessandro, U.; Rosenthal, P.J. and Talisuna, A.O. (2009):** " Effectiveness of quinine versus artemether-lumefantrine for treating uncomplicated falciparum malaria in Ugandan children: randomised trial" BMJ 339: 2763
- Agnew, A. (1995):**The relationship between worm burden and levels of a circulating antigen (CAA) of 5 species of *Schistosoma* in mice. Parasitology 111 : 67–76.
- Allan, D.; Payares, G. and Evans, W.H. (1987) :**The phospholipid and fatty acid composition of *Schistosoma mansoni* and of its purified tegumental membranes, Mol. Biochem. Parasitol. 23 : 123–128.
- Amer, M.(1982):** Cutaneous schistosomiasis. International Journal of Dermatology, 21: 44-46.
- Andrade, Z.A.(1965):** Hepatic schistosomiasis; morphological aspects. In: progress in liver disease. ed. Pepper, H. and Schaffner, E. 2: 222-242.
- Andrade, Z.A. and Warren, K.S. (1964):** Mild prolonged schistosomiasis in mice: alteration in host response with time and the development of portal fibrosis. Trans R Soc Trop Med Hyg 58 : 53.
- Andrews, P. (1985):** Praziquantel: mechanisms of anti-schistosomal activity. Pharmacol. Ther. 29:129-156 .

- Andrews, P.; Thomas, H. and Weher, H.(1980):** The in vitro uptake of ^{14}C praziquantel by cestodes, trematodes and nematodes. *J. Parasitol.*, 66:920-925.
- Araujo, M.I. (1994):** The absence of gamma-interferon production of *S. mansoni* antigens in patients with schistosomiasis, *Braz. J. Med. Biol. Res.* 27 : 1619-1625.
- Arauo, M.I. (1996):** Evidence of a T helper type 2 activation in human schistosomiasis, *Eur. J. Immunol.* 26 : 1399-1403.
- Barnes, K.I.; Chanda, P. and Barnabas, G. (2009):**" Impact of the large-scale deployment of artemether/lumefantrine on the malaria disease burden in Africa: case studies of South Africa, Zambia and Ethiopia". *Malar J.* 8 (1): S8.
- Barreto, M.L.; Smith, D.H. and Sleight, A.C. (1990):** Implications of faecal egg count variation when using the Kato-Katz method to assess *Schistosoma mansoni* infections, *Trans. R. Soc. Trop. Med. Hyg.* 84 : 554-555.
- Basch, P.F. and Basch, N. (1982):** *Schistosoma mansoni*: scanning electron microscopy of schistosomula, adults and eggs grown "in vitro". *Parasitology* 85: 333-338.
- Beaver, P.C.; Juny, P.C. and Cupp, E.W.(1984):** General considerations of schistosomes. In *Clinical Parasitology*. 19th ed. Lea and Febiger, Philadelphia, 27:415.
- Becker, B.; Mehlhorn, H.; Andrews, P.; Thomas, H. and Eckert, J.(1980):** Light and electron microscopic studies on the effect of praziquantel on *Schistosoma mansoni*, *Dicrocoelium dendriticum*, and *Fasciola hepatica* (Trematoda) in vitro. *Z. Parasitenkd.* 63: 113-128.
- Beltran, S. and Boissier, J. (2008):** "*Schistosoma* monogamy: who, how, and why?". *Trends in Parasitology* 24 (9): 386-91.
- Bergquist, R. (1996):** Prospects of vaccination against schistosomiasis. *Scand J Infect Dis Suppl* 76 : 60.
- Bergquist, M.V.; Johansen, J. and Utzinger, J. (2009):** Diagnostic dilemmas in helminthology: what tools to use and when?, *Trends Parasitol.* 25 : 151-156.
- Bergwerff, A. A. and Van Dam, G.J. (1994) :**The immunologically reactive O-linked polysaccharide chains derived from circulating

- cathodic antigen isolated from the human blood fluke *Schistosoma mansoni* have Lewis-X as a repeating unit. Eur. J. Biochem. 225 : 467–482.
- Berry, M.; Mukhopadhyay, V.S.; Suri, S. and Pawa, S.(2004):** Diagnostic radiology, gastroenterology and hepatobiliary imaging. 2nd : 256.
- Bilharz, T.M.(1852):** Fernere Beobachtungun uber das die Pfortader des Menschen bewohnende *Distomum Haematobium* und sein verhaltniss Zu gewissen pathologischen Bildungen. Z. Wissensch. Zool., 4:74. Quoted from **Jordan, P.; Wabbe, G. and Sturrock, E.(1993):** "Human schistosomiasis" p.1. University Press, Cambridge, UK.
- Boros, D.I.(1989):** Immunopathology of *S.mansoni* infection. Clinical Microbiology Reviews, 2(3): 250-269.
- Braschi, S.; Borges, W.C. and Wilson, R.A. (2006):** "Proteomic analysis of the schistosome tegument and its surface membranes". Memórias Do Instituto Oswaldo Cruz 101 (1): 205–12.
- Brindley, P.J.; Strand, M.; Norden, A.P. and Sher, A.(1989):** Role of host antibody in the chemotherapeutic action of praziquantel against *Schistosoma mansoni*: identification of target antigens.
- Brouwers, J.F.H.M.; Versluis, C.; Van Golde ,L.M.G. and Tielens, A.G.M. (1998):** 5-Octadecenoic acid: evidence for a novel type of fatty acid modification in schistosomes, Biochem. J. 334 : 315–319.
- Bruun, B. and Aagaard-Hansen, J. (2008):** The Social Context of Schistosomiasis and its Control: An Introduction and Annotated Bibliography. World Health Organization, Geneva.
- Caffrey, C.R. (2007):** Chemotherapy of schistosomiasis: present and future, Current Opinion in Chemical Biology 11 : 433–439.
- Caulfield, J.P. and Cianci, C.M. (1985):** Human erythrocytes adhering to schistosomula of *Schistosoma mansoni* lyse and fail to transfer membrane components to the parasite, J. Cell Biol. 101 : 158–166
- Caulfield, J.P.; Korman, G.; Butterworth, A.E.; Hogan, M. and David, J.R. (1980):**The adherence of human neutrophils and eosinophils to schistosomula: evidence for membrane fusion between cells and parasites, J. Cell Biol. 86 : 46–63.
- (CDC): Center for Disease Prevention and control(2009):**www.dpd.cdc.gov/dpdx.

- Cheever, A.W.; Maacedoni, J.G.; Cheever, E.A. and Mosimann, J.E.(1992): Persistence of eggs and hepatic fibrosis after treatment of infected mice. *Am. J. Trop. Med. Hyg.*,46(6):752-758.
- Cheng, G.F.; Lin, J.J.; Feng, X.G.; Fu, Z.Q.; Jin, Y.M.; Yuan, C.X.; Zhou, Y.C. and Cai, Y.M. (2005): Proteomic analysis of differentially expressed proteins between the male and female worm of *Schistosoma japonicum* after pairing, *Proteomics* 5 : 511–521.
- Chitsulo, L.;Engels, D.;Montresor, A. and Savioli, L. (2000): The global status of schisto- somiasis and its control. *Acta Trop.* 77: 41–51.
- Cioli, D. (2000): Praziquantel: is there real resistance and are there alternatives?, *Curr. Opin. Infect. Dis.* 13 : 659–663.
- Cioli, D.; Pica-Mattoccia, L. and Archer, S. (1993): Drug resistance in schistosomiasis. *Parasitol. Today*, 9:162.
- Cioli, D.; Botros, S.S.; Wheatcroft-Francklow, K.; Mbaye, A.; Southgate, V.; Tchuem Tchuenté, L.A.; Pica-Mattoccia, L.; Troiani, A.R.;El-Din S.H.;Sabra, A.N.;Albin ,J.; Engels ,D. and Doenhoff, M.J. (2004):Determination of ED50 values for praziquantel in praziquantel-resistant and -susceptible *Schistosoma mansoni* isolates, *Int. J. Parasitol.* 34 : 979–987.
- Cioli, D.;Valle, C.; Angelucci, F. and Miele, A.E. (2008): Will new antischistosomal drugs finally emerge?, *Trends Parasitol.* 24 : 379–382.
- Clegg, J. A. (1965): In vitro cultivation of *Schistosoma mansoni*. *Exp. Parasitol.* 16: 133-147.
- Coelho, P.M.Z.; Gazzinelli, G. and Pellegrino, J.(1980): *S.mansoni*: Host antigen occurrence on worms recovered from laboratory vertebrate animals. *Parasitol.*, 81: 349-354.
- Coles, G.C.; Bruce, J.I.; Kinoti, G.K.; Mutahi, W.T.; Dias, E.P. and Katz, N.(1986): Drug resistance in schistosomiasis. *Trans. R. Soc. Trop. Med. Hyg.*, 80: 347.
- Combes, C. (1990): Where do human schistosomes come from? An evolutionary approach. *Trends Ecol Evol* 5: 334-336.
- Conor, A.(2007):Current opinion in chemical biology 11:433-439.
- Contigli, C.(1999):Phenotype and cytokine profile of *Schistosoma mansoni* specific T cell lines and clones derived from

- schistosomiasis patients with distinct clinical forms, Clin. Immunol. 91 : 338–344
- Contractor, Q.Q.; Benson, L.; Contractor, T.Q. and Kher, Y.R. (1990):** Laparoscopic appearance of liver in *S. mansoni* infection. Endoscopy, 22: 137.
- Cook, G.C.(1996):** Manson's Tropical Diseases, 20th ed. WB Saunders Company LTD., England, 3:1413.
- Correnti, J.M.; Brindley ,P.J. and Pearce ,E.J. (2005):**Long-term suppression of cathepsin B levels by RNA interference retards schistosome growth, Mol. Biochem. Parasitol. 143: 209–215..
- Coura, J.R. and Amaral, R.S. (2004) :**Epidemiological and control aspects of schistosomiasis in Brazilian endemic areas, Mem. Inst. Oswaldo Cruz 99 (1) : 13–19
- Cousin, C.E and Dorsey, C.H.(1991):** Nervous system of *Schistosoma mansoni* cercariae: Organization and fine structure. Parasitol. Res., 2:132.
- Crompton, D.W. (1999):**How much human helminthiasis in the world? The Journal of Parasitology 85 (3): 397–403 .
- Davis, A.(1982):** Management of the patients with schistosomiasis. Quoted from **Jordan, P. and Webbe, G** eds: Schistosomiasis. Epidemiology, Treatment and control. William Heinemann Medical Books LTD., London, :184.
- Davis, A.(1986):** Recent advances in schistosomiasis. Quarterly Journal of Medicine (New series) 58(95):110.
- Davis, A.(1993):** Antischistosomal drugs and clinical practice: "In Human Schistosomiasis". Quoted from (**Jordan, P.; Wabbe, G. and Sturrock, R.F. eds**): 367. University Press, Cambridge, UK
- Davis, A.(1996):** Schistosomiasis. In Manson's Tropical Diseases. 20th ed., (Cook, C.G. ed) WB Saunders Company LTD, London, 30: 1551-1559.
- Davis, A. and Wegner, D.H.G.(1979):** Multicenter trials of praziquantel in human schistosomiasis, design and techniques, Bull. W.H.O., 57: 767.
- De Jesus, A.R.(2002):**Clinical and immunologic evaluation of 31 patients with acute *schistosomiasis mansoni*, J. Infect. Dis. 185: 98–105

- De Jesus, A.R.(2004):** Association of type 2 cytokines with hepatic fibrosis in human *Schistosoma mansoni* infection, *Infect. Immun.* 72 : 3391–3397.
- De Jesus, A.R. ; Araujo, I. O.; Becelar, A.; Magalhães, E.; Pearce, D.; Harn, M. and Carvalho, E.M. (2000):** Human immune responses to *Schistosoma mansoni* vaccine candidate antigens. *Infect Immunol* 68 : 2797.
- De-Jonge, N. (1988):** Detection of circulating anodic antigen by ELISA for seroepidemiology of schistosomiasis *mansoni*. *Trans. R. Soc. Trop. Med. Hyg.*, 82: 591.
- De Jonge ,N. (1989):** Time-resolved immunofluorometric assay (TR-IFMA) for the detection of schistosome circulating anodic antigen. *Trans. R. Soc. Trop. Med. Hyg.* 83 : 659–663.
- De Vlas, S.J. and Gryseels, B. (1992):** Underestimation of *Schistosoma mansoni* prevalences, *Parasitol. Today* 8 : 274–277.
- De Vlas, S.J.; Gryseels, B.;Van Oortmarssen, G.J.; Polderman, A.M. and Habbema, J.D.F. (1993):** A pocket chart to estimate true *Schistosoma mansoni* prevalence, *Parasitol. Today* 9 : 305–307 .
- De Vlas, S.J. ;Engels, D.; Rabello, A.L.T. ;Oostburg, B.F.J.; Van Lieshout, L.; Polderman, A, M.; Van Oortmarssen, G.J.; Habbema, J.D.F. and Gryseels, B. (1997):** Validation of a chart to estimate true *Schistosoma mansoni* prevalences from simple egg counts, *Parasitology* 114 : 113–121.
- Dean, L.L. and Podesta, R.B. (1984):** Electrophoretic patterns of protein synthesis and turnover in apical plasma membrane and outer bilayer of *Schistosoma mansoni*, *Biochim. Biophys. Acta* 799 : 106–114.
- Deelder, A.M. (1976):** *Schistosoma mansoni*: demonstration of two circulating antigens in hamsters. *Exp. Parasitol.* 40 : 189–197.
- Deelder, A.M. (1989):** Quantitative determination of circulating antigens in human *schistosomiasis mansoni* using an indirect haemagglutination assay. *Am. J. Trop. Med. Hyg.* 40 : 50–54 .
- Deelder, A.M.; De Jonge, N.; Boerman ,O.C.; Fillie ,Y.E.; Hilberath, G.W.; Rotmans, J.P.; Gerritse ,M.J. and Schut, D.W. (1998):** Sensitive determination of circulating anodic antigen in *Schistosoma mansoni* infected individuals by an enzyme-linked immunosorbent assay using monoclonal antibodies, *Am. J. Trop. Med. Hyg* 30:34-35.

- Dias, L.C.; Pedro, R. and Deberaldini, E.R. (1982):** Use of praziquantel in patients with schistosomiasis *mansoni* previously treated with oxamniquine and/or hycanthone: Resistance of *Schistosoma mansoni* to schistosomicidal agents. Trans. R. Soc. Trop. Med. Hyg., 76: 652.
- Doenhoff, M.J.; Cioli, D. and Utzinger, J. (2008):** Praziquantel: mechanisms of action, resistance and new derivatives for schistosomiasis, Current Opinion in Infectious Diseases 21 : 659–667.
- Doetze, J.; Satoguina, G.; Burchard, T. ;Rau, C.; Loliger, B. Fleischer and Hoerauf (2000) :**Antigen-specific cellular hyporesponsiveness in a chronic human helminth infection is mediated by T(h)3/T(r)1-type cytokines IL-10 and transforming growth factor-beta but not by a T(h)1 to T(h)2 shift, Int. Immunol. 12: 623–630.
- Dorn, A.; Vippagunta, S. R.; Matile, H.; Jaquet, C.; Vennerstrom, J. L. and Ridley, R. G. (1998):** An assessment of drug-haematin binding as a mechanism for inhibition of haematin polymerisation by quinoline antimalarials. Biochemical Pharmacology 55, 727–736.
- Duvall, R.H. and Dewitte, W. B. (1967):**An improved perfusion technique for recovering adult schistosomes from laboratory animals. Am. J. Trop. Med. Hyg. 16: 483-486.
- El-Gendy, S.D.; Osman, A.M. and Al-Sherbiny, M.M.(1999):** Epidemiology and immuodiagnosis of *schistosomiasis haematobium* in low endemic area in Egypt. J. Egypt. Soc. Parasitol., 29(1): 229-246.
- Elwi, A.M. (1976):** Pathology of schistosomiasis. Published by National Information, Documental Center, Dokki, Cairo, Egypt, 54-75.
- Engels, D. ; Sinzinkayo ,E. and Gryseels, B. (1996):** Day-to-day egg count fluctuation in *Schistosoma mansoni* infection and its operational implications, Am. J. Trop. Med. Hyg. 54 : 319–324.
- Engels, D.; Nahimana, S. ; De Vlas ,S.J. and Gryseels, B. (1997a):** Variation in weight of stool samples prepared by the Kato–Katz method and its implications, Trop. Med. Int. Health 2: 265–271.
- Engels, D.; Nahimana, S. ; de Vlas ,S.J. and Gryseels, B. (1997b):** Intraspecimen fecal egg count variation in *Schistosoma mansoni* infection, Am. J. Trop. Med. Hyg. 57 : 571–577.

- Engels, D. ;Savioli, L.; Stansfield, S.; Bundy, D.A.; Mitchell ,A. and Bhatia, R. (2002): Schistosomiasis and soil-transmitted helminth infections control efforts. Trans R Soc Trop Med Hyg 9(6) :577–579.
- Eraky, M.A. and Hammam, O.A. (2010): Evaluation of early efficacy of mefloquine on Egyptian *Schistosoma mansoni* strain in mice. Egypt. Journal of medical science 31(1):447-461.
- Fallon, P.G. and Doenhoff, M. J. (1994): Drug-resistant schistosomiasis: resistance to praziquantel and oxamniquine induced in *Schistosoma mansoni* in mice is drug specific. Am. J. Trop. Med. Hyg. 51:83-88 .
- Farid, Z.; Higashi, G.I.; Bssily, S.; Trabolsi, B. and Watten, R.H.(1980): Treatment of advanced hepatosplenic schistosomiasis with oxamniquine. Trans. R.Soc. Trop. Med. Hyg., 74: 400.
- Fenwick, Savioli, L.; Engels, D.; Bergquist, N.R. and Todd, M.H. (2003): Drugs for the control of parasitic diseases: current status and development in schistosomiasis, Trends Parasitol. 19 .
- Foster, R.(1973): The principle development of oxamniquine. Rev. Inst. Med. Trop. Sao Paulo, 15: 1.
- Fripp, P.J. (1967): The sites of (1–14C) glucose assimilation in *Schistosoma haematobium*, Comp. Biochem. Physiol. 23 : 893–898.
- Furlong, S.T. and Caulfield, J.P. (1988): *Schistosoma mansoni*: sterol and phospholipid composition of cercariae, schistosomula, and adults, Exp. Parasitol. 65: 222–231.
- Gatlin, M.R.; Black ,C.L.; Mwinzi, P.N.; Secor, W.E.; Karanja, D.M. and Colley, D.G. (2009): "Association of the gene polymorphisms IFN-gamma +874, IL-13 -1055 and IL-4 -590 with patterns of reinfection with *Schistosoma mansoni*". PLoS Neglected Tropical Diseases 3 (2): 375.
- Genovese, R. F.; Newman, D. B. and Brewer, T. G. (2000): Behavioral and neural toxicity of the artemisinin antimalarial, arteether, but not artesunate and artelinate, in rats. Pharmacol. Biochem. Behav. 67:37-44.
- Gobert, D.J.; Stenzel, D.P.; McManus and Jones ,M.K. (2003): The ultrastructural architecture of the adult *Schistosoma japonicum* tegument, Int. J. Parasitol. 33 :1561–1575.
- Goes, A.M. ; Gazzinelli, G.; Rocha, R.; Katz ,N. and Doughty, B.L., (1991): Granulomatous hypersensitivity to *Schistosoma mansoni*

- egg antigens in human schistosomiasis. III. In vitro granuloma modulation induced by immune complexes. *Am J Trop Med Hyg* 44 : 434.
- Golan, C.S. ;Brown, C.M. ;Cianci, S.T.; Furlong, S.T. and Caulfield, J.P. (1986):** Schistosomula of *Schistosoma mansoni* use lysophosphatidylcholine to lyse adherent human red blood cells and immobilize red cell membrane components, *J. Cell Biol.* 103: 819–828.
- Goldsmith, R.S. (1996):** Current Medical Diagnosis and Treatment, 35th ed. Lange medical book 918-980.
- Grogan, N. (1998):** Antigen-specific proliferation and interferon- γ and interleukin-5 production are down-regulated during *Schistosoma haematobium* infection, *J. Infect. Dis.* 177: 1433–1437.
- Gryseels, B.; Polman, K.; Clerinx, J. and Kestens, L., (2006):** Human schistosomiasis, *Lancet* 368 : 1106–1118.
- Gunderson, S.G. (1992):** Magnetic bead antigen capture enzyme-linked immunoassay in microtitre trays for rapid detection of schistosomal circulating anodic antigen. *J. Immunol. Methods* 148 : 1–8.
- Hagan, P. and Abath, F.G. (1992):** Recent advances in immunity to human schistosomiasis. *Mem. Inst Oswaldo. Cruz.*, 87(4): 95-98.
- Hang, I.M.; Warren, K.S. and Boros, D.L. (1974):** *S. mansoni* antigenic secretions and the etiology of egg granuloma in mice. *Exp. Parasitol.*, 35:288-298.
- Harder, A. (2002):** Treatment procedures and preventive controls for schistosomiasis. *Parasitology Today*. 16:364-366.
- Hardman, J.G; Limbird, L.E.; Molinoff, P.B.; Ruddon, R.W and Gilman, A.G. (1996):** Goodman and Gilman's the pharmacological Basis of Therapeutics. 9th ed. International Edition. McGraw- Hill
- Hassan, M.M.; El-motaiem, M.; Afifi, H.; Abaza, B.; El-Shafei, M. and Massoud, A.M. (2003):** In vitro effect of mirazid on *Schistosoma mansoni* worms. *J. Egypt. Soc. Parasitol.*, 33(3): 999-1008.
- He, Y.X.; Hu, Y.Q and Yu, Q.F. (1992):** Sensitivity of different isolates of *S. japonicum* from China to praziquantel. *Asian. J. Trop. Med. Publ. Hith.*, 23(2):261-263.
- Hennessy, M. and Spiers, J.P. (2007) :** A primer of the mechanics of P-glycoprotein in the multidrug transporter, *Pharmacol Res* 55.

- Higashi, G.I. and Farid, Z.(1979):** Oxamniquine fever-drug induced immune complex reaction. British Medical Journal, 2: 830.
- Hockley, D.J.(1973):** Ultrastructure of the tegument of *Schistosoma*, Adv. Parasitol. 11 : 233–305.
- Hockley, D.J. and McLaren, D.J. (1971):** The outer membrane of *Schistosoma mansoni*, Trans. R. Soc. Trop. Med. Hyg. 65 : 432.
- Hockley, D.J. and McLaren, D.J. (1973):** *Schistosoma mansoni*: changes in the outer membrane of the tegument during development from cercaria to adult worm, Int. J. Parasitol. 3 : 13–25.
- Holland, J. C.; Pellegrino, J. and Cozinelli, F. (1974):** Infection of mice with cercariae and schistosomula of *Schistosoma mansoni* by i.v. and s.c. routes. Rev. Inst. Med. Trop. S. Paulo. 16: 132-146.
- Hotez, J.; Fenwick, A.; Savioli, L. and Molyneux, D.H. (2009):** Rescuing the bottom billion through control of neglected tropical diseases, Lancet 373 : 1570–1575.
- Hotez, P.J. and Fenwick, A. (2009):** Schistosomiasis in Africa: an emerging tragedy in our new global health decade. PLoS Negl. Trop. Dis. 3:485.
- Hussein, M.R.; Abu-Dief, E.E.; El Hady, H.A.; Mahmoud, S.S. and Salah, E.M.(2005):** Quantitative comparison of infected *schistosomiasis mansoni* and *haematobium*: animal model analysis of the granuloma cell population. J. Egypt. Soc. Parasitol., 35(2): 467-76.
- Ismail, M.M.; Atta, A.A.; El-Raziky, E.H.; El-Ridi, A.M.S.; Attia, M.M.; El-Gamal, R.L. and El-Khashab, M.(1986):** Enzyme linked immunosorbent assay for the evaluation of praziquantel therapy in different groups of bilharzial patients. J. Egypt. Soc. Parasitol., 16(1): 269.
- Jaureguierry, S.; Perez, L.; Paris, L.; Bricaire, E.; Danis, M. and Gaumes, E.(2005):** Invasive schistosomiasis. Presse. Med., 34(21): 1641-5.
- Jha, S.; Kumar, R. and Kumar, R. (2006):** "Mefloquine toxicity presenting with polyneuropathy—a report of two cases in India". Trans R Soc Trop Med Hyg 100 (6): 594–96.
- Jiang, P.J.; Skelly, C.B. ;Shoemaker and Caulfield, J.P. (1996):** *Schistosoma mansoni*: the glucose transport protein SGTP4

- is present in tegumental multilamellar bodies, discoid bodies, and the surface lipid bilayers, *Exp. Parasitol.* 82 : 201–210.
- Jiraungkoorskul, W. ; Sahaphong, S.; Sobhon, P. ; Riengrojpitak, S.; and Kangwanrangsan, N. (2005):** Effects of praziquantil and artesunate on the adult *S.mekongi* harboured in mice. *Parasitology International* 54, 177–183.
- Johansen, M.V. ; Sithithaworn, P.; Bergquist ,R. and Utzinger, J. (2010):** Towards improved diagnosis of zoonotic trematode infections in Southeast Asia, *Adv. Parasitol.* 73 : 171–195.
- Jones, M.K.; Gobert, G.N.; Zhang, L.; Sunderland ,P. and McManus D.P. (2004):** The cytoskeleton and motor proteins of human schistosomes and their roles in surface maintenance and host–parasite interactions, *Bioessays* 26 : 752–765.
- Kalapothakis, E.; Hirst, E. and Evans, W.H. (1988):** Perturbations of the topography of the tegument of adult *Schistosoma mansoni*. A scanning electron microscope study. *Braz J Med Biol Res* 21: 961–969.
- Katz, N. ; Chaves, A. and Pellegrino, J. (1972):** A simple device for quantitative stool thick-smear technique in schistosomiasis mansoni, *Rev. Inst. Med. Trop. Sao Paulo* 14 : 397–400
- Katz, N.; Rocha, R.S.; Desouza, C.P.; Filho, P.; Bruce, J.L.; Coles, G.C. and Kinoti, G.K. (1991):** Efficacy of alternating therapy with oxamniquine and praziquantel to treat *Schistosoma mansoni* in children following failure of first treatment. *Am. J. Trop. Med. Hyg.*, 44:509.
- Keiser, J. and Utzinger, J. (2007):** Artemisinins and synthetic trioxolanes in the treatment of helminth infections. *Curr Opin Infect Dis* 20: 605–612
- Keiser, J.; Chollet, J. ;Xiao, S.H.; Mei, J.Y.; Jiao, P.Y.; Utzinger, J. and Tanner, M. (2009):** Mefloquine—an aminoalcohol with promising antischistosomal properties in mice, *PLoS Negl. Trop. Dis.* 3 : 350.
- Keiser, J.; Vargas, M. and Doenhoff, J. (2010):** Activity of Artemether and Mefloquine against Juvenile and Adult *Schistosoma mansoni* in Athymic and Immunocompetent NMRI Mice. *Am. J. Trop. Med. Hyg.*, 82(1): 112–114
- Khayyal, M.T. (1965):** Significance of worm shift in experimental chemotherapy of schistosomiasis mansoni with the emphasis on the action of anaesthetics. *Nature*, 205: 1331–1332.

- Khoo, S. ;Sarda, X.; Xu, J.P. ;Caulfield, M.R. ;McNeil, S.W. ;Homans, H.R. ;Morris and Dell, A. (1995):** A unique multifucosylated-3GalNAc beta 1-4GlcNAc beta 1-3Gal alpha 1-motif constitutes the repeating unit of the complex O-glycans derived from the cercarial glycocalyx of *Schistosoma mansoni*, J. Biol. Chem. 270 : 17114–17123.
- King, C.H. (2007):** Lifting the burden of schistosomiasis– defining elements of infection- associated disease and the benefits of King antiparasite treatment. J Infect Dis 196:653– 655
- King, C.H. and Dangerfield-Cha, M.(2008):** The unacknowledged impact of chronic schistosomiasis, Chronic Illness 4 : 65–79.
- Kohn, A.; López-Alvarez, M.L. and Katz, N. (1982):** Transmission and scanning electron microscopical studies in the tegument of male *Schistosoma mansoni* after oxamniquine treatment. Ann Parasit Hum Comp 57: 285-291.
- Kojima, S. (1998):** Schistosomiasis. In Topley and Wilson Microbiology and Microbial infections. 9th ed, Oxford University prss. Inc. New York, USA. 5:479-506.
- Kongs, A.; Marks, G.; Verle, P. and Van der Stuyft, P. (2001):** The unreliability of the Kato–Katz technique limits its usefulness for evaluating *S. mansoni* infections, Trop. Med. Int. Health 6 : 163–169.
- Kruatrachue, M.; Upatham, E.S.; Sahaphong, S. and Riengrojpitak, S. (1979):** Scanning and transmission electronmicroscopy of Mekong *Schistosoma* eggs and adults. S Asian J Trop Med Publ Hlth 10: 85-96.
- Kusel ,J.R. and Mackenzie, P.E. (1975):**The measurement of the relative turnover rates of proteins of the surface membranes and other fractions of *Schistosoma mansoni* in culture, Parasitology 71 : 261–273.
- Kusel ,J.R.; Mackenzie ,P.E. and McLaren, D.J. (1975):** The release of membrane antigens into culture by adult *Schistosoma mansoni*, Parasitology 71 : 247–259.
- Lambertucci, J.R. (1993):** *S. mansoni*: Pathological and clinical aspects. “In Human Schistosomiasis”. (Jordan, P.; Webbe, G. and Sturrock, R.F. eds) 3rd ed., CAB International, Walling ford, Oxon, U.k. p:33-85.

- Lambertucci, J.R.; Serufo, J.C.; Gerspacher, R.L.; Rayes, A.A.; Teixeira, R.; Nobre, V.; Antunes, C.M.(2000):** *S. mansoni* assessment of morbidity before and after control. *Acta. Trop.*, 77(1): 101-109.
- Lan, L.B.; Ayesh, S.; Lyubimov, E.; Pashinsky, I. and Stein, W.D. (1996):** Kinetic parameters for reversal of the multidrug pump as measured for drug accumulation and cell killing, *Cancer Chemother Pharmacol* 38: 181-190.
- Lanfredi ,R.M.; De Souza ,W. and Gomes, D.C. (1995):** Comparative study of four species of *Trichuris* Roederer, 1761 (Nematoda, Trichurinae) by scanning electron microscopy. *Mem Inst Oswaldo Cruz* 90: 489-496.
- Le, W. J.; You, J. Q.; Yang, Y. Q.; Mei,J. Y.; Guo,H. F.; Yang,H. Z. and Zhang, C. W.(1982):** Studies on the efficacy of artemether in experimental schistosomiasis. *Acta Pharmaceut. Sin.* 17:187-193. (In Chinese.)
- Le, W. J.; You, J. Q. and Mei,J. Y. (1983):** Chemotherapeutic effect of artesunate in experimental schistosomiasis. *Acta Pharmaceut. Sin.* 18:619.
- Leiper, R.T. (1918):** researches on Egyptian bilharziasis. *The Journal of The Royal Army Medical Corps*, 30:235-242. Quoted from **Woodruff, A.W.(1979):** *Medicine in the tropics*. 1st ed. Churchill Livingstone Edinburgh, London, p.173.
- Leitch, B.; Probert, A.J. and Runham, N.W. (1984):** The ultrastructure of the tegument of adult *Schistosoma haematobium*, *Parasitology* 89 : 71-78
- Liang, U. S.; Bruce,J. I. And Boyd, D. A. (1987):** Laboratory cultivation of *Schistosoma* vector snails and maintenance of *schistosoma* life cycle. In *Proceedings of the 1st Sino-American Symposium on biotechnology and parasitic diseases*, Department of Parasitology, National Yangming Medical College, Taipei, Taiwan 1: 34- 48.
- Lima, S.F.; Vieira, L.Q.; Harder, E.R. and Kusel, J.R. (1994):** Effects of culture and praziquantel on membrane fluidity parameters of adult *Schistosoma mansoni*. *Parasitology*, 19: 57-64.
- Loverde, P.T. and Chen,. L (1991):** "Schistosoma female reproductive development". *Parasitology Today* 7 (11): 303-8.

- MacGregor, A.N. ; Kusel, J.R. ; and Wilson, R.A. (1988):** Isolation and characterisation of discoid granules from the tegument of adult *Schistosoma mansoni*, Parasitol. Res. 74 : 250–254
- Machado-Silva, J.R.; Galvão, C.; Oliveira, R.M.F.; Presgrave, O.A.F. and Gomes, D.C. (1995):** *Schistosoma mansoni* Sambon, 1907: Comparative morphological studies of some Brazilian strains. Rev Inst Med Trop São Paulo 37: 441-447.
- Maguire, J.D.; Krisin, H.; Marwoto, H.; Richie, T.L.; Fryauff, D.J. and Baird, J.K. (2006):** "Mefloquine is highly efficacious against chloroquine-resistant *Plasmodium vivax* malaria and *Plasmodium falciparum* malaria in Papua, Indonesia". Clin Infect Dis 42 (8): 1067–72.
- Maizels, R.M. ; Bundy, D.A. ; Selkirk, M.E. ; Smith D.F. and Anderson, R.M. (1993):** Immunological modulation and evasion by helminth parasites in human populations, Nature 365 : 797–805.
- Makanga, M. and Krudsood, S. (2009):** The clinical efficacy of artemether/lumefantrine (Coartem)]. Malar J. 8 (1): S5.
- Malaquias, L.C.(1997):** Cytokine regulation of human immune response to *Schistosoma mansoni*: analysis of the role of IL-4, IL-5 and IL-10 on peripheral blood mononuclear cell responses, Scand. J. Immunol. 46 : 393–398.
- Manneck, T.; Haggenmüller, Y. and Keiser, J. (2010):** Morphological effects and tegumental alterations induced by mefloquine on schistosomula and adult flukes of *Schistosoma mansoni*, Parasitology 137 : 85–98.
- Manson, P.(1902):** Report of a case of bilharzias from the west Indies. J. Trop. Med., 5: 384. Quoted from **(Jordan, P.; Webbe, G. and Sturrock, R.E.(1993):** Human Schistosomiasis., University Press Cambridge, UK.
- Markell, E.K.; Voge, M. and John, D.T. (1999):** Medical Parasitology. 8th ed., Saunders Company.
- Massoud, A.M.; El-Ebiary, F.H. and Abdel-Salam, N.F.(2004):** Effect of myrrh extract on the liver of normal and bilharzially infected mice; An ultrastructural study. J. Egypt. Soc. Parasitol., 34(1): 1-21.
- McLaren, D.J.(1980):** *Schistosoma mansoni*: The parasite surface in relation to host immunity. John Willey & Sons, Ltd., England, 229.
- McLaren, D.J. and Hockley, D.J. (1977):** Blood flukes have a double outer membrane. Nature 269: 147-149.

- McLaren, D.J. ; Clegg, J.A. and Smithers, S.R. (1975):** Acquisition of host antigens by young *Schistosoma mansoni* in mice: correlation with failure to bind antibody in vitro, *Parasitology* 70 : 67–75
- Mc Manus, D.P. (1999):** The search for a vaccine against schistosomiasis—a difficult path but an achievable goal. *Immunol Rev* 171 : 149.
- Meuleman, E.A.; Holzmman, P.J. and Peet, R.C.(1980):** The development of daughter sporocysts inside the mother sporocyst of *Schistosoma mansoni* with special reference to the ultrastructure of the body wall. *Z Parasitenkd.*;61(3):201-12.
- Medhat, A.; Zarzour, A. and Nafeh, M.(1997):** Evaluation of an ultrasonographic score for urinary bladder morbidity in *schistosomiasis haematobium* infection. *AM. J. Trop. Med. Hyg.*, 57: 16-19.
- Mehlhorn, H.; Frenkel, J.K. Becher, B.; Andrews, P. and Thomas, H. (1983):** Light and electron microscopic studies on *Schistosoma mansoni* granulomas of mouse livers following treatment with praziquantel. *Tropenmedizin Und Parasitologie.* 33:299-239.
- Melman, S.D.; Steinauer M.L., ;Cunningham, C. ;Kubatko, L.S.; Mwangi, I.N.; Wynn, N.B.; Mutuku M.W, ;Karanja, D.M.S.; Colley, D.G.; Black, C.L.;Secor, W.E.; Mkoji, G.M. and Loker, E.S. (2009):** Reduced susceptibility to praziquantel among naturally occurring Kenyan isolates of *Schistosoma mansoni*, *PLoS Neglected Tropical Diseases* 3 :504
- Meyer, F. ; Meyer, H. and Bueding, E. (1970):** Lipid metabolism in the parasitic and free-living flatworms, *Schistosoma mansoni* and *Dugesia dorotocephala*, *Biochem. Biophys. Acta* 210 : 257–266.
- Middleton, W.; Kurtz, A.B. and Hertzberg, B.S.(2004):** Ultrasound; the requisites. 2nd ed. 66:67-179.
- Miller, F.H.; Tulloch, G.S. and Kuntz, R.E. (1972):** Scanning electron microscopy of integumental surface of *Schistosoma mansoni*. *J Parasitol* 58: 693-698.
- Moffat, D. and Kusel ,J.R. (1992):** Fluorescent lipid uptake and transport in adult *Schistosoma mansoni*, *Parasitology* 105 : 81–89.
- Montenegro, S.M. (1999):** Cytokine production in acute versus chronic human *schistosomiasis mansoni*: the cross-regulatory role of interferon-gamma and interleukin- 10 in the responses of

- peripheral blood mononuclear cells and splenocytes to parasite antigens, *J. Infect. Dis.* 179 : 1502–1514: 133–136.
- Montresor, A. ; Crompton, D.W.T.; Bundy, D.A.P. ; Hall, A. and Savioli, L. (1998):** Guidelines for the evaluation of soil-transmitted helminthiasis and schistosomiasis at community level, World Health Organization, Geneva .
- Mordi, M.N.; Mansor, S.N.; Navaratnam, V. and Wernsderfer, W.H.(1997):** Single dose pharmacokinetics of oral artemether in healthy Malaysian volunteers *Br J clin. Pharmacol* 5:43-363.
- Morris, G.P. and Threadgold, L.T. (1967):** Ultrastructure of the tegument of adult *Schistosoma mansoni*. *J Parasitol* 54: 15-27.
- Morris, G.P. and Threadgold, L.T. (1968):** Ultrastructure of the tegument of adult *Schistosoma mansoni*, *J. Parasitol.* 54 : 15–27.
- Mwatha, J.K. (1998):** High levels of TNF, soluble TNF receptors, soluble ICAM-1, and IFN-gamma, but low levels of IL-5, are associated with hepatosplenic disease in human *schistosomiasis mansoni*, *J. Immunol.* 160 : 1992–1999.
- N'Goran, E. K.; Utzinger, J.; Gnaka, H. N.; Yapi ,A.; N'Guessan, N. A.; Kigbafori, S. D.; Lengeler, C.; Chollet, J.; Xiao, S. H. and Tanner, M. (2003):** Randomized, double-blind, placebo-controlled trial of oral artemether for the prevention of patent *Schistosoma haematobium* infections. *Am. J. Trop. Med. Hyg.* 68:24-32.
- Nishimura, K. and Hung, T. (1997):** Current views on geographic distribution and models of infection of neurohelminthic diseases, *J. Neurol. Sci.* 145 : 5–14.
- Nosten, F. and Brasseur. P. (2002):**Combination therapy for malaria: the way forward? *Drugs* 62:1315-1329.
- Obeng, B.B. ; Aryeetey, Y.A. ; de Dood, C.J. ; Amoah, A.S. ; Larbi, I.A. ; Deelder, A.M. ; Yazdanbakhsh, M.; Hartgers, F.C. ; Boakye, D.A. ; Verweij, J.J. ; van Dam G.J. and van Lieshout L. (2008):** Application of a circulating-cathodic-antigen (CCA) strip test and real-time PCR, in comparison with microscopy, for the detection of *Schistosoma haematobium* in urine samples from Ghana, *Ann. Trop. Med. Parasitol.* 102 : 625–633.
- Ofori-Adjei, D.; Adjepon-Yamoah, K.K. and Lindstorm, B.(1988):** Oral praziquantel kinetics in normal and *Schistosoma haematobium* infected subjects. *Therapeutic Drug Monitoring*, 10:45.
- Okabe, K. and Tanaka, T. (1961):** Urine precipitation reaction for *schistosomiasis japonica*. *Kurume Med. J.* 8 : 24–29.

- Oliveira, M.F.; d'Avila, J.C. and Torres, C.R. (2000): "Haemozoin in *Schistosoma mansoni*". Molecular and Biochemical Parasitology 111 (1): 217-21.
- Oliveira, G.; Rodrigues, N. B.; Romanha, A. J.; Bahia, D. (2004): Genome and genomic of Schistosomes Canadian Journal of Zoology 82 (2): 375-390 .
- Otubanjo, O.A.(1981): Astiban-induced damage to tegument and the male reproductive system. Exp. Parasitol. 52: 161-170.
- Pearce, E.J. and MacDonald, A.S. (2002) : The immunobiology of schistosomiasis, Nat. Rev. Immunol. 2 : 499-511.
- Peeling, R.W. ; Smith, P.G. and Bossuyt, P.M. (2006): A guide for diagnostic evaluations, Nat. Rev. Microbiol. 4 (9) pp. S2-6
- Pellegrino, J. and Faria, J.(1965): The oogram method for the screening of drugs in *schistosomiasis mansoni*. AM. J. Trop. Med. Hyg., 14(3):363-369
- Perez,H. and Terry,R.J. (1973): The killing of adult *Schistosoma mansoni* in vitro in the presence of antisera to host antigenic determinants and peritoneal cells, Int. J. Parasitol. 3 : 499-503.
- Perez, H.; Clegg, I. A. and Smithers, S. R. (1975): Acquired immunity to *Schistosoma mansoni* in the rat: Measurement of immunity by the lung recovery technique. Parasitol. 69: 349-359
- Peters, A. P. and Warren, K. S. (1969): A rapid method of infecting mice and other laboratory animals with *S. mansoni* subcutaneous injection. J. Parasitol. 55: 558-563.
- Pica-Mattoccia, L. and Cioli, D. (2004): Sex- and stage-related sensitivity of *Schistosoma mansoni* to in vivo and in vitro praziquantel treatment. International Journal for Parasitology 34: 527-533.
- Polderman, A.M.; Gryseels, B.; Gerolod, J.L.; Mpania-Kayitseshonga, S. and Manshanda, J.P.((1984): Side effects of praziquantel in the treatment of *Schistosoma mansoni* in Maniema, Zaire. Trans. R. Soc. Trop. Med. Hyg., 78: 752.
- Pontes, L.A.; Dias-Neto, E. and Rabello, A. (2002): Detection by polymerase chain reaction of *Schistosoma mansoni* DNA in human serum and feces, Am. J. Trop. Med. Hyg. 66 : 157-162.
- Pontes, L.A. ; Oliveira, M.C. ; Katz, N. ; Dias-Neto ,E. and Rabello, A. (2003): Comparison of a polymerase chain reaction and the

- Kato-Katz technique for diagnosing infection with *Schistosoma mansoni*, Am. J. Trop. Med. Hyg. 68 : 652-656. 1996: 319-324.
- Price, R.; van Vugt, M. ; Phaipun, L.; Luxemburger, C. ; Simpson, J.; McGready, R. ; ter Kuile, F.; Kham, A. ; Chongsuphajaisiddhi, T.; White, N. J. and Nosten, F. (1999): Adverse effects in patients with acute falciparum malaria treated with artemisinin derivatives. Am. J. Trop. Med. Hyg. 60:547-555.
- Rabello, A.L.; Rocha, R.S.; De-Oliveira, J.P.; Katz, N. and Lambertucci, J.R.(1992): Stool examination and rectal biopsy in the diagnosis and evaluation of therapy of schistosomiasis *mansoni*. Rev. Inst. Med. Trop. Sao Paulo, 34(6): 601.
- Race, G.J.; Martin, J.H.; Moore, D.V. and Larsh, J.E. (1971): Scanning and transmission electronmicroscopy of *Schistosoma mansoni* eggs, cercariae and adults. Am J Trop Med Hyg 20: 914-927.
- Raso, A. ;Tafuri, N. ; Monteiro, E.R. and Tafuri, W.L. (2006): The tumoral form of cerebellar schistosomiasis: case report and measure of granulomas, Rev. Soc. Bras. Med. Trop. 39 : 283-286.
- Redman, C.A.; Robertson, A.; Smith, J.M. and Miller, S.C.(1996): Praziquantel: An urgent and exiting challenge. Parasitol. Today., 12:14.
- Retick, A.B.; Darracott-Voughan, E. and Wein, A.J.(2002): Parasitic diseases of the urinary tract. Cambell's urology, 8th ed. Saunders Press, :763-780.
- Rey, A. and Luís, S. (1991): Parasitology. Rio de Janeiro, RJ: Editora Guanabara Koogan S.A. 351:62.
- Reynolds, S.R.; Dahl, C.E. and Harn, D.A., (1994): T and B epitope determination and analysis of multiple antigenic peptides for the *Schistosoma mansoni* experimental vaccine triose-phosphate isomerase. J Immunol 152 : 193
- Roberts, S.M. ; Aitken, R.; Vojvodic ,M.; Wells, E.and Wilson, R.A. (1983): Identification of exposed components on the surface of adult *Schistosoma mansoni* by lactoperoxidase-catalysed iodination, Mol. Biochem. Parasitol. 9 : 129-143.
- Rogers, M.V. and McLaren, D.J. (1987): Analysis of total and surface membrane lipids of *Schistosoma mansoni*, Mol. Biochem. Parasitol. 22 : 273-288

- Rogers, R.A.; Jack, R.M. and Furlong, S.T.(1993): Lipid and membrane protein transfer from human neutrophils to schistosomes is mediated by ligand binding, J. Cell Sci. 106 :485-491.
- Rogers, S.H. and Bueding, E. (1975) : Anatomical localization of glucose uptake by *Schistosoma mansoni* adults, Int. J. Parasitol. 5 : 369-371.
- Rollinson, D. and Southgate, V.(1987): The genus *Schistosoma*: a taxonomic appraisal. In: D. Rollinson and A.J. Simpson, Editors, The Biology of Schistosomes, from Genes to Latrines, Academic Press, London : 1-49.
- Ross, A. G. P.; Bartley, P. B.; Sleight, A. C.; Olds, G. R.; Li, Y. S. ; Williams, G. M. and McManus, D. P. (2002):Schistosomiasis. N. Engl. J. Med. 346:1212-1220
- Rumjanek, F.D. ;Campos, E.G. and Afonso, L.C. (1988): Evidence for the occurrence of LDL receptors in extracts of schistosomula of *Schistosoma mansoni*, Mol. Biochem. Parasitol. 28 : 145-152.
- Ruppel, A. and McLaren, D.J. (1986) :*Schistosoma mansoni*: surface membrane stability in vitro and in vivo, Exp. Parasitol. 62 : 223-236.
- Sabah, A. A.; Fletcher, C. ; Webbe, G. and Doenhoff, M. J. (1986): *Schistosoma mansoni*: chemotherapy of infections of different ages. Exp. Parasitol. 61:294-303.
- Saif, M.; Gaber, A.; Hassanein, G. and Khamis, S.(1978): Efficacy of oxamniquine in treatment of *S. mansoni* in close community on Egypt. J. Egypt. Med. Ass., 61:427.
- Sambon, L.W.(1907): Remarks on *Schistosoma mansoni*. Am. J. Trop. Med. Hyg., 10:303. Quoted from, Jordan, P.; Webbe, G. and Sturrock, R.E. (1993): Human Schistosomiasis, University Press Cambridge, UK. P.1.
- Samuelson, J.C. ; Caulfield , J.P. and David, J.R. (1982): Schistosomula of *Schistosoma mansoni* clear concanavalin A from their surface by sloughing, J. Cell Biol. 94 : 355-362.
- Saunders, N.; Wilson, R.A. and Coulson, P.S. (1987): The outer bilayer of the adult schistosome tegument surface has a low turnover rate in vitro and in vivo, Mol. Biochem. Parasitol. 25 : 123-131.
- Schutte, C.H.J.(1974): Studies on the South African strains of *S. mansoni*. Morphology of miracidium. South African Journal of Science, 70: 299-302.

- Senft, A.W.; Gibler, W.B. and Knopf, P.M. (1978): Scanning electron microscope observations on tegument maturation in *Schistosoma mansoni* grown in permissive and non-permissive hosts. *Am J Trop Med Hyg* 27: 258-266.
- Shao, Y.M.; Ayesb, S. and Stein, W.D. (1997): Mutually co-operative interactions between modulators of P-glycoprotein, *Biochim Biophys Acta* 1360 : 30-38.
- Silk, M.H.; Spence, I.M. and Gear, J.H.S. (1969): Ultrastructural studies of the blood fluke—*Schistosoma mansoni*. I. The integument, *S. Afr. Med. Sci.* 34 : 1-10.
- Silveira, A.M. (2004): Human *schistosomiasis mansoni*: intensity of infection differentially affects the production of interleukin-10, interferon-gamma and interleukin-13 by soluble egg antigen or adult worm antigen stimulated cultures, *Trans. R. Soc. Trop. Med. Hyg.* 98 : 514-519.
- Simpson, A.J.; Singer, D.; McCutchan, T.F.; Sacks, D.L. and Sher (1983): Evidence that schistosome MHC antigens are not synthesized by the parasite but are acquired from the host as intact glycoproteins, *J. Immunol.* 131 : 962-965.
- Sinclair, D.; Zani, B.; Donegan, S.; Olliaro, P. and Garner, P. (2009): "Artemisinin-based combination therapy for treating uncomplicated malaria". *Cochrane Database Syst Rev.* 8 : 3.
- Skelly, P.J. and Shoemaker, C.B. (1996): Rapid appearance and asymmetric distribution of glucose transporter SGTP4 at the apical surface of intramammalian-stage *Schistosoma mansoni*, *Proc. Natl Acad. Sci. USA* 93 : 3642-3646.
- Skelly, P.J. and Shoemaker, C.B. (2001): The *Schistosoma mansoni* host-interactive tegument forms from vesicle eruptions of a cyton network, *Parasitology* 122 : 67-73.
- Skelly, P.J.; Pfeiffer, R.; Verrey, F. and Shoemaker, C.B. (1999): SPRM1lc, a heterodimeric amino acid permease light chain of the human parasitic platyhelminth, *Schistosoma mansoni*, *Parasitology* 119 : 569-576.
- Smith, J.H.; Reynolds, E.S. and von Lichtenberg, F. (1969): The integument of *Schistosoma mansoni*, *Am. J. Trop. Med. Hyg.* 18 : 28-49.
- Smith, T.M. and Brooks Jr, T.J. (1969): Lipid fractions in adult *Schistosoma mansoni*, *Parasitology* 59 : 293-298.

- Smithers, S.R.; Terry, R.J. and Hockley, D.J. (1969):** Host antigens in schistosomiasis, Proc. R. Soc. Lond. B Biol. Sci. 171 : 483–494.
- Snyder, S.D. and Eric, S.L.(2000):** Evolutionary relationship among the schistosomatidae (Platyhelminthes: Digenea) and an Asian origin for *Schistosoma*. J. Parasitol., 86(2): 283-288.
- Steinmann, P. ;Keiser, J.; Bos, R.; Tanner ,M. and Utzinger, J. (2006):** Schistosomiasis and water resources development: systematic review, meta-analysis, and estimates of people at risk, The Lancet Infectious Diseases 6 : 411–425.
- Stothard, J.R.; Chitsulo, L.; Kristensen, T.K. and Utzinger, J. (2009):** Control of schistosomiasis in sub-Saharan Africa: progress made; new opportunities and remaining challenges. Parasitology 136:1665–1675.
- Sturrock, R.F.(1987):** Biology and ecology of human schistosomes. Baillier's Clinical Tropical Medicine and Communicable Disease, 2(2): 249.
- Symmer's, W.S.C. (1904):** Note on a new form of liver cirrhosis due to the presence of ova of Bilharzia. J. Pathol. Bact., 9: 237-239.
- Taylor, M.D.; LeGoff, L.; Harris, A.; Malone, E.; Allen, J.E. and Maizels, R.M. (2005):**Removal of regulatory T cell activity reverses hyporesponsiveness and leads to filarial parasite clearance in vivo, J. Immunol. 174 : 4924–4933.
- Ten Hove, R.J.; Verweij, J.J.; Vereecken, K.; Polman, K.; Dieye, L. and van Lieshout, L. (2008):** Multiplex real-time PCR for the detection and quantification of *Schistosoma mansoni* and *S. haematobium* infection in stool samples collected in northern Senegal, Trans. R. Soc. Trop. Med. Hyg. 102 : 179–185.
- Terry, R.J.(1994):** Human immunity to schistosomes: Concomitant immunity. Parasitol. Today, 10(10): 377.
- Uglen, G.L. and Read, C.P. (1975):** Sugar transport and metabolism in *Schistosoma mansoni*, J. Parasitol. 61: 390–397.
- Utzinger, J. and Keiser, J. (2004):** Schistosomiasis and soil-transmitted helminthiasis: common drugs for treatment and control, Expert Opinion on Pharmacotherapy 5 : 263–285.
- Utzinger, J.; Booth ,M.; N'Goran, E.K.; Muller, I.; Tanner, M. and Lengeler, C. (2001):**Relative contribution of day-to-day and intra-specimen variation in faecal egg counts of *Schistosoma mansoni* before and after treatment with praziquantel. Parasitology, 122(5):537-544

- Utzinger, J.; Chollet, J.; Tu, Z. W.; Xiao, S. H. and Tanner, M. (2002):** Comparative study of the effects of artemether and artesunate on juvenile and adult *Schistosoma mansoni* in experimentally infected mice. *Trans. R. Soc. Trop. Med. Hyg.* 96:318-323.
- Utzinger, J.; Raso, G.; Brooker, S.; de Savigny, D.; Tanner, M.; Ornbjerg, N.; Singer, B.H. and N'Goran, E.K. (2009):** Schistosomiasis and neglected tropical diseases: towards integrated and sustainable control and a word of caution. *Parasitology* 136, 1859–1874.
- Van Agtmael, M. A.; Eggelte, T. A. and van Boxtel, C. J. (1999):** Artemisinin drugs in the treatment of malaria: from medicinal herb to registered medication. *Trends Pharmacol. Sci.* 20:199-205.
- Van der Kleij, D. and Yazdanbakhsh, M. (2003) :** Control of inflammatory diseases by pathogens: lipids and the immune system, *Eur. J. Immunol.* 33 : 2953–2963.
- Van Etten, L. (1994):** Rapid diagnosis of schistosomiasis by antigen-detection in urine with a reagent strip. *J. Clin. Microbiol.* 32 : 2402–2406.
- Van Lieshout, L. (1995) :** Analysis of worm burden variation in human *Schistosoma mansoni* infections by determination of serum levels of circulating anodic antigen and circulating cathodic antigen. *J. Infect. Dis.* 172 : 1336–1342.
- Van Lieshout, L. (2000):** Immunodiagnosis of schistosomiasis by determination of the circulating antigens CAA and CCA, in particular in individuals with recent or light infections, *Acta Trop.* 77 : 69–80 .
- Van Nassauw, L. ; Toovey, S.; Van, O.p.; den Bosch, J.; Timmermans, J.P. and Vercruysse, J. (2008):** Schistosomicidal activity of the antimalarial drug, mefloquine, in *Schistosoma mansoni*-infected mice, *Travel Medicine and Infectious Disease* 6 : 253–258.
- Viana, I.R.C. (1994):** Interferon- γ production by peripheral blood mononuclear cells from residents of an area endemic for *Schistosoma mansoni*, *Trans. R. Soc. Trop. Med. Hyg.* 88 : 466–470.
- Von Lichtenberg, L. (1987):** Consequences of infections with Schistosomes. In: D. Rollinson and A.J. Simpson, Editors, *The*

- Biology of Schistosomes, from Genes to Latrines, Academic Press, San Diego : 185–232.
- Walter ,R.D.; Wittich, R.M. and Kuhlow, F. (1987): Filaricidal effect of mefloquine on adults and microfilariae of *Brugia patei* and *Brugia malayi*. *Trop Med Parsitol.* 38:55-6.
- Wang, L.D.; Chen, H.G. ; Guo, J.G.; Zeng, X.J. ;Hong, X.L.; Xiong, J.J. ;Wu, X.H. ;Wang, X.H. ;Wang, L.Y. ;Xia, G. ;Hao, Y. ;Chin D.P. and Zhou, X.N. (2009): A strategy to control transmission of *Schistosoma japonicum* in China, *N. Engl. J. Med.* 360 : 121–128.
- Warren, K.S. (1973): The pathology of schistosome infections. *Helminthological. Abstracts* 42:591.
- Warren, K.S. (1978): Hepatosplenic schistosomiasis. A great neglected disease of the liver. *Gut.* 19:572-577.
- Warren, K.S. and Mahmoud, A.A.F. (1976): Targeted mass treatment: A new approach to thee control of schistosomiasis. Reprinted from *Transaction of the Association of American Physicians*, 39: 195.
- White, N.J. (1999): Antimalarial drug resistance and combination chemotherapy. *Philos. Trans. R. Soc. London Ser. B Biol. Sci.* 354:739-749.
- White, N.J. (2008): Qinghaosu (artemisinin): the price of success, *Science* 320 : 330–334.
- Wiest, P.M.; Tartakoff, A.M.; Aikawa, M. and Mahmoud ,A.A. (1988): Inhibition of surface membrane maturation in schistosomula of *Schistosoma mansoni*, *Proc. Natl Acad. Sci. USA* 85 : 3825–3829.
- Williams, M.E.(1994): Leukocytes of patients with *Schistosoma mansoni* respond with a Th2 pattern of cytokine production to mitogen or egg antigens but with a Th0 pattern to worm antigens, *J. Infect. Dis.* 170 : 946–954.
- Wilson, R.A. and Barnes, P.E. (1974a): An in vitro investigation of dynamic processes occurring in the schistosome tegument, using compounds known to disrupt secretory processes, *Parasitology* 68 : 259–270.
- Wilson, R.A. and Barnes, P.E. (1974b): The tegument of *Schistosoma mansoni*: observations on the formation, structure and composition of cytoplasmic inclusions in relation to tegument function, *Parasitology* 68 : 239–258.

- Wilson, R.A. and Barnes, P.E. (1977): The formation and turnover of the membranocalyx on the tegument of *Schistosoma mansoni*, *Parasitology* 74 : 61–71.
- Wilson, R.A. and Coulson, P.S. (2009): "Immune effector mechanisms against schistosomiasis: looking for a chink in the parasite's armour". *Trends in Parasitology* 25 (9): 423–31.
- World Health Organization (1985): The control of schistosomiasis: report of a WHO expert committee. WHO Tech. Rep. Ser. 728: 1–114.
- World Health Organization (2002): WHO Expert Committee. Prevention and control of schistosomiasis and soil-transmitted helminthiasis. World Health Organ Tech Rep Ser., 912:1–57.
- World Health Organization (2010): Increasing access to health workers in remote and rural areas through improved retention: global policy <http://whqlibdoc.who.int/ng>.
- Xiao, S.H. and Catto, B.A. (1989): In vitro and in vivo studies of the effect of artemether on *Schistosoma mansoni*. *Antimicrob Agents Chemother* 33 : 1557–1562.
- Xiao, S.H.; Shi, Z.G.; Zhuo, S.J.; Wang, C.Z.; Zhang, Z.G.; Chu, B.; Zheng, J. and Chen, M.G. (1996): Field studies on the preventive effect of oral artemether against schistosomal infection. *Chin. Med. J.* 109: 272–275.
- Xiao, S. H.; Hotez, P. J. and Tanner, M. (2000): Artemether, an effective new agent for chemoprophylaxis against schistosomiasis in China: its in vivo effect on the biochemical metabolism of the Asian schistosome. *Southeast Asian J. Trop. Med. Public Health* 31:724–732.
- Xiao, S.H. ;Shen, B.G.; Chollet, J.; Utzinger, J. and Tanner, M. (2001): Tegumental alterations in juvenile *Schistosoma haematobium* harboured in hamsters following artemether treatment. *Parasitol Int* 50: 175–183.
- Xiao, S. H., Tanner, M.; N'Goran, E. K.; Utzinger, J.; Chollet, J.; Bergquist, R.; Chen, M. G. and Zeng, J. (2002): Recent investigations of artemether, a novel agent for the prevention of schistosomiasis japonica, mansoni and haematobia. *Acta Trop.* 82:175–181.
- Xiao, S.H.; Keiser, J.; Chollet, J.; Utzinger, J.; Dong, Y.; Endriss, Y.; Vennerstrom, J.L. and Tanner, M. (2007): In vitro and in vivo activities of synthetic trioxolanes against major human

- schistosome species, Antimicrob. Agents Chemother. 51 : 1440–1445.
- Xu, X. and Caulfield, J.P. (1992):** Characterization of human low density lipoprotein binding proteins on the surface of schistosomula of *Schistosoma mansoni*, Eur. J. Cell Biol. 57 : 229–235.
- Yang, W.; Jackson, D.C.; Zeng, Q. and Mc-Manus, D.P.(2000):** Multiepitope schistosome vaccine candidates tested for protective immunogenicity in mice. Vaccine, 19(1): 103-113.
- Zhang, C.; Xio, S.; Utizinger, J.; Chollet, J.; Keiser, J. and Tanner, M.(2009):** Histopathological and changes in adult *S.jabonicum* harbored in mice treated with single dose of mefloquine .Parasitology Research 104:1407-1416.
- Zhou, Y. and Podesta, R.B. (1989):** Effects of serotonin (5HT) and complement C3 on the synthesis of the surface membrane precursors of adult *Schistosoma mansoni*, J. Parasitol. 75 : 333–343.
- Zouain, C.S.; Gustavson, S. ; Oliveira ,S.C. ; Azevedo, V. ; Alves, J.B. and Goes , A.M. (2001):**The role of IL-10 and IgG1 in the protection and granulomatous response in *Schistosoma mansoni* P24-immunized mice. Vaccine 19 : 1218.