

REFERENCES

Andrew CP, Martha KT, Edmund SS, et al. (2007): Urologic Imaging without X-rays: Ultrasound, MRI, and Nuclear Medicine Article Last Updated: Jan 4.

Auge BK, Munver R, Kourambas J, et al (2002): Endoscopic management of symptomatic caliceal diverticula: a retrospective comparison of percutaneous nephrolithotripsy and ureteroscopy. J Endourol; 16:557–63.

Bates JA. (2004): Abdominal ultrasound. How, Why and When, 7: 154-157.

Bellman GC, Pardalidis NP, Smith AG. (1995): Management of hemorrhage after percutaneous renal surgery. J Urol 1995.

Buchholz NP. (2001): Double infundibular obliteration with abscess formation after percutaneous nephrolithotomy. Urol Int; 66:46.

Cadeddu JA, Arrindell D, Moore RG. (1997): Near fatal air embolism during percutaneous nephrostomy placement. J Urol; 158:1519.

Chien GW and Bellman GC. (2002): Blind percutaneous renal access. J Endourol; 16:93-96

Cooper CS and Hedican SP. (2003): Mini-percutaneous ultrasonic nephrolithotripsy for pediatric Staghorn calculi. J Urol; 170: 1336–1337.

- Cuellar DC and Averch TD. (2004):** Holmium laser percutaneous nephrolithotomy using a unique suction device. J Endourol; 18: 780.
- Curtis R, Thorpe AC and Marsh R. (1997):** Modification of the technique of percutaneous nephrolithotomy in the morbidly obese patient. Br J Urol; 79:138–40.
- De la Rosette, Tsakiris P. Ferrandino M, et al. (2008):** Beyond Prone Position in Percutaneous Nephrolithotomy: A Comprehensive Review Eur Urol 54; 1262–1269.
- De Sio M, Autorino R., Quarto G. et al. (2008):** Modified supine versus prone position in percutaneous nephrolithotomy for renal stones treatable with a single percutaneous access: a prospective randomized trial. Eur Urol 54; (196 - 203) .
- Denby S, Fracs and Villis M. (2007):** Percutaneous nephrolithotomy in the supine position: A neglected approach? J. Endo. Vol. 21, N. 12.
- Denstedt JD, Clayman RV and Picus DD. (1991):** Comparison of endoscopic and radiological residual fragment rate following percutaneous nephrolithotripsy. JUrol; 145:703.
- Desai MR, Kukreja RA, Desai MM, et al. (2004):** A prospective randomized comparison of type of nephrostomy drainage following percutaneous nephrostolithotomy: Large bore versus small bore versus tubeless. J Urol; 172:565-7.
- Domenech A, Vivaldi B, Diaz C et al. (2008):** complications in percutaneous nephrolithotomy: A comparative study between the

supine and prone positions using the modified claviens system. Urology 72 (Suppl. 5A), November.

Drake RL, Vogl W, and Mitchell AW. (2007): Gray's anatomy; 4:272-293.

El-Assmy AM, Shokeir AA, El-Nahas AR, et al. (2007): Outcome of percutaneous nephrolithotomy: effect of body mass index. Eur Urol; 52 :199–205.

El-Assmy AM, Shokeir AA, Mohsen T, et al. (2007): Renal access by urologist or radiologist for percutaneous nephrolithotomy: Is it still an issue? J Urol; 178: 916 – 20.

Evans CP and Stoller ML. (1993): The fate of the iatrogenic retroperitoneal stone. J Urol; 150:827.

Fernstrom I and Johansson B. (1976): Percutaneous pyelolithotomy: A new extraction technique. Scand J Urol Nephrol; 10:257. Quoted from Vicentini et al., 2009.

Fuchs GJ, Moody JA, Gutierrez-Aceves J, et al. (2001): Complications of percutaneous renal surgery: prevention, recognition and management. Quoted from Smith et al., (2006) 20: 160-166.

Gerspach JM, Bellman GC, Stoller ML, Fugelso P. (1997): Conservative management of colon injury following percutaneous renal surgery. Urology; 49: 831. Quoted from Smith et al., (2006) 20: 160-166.

- Giblin JG, Lossef S and Pahira JJ. (1995):** A modification of standard Percutaneous nephrolithotripsy technique for the morbidly obese patient. *Urology*; 46:491–3. Quoted from El-Assmy et al., (2007).
- Giusti G, Piccinelli A, Taverna G, et al (2007):** Miniperc? No, thank you! *Eur Urol*; 51:810.
- Heidenreich A, Desgrandschamps F, Terrier F. (2002):** Modern approach of diagnosis and management of acute flank pain: review of all imaging modalities. *EurUrol*; 41:351.
- Hellawell GO, Mutch SJ, Thevendran G, et al. (2005):** Radiation exposure and the urologist: what are the risks? *J Urol*; 174:948.
- Hernandez DJ, Sinkov VA, Roberts WW, et al. (2001):** Measurement of bio-impedance with a smart needle to confirm percutaneous kidney access. *J Urol*; 166: 1520 – 1523.
- Holman E, Pasztor I, Khan AM. (1993):** Laparoscopically controlled and assisted percutaneous transperitoneal nephrolithotomy in a pelvic dystopic kidney. *J Endourol*; 7(4): 303–30.
- Hopper KD and Yakes WF. (1990):** The posterior intercostal approach for percutaneous renal procedures: risk of puncturing the lung, spleen, and liver as determined by CT. *AJR Am J Roentgenol*; 154:115.
- Hopper KD, Sherman JL, Luethke JM, et al. (1987):** The retrorenal colon in the supine and prone position. *Radiology* 162: 443- 446.

- Hopper KD, Sherman JL, Williams MD, et al. (1987):** The variable anteroposterior position of the retroperitoneal colon to the kidneys. *Invest. Radiol.* 22: 298-302.
- Jackman SV, Potter SR, Regan F, et al. (2000):** Plain abdominal x-ray versus computerized tomography Screening: sensitivity for stone localization after non-Enhanced spiral computerized tomography. *J Urol*; 164:308.
- Kerbl K, Picus DD and Clayman RV. (1994):** Clinical experience with the Kaye nephrostomy tamponade catheter. *Eur Urol*; 25:94
- Kidd CR and Conlin MJ. (2003):** Ureteroscopically assisted percutaneous renal access. *Urology*; 61:1244-1245.
- Kim HL and Schulman P. (2004):** The Paky, Hermes, Aesop, Zeus, and da Vinci robotic systems. *Urol Clin North Am*; 31:659-669.
- Kim IY and Clayman RV. (2006):** Surgical Renal Anatomy AUA Update Series volume (25) Lesson 40; 361- 367.
- Kim SC, Tinmouth WW, Kuo RL, et al. (2005):** using and choosing a nephrostomy tube after percutaneous nephrolithotomy for large or complex stone disease: A treatment strategy. *J Endourol*; 19 (3): 348 – 352.
- Kondas J, Szentgyorgyi E, Vaczi L and Kiss A. (1994):** Splenic injury: a rare complication of percutaneous nephrolithotomy. *Int Urol Nephrol*; 26:399. Quoted from Shah and Assimos (2006) in Smith's Textbook Of Endourology 20: 160 - 166.

- Koolpe HA and Lord B. (1990):** Eccentric nephroscopy for the incarcerated nephrostomy. *Urol Radiol*; 12:96.
- Krombach GA, Mahnken A, Tacke J, et al. (2001):** US-guided nephrostomy with the aid of a magnetic field-based navigation device in the porcine pelvicaliceal system. *J Vasc Interv Radiol* ; 12:623-628.
- Kuo RL, Paterson RF, Siqueira TM, et al. (2004):** In vitro assessment of lithoclast ultra intracorporeal lithotripter. *J Endourol*; 18: 153.
- Lam HS, Lingeman JE, Mosbaugh PG, et al. (1992):** Evolution of the technique of combination therapy for staghorn calculi: a decreasing role for extracorporeal shock wave lithotripsy. *J Urol*; 148:1058.
- Leder RA and Nelson RC. (2001):** Three-dimensional CT of the genitourinary tract. *J Endourol*; 15:37.
- Lee WJ, Smith AD, Cubelli V, et al. (1987):** Complications of percutaneous nephrolithotomy. *AJR Am J Roentgenol*; 148:177. Quoted from Shah and Assimos (2006) in Smith's Textbook Of Endourology 20: 160 - 166.
- Lemaitre L, Mestdagh P, Marecaux-Delomez J, et al. (2000):** Percutaneous nephrostomy: Placement under laser guidance and real-time CT fluoroscopy. *Eur Radiol* ; 10:892-895
- Lu HF, Shekarriz B, and Stoller ML. (2002):** Donor-gifted allograft urolithiasis: early percutaneous management. *Urology*; 59:25.

- Manohar T, Jain P and Desai M. (2007):** Supine percutaneous nephrolithotomy: effective approach to high – risk and morbidly obese patients. J Endourol 2007;21:44–9.
- Man-Tat NG, Wai-Ho S, Chi-Wai Ch, et al. (2004):** Supine position is safe and effective for percutaneous nephrolithotomy. J. Endo. Vol. 18, N. 5.
- Matlaga BR, Shah OD, Zagoria RJ, et al. (2003):** Computerized tomography guided access for percutaneous nephrostolithotomy. J Urol; 170:45.
- McAleer SJ and Loughlin KR. (2004):** Nephrolithiasis and pregnancy. Curr Opin Urol; 14:123.
- Meretyk S, Albala DM, Clayman RV, et al. (1992):** Endoureterotomy for treatment of ureteral strictures. J Urol; 147:1502.
- Miller NL, Matlaga BR, and Lingeman JE. (2007):** Techniques for fluoroscopic percutaneous renal access. J Urol; 178:15-22.
- Munver R, Delvecchio FC, Newman GE, et al. (2001):** Critical analysis of supracostal access for percutaneous renal surgery. J Urol; 166:1242.
- Mutazindwa T and Hussein T. (1996):** Imaging in acute renal colic: the intravenous urogram remains the gold standard. Eur J Radiol; 23: 238.
- Neto EA, Mitre AI, Gomes CM, et al. (2007):** Percutaneous nephrolithotripsy with the patient in a modified supine position. JUrol; 178 : 165-8.

- Ng MT, Sun WH Cheng CW et al. (2004):** Supine position is safe and effective for percutaneous nephrolithotomy. *J Endourol*; 18: 469-74.
- Norris TG. (2002):** Radiation safety in fluoroscopy. *Radiol Technol*; 73:511.
- Park S, Maxwell V, Meng, et al. (2006):** Percutaneous Nephrolithotomy Technical Aspects. In urinary stone disease. Edited by; Stoller M, and Meng M. (2006).38; 609-42
- Park S, Margaret S. and Pearle (2006):** Imaging for Percutaneous Renal Access and Management of Renal Calculi. *Urol Clin Nam* 33; 353–364.
- Patel A and Fuchs GJ. (1998):** Air travel and thromboembolic complications after percutaneous nephrolithotomy for staghorn stone. *J Endourol* 12: 68 – 72.
- Patel A, Soonawalla P, Shepherd SF, et al. (1996):** Long-term outcome after percutaneous treatment of transitional cell carcinoma of the renal pelvis. *J Urol* 1996; 155:868.
- Pearle MS, Corwin TS and Mullican MA. (2000):** Comparison of intra operative chest fluoroscopy, chest radiography and chest CT for detecting hydropneumothorax after percutaneous nephrostolithotomy . *J Endourol*; 14:A86.
- Pearle MS, Watamull LM and Mullican MA. (1999):** Sensitivity of noncontrast helical computerized tomography and plain film radiography compared to flexible nephroscopy for detecting

residual fragments after percutaneous nephrostolithotomy. J Urol; 162: 23.

Pietrow PK, (2006): Percutaneous Nephrolithotomy Indications and Outcomes in Current Clinical Urology, In urinary stone disease. Edited by; Stoller M, and Meng M. (2006).38; 609-42.

Pietrow PK, Auge BK, Zhong P, et al. (2003): Clinical efficacy of a combination pneumatic and ultrasonic lithotrite. J Urol; 169:1247.

Portis AJ, Laliberte MA, Drake S, et al. (2006): Intraoperative fragment detection during percutaneous nephrolithotomy: Evaluation of high magnification rotational fluoroscopy combined with aggressive nephroscopy. J Urol; 175:162-5.

Preet S, and Raj MP. (2007): Ultrasound-Guided Genitourinary Interventions: Ultrasound Clin: 2, 115–120.

Raj GV, Auge BK, Weizer AZ, et al. (2003): Percutaneous management of calculi within horseshoe kidneys. J Urol; 170:48.

Rana AM, Bhojwani JP, Junejo NN, et al. (2008): Tubeless PCNL with patients in supine position: procedure for all seasons? With comprehensive technique. Urology;71:581–5.

Roberts S, Bolton DM and Stoller ML. (1994): Hypothermia associated with percutaneous nephrolithotomy. Urology; 44:832.

Sakurai H, Mitsuhashi N, Harashima K, et al. (2004): CT- fluoroscopy guided interstitial brachtherapy with image-based treatment planning forum unrespectable locally recurrent rectal carcinoma. Brachytherapy; 3:222.

- Sampaio FJB. (2000):** Renal anatomy: endourologic considerations. Urol clin North Am; 27 : 585 – 607.
- Scoffone CM, Cracco CM, Poggio M, et al. (2008):** Endoscopic combined intrarenal surgery in Galdakao-Modified supine valdivia position for urolithiasis: a possible new PCNL standard. Eur Urol in Press.
- Sengupta S, donnellan S and Vincent M, et al. (2000):** CT Analysis of Caliceal Anatomy in the supine and prone positions; Journal Of Endourology Volume 14, Number 7; 555- 557.
- Shah O and Assimos DG. (2006):** Complications of Percutaneous Renal Surgery in Smith's Textbook Of Endourology 20: 160 - 166.
- Shoma AM, Eraky I, El-Kenawy MR and El-Kappany HA.(2002):**Percutaneous nephrolithotomy in the supine position: technical aspects and functional outcome compared with the prone technique. Urology; 60:388–92.
- Skolarikos A, Alivizatos G and de la Rosette J. (2005):** Percutaneous Nephrolithotomy and its Legacy, EurUrol 47: 22–28
- Smith AD, Badlani GH, Bagley, DH, et al. (2006):** Smith's Textbook Of Endourology; 12: 79-101.
- Smith AD, Badlani, GH, Bagley, DH et al. (2006):** Smith's Textbook Of Endourology; 20: 161.
- Soble JJ and Streem SB. (1998):** Prospective randomized comparison of percutaneous nephrostomy tract dilation techniques. J Endourol; 12:S115A.

- Stoller M, and Meng M. (2006):** Urinary stone disease 32; 621 – 630.
- Stoller ML, Wolf JS Jr and St Lezin MA. (1994):** Estimated blood loss and transfusion rates associated with percutaneous nephrolithotomy. J Urol; 152:1977.
- Teichman JM, Long RD and Hulbert JC. (1995):** Long-term renal fate and prognosis after staghorn calculus management. J Urol. 153-1403. Quoted from Smith et al., (2006) 20 :160 – 166.
- Thiruchelvam N, Mostafid H, Ubhayakar G. (2005):** Planning percutaneous nephrolithotomy using multidetector computed tomography urography, multi-planar reconstruction and three-dimensional reformatting. BJU Int; 95:1280.
- Valdivia J. GJ. Valle gerhold, J.A. Lopez lopez, et al. (1998):**Technique And Complications Of Percutaneous Nephroscopy: Experience With 557 Patients In The Supine Position The Journal Of Urology Vol. 160. 1975-1978.
- Valdivia JG. (2009):** Complete Supine Percutaneous Nephrolithotripsy Comparison With the Prone Standard Technique: The Time for Change From Prone to Supine Position Has Come! UroToday International Journal on the internet.
- Vicentini FC, Gomes CM, Danilovic A, et al.(2009):**Percutaneous nephrolithotomy: Current concepts. Indian J Urol; 25:4-10.
- Wein, Kavoussi, Novick P and Peters (2007):** Campbell, Walsh Urology. Ninth edition 46:1526-1565.

Wong C and Leveillee RJ. (2002): Single upper pole percutaneous access for treatment of greater than or equal to 5-cm complex branched staghorn calculi: is shockwave lithotripsy necessary? J Endourol; 16: 477.

Woodside JR, Stevens GF, Stark GL, et al. (1985). Percutaneous stone removal in children. J Urol; 134(6): 1166–1167.

Zeren S, Satar N, Bayazit Y, et al . (2002):Percutaneous nephrolithotomy in the management of pediatric renal calculi. J Endourol; 16(2): 75–78.

Zhou X, Gao X, Wen J et al. (2008): Clinical value of minimally Invasive percutaneous nephrolithotomy in the supine position under the guidance of real-time ultrasound: report of 92 cases.Urol Res; 36:111–4.