

REFERENCES

REFERENCES

Barger, A.C. and Herd, J.A. (1971) :

The renal circulation.

New Engl. J. Med., 35 : 43.

Barrier Steven L. (1982) :

Nephrotoxicity of cephalothin-aminoglycoside
interactions.

Arch. Intern. Med., 142 : 1754.

Barza, M.; Pinn, V.; Tang Vay, P. and Murray, T. (1978) :

Nephrotoxic of cephalosporins and aminoglycoside
alone and a combination in a rat model.

J. Antimicrob. Chemother., 4 supp : 59.

Becker, R.F.; Wilson, J.W. and Gehwerler, J.A. (1971) :

The urinary system. In Becker, R.F.; Wilson, J.W.
and Gehwerler, J.A. (Ed.). The anatomical basis
of medical practice.

Williams, Baltimore, p. 35.

Bennett, W.M.; Luft, F. and Porter, G.A. (1980) :

Pathogenesis of renal failure due to aminoglycoside.

Am. J. Med., 69 : 767.

Bennett, W.M. (1981) :

Aminoglycoside. In Klaher, S. and Mossry, S.G.
(Ed.). Contr. Nephrology.
Plenum Medical Book, New York, p. 466.

Bennett, W.M.; Plamp, G.E.; Elliot, W.C.; Parker, R.A. and
Proter, G.A. (1982) :

Effect of basic aminoacid and aminoglycoside on
³H gentamicin uptake in cortical slices of rat
and human kidney.
J. Lab. Clin. Med., 99 : 156.

Bennett, W.M.; Elliot, W.C.; Houghton, D.C. and Porter, G.A.
(1982) :

Reduction of experimental gentamicin nephrotoxi-
city in rat, dietary calcium loading.
Antimicrob. Agent Chemother., 22 : 508.

Berndt, W.O. (1981) :

Use renal function tests in the evaluation of
nephrotoxicity effect. In Cumming N.B. and Klahr,
S. (Ed.). Chronic renal diseases.
Plenum Medical Book, London, p. 14.

Betts Robert, F. (1985) :

The amikain surveillance studies. Part I Empiric
therapy.
Challenges and Choeses, p. 3-4.

- Blaser, T.; Ruttiman, S.; Bhend, H. and Luthy, R. (1983) :
Increase of amikacin half life during therapy.
Antimicrob. Agent Chemother., 23 : 888.
- Bloom, W. and Fawcett, D.W. (1975) :
The urinary system. In Bloom, W. and Fawcett,
D.W. (Ed.). A text book of histology.
W.S. Saunders, London, p. 766.
- Bowman, R.L.; Silverblatt, F.J. and Kaloyands, G.J. (1977) :
Comparison of the nephrotoxicity of netilmicin
and gentamicin in rat.
Antimicrob. Agent Chemother., 12 : 474.
- Brater, D.C. and Chennavasin, P. (1982) :
Antibiotic. In Brater, D.C. and Chennavasin, P.
(Ed.). Drug use in renal diseases.
Adis Health Science Press, Tokyo, p. 90.
- Brenner, B.M. and Lazarus, J.M. (1983) :
Aminoglycoside nephrotoxicity. In Brenner, B.M.
and Lazarus, J.M. (Ed.). Acute renal failure.
W.B. Saunders, London, p. 284.
- Brochner, M.J. and Rodbro, P. (1976) :
Selection of routine methods for determination of
glomerular filtration rate in adult patients.
Scand. J. Clin. Invest., 36 : 35.

Bryan, L.E. and Van Den Elzen, H.M. (1977) :

Effect of membrane energy mutation.

Antimicrob. Agent Chemother., 12 : 163.

Bryan, L.E. and Kwan, S. (1984) :

Mechanism of aminoglycoside resistance of anaerobic bacteria.

J. Antimicrob. Chemother., 8 : 201.

Carlton, E.C. (1982) :

Renal function test. In Harrison, H.; Gittes, R.

F.; Porlmitter, A.D. and Stomey, T.A. (Ed.).

Chambbell Urology.

W.B. Saunders, London, p. 109.

Chan, R.A.; Benner, E.J. and Hoeprich, P.D. (1972) :

Gentamicin ototoxicity and nephrotoxicity.

Ann. Intern. Med., 76 : 773.

Collar, V.U.; Lietman, P.S. and Mitch, W.E. (1979) :

Evidence of luminal uptake of gentamicin in perfused rat kidney.

J. Pharmacol. Exp. Ther., 210 : 701.

Cone, L.A. (1982) :

A survey of prospective, controlled clinical trials of gentamicin, tobramycin, amikacin and netilmicin.

Clinical Therapeutic, 5 : 155.

Cronin, R.E.; Bulger, R.E.; Southen, P. and Henrich, W.L.

(1980) :

Natural history of aminoglycoside in the dog.

J. Lab. Clin. Med., 95 : 114.

Cross, A.S.; Opal, M.S. and Kopecko, D.J. (1983) :

Progressive increase in antibiotic resistance of Gram-negative bacterial isolates. Walter Reed Hospital 1976-1980. Specific analysis of gentamicin, tobramycin and amikain resistance.

Arch. Intern. Med., 143 : 2075.

Culter Ralphe, Gyselynck, A.M.; Fleet, P. and Forrey, A.

(1972) :

Correlation of serum creatinine concentration and gentamicin half life.

J. A. M. A., 21 : 1037.

Dahlager, J.I. (1980) :

The effect of netilmicin and other aminoglycoside on renal function.

Scand. J. Infect. Dis., 23 : 96.

Dahlager, J.I. (1981) :

The role of the aminoglycoside in modern clinical practice.

A doctor to doctor round table discussion hold in Nice, France, December, p. 35.

Davies, D.V. and Coupland, R.E. (1967) :

Kidneys. In Davies, D.V. and Coupland, R.E. (Ed.)

Gray's Anatomy.

Longmans, London, p. 211.

Davis, J. and Courvalin, P. (1977) :

Mechanism of resistance aminoglycoside.

Am. J. Med., 23 : 621.

Dellinger, P.; Murphy, T.; Pinn, V.; Barza, M. and Weinstein,

L. (1976) :

The protective effect of cephalothin against gentamicin induce nephrotoxicity in rats.

Antimicrob Agent Chemother., 9 : 172.

Duart, C.G.; Elveback, L.R. and Liedtke, R.R. (1985) :

Creatinine. In Duart, C.G. (Ed.). Creatinine renal function tests.

Little Brown, p. 1.

Dukes, M.G. and Elis, J. (1982) :

Nephrotoxicity of aminoglycoside. In Dukes, M.G. and Elis, J. (Ed.). Side effect of drug annual 6,

Excerpta Medica, Oxford, p. 48.

- Edward, G.Q.; Smith, G.R.; Baughman, K.L.; Rogers, J.F. and
Lietman, P.S. (1976) :
Concentration of gentamicin and amikain in human
kidney.
Antimicrob. Agent Chemother., 9 : 925.
- El-Hawary, M.B.; Khayyal, M.T. and Isaak, Z. (1985) :
Antibiotics. In El-Hawary, M.B.; Khayyal, M.T.
and Isaak, Z. (Eds.). Handbook of pharmacology.
The Scientific Book Centre, Cairo, p. 618.
- Elliot, W.C.; Houghton, D.C.; Gilbert, D.N.; Humler, B.J. and
Bennett, W.M. (1982) :
I- Degree and permanence of acquired insensitivity.
Lab. Clin. Med., October, p. 501.
- Elliot, W.C.; Houghton, D.C.; Gilbert, D.N.; Humler, B.J. and
Bennett, W.M. (1982) :
II- definition of condition necessary to induce
acquired insensitivity.
Lab. Clin. Med., October, p. 513.
- Evan, A.P.; Huser, J.; Arasthi, P.S.; Rankin, L.I. and Luft,
F.C. (1979) :
Gentamicin induce glomerular injury.
J. Antimicrob. Agent Chemother., 19 : 703.

Fahmy, M.F. (1986) :

Aminoglycoside therapy in renal failure. In Fahmy, M.F. (Ed.). Collective urology. Cairo Univ., p. 469.

Feldman Stuart, Meng Ynn Wany and George, J. Kaloyanids (1982) :

Aminoglycoside induce a phospholipidosis in the renal cortex of the rat. *J. Pharmacol. Exper. Ther.*, 220 : 518.

Ganong, W.F. (1983) :

Renal function. In Ganong, W.F. (Ed.). Review of medical physiology. Los Altos, California, p. 532.

Gardner, E.; Gray, D.T. and Orhilly, R. (1975) :

Kidney, ureter, suprarenal gland. In Gardner, E. and Orhilly, R. (Ed.). A regional study of human structure. W.B. Saunders, London, p. 406.

Gilbert, D.N.; Starr, P.P.; Bennett, W.M.; Houghton, D.C. and

Proter, G. (1978) :

Comparative nephrotoxicity of gentamicin and tobramycin. *Antimicrob. Agents Chemother.*, 13 : 34.

Gray, N.E.; Buzzeeo, L.; Saloki, J. and Eisenger, R.B. (1976) :

Gentamicin associated acute renal failure.

Arch. Intern. Med., 136 : 1101.

Ham, A.W. and Cormack, D.H. (1979) :

The urinary system. In Ham, A.W. and Cormack, D.H.
(Ed.). *Hisology*.

J.B. Lippincott, Toronto, p. 753.

Hamdi, H.; Abdel, R.Y.; Malek, A.; El-Bagory, I.; Ibrahim, Z.

and Macrom, F. (1982) :

Kidney. In Hamdi, H.; Abdel, R.Y.; Malek, A.; El-

Bagory, I.; Ibrahim, Z. and Macrom, F. (Ed.). *Kidney
fundamentals of human physiology*.

El-Naser Modern Bookshop, Cairo, p. 1.

Higgines, L. and Kastnass, S. (1967) :

Antimicrobial agent. In Goodman and Gilman (Ed.).

Goodman and Gilman's the pharmacological basis of
therapeutic.

Macmillan, Toronto, p. 1102.

JosEpivitz Christine, Pastorza Munoz, E.; Tunmerman, D.; Scott,

M.; Feldman, S. and Kaloyandes, G.J. (1982) :

Inhibition of gentamicin uptake in rat renal cortex
in vivo by aminoglycoside and organic polycations.

J. Pharmacol. Exper. Ther., 223 : 97.

Kahlmeter, G. (1980) :

Clinical pharmacokinetics : Netilmicin.

Scand. J. Inf. Dis., 23 : 74.

Kahlmeter, G. (1981) :

The role of aminoglycoside in modern clinical practice.

A doctor to doctor round table discussion held in Nice, France, December, p. 25.

Kahlmeter, G. and Dahlager, J.I. (1984) :

Aminoglycoside nephrotoxicity.

J. Antimicrob. Chemother., 13 : 9.

Kalsyanide, G.J. and Munoz, E.P. (1980) :

Aminoglycoside nephrotoxicity.

Kidney Int., 18 : 575.

Kaplan, L.A. (1984) :

Enzymes. In Kaplan, L.A. and Presce, A.J. (Ed.).

Clinical chemistry.

C.V. Mosby, Toronto, p. 1116.

Klastersky, J.; Hensgens, C.; Henri, A. and Daneau, D. (1974) :

Comparative clinical study of tobramycin and gentamicin.

Antimicrob. Agent Chemother., 5 : 133.

Kuhar, M.J.; Mak, L.L. and Lietman, P.S. (1979) :

Autoradiographic localization of H^3 gentamicin in proximal tubule of mice.

Antimicrob. Agent Chemother., 15 : 131.

Lerner, A.N. (1983) :

The role of aminoglycoside in modern clinical practice.

A doctor to doctor round table discussion held in Nice, France, December, 23 : 1124.

Lerner, A.Martin (1983) :

Randomized controlled trial the comparative efficacy, auditory, toxicity and nephrotoxicity of tobramycin and netilmicin.

Lancet, 21 : 1123.

Lipsky, J.J.; Cheno, I.; Sacktor, B. and Litman, P.S. (1980) :

Gentamicin uptake by renal tubule brush border membrane vesicle.

J. Pharmacol. Exper. Ther., 215 : 119.

Lockhart, R.D.; Hamilton, G.F. and Fyfe, F.W. (1970) :

The urinary system. In Lockhart, R.D.; Hamilton, G.F. and Fyfe, F.W. (Ed.). Anatomy of the human body.

Faber Limited, London, p. 45.

Lode, H. (1981) :

The role of the aminoglycoside in modern clinical practice.

A doctor to doctor round table discussion held in Nice, France, December, p. 40.

Luft, F.C.; Rankin, L.I.; Sloan, K.S. and Yan, M.N. (1978) :

Recovery from aminoglycoside nephrotoxicity with continued drug administration.

Antimicrob. Agent Chemother., 3 : 201.

Luft, F.C. and Evans, A.P. (1980) :

Comparative effect of tobramycin and gentamicin on glomerular ultrastructure.

J. Inf. Dis., 142 : 910.

Luft, F.C. (1985) :

Pathogenesis of aminoglycoside nephrotoxicity.

In Hook, J.B. (Ed.). Toxicology of the kidney.

Raven Press, New York, p. 407.

Luthy, R. (1981) :

The role of aminoglycoside in modern clinical practice.

A doctor to doctor round table discussion held in Nice, France, December, p. 36.

Malis, C.D.; Racusen, L.C.; Salez, K. and Welton, A. (1984) :

Nephrotoxicity of lysine and of a single dose of aminoglycoside in rats given lysine.

J. Lab. Clin. Med., May, 660.

Moffat, D.V. (1982) :

Development of urogenital system. In Chisholm, G.D. and Williams, D.I. (Ed.). Scientific foundations of urology.

William Heimmann, London, p. 130.

Moore, R.D.; Smith, C.R.; Lipsky, J.J.; Mellits, E.D. and

Fietman, P.S. (1984) :

Risk of nephrotoxicity in patient treated with aminoglycoside.

Ann. Intern. Med., 101 : 352.

Moore, R.D. and Smith, G.R. (1985) :

Aminoglycoside nephrotoxicity and age.

Ann. Intern. Med., 102 : 352.

Morin, J.P.; Viotle, G.; Vandewalle, A.; Hoof, V.; Tultens, P.

and Fillastre, J.P. (1980) :

Gentamicin induced nephrotoxicity.

Kidney Int., 18 : 585.

Nanra, S. Ranjit (1986) :

Drug-induced renal disease, renal disorders.
Med. Int., Middle Eastern ed., Sept., p. 1366.

Napolitani, D.F (1970) :

Urinary system. In Napolitani, D.F. (Ed.).
Human anatomy.
J.B. Lippincott Co., Toronto, p. 131.

Pepperell, R.J. (1983) :

Drug use in pregnancy.
Med. Int., November, 35 : 1636.

Perrin, P. and Gille, Y. (1983) :

Prophylactic antibiotic in urology. In Francois,
B. and Perrin, P. (Ed.). Urinary infection.
Butterworths, London, p. 221.

Quarum, M.L.; Houghton, D.C.; Gilbert, D.N.; Carron, D.A. and

Bennett, W.M. (1984) :

Increasing dietary calcium moderates experimental
gentamicin nephrotoxic.
J. Lab. Clin. Med., 6 : 104.

Resnick, J.B. and Hamburger, R.J. (1982) :

Clinical assessment of renal function. In Resnick,
J.B. and Hamburger, R.J. (Ed.). Nephrology.
J. B. Lippincott, Toronto, p. 30.

Richterich, R. and Colombo, J.P. (1981) :

Kidneys. In Richterich, R. and Colombo, J.P. (Ed.).

Clinical chemistry.

John Wiley, Toronto, p. 673.

Sande, M.A. and Mandell, G.L. (1980) :

Aminoglycoside. In Goodman and Gilman (Ed.).

Goodman and Gilman's the pharmacological basis
of therapeutic.

Macinillans, Toronto, p. 1162.

Sande, M.A. and Mandell, G.L. (1986) :

Aminoglycoside. In Goodman and Gilman (Ed.).

Goodman and Gilman's the pharmacological basis
of therapeutic.

Macinillans, London, p. 1159.

Schentage, J.J.; Lasezkay, G.; Cumbo, J.J.; Plant, M.E. and

Jusko, W.J. (1978) :

Accumulation pharmacokinetics of tobramycin.

Antimicrob. Agent Chemother., 13 : 649.

Schentage, J.J. and Plant, M.E. (1980) :

Patient treated with aminoglycoside.

Kidney Int., 17 : 654.

Schotteluum, B.A. and Schotteluum, D.D. (1978) :

Renal physiology. In Schotteluum, B.A. and
Schotteluum, D.D. (Ed.). Textbook of physiology.
C.V. Mosby, Saint Louis, p. 540.

Scott, D.F.; Wood, D.O.; Brownell, G.H.; Carter, M.J. and

Best, G.C. (1978) :

Aminoglycoside modification by gentamicin-resis-
tance isolate of *Staphylococcus aureus*.
Antimicrob. Agent Chemother., 13 : 641.

Shepard, R.S. (1971) :

Clinical evaluation of kidney function. In Shepard,
R.S. (Ed.). Human physiology.
J.B. Lipincott, Toronto, p. 377.

Smith, G.R. (1977) :

Experience points of gentamicin.
New Engl. J. Med., 12 : 296.

Smith, G. R.; Lipsky, J.J.; Laskin, O.L.; Helimann, D.B.; Mellits,

D.; Longstreth, J. and Lietman, P.S. (1980) :

Double blind comparison of nephrotoxicity of genta-
micin and tobramycin.
New Engl. J. Med., 15 : 1106.

Smith, S.T. (1986) :

Basic physiology, analytical procedures. In

Smith, S.T. (Ed.). Clinical chemistry, principles.

W.B. Saunders, Toronto, p. 413.

Spyker, D.A. and Cuerrant, R.L. (1977) :

Gentamicin dosage.

Ann. Int. Med., 80 : 357.

Thompson, F.D. (1982) :

Laboratory and biochemical test. In Chisholm, G.

D. and Williams, D.I. (Ed.). Scientific founda-

tions of urology.

Williams, London, p. 76.

Tietz, N.M. (1986) :

Nitrogen metabolites and renal function. In Tietz,

N.M.; Bhaganvan, N.V.; Conn, R.B.; Pruden, E.L. and

Whitery, S.R. (Ed.). Textbook of clinical chemistry.

W.B. Saunders, London, p. 1268.

Tight, J.R. (1982) :

Histology and ultrastructure. In Chisholm, G.D.

and Williams, D.I. (Ed.). Scientific foundations

of urology.

William Heimann, London, p. 20.

Tobias, G.J.; Langulin, M.C. and Hopper, J. (1962) :

Endogenous creatinine clearance.

New Engl. J. Med., 266 : 317.

Valtin, H. (1979) :

Renal function assessment. In Valtin, H. (Ed.).

Renal dysfunction.

Little Brown Co., Boston, p. 205.

Whelton, A. and Salez, K. (1981) :

Mechanism of drug nephrotoxicity. In Zurkzoglu, W.; Papadimitriou, M.; Pyrpasopolus, M.; Sions, M. and Zambolis, C. (Ed.). Eighth International congress of nephrology.

Thessloniki, Greece, p. 144.

Whelton, A. and Salez, K. (1982) :

Aminoglycoside nephrotoxicity.

J. Lab. Clin. Med., 90 : 148.

Wilson, A.; Schild, H.O. and Modell, W. (1979) :

Aminoglycoside. In Wilson, A.; Schild, H.O. and Modell, W. (Ed.). Applied pharmacology.

Churchill Livingstone, p. 486.

Wootton, I.D. and Freeman, H. (1982) :

Kidneys. In Wootton, I.D. and Freeman, H. (Ed.).

Microanalysis in medical biochemistry.

Churchill Livingstone, London, p. 48.

Yasuhara Hajime; Kobayaski; Sakamoto, D. and Kamijo, K. (1982) :

Pharmacokinetic of amikain.

J. Clin. Pharmacol., 22 : 403.