

References

- 1- **Adcock LM and Papile L (2008):** Perinatal asphyxia. In: Cloherty JP, Eichewald EC, Stark AR (Eds). Manual of neonatal care (6th ed). Lippincott Williams and Wilkins. Philadelphia; 27C:518-28.
- 2- **Aggarwal R, Jain A, Deorari AK et al. (2008):** "Post-resuscitation management of asphyxiated neonates". Indian J Pediatr; 75:175-80.
- 3- **Aggarwal R, Deorari AK and Paul VK. (2001):** Post-resuscitation management of asphyxiated neonates. Ind J Pediatr; 63(12): 1149.
- 4- **Allemand F, Reale F, Sposato M and Allemand A. (2009):** Perinatal Hypoxic-Ischemic Encephalopathy: epileptic and paretic outcome at one year of age. Ital J Pediatr; 35:14.
- 5- **Al-Naqeeb N, Edwards AD, Cowan F and Azzopardi D (1999):** Assessment of neonatal encephalopathy by amplitude-integrated electroencephalography. Pediatr; 103:1263-71.
- 6- **Amato M and Donate F (2000):** Update on perinatal hypoxic insult: mechanism, diagnosis, and intervention. Eur J Pediatr Neurol; 4(5): 203-9.
- 7- **Ambalavanan N, Carlo WA, Shankaran S, Bann CM, Emrich L, et al.(2006):** Predicting Outcomes of Neonates Diagnosed With Hypoxic Ischemic Encephalopathy. Pediatr; 118(5): 2084-93.
- 8- **American Academy of Pediatrics (1992):** "Relation between perinatal factors and neurological outcome". In Guidelines for Perinatal Care. 3rd ed. ELK Grove Village, III: American Academy of Pediatrics; 221-34.

References

- 9- **American Academy of Pediatrics and American College of Obstetrics and Gynecology (1996):** Committee on fetus and newborn, and Committee on obstetric practice: Use and abuse of the Apgar score. *Pediatr*; 98:141-42.
- 10- **Anast CS, Winnacker JL, Forte LR et al. (1976):** Impaired release of parathyroid hormone in magnesium deficiency. *J Clin Endocrinol Metab.*;42:707-17.
- 11- **Andrew W and Marianne T (2002):** Clinical trials of treatments after perinatal asphyxia. *Curr Opin Pediatr*; 14(6): 664-8.
- 12- **Apgar V (1953):** A proposal for a new method of evaluation of the newborn infants. *Anesth Analg*; 32:260-7.
- 13- **Arvin KL, Han BH, Du Y, et al. (2002):** Minocycline markedly protects the neonatal brain against hypoxic ischemic injury. *Ann Neurol*; 52:54-61.
- 14- **Atsmon J and Dolev E (2005):** Drug induced hypomagnesemia: scope and management. *Drug Safety*; 28:763-88.
- 15- **Attia N, Taman K, Abdel-Rahman S and Faris L (1997) :** Role of SPECT and Cranial Sonography in the Evaluation of Brain Insult in Perinatal Asphyxia. *Egy J Pediatr*; 14(4): 619-63.
- 16- **Attia NI, Ali GS and Mahmoud MN (2002):** Lipid Peroxidation In Hypoxic-Ischemic Encephalopathy. *Egy J Pediatr*; 3(1).
- 17- **Aurora S and Snyder EY (2004):** Perinatal asphyxia. In: Cloherty JP, Eichenwald EC and Stark AR. (Eds). *Manual of Neonatal Care*. (15th ed). Lippincott Williams and Wilkins. Philadelphia; (27): 536-55.

References

- 18- **Aydin A, Gene K, Akhisaroglu M, et al. (2003):** Erythropoietin exerts neuroprotective effect in neonatal rat model of hyoxic-ischemic brain injury. *Brain Dev*; 25:494-8.
- 19- **Baker SB and Worthley LI (2002):** The Essentials of Calcium, Magnesium and phosphate metabolism. Part I physiology. *Crit care Resusc*; 4:301-6.
- 20- **Barnett A, Mercuri E, Rutherford M, Haataja L, Handerson S, et al. (2002):** Neurological and perceptual motor outcome at 5-6 years of age in children with neonatal encephalopathy: relationship with neonatal brain MRI. *Neuropediatr*; 33:242-8.
- 21- **Benson JE, Bishop MR and Cohen HL (2002):** Intracranial neonatal neurosonography: an update. *Ultra Q*; 18(2): 89-114.
- 22- **Berger R and Garnier Y (1999):** "Pathophysiology of perinatal brain damage". *Brain Res Rev*; 30(2): 107-34.
- 23- **Berger R and Garnier Y (2001):** Perinatal brain injury. *J Perinat Med*; 29(1): 85-6.
- 24- **Blood AB, Hunter CJ and Power GG (2003):** Adenosine mediates decreased cerebral metabolic rate and increased cerebral flow during acute moderate hypoxia in the near term fetal sheep. *L physiolo*; 553(3): 935-45.
- 25- **Boichot C, Walker PM, Durand C, et al. (2006):** Term neonate prognoses after perinatal asphyxia: contributions of MR imaging, MR spectroscopy, relaxation times, and apparent diffusion coefficients. *Radiol*; 239: 839-48.
- 26- **Boo NY, Hafidz H, Nzwawi HM, Cheah FC, Fedzil YJ, et al. (2005):** comparison of serum cardiac troponin T and creatine kinase

References

- MB isoenzyme mass concentrations in asphyxiated term infants during the first 48 hrs of life. *J pediatri* 41(7): 331-7.
- 27- **Borruto F, Comparetto C, Weher E and Tresser A (2006):** Screening of Fetal Distress by Assessment of Umbilical Cord Lactate. *Clin Exp Obstet Gynecol*; 33(4): 219-22.
- 28- **Broner CW, Stidhalm GL, Westenkirchner DF. et al. (1990):** Hypermagnesemia and hypocalcemia as predictors of high mortality in critically ill pediatric patients. *Crit care Med*; 18:921-8.
- 29- **Brus F, Van Oeveren W, Okken A, et al. (1997):** Number and activation of circulating polymorphonuclear leukocytes and platelets are associated with neonatal respiratory distress syndrome severity. *Pediatr*; 99:672-80.
- 30- **Bushinsky DA and Monk RD (1998):** Electrolyte quintet: calcium. *Lancet*; 352:306-11.
- 31- **Cadell JL and Reed GF (1989):** Unreliability of plasma magnesium value in asphyxiated neonates. *Magnesium*; 8: 11-6.
- 32- **Caravale B, Allemand F and Libenson MH (2003):** Factors Predictive of Seizures and Neurologic Outcome in Perinatal Depression. *Pediatr Neurol*; 29(1):18-25.
- 33- **Celtik C, Acunas B, Oner N and Pala O (2004):** Neuron specific enolase as a marker of the severity and outcome of hypoxic-ischemic encephalopathy. *Brain Dev*; 26(6): 398-402.
- 34- **Chamnana-Vanaki J and Perlman JM (2000):** Management of perinatal HIE. *Rec Adv Pediatr*; 18(3): 33.

References

- 35- ***Chamnanavanaki and Perlman JM (2000):*** Management of perinatal hypoxic ischemic encephalopathy. *Recent Advances in Pediatrics*; 18(3):33-46.
- 36- ***Changlian Z, Falin X, Xiaoyang W, Masahiro S, Yasu U, et al. (2003):*** Different apoptotic mechanisms are activated in male and female brains after NHIE. *J Neuro Chem*; 96:1016-9.
- 37- ***Chao CP, Zaleski CG and Patton AC (2006):*** Neonatal Hypoxic Ischemic Encephalopathy: Multimodality Imaging Findings. *Rad Graph*; 26:S159-S72.
- 38- ***Chapman IA and Stoll BJ (2007):*** Hypoxia-ischemia. In: Behrman RE, Kliegman RM and Jenson HB (Eds). *Nelson Textbook of Pediatrics*. (18th ed). WB Saunders. Philadelphia; 718-20.
- 39- ***Chen J, Hunang H, Zheng K, Xia R, Chen J, et al. (2007):*** Early application of nerve growth factor affects serum inflammatory cytokine levels in neonatal hypoxic ischemic encephalopathy. *Neur Reg* 2(7): 737-9.
- 40- ***Chen ZL, He RZ, Peng Q, Guo KY, Zhang YQ, et al. (2009):*** Prenatal risk factors for neonatal asphyxia: how risk for each? *Zhongguo Dang Dai Er Ke Za Zh*; 11(3): 161-5.
- 41- ***Cheong JL, Cady EB, Penrice J, Wyatt JS, Cox IJ, et al. (2006):*** Proton MR Spectroscopy in Neonates with Perinatal Cerebral Hypoxic-Ischemic Injury: Metabolite Peak-area Ratios, Relation Times and Absolute Concentrations. *Am J Neuroradiol*; 27(7):1546-54.

- 42- Correa F, Enriquez G, Possello J, Lucaya J, Piqueras J, et al. (2004):** Posterior Fontanelle sonography: An Acoustic Window into the Neonatal; 25(7): 1274-82.
- 43- C.P. Panayiotopoulos. (2010):** MD, PhD, FRCP, A clinical guide to epileptic syndrome and their treatment. Second edition. London: Springer; Epilepsia; 42: 1212-8.
- 44- Cuzzocrea S (2004):** Effect of Inhibitors of Nitric Oxide in Animal Models and Future Directions for Therapy in inflammatory Disorders. Current Medicinal Chemistry-Anti-Inflammatory & Anti-Allergy Agents; 3(3):261-70.
- 45- Dai H and Luo YH (2009):** Changes of electroencephalographic levels in neonates with hypoxic ischemic-encephalopathy. Zhongguo Dang Dai Er Ke Zhi; 11(3): 173-6.
- 46- Deguchi K, Oguchi K and Takashima S (1997):** Characteristic neuropathology of leukomalacia in extremely low birth weight infants. Pediatr Neurol; 16(4): 296-300.
- 47- Delivoria- Papdopoulos M and Mishra DP (1998):** Mechanisms of cerebral injury in perinatal asphyxia and strategies for prevention. J Pediatrics 132: 305-45.
- 48- Demenezes MS (2006):** hypoxic ischemic brain injury in the newborn. URL: <http://www.eMedicine.com>.
- 49- Depp R (1995):** "Perinatal asphyxia: assessing its causal role and timing". Semin Pediatr Neurol; 2(1): 3-36.
- 50- Eichenwald EC and Stark AR. Eds.: Manual of Neonatal care (5th – edition). Lippincott Williams and Wilkins. Philadelphia; PP. 643-650**

References

- 51- **El-Ayouty M, Abdel-Hady H, El-Mogy S, Zaghlol H, El-Beltagy M, et al. (2007)** : Relationship between electro-encehalography and magnetic resonance imaging findings after hypoxic-ischemic encephalopathy at term. *Am J Perinatol*; 24(8): 467-73.
- 52- **Engle WD, Laptook R and Perlman JM (1999)**: Acute change in arterial carbon dioxide tension and acid – base status, and early neurologic characteristics in term infants following perinatal asphyxia. *Resuscitation*; 42(1): 11-7.
- 53- **Ferriero DM (2004)**: Neonatal brain injury. *N Engl J Med*; 351:1985-95.
- 54- **Fleming CR, George L, Stoner GL et al. (1996)**: The importance of urinary magnesium values in patients with gut failure. *Mayo Clin Proc*; 71:21-4.
- 55- **Forfar and Arneil (2003)**: Hypoxic ischemia. In: McIntosh N, Helms PJ and Symth R. (Eds). *Textbook of Pediatrics*. (6th ed). Churchill Livingstone; P693-9.
- 56- **Furtakul S, Praisuwanna P and Thaitumyanon P (2006)**: Risk Factors for hypoxic ischemic encephalopathy in asphyxiated newborn infants. *J Med Assoc Thai*; 89(3): 322-8.
- 57- **Gabbe SG, Niebyl JR and Simpson JL (2002)**: Normal and Problem Pregnancies. (4th ed). Churchill Livingstone.; 661-7.
- 58- **Gathwala G (2001)**: Neuronal protection with magnesium. *Indian J Pediatr* 68 (5):417-9.

References

- 59- **Gazzolo D, Marinoni E, Di Iorio R, Braschetti M, Kornacka M, et al. (2004):** urinary S100B protein measurements: a tool for early identification of hypoxic-ischemic encephalopathy in asphyxiated fullterm infants. *Crit Care Med*; 32(1): 131-6.
- 60- **Gluckman PD, Wyatt JS, Azzopardi D, et al. (2005):** Selective head cooling with mild systemic hypothermia after neonatal encephalopathy: multicenter randomized trial. *Lancet*; 365:663-70.
- 61- **Gomella TL, Cunningham MD, Eyal FG and Zenk K (2004):** Perinatal asphyxia. In: Gomella TL, Cunningham MD, Eyal FG and Zenk KE. (Eds). *Neonatology; Management, procedures, On-call problems, Diseases and Drugs*. (5th edition), Lange,; Pp. 512-23.
- 62- **Gomella TL, Cunningham MD, Eyal FG, Zenk KE et al. (2004 a):** Commonly used medications, Magnesium Sulphate. In *Clinical manual*. Gomella et al (eds), 5th edition. Mc Graw – Hill companies, p. 620.
- 63- **Gomella TL, Cunningham MD, Eyal FG, Zenk KE et al. (2004):** Disorders of calcium and magnesium metabolism. In *Clinical manual*. Gomella et al (eds) 5th edition. Mc Graw – Hill Companies. P. 563-71.
- 64- **Gomes JA, Stevens RD, Lewin JJ, Mirski MA and Bhardwaj A (2005):** Glucocorticoid therapy in neurological critical care. *Crit Care Med*; 33 (6): 1214-24.
- 65- **Graham EA, Ruis KA, Hartman AI, Northiugton FJ and Fox HE (2008):** A systematic review of the role of intrapartum hypoxia – ischemia in the causation of neonatal encephalopathy. *Am J Obstet Gynecol*; 199(6):585-6.

References

- 66- **Greenamyre JT, Porter RHP (1994):** Anatomy and physiology of glutamate in the CNS. *Neurology* 44:7-12.
- 67- **Greenbaum LA (2004):** Electrolyte and acid base disorders. In: Nelson Textbook of Pediatrics, Behrman RE, Kliegman RM, Jensen HB (eds) 17th ed. Philadelphia; El Sevier, p. 208-19.
- 68- **Gupta BD, Sharma P, Bagla J, Parakh M and Soni JP (2005) :** Renal Failure in Asphyxiated Neonates. *Ind Pediatr*; 42(17):928-34.
- 69- **Haan HH and Hasaart TH (1995):** "Neuronal death after perinatal asphyxia". *Eur J Obstet Gynecol Reprod Biol*; 61(2): 123-7.
- 70- **Hagberg B, Hagberg G, Beckung E and Uvebrant P (2001):** Changing panorama of cerebral palsy in Sweden. VIII. Prevalence and origin in the birth year period 1991-94. *Acta prdiatr*; 90:271-7.
- 71- **Hall RT, Hall FK and Daily D (1998):** High-dose Phenobarbital therapy in term newborn infants with severe perinatal asphyxia: a randomized, prospective study with three year follow-up. *J pediatri*; 132:345-8.
- 72- **Hankins GD and Speer M (2003) :** "Defining the pathogenesis and pathophysiology of neonatal encephalopathy and cerebral palsy". *Obstet. Gynecol*; 102(3): 628-36.
- 73- **Hasan Tekgul, Kimberlee Gauvreau, Janet Soul., et al. (2006):** The current etiological profile and neuro-developmental outcome of seizures in term newborn infants, *MPHa (Pediatrics)*. Vol. 117, No4, April 2006, PP.1270-1280 (DOI: 10.1542/PEDS.2005-1178).

References

- 74- **Hassanein SM, Moustafa MF, AfifyHM, Abdel-latif MM and El-Houshy LM (2002):** Value of Lactate Dehydrogenase (LDH), Aspartate Aminotransferase (AST), and Total Creatine Kinase (TCK) in predicting Sequelae of perinatal Asphyxia. *ESNP*;3(1).
- 75- **Hellström-Westas L, Rosen I, De-Vries LS and Greisen G (2006):** Amplitude-integrated EEG Classification and Interpretation in Preterm and Term Infants. *Neurology*; 7(2):c76-c87.
- 76- **Higgins RD, Touse NK, Periman J, et al. (2006):** Hypothermia and perinatal asphyxia: executive summary of the National Institute of Child Health and Human Development workshop, *J Pediatr*; 148 (2): 170 - 5.
- 77- **Hoffman DJ, Marro PJ, McGowan JE, Mishra OP, Delevoria-Papadopoulos M et al. (1994):** Protective effect of Mg sulfate infusion of NMDA receptor binding characteristics during cerebral cortical hypoxia in the newborn piglet. *Brain Res*; 664: 144-9.
- 78- **Huppi PS (2002) :** Advances in postnatal neuroimaging : relevance to pathogenesis and treatment of brain injury. *Clin Perinatol*; 29:827-56.
- 79- **Huttner KM (2004):** Hypocalcemia, hypercalcemia and hypermagnesemia. In *Manual of Neonatal Care*. Cloherty JP, Eichenwald EC, Stark AR (eds) 8th ed. Lippincott Williams and Wilkins, Philadelphia pp 579-88.
- 80- **Hypoxic – Ischemic injury To the Neonatal Brain. AJNR Am J. neuroradiol**, 25(3): 263 - 8.

References

- 81- **Ichiba H, Tamai H, Negishi H, et al. (2002):** Randomized controlled trial of magnesium sulfate for severe birth asphyxia. *Pediatr Int*; 44: 505 - 9.
- 82- **Ilves P, Kiisk M, Soopold T, et al. (2000):** Serum total magnesium and ionized calcium concentrations in asphyxiated term newborn infants with hypoxic-ischemic encephalopathy. *Acta Paediatr*; 89 : 680-5
- 83- **Ingemarsson I (2003):** Gender aspects of preterm birth. *Int J Obst Gyn*; 110(20): 34-8.
- 84- **Ito BA, Al-Hawsawi ZM and Khan AH. (2003):** Hypoxic ischemic encephalopathy: Incidence and risk factors in North Western Saudi Arabia. *Saudi Med J*; 24(2):147-53.
- 85- **Jacobs S, Hunt R, Tarnow –Mordi W, Inder T and Davis P (2007):** Cooling for newborns with hypoxic ischemic encephalopathy. *Coch Data Syst Rev* ;(4): CD-00311.
- 86- **Johnston BM, Mallard EC, Williams CE and Gluckman PD (1996):** Insulin-like growth factor-1 is a potent neuronal rescue agent after hypoxic-ischemic injury in fetal lambs. *J Clin Invest*; 97: 300-8.
- 87- **Johnston MV (1993):** Cellular alterations associated with perinatal asphyxia. *Clin Invest Med*; 16:122-32.
- 88- **Kanik E, Ozer EA, Bakiler AR, Aydinlioglu H, Dorak C, et al. (2009):** Assessment of myocardial dysfunction in neonates with hypoxic-ischemic encephalopathy: is it a significant predictor of mortality? *J Mat-Fet Neo Med*: 22(3):239-42.

References

- 89- **Karlsson M, Blennow M, Nemeth A and Winblad B (2007):** Dynamics of hepatic enzyme activity following birth asphyxia. *Acta Paediatr*; 95(11): 1405 - 11.
- 90- **Key LL and Carpenter TO (1990):** Metabolism of calcium, phosphorus and other divalent ions. In: *Pediatric textbook of fluids and electrolytes*. Ickikawa I (ed): Baltimore, Williams and Wilkins, p.98.
- 91- **Khashaba MT, Shouman BO, Shaltout AA, et al. (2006):** Excitatory amino acids and magnesium sulfate in neonatal asphyxia. *Brain Dev*; 28 (6): 375-9.
- 92- **Klinger G, Beyene J, Shah P and Perlman M (2005):** Do hyperoxemia and hypocapnia add the risk of brain injury after intrapartum asphyxia? *Arch Dis Child Fet Neo*. Ed; 90: 49 - 52.
- 93- **Langley WF and Mann D (1991):** Central nervous system magnesium deficiency. *Arch Intern Med*; 151: 593-6.
- 94- **Laptook AR and Corbett RJ (2002):** The effects of temperature on hypoxic-ischemic brain injury. *Clin perinatol*; 29: 623 - 649.
- 95- **Le- Strange E, Saeed N, Cowan FM, Edwards AD and Rutherford MA (2004):** MR imaging Quantification of Cerebellar Growth Following Hypoxic-Ischemic Injury to the Neonatal Brain. *AJNR Am J Neuroradiol*; 25 (3): 263-8.
- 96- **Legido A, Katsetos CD, Mishra OP and Delivoria – Papadopoulos M (2000):** Perinatal Hypoxic Ischemic Encephalopathy: Current and Future Treatments. *Int Pediatr*. 15 (3): 143-51.

References

- 97- **Leuthner SR and Das UG (2004):** Low Apgar scores and the definition of birth asphyxia. *Pediatr Clin Nor Am*; 51 (3): 737- 45.
- 98- **Lin HC, Su BH, Lin TW and Tsai CH (2004):** Risk factors of mortality in meconium aspiration syndrome. *Acta Pediatr*; 45(1):30-4.
- 99- **Levene M, Blennow M, Whitelaw A et al. (1995):** Acute effects of two different doses of magnesium sulphate in infants with birth asphyxia. *Arch Dis Child*; 73:F174-F7.
- 100- **Levene MI, Evans DJ, Mason S, Brown J et al. (1999):** An international network for evaluating neuroprotective therapy other severe asphyxia. *Semin Perinatal*; 23:226-33.
- 101- **Levin BS and Coburn JW (1984):** Magnesium, the mimic/antagonist of calcium. *N Eng J Med*; 310:1253-5.
- 102- **Liu Z, Xiong T and Meads C (2006):** Clinical effectiveness of treatment with hyperbaric oxygen for neonatal hypoxic ischaemic encephalopathy: systematic review of Chinese literature. *BMJ*; 333 (7564): 374.
- 103- **Loh SF, Woodworth A and Yeo GSH (2008):** Umbilical cord blood gas analysis at delivery. *Sing Med j*; 49: issue 05.
- 104- **Mahura IS (2003):** Cerebral ischemia – hypoxia and biophysical mechanisms of neurodegeneration and neuroprotection effects. *Fiziol Zh*; 49:7-12.

References

- 105-Makikallio K, Vuolteenaho O, Jouppila P and Reasanen J (2000):** Association of severe placental insufficiency and systemic venous pressure rise in the fetus with increased neonatal cardiac troponin T levels. *Am J Obstet Gynecol*; 183: 726 -31.
- 106-McAlmon KP (2004) :** Necrotizing enterocolitis. In: Cloherty JP, Eichenwald EC and Stark AR. (Eds). *Manual of Neonatal Care*. (5th edition). Lippincott Williams and Wilkins. Philadelphia; Pp. 643-50.
- 107-Marcus JC, Valencia GB, Altura BT, et al. (1998):** Serum ionized magnesium in premature and mature term infants. *Pediatr Neurol*; 18(4): 311-4.
- 108-Marret S, Gressens P, Gadisseux JF, Evrand P (1995):** Prevention by magnesium of excitotoxic neuronal death in the developing brain: an animal model for clinical intervention studies. *Dev Med Child Neurol*; 37:473-84.
- 109-Martin HE (1969):** Clinical magnesium deficiency. *Ann NY Acad Sci*; 162:891-900.
- 110-McClure MM, Threlkeld SW and Fitch RH (2007):** Auditory processing and learning / memory following erythropoietin administration in neonatally hypoxic – ischemic injured rats .*Brain Res*; 9; 1132 (1): 203- 9.
- 111-McDonald JW, Silverstein FS and Johnson MV (2000):** Physiology and pathophysiology rules of brain injury in perinatal asphyxia .*Brain Res Rev*; 15: 41.
- 112-Mclean RM (1994):** Magnesium and its therapeutic uses: a review. *Am J Med*; 96: 63 - 76.

References

- 113-Mehta KC, Kalkwarf HJ, Mimouni F, et al. (1998):** Randomized trial of magnesium administration to prevent hypocalcemia in infants of diabetic mothers. *J Perinatol*; 18(5): 352-6.
- 114-Ment LR, Bada HS, Barnes B, Great PE, Hirtz D, et al. (2004):** Neuroimaging of the neonate .*Neuroradiology*; 22- 36.
- 115-Miller SP, Ramaswamy V, Michelson D, et al. (2005):** "Patterns of brain injury in term neonatal encephalopathy".*J Pediatr*;146(4):453-60
- 116-Mishra O and Delivoria Papadopoulos M (1999):** Cellular mechanisms of hypoxic injury in the developing brain. *Brain Res Bull*; 48:233-8.
- 117-Morrisett RA, Mott DD, Lewis DV, Wilson WA, Swartzwelder HS (1990):** Reduced sensitivity of NMDA component of synaptic transmission to magnesium in hippocampal slices from immature rats. *Dev Brain Res*. 86:257-62.
- 118-Moster D, Lie RT, Irgens LM et al. (2001):** "The association of Apgar score with subsequent death and cerebral palsy: A population based study in term infants". *J Pediatr*; 138:798-803.
- 119-Mousa SA, Ritger RC and Smith RD (1992):** Efficacy and safety of desferoxamine conjugated to hydroxyethyl starch. *J Cardiovasc Pharmacol*; 19: 425 - 9.
- 120-Nelson N, Finnstrom O, Larsson L (1989):** Plasma ionized calcium, phosphate and magnesium in preterm and small for gestational age infants. *Acta Paediatr Scand*; 78:351-7.

References

- 121-Nikas I, Dermentzoglou V, Theofanopoulou M and Theofanopoulou V (2008):** Parasagittal Lesions and ulegyria in hypoxic ischemic encephalopathy: neuroimaging findings and review of the pathogenesis. *J Child Neurol*; 23 (1): 51-8.
- 122-Nordin BE (1997):** Calcium and osteoporosis. *Nutrition*; 13(7-8): 664-86.
- 123-Nowak L, Bregestovski P, Asher P Herbert A, Prochiantz A et al, (1984):** Magnesium gates glutamate activated channels in mouse central neurons. *Nature*; 307:462-5.
- 124- Ojha RK, Singh SK, Batra S, Sreenivas V and Puliyl JM (2006):** Lactate: Creatinine Ratio in Babies with Thin Meconium Staining of Amniotic Fluid .*BMC Peiatr*; 6: 13- 20.
- 125- Olusegeun J Adebami. (2010):** Study to determine the pattern and outcome of neonatal seizures in Osogbo, South-Western Nigeria. Department of Pediatrics and child health, S A Journal of child health. Vol. 4, No.2 article.
- 126-Oren H, Duman N, Abacioglu H, et al. (2001) :** Association between serum macrophage colony – stimulating factor levels and monocyte and thrombocyte counts in healthy, hypoxic, and septic term neonates .*Pediatr*; 108- 329 -32.
- 127-Palmer C (1995):** Hypoxic – ischemic encephalopathy: Therapeutic approaches against microvascular injury, and role of neutrophils, PAF, and free redicals. *Clini Perinatol*; 22(2): 481- 517.
- 128- Palmer J (2004):** Perinatal Hypoxic-Ischemic Disease URL : <http://www.google.com>.

References

- 129- Paul D' orazio. (2008):** Ph. D. Fundamentals of clinical chemistry. Electrochemistry and chemical sensor. Fourth edition, P. 85.
- 130- Park SS, Chung SH, song JH, Kim SK, Kim SD, et al. (2007) :** The correlation between the Severity of Hypoxic Ischemic Encephalopathy and the Develompent of Acute Renal Failure in Asphyxiated Neonates .J Kor soc Pediatr Nephrol; 11 (1) : 32 -40 .
- 131- Partridge SC, Mukherjee P, Henry R, Miller SP, Berman JI, et al. (2004) :** Diffusion tensor imaging : serial quantitaion of white matter tract maturity in premature newborns .Neuroimage; 22 : 1302-14.
- 132- Perlman JM (2006):** Intervention strategies for Neonatal Hypoxic Ischemic Cerebral Injury. J Clin ther; 28(9):1353 - 65.
- 133- Peeters-Scholte C, Braun K, Koster J, Blomgren K, Buonocore G, et al. (2003):** Effect of allopurinol and desferoxamine on reperfusion injury of the brain in newborn piglets after hypoxia ischemia. Pediatr Res; 54: 516 - 22.
- 134- Petersen S, Christensen NC, Fogh-Andersen N. (1981):** Effect on serum Calcium of 1 α -hydroxy-Vitamin D₃ supplementation in infants of low birth weight, infants with perinatal asphyxia and infants of diabetic mothers. Acta Paediatr Scand; 70: 897-901.
- 135- Prentice A (2000):** Calcium in pregnancy and lactation. Annu Rev Nutr; 20:249-72.
- 136- Pressler MR, Boylan GB, Morton M, Binnie CD and Rennie JM (2001):** Early serial EEG in HIF. Clin Neurophysiol; 112(1): 31

References

- 137-Qian J, Zhou D, & Wang YW (2009):** Umbilical artery blood s100 β protein: a tool for the early identification of neonatal hypoxic – Ischemic encephalopathy. *Eur J Pediatr*; 168(1): 71 -7.
- 138-Raju TNK (2006):** Hypoxic Ischemic Encephalopathy. URL: [http // www. eMedicine.com](http://www.eMedicine.com)
- 139-Reddy S, Dutta S and Narang A (2008) :** Evalution of lactate Dehydrogenase, Creatine Kinase and Hepatic Enzymes for the Retrospective Diagnosis of Perinatal Asphyxia Among Sick Neonates. *Ind pediatr*; 45: 144 - 7.
- 140-Ringer SA (2004):** Resuscitaion in the delivery room .In: Cloherty JP, Eichenwald WC and Strak AR (Eds.). *Manual of Neonatal Care*. (5th edition). Lippincott Williams & Wilkins. Phildelphia; 57-69.
- 141-Rosemary D and Higgins MID (2005):** Hypoxic ischemic Encephalopathy and Hypothermia: A Critical Look, by The American College of Obstetricians and Gynecologists. Published by Lipioncott Williams & Wilkins; 106: 6.
- 142-Rodd C and Goodyer P (1999):** Hypercalcemia of the newborn. Etiology, evaluation and management. *Pediatr Nephrol*; 13(6):542-7.
- 143-Rodriguez MJ, Ursu G, Bernal F, et al. (2001):** Perinatal human hypoxic ischemic vulnerability correlates with brain calcification. *Neurobiol Di*; 8(1): 59-68.
- 144-Roelant – Vanrijn AM, Nikkels PG, Groenendall F, Vander Ground J, Barth PG, et al. (2001):** Neonatal diffusion weighted MR imaging: relation with histopathology or follow up MR examination .*Neuropediatr*; 32:286 - 94.

References

- 145-Ryan MP (1987) :** Diuretics and potassium / magnesium depletion: directions for treatment. *Am J Med*; 82:38-47.
- 146-Sameshima H, Ota A, Ikenove T et al. (1999):** Pretreatment with magnesium sulfate protects against hypoxic ischemic brain injury but post asphyxial treatment worsens brain damage in seven-day-old rats. *Am J Obstet Gynecol*; 180:725-30.
- 147-Sangkae C and Perman PM (2000):** Management of perinatal HIE. *Rec adva pediatri*; 8(3): 33.
- 148-Sarnat HB and Sarnat MS (1976):** Neonatal encephalopathy following fetal distress: A clinical and electroencephalographic study. *Arch Neuro*; 33: 696 - 705.
- 149-Shalak L and Perlman JM (2004):** Hypoxic –ischemic braing injury in the infant: current concepts. *Early Hum Dev*: 80: 125 - 41.
- 150-Shalak LF, Laptook AR, Velophi SC and Perlman M (2003):** a-EEG coupled with an early neurologic examination enhances prediction of term infants at risk for persistent encephalopathy. *Pediatr*; 111: 351 - 7.
- 151-Shankaran S, Laptook AR, Ehrenkranz RA, et al. (2005):** whole – body hypothermia for neonates with hypoxic - ischemic encephalopathy. *N Eng1 J Med*; 353 (15): 1574 - 84.
- 152-Shaw PS, Ohlsoon A and Perlman M (2007):** Hypothermia to treat neonatal hypoxic ischemic encephalopathy. *Arch pediatri Adolesc Med*; 161 (10): 951 - 8.

References

- 153-Singh D, Kumar P, Majumdar S and Narang A (2004):** Effect of Phenobarbital On Radicals In Neonates with Hypoxic ischemic encephalopathy - A Randomized Controlled Trial. *J. Perinat Med*; 32 (3): 278 - 81.
- 154-Stark EJ and Joanna SJ (2004):** Cerebral artery Doppler ultrasonography for prediction of outcome after perinatal asphyxia. *J Ultra Med*; 13: 595 - 600.
- 155-Stoll BJ and Kleigman RM (2004):** The fetus and the neonatal infant. In: Behrman RE, Kleigman RM and Jenson HB, (Eds) *Nelson Textbook of pediatrics*. (17th edition). WB: Saunders Co., London; 523 - 86.
- 156-Stoll BJ and Piazza AJ (2007):** Digestive system disorders. In: Behrman RE, Kliegman RM and Jenson HB, (Eds). *Nelson Textbook of Pediatrics*. (18th edition). WB: Saunders: Philadelphia. 753-66.
- 157-Tan WKM, Williams CE and Mallard CE (1994):** Monosialoganglioside GM1 treatment after a hypoxic - ischemic episode reduces the vulnerability of the fetal sheep brain to subsequent injuries *Am J Obstet Gynecol*, 170: 663 - 70.
- 158-Taniguchi H, Mohri I, Okabe – Arahori H, Aritake K, Wada K, et al. (2007):** Prostaglandin D2 Protects Neonatal Mouse Brain from hypoxic ischemic Injury. *J neurosci*; 27 (16): 4303 - 12.
- 159-Tekgul H, Yalaz M, Kutukeuler N, Ozbek S, Kose T, et al. (2004):** Value of biochemical markers for outcome in term infants with asphyxia. *Pediatr Neurol*; 31 (5): 326 - 32.

References

- 160-Theocharis P, Giapros V, Tsampoura Z, Basioti M, Andronikou S, et al. (2011) :** Neonatal Intensive Care Unit, University of Ioannina, Greece.
- 161-Toet MC, Meij VW, De-Vries LS, Uiterwall CS and Van Huffelen AC (2002):** Comparison between simultaneously recorded amplitude-integrated EEG (cerebral function monitor) and standard EEG in neonates. *Pediatr*, 109: 772 - 9.
- 162-Tong GM and Rude RK (2005):** Magnesium deficiency in critical illness. *J Intensive Care Med*; 20:3-17.
- 163-Tsang RC (1992):** Calcium, phosphorus and magnesium metabolism. In *Fetal and neonatal physiology*. RA polin and WW Fox (eds). Philadelphia. Saunders.
- 164-Van Hook JW (1991):** Hypermagnesemia. *Crit Care Clin*, 7:215-23.
- 165-Vannucci RC and Palmer C (1997):** Hypoxia-ischemia: Pathogenesis and Neuropathology. In: *Neonatal Perinatal Medicine, Diseases of the fetus and Infant*. Fanaroff AA. And Martin, RJ (eds). 6th edition. Philadelphia, Mosby, pp. 856-77.
- 166-Vannucci RC and Perlman JM (1997):** Intervention for perinatal hypoxic- ischemic encephalopathy. *Pediatr*: 100: 1004 - 14.
- 167-Vannucci RC, Towfighi J and Vannucci SJ (2004):** Secondary energy failure after cerebral hypoxia - ischemia in the immature rat. *J Cereb Blo Flo Metab*, 24(10): 1090 - 7.
- 168-Vannucci RC (1993):** "Mechanisms of perinatal hypoxic-ischemic brain damage". *Semin Perinatol*; 17(5): 330-7.

References

- 169-Verkalan MT (2009):** The chilling Details: Hypoxic – ischemic Encephalopathy. *J perin Neo – Nurs*; 23(1) : 59-68.
- 170-Volpe JJ (2008):** Hypoxic –ischemic encephalopathy: biochemical and physiological aspects In : volpe JJ, ed. *Neurology of the Newborn*. Philadelphia, PA: Elsevier Saunders. 331- 94.
- 171-Volpe JJ (2008):** Hypoxic-ischemic encephalopathy: clinical aspects. In: Volpe JJ, ed. *Neurology of the Newborn*. Philadelphia PA. Elsevier Saunders; 400 - 80.
- 172-Volpe JJ (1995):** Hypoxic ischemic encephalopathy. In *Neurology of the newborn*. Johns HC, Richards HK and Bucknal RM (eds). 3rd edition. W.B. Saunders Co., Philadelphia, 250-79.
- 173-Volpe JJ (2001):** Hypoxic ischemic encephalopathy. In *Neurology of the Newborn*. Volpe JJ (eds), 4th edition. W.B. Saunders, Philadelphia, 217-94.
- 174-Wadhawan E, Vohr BR, Fanaroff AA, Pertitt RL Duara S, et al. (2003):** Does labor influence neonatal and neurodevelopmental outcomes of extremely low birth weight infants who are born by cesarean delivery? *Am J obstet Gynecol*; 189(2): 501 - 6.
- 175-Whang R, Oei TO, Aikawa JK, et al. (1984):** Predictors of clinical hypomagnesemia: hypokalemia hypophosphatemia, hyponatremia, and hypocalcemia. *Arch Intern Med*; 144: 1794 - 6.
- 176-Whang R (1987):** Magnesium deficiency pathogenesis, prevalence, and clinical implications. *Am J Med*; 82(3A): 24 - 9.

References

- 177-Whit RE and Hartzell HC (1989):** Magnesium ions in cardiac function: regulator of ion channels and second messengers. *Biochem Pharmacol*; 38: 859 - 67.
- 178-Wood NI, Barone FC, Benham CD et al. (1997):** The effects of SB 206284A, a novel neuronal calcium – channel antagonist, in models of cerebral ischemia. *J Cereb Blood Flow Metab*; 17: 421 - 9.
- 179-Wu WY, Backstrand HK, Zhao S, Fullerton JH and Johnston SC (2004):** Declining diagnosis of birth asphyxia in California 1991-2000. *Pediatr*, 114: 1584 - 90.
- 180-Yoneda S, Ibara S, Kobayashi K, Kato E, Maruyama Y, Maruyama H, Sumida Y, Sunami R, Sakai M, Ikenoue T, Saito S et al. (2004):** Low adjusted serum ionized calcium concentration shortly after birth predicts poor outcome in neonatal hypoxic– ischemic encephalopathy. *J Obstet Gynecol Res Vol.* 31, no. 6:57-64.
- 181-Zanelli SA, Naylor M, Dobbins N, et al. (2008):** Hypoxic ischemic encephalopathy. Saunders: Elsevier. eMedicine. *Pediatr*; 1-9.
- 182-Zanelli SA, Naylor M, Dobbins N, et al. (2008):** Implementation of a "Hypothermia for HIE program: 2- year experience in a single NICU. *J perinatol*; 2(28): 171- 5.
- 183-Zapater P, Moreno J, Horga J (1997):** Neuroprotection by the novel calcium anagonist PCA50938, nimodipine and flunarizine, in gerbil global brain ischemia. *Brain Res*; 772:57-62.

References

- 184-Zarifi MK, Astrakas LG, Poussaint TY, Plessis AA, Zurakowsk D, et al. (2002) :** Prediction of Adverse outcome with cerebral lactate level and asphyxia Diffusion coefficient in infants With perinatal Asphyxia. Radiol; 225 (3) : 859 - 70.
- 185-Ziegler R (2001):** Hypercalcemic crisis. J Am Soc Nephrol 12 (Suppl. 17): S3 - S9.
- 186-Zolaga GP (1992):** Hypocalcemia in critically ill patients. Crit Care Med; 20: 251-62.